CASE REPORT

Pelvic *Actinomyces israelii* abscess: a differential diagnosis of a pelvic massAjit Kaur Dhillon,¹ Neil Fairlie,² Guy Finch³¹Department of Vascular Surgery, Derby Hospital NHS Foundation Trust, Derby, UK²Department of Radiology, Northampton General Hospital, Northampton, UK³Department of General Surgery, Northampton General Hospital, Northampton, UK

Correspondence to

Ajit Kaur Dhillon, ajitdhillon@doctors.org.uk

Accepted 9 November 2015

SUMMARY

A 52-year-old woman was admitted to our district general hospital, with a rapidly growing, increasingly tender mass in the right iliac fossa, difficulty mobilising with a fixed flexion deformity of the right hip and 15 kg weight loss in 5 months. Her 8-month long surgical history for investigation of a pelvic mass stemmed from the removal of an intrauterine device. It included radiological and surgical investigations. We report the second case in the literature, of a patient presenting with abdominal wall abscess, psoas abscess and hydronephrosis as a long-term complication of *Actinomyces israelii* infection of the ipsilateral ovary with a favourable outcome having excluded ovarian malignancy.

BACKGROUND

Actinomycosis is a chronic suppurative and granulomatous, fibrosing infection caused by Gram-positive bacteria, commonly *Actinomyces israelii*. It is a commensal organism of the oral, gastrointestinal and genital tracts, and is characteristically a mimicker of malignancy. Infection commonly supervenes on tissue trauma, for example, following surgery, or by the presence of a foreign body such as an intrauterine device.¹

The bacterium penetrates the damaged mucosa and initiates an inflammatory response leading to the formation of an inflammatory pseudotumour and abscess. The difficulty in diagnosis arises because the abscess grows slowly with potential to cause fistula or perforation. The affected patient will usually present with an indolent illness, and non-specific signs and symptoms, all of which delay definitive diagnosis.^{1 2} Definitive treatment depends on many factors, including location of infection and duration of symptoms, but the cornerstone is antibiotic therapy based on a β lactam inhibitor and a β lactamase inhibitor combination. Surgery may be necessary to reduce the burden of infected tissue and to aid diagnosis, to exclude malignancy. Diagnosis of actinomycosis is primarily based on the presence of Gram-positive filaments and sulfur granules on histology since microbiology cultures are negative in a high proportion of cases.

CASE PRESENTATION

A 52-year-old woman was admitted to our district general hospital as an emergency, with abdominal pain and difficulty mobilising due to an increasingly painful mass in the right iliac fossa (RIF), and fixed flexion deformity of the ipsilateral hip. She had

non-specific symptoms with episodes of fever, night sweats, anorexia and unintentional weight loss of 15 kg in the previous 5 months. Other symptoms included urinary frequency, without dysuria, and variable bowel habit.

Her medical history was unremarkable except for menopause. However, the events following removal of a copper intrauterine coil 8 months earlier, after the coil had been in situ for 20 years, were significant. Following symptoms of vague pelvic discomfort, a pelvic ultrasound scan taken 7 months earlier reported a complex 60×38×44 mm³ mass with fluid and solid elements within it in the left ovary and an irregular 50×39×46 mm³ mass with a thickened endometrium (4.8 mm) in the right ovary. A Pipelle biopsy, carried out 6 months earlier, reported severe endometritis containing colonies of *A. israelii*, treated with a 7-day course of oral co-amoxiclav. The CA125 was elevated at 95 Units/mL. Left salpingo-oophorectomy and adhesiolysis biopsies of the peritoneum were performed at an elective diagnostic laparoscopy 5 months earlier. The histology reported only inflammatory changes without evidence of intraperitoneal actinomycosis or tuberculous infection, and excluded ovarian malignancy.

At follow-up with the gynaecologists 3 months earlier, the patient reported a shooting pain from within the vagina radiating to the umbilicus, associated with nocturia. In the meantime, she had been diagnosed with right-sided sciatica by the general practitioner. On examination, a tender mass in the RIF was noted. The current CT scan was compared with CT scans carried out 6 and 2 months earlier. The main difference between the scans from 6 and 2 months earlier and the current scan was the development of right side hydronephrosis (figures 1 and 2) and 'marked inflammation in the pelvis, surrounding the caecal pole, adherent to the right adnexal mass'. A retrograde J-J stent was placed into the right ureter for presumed malignant ureteric obstruction. Colonoscopy failed due to poor bowel preparation and a repeat colonoscopy was planned for a later date. Similarly, an ultrasound-guided biopsy of the inflamed area in the pelvis was also being planned, but the patient was entered into the current admission before either investigation could be performed.

On examination, she was tachycardic (100 bpm), hypotensive (90/60 mm Hg) and afebrile, but remained alert and orientated. She had a midline laparotomy scar from a surgery many years earlier at a different hospital; the details were unknown to the patient. A firm tender area of erythema was

CrossMark

To cite: Dhillon AK, Fairlie N, Finch G. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-211595

Figure 1 CT scan 6 months earlier, sagittal view showing no hydronephrosis and no pelvic mass.

seen around the previous laparoscopy port site low in the RIF and associated with the right inguinal lymphadenopathy. Serology showed haemoglobin of 61 (125–150 g/dL), white cell

Figure 2 CT scan 2 months earlier, sagittal view showing iliopsoas mass with hydronephrosis.

Figure 3 CT scan during current emergency admission, sagittal view showing the ureteric stent in situ and compression from the iliopsoas mass.

count $9.1 \times 10^9/L$ ($4.0-10.00 \times 10^9/L$), platelet count $497 \times 10^9/L$ ($150-400 \times 10^9/L$) and C reactive protein (CRP) 265 mg/L, and the estimated glomerular filtration rate was reduced to 73 mL/min. The rest of the patient's profile was normal. Plain radio-graphs of the chest and abdomen were unremarkable.

This was considered to be a complication of an ongoing working diagnosis of pelvic malignancy. After transfusion of red blood cells and a gynaecological review, a CT scan of the abdomen and pelvis was repeated. It showed 'persistent right hydronephrosis despite the presence of the ureteric stent', and 'inflammatory change extending from the right groin into the right iliopsoas and right flank, and anteriorly to the skin (figures 3 and 4). Little central fluid content was noted and percutaneous drainage was unlikely to be helpful.' On consideration of the patient's clinical presentation and the diagnostic uncertainty, the decision was made to perform a laparotomy.

Figure 4 CT scan during current emergency admission, axial view showing the psoas abscess pointing through the anterior abdominal wall.

At laparotomy, a right flank, 5.5 cm retroperitoneal abscess without obvious intraperitoneal communication was found, with fibrosis over the iliacus and psoas muscle extending over the pelvic brim. The right ureter was greatly dilated (15 mm) despite the presence of a ureteric stent; the right ovary appeared normal, but the Fallopian tube was tortuous within an inflammatory mass containing the caecum and terminal ileum, in the pouch of Douglas. Although most of the tissues were thickened they did not appear to be malignant. The decision was made to carry out a right oophorectomy, an ileo-caecectomy, multiple retroperitoneal biopsies, and incision and drainage of the retroperitoneal abscess. All the excised tissue appeared fibrotic and felt solid; it had an off-white colouring.

DIFFERENTIAL DIAGNOSIS

- ▶ Ovarian malignancy
- ▶ Caecal malignancy

OUTCOME AND FOLLOW-UP

The diagnosis of *A. israelii* in the right ovarian tissue, small and large intestines was confirmed by histology showing the characteristic yellow sulfur granules containing filaments after Gram stain and treatment with silver stain. Peritonitis was reactive only. Intra-abdominal tuberculosis was excluded.

The pathologist suggested the source of the abscess was most likely the right ovary, but clinically there was stronger suspicion that this was an ascending infection from the endometritis secondary to *A. israelii* identified on Pipelle biopsy, which had been undertreated at that time. It would, however, be difficult to substantiate these suspicions especially on CT scan. Progress of pathology may have been easier to show had a CT scan been carried out at baseline, since direct comparison with the baseline ultrasound and subsequent CT scans is difficult. The patient was discharged home on a 6-month course of doxycycline after 15 days of recovery in hospital. There were concerns about the patient developing an allergy to penicillin administered preoperatively and postoperatively, hence doxycycline was started on confirmation of diagnosis.

An outpatient review reported a very well woman, with a repeat CT scan showing a complete resolution of the previous pathology, apart from residual hydronephrosis. Serology was also much improved, with CRP <5 mg/L, white cell count $3.9 \times 10^9/L$, haemoglobin 127 g/L and platelets $245 \times 10^9/L$.

DISCUSSION

Actinomycosis is a chronic suppurative and granulomatous fibrosing infection caused by Gram-positive bacteria, commonly *A. israelii*. It is a commensal organism of the oral, gastrointestinal and genital tracts, and is characteristically a mimicker of malignancy. Infection commonly supervenes on tissue trauma, for example, following surgery, or by the presence of a foreign body such as an intrauterine device.¹ The bacterium penetrates the damaged mucosa and initiates an inflammatory response leading to the formation of an inflammatory pseudotumour and abscess. It spreads by continuity along the tissue planes rather than taking haematogenous or lymphatic routes. Interestingly, it rarely presents with intraluminal pathology, hence investigations such as endoscopy and barium enema merely show external compression. The difficulty in diagnosis arises because the abscess grows slowly, with potential to cause fistula or perforation. The affected patient will usually present with an indolent illness, and non-specific signs and symptoms, all of which delay definitive diagnosis. Radiological investigations such as CT,

ultrasound and MRI, confirm the presence of a pseudotumour, but the differential diagnosis still includes malignancy.

Microbiological cultures are negative in 76% of cases such that the preoperative diagnosis based on fluid culture yields a diagnosis in 10% of cases. Again, primarily because of a low level of suspicion and difficulties in identifying actinomycosis routinely, diagnosis is confirmed by cultures and histology.^{1–4} Presence of Gram-positive filaments and sulfur granules on histology are strongly suggestive of actinomycosis. Sulfur granules are colonies of organisms that appear as round or oval basophilic masses with eosinophilic terminal ‘clubs’ on staining with H&E.⁵

Treatment ultimately depends on the timing of the diagnosis, location of infection and severity of presenting symptoms. The cornerstone of treatment remains antibiotics, with the choice of regimen taking into account the site affected, other co-pathogens and, ideally, the specific strain of actinomycosis. Typically, a first-line regimen would consist of a β lactam and a β lactamase inhibitor. For patients who are allergic to penicillin, doxycycline, clindamycin and erythromycin are suitable alternatives.^{2–5} Surgical resection of infected tissue may be necessary, especially to exclude malignancy, and to drain abscesses that are deemed unsuitable for percutaneous drainage. Surgery alone has been shown to be of little benefit.^{1–4} By reducing the infective load, antibiotics may exert a greater effect. Importantly, the need for surgical intervention needs to be considered on an individual basis.⁵ The duration of antibiotic therapy depends on the initial infective burden, whether surgical resection has been carried out, the system affected and the patient’s initial response to therapy. Cure of pelvic actinomycosis has been reported after surgical resection and 3 months of antibiotics, but there are other reports of cure following 1–2 months of antibiotics. Prolonged courses are needed to overcome the poor penetration of antibiotics into the fibrotic tissue and avascular planes. Follow-up at regular intervals, both clinical and radiological is vital.^{2–5}

Learning points

- ▶ A high index of suspicion is required in patients who have had an intrauterine device in situ or recent abdominopelvic surgery, and who present with vague symptoms such as lower abdominal pain, urinary symptoms and possibly a leucocytosis.
- ▶ Close liaison with the microbiologists is important to carry out extended cultures and histological analysis of tissue and fluid to exclude presence of yellow sulfur granules and filaments, characteristic of actinomycosis species.
- ▶ In patients referred with a pelvic mass that does not progress as expected for a malignancy, consider presence of an atypical infection such as actinomycosis.

Acknowledgements The authors wish to thank Ms Pip Dival, clinical librarian, for her help in the literature search and retrieval.

Contributors AKD prepared the manuscript. NF reported on radiological imaging. GF was responsible for concept design and joint writing of the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Marella VK, Hakimian O, Wise GJ, *et al.* Pelvic actinomycosis, urologic perspective. *Int Braz J Urol* 2004;30:367–76.
- 2 Choi MM, Baek JH, Lee JN, *et al.* Clinical features of abdominopelvic actinomycosis: report of twenty cases and literature review. *Yonsei Med J* 2009;50:555–9.
- 3 Pitot D, De Moor V, Demetter P, *et al.* Actinomycotic abscess of the anterior abdominal wall: a case report and literature review. *Acta Chir Belg* 2008;108:471–3.
- 4 de Feiter PW, Soeters PB. Gastrointestinal actinomycosis: an unusual presentation with obstructive uropathy. *Dis Colon Rectum* 2001;44:1521–5.
- 5 Wong VK, Turmezei TD, Weston VC. Clinical review: actinomycosis. *BMJ* 2011;343:d6099.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow