Management of Mild Traumatic Brain Injury Symptoms in a 31-Year-Old Woman Using Cervical Manipulation and Acupuncture: A Case Report

Danielle M. Gergen DC*

Chiropractor, Oxboro Family Chiropractic, Bloomington, MN

Received in 21 April 2015; received in revised form 5 August 2015; accepted 5 August 2015

Key indexing terms:

Brain injuries;
Brain concussion;
Neck pain;
Manipulation,
chiropractic;
Acupuncture points

Abstract

Objective: The objective was to describe chiropractic and acupuncture care of a patient with acute mild traumatic brain injury (mTBI) symptoms.

Clinical Features: A 31-year-old woman had acute neck pain, headache, dizziness, nausea, tinnitus, difficulty concentrating, and fatigue following a fall. She was diagnosed at an urgent care facility with mTBI immediately following the fall. Pharmaceutical intervention had been ineffective for her symptoms.

Intervention and Outcome: The patient was treated with chiropractic adjustments characterized as high velocity, low amplitude thrusts directed to the cervical spine and local acupuncture points in the cervical and cranial regions. The patient received care for a total of 8 visits over 2.5 weeks with resolution of concussive symptoms.

Conclusion: This patient with mTBI responded favorably to a conservative treatment protocol with the combination of chiropractic and acupuncture care.

© 2015 National University of Health Sciences.

Introduction

Mild traumatic brain injury (mTBI) has gained media attention over the effects of repetitive head injuries on brain development and health in both athletes and trauma patients. Approximately 75% of the 1.7 million brain

injuries in the United States are classified as concussions or mTBI annually.¹ This figure reflects the sharp increase in mTBIs incidence from 521.0 reported injuries per 100,000 people to 823.7 per 100,00 from 2001 to 2010.² The mechanism of injury is largely attributed to local head trauma from biomechanical forces.³ Currently, there is a large focus on a young, athletic population with injuries from contact sports, although the majority of mTBIs occur after traumatic falls and motor vehicle crashes.⁴ As a result of the trauma, current evidence

* Oxboro Family Chiropractic, 9864 Lyndale Ave S, Bloomington, MN 55420. Tel.: +1 952 948 9225; fax: +1 952 888 6011.

E-mail address: d.m.gergen@gmail.com.

suggests that there is a rapid hyperglycolysis in the neural tissue followed by a prolonged period of hypoglycolysis, altered permeability of the blood brain barrier, increased inflammation, and a rise in interleukin levels.⁵ Symptoms arising from these cellular events include nausea, dizziness, headache, tinnitus, acute pain, and cognitive deficits.⁶ Symptoms typically resolve over weeks to months, although 36% of patients diagnosed with mTBI continue to report persisting symptoms for 6 months or longer.⁷ In addition, the Centers for Disease Control and Prevention reported that as many as 15% of patients with diagnosed mTBI experience persistent, disabling problems associated with concussive symptoms.⁸

Consistently, the literature reports rest as the most common treatment method for mTBI symptoms.^{5,9–13} Symptom education and material on future prevention are also commonly given as postconcussive treatment especially in an athletic population where return to play is a major focus.^{6,13} Although rest and education have been found to reduce some mTBI symptoms, these can be insufficient as a treatment strategy in an adult, working population where symptoms often can be triggered by occupational duties and can greatly impact daily activities. Pharmaceutical treatment options, such as DDAVP, have demonstrated little positive effect on symptoms.^{5,13,14} Self-management with over-the-counter options like nonsteroidal anti-inflammatory drugs has also been shown to be ineffective in symptom reduction despite the inflammatory cascade that occurs during mTBI.⁵

Chiropractic management of mTBI has a prominent focus on an athletic population within the literature. Multiple reviews outline the high incidence of mTBI and indicated that chiropractors as first-contact providers should have awareness of symptoms, educational materials for patients, and stress rest before a return to play or activities.^{1,15,16} In addition, manual adjustments both with and without soft tissue therapies have been shown to be effective in the treatment of some mTBI symptoms including somatic pain related to mTBI, whiplash, headache, migraine, loss of motion in the cervical spine, and additional sequelae from trauma.^{17–20}

In Traditional Chinese Medicine (TCM), it is theorized that external trauma such as in mTBI often causes a TCM condition known as *blood stasis* which causes pain and dysfunction in the injured area and can be alleviated with TCM treatments such as acupuncture. Acupuncture has been found to have longer-lasting therapeutic effect for acute headache after trauma when compared with pharmaceutical options.^{21–25} Local needling produces analgesic effects in areas of injury

by locally increasing the blood supply through the release of vasodilatory neuropeptides and thereby increasing local oxygen and cytokines along with reducing inflammation.^{26,27} Acupuncture also has been indicated as having a positive therapeutic effect on other mTBI-related symptoms such as headaches, neck pain, and nausea.^{28–31}

Because of the prevalent nature of mTBI injuries, there is need for additional treatment protocols to be available for injuries sustained by adults in nonathletic situations. The purpose of this case report is to describe the management of an adult patient with mTBI injury through the use of manual cervical chiropractic manipulation, soft tissue therapy, and acupuncture.

Case Report

A 31-year-old woman presented to the chiropractic clinic for treatment of acute mTBI symptoms after a traumatic fall 3 weeks prior. She reported that she slipped on ice and landed prone, striking her face on the ground. She lost consciousness for a period of 3–4 minutes. The patient presented to urgent care with symptoms of acute neck pain, frontal and occipital headaches, nausea, dizziness, tinnitus, fatigue, and cognitive “foginess.” She was diagnosed with mTBI and prescribed Flexeril and Vicodin. Medical intervention temporarily alleviated her symptoms, but she found the relief to be short lived.

The patient described her physical and cognitive symptoms as unchanged from her initial presentation to urgent care. She reported constant dull neck pain with intermittent episodes of sharp pain into her arms bilaterally and upper back associated with movement. The symptoms made it difficult to sleep, perform her occupational duties, and participate in physical activity. She noted her neck pain to be 9/10 on a verbal Numeric Pain Scale (NPS) and headache pain to be 9/10 NPS upon initial evaluation.

On examination, the patient had a loss of active cervical rotation 5°–10° bilaterally and was noted to have static rotational changes to her occiput. Her neurologic examination was nonfocal, and she had no evidence of cranial trauma. Vitals were within normal limits. Plain radiographs of the cervical spine demonstrated a loss of cervical lordosis and a mild right lateral convexity of C4–C7, with minimal signs of vertebral body elongation inferiorly on C4, C5, and C6. Result of Romberg test was negative on orthopedic examination. Local pain and tension were noted during cervical

compression, foraminal compression test, and shoulder depression. Motion palpation found rotational restriction of C3 on the left and C7 on the right. The patient did not have a personal or family history of migraine, neck pain, or significant medical history. The working diagnosis was mTBI with cervical sprain/strain, cervicogenic headache, segmental dysfunction of the cervical spine, and muscle spasm.

Treatment visits consisted of high-velocity, low-amplitude manipulation of C3 and C7. Trigger point therapy was applied to the upper trapezius and suboccipital muscles bilaterally for 10-15 seconds per point. Therapeutic stretching of the upper trapezius muscles was augmented with intersegmental traction for 10 minutes. Cryotherapy with ice packs was applied for the first treatment only. Treatment frequency was 3 visits per week for 2 weeks. After 6 treatments of manual manipulation to the cervical region and soft tissue therapies, the patient reported decreased pain levels in the neck from 9/10 NPS to 2/10 NPS. Headache intensity decreased from 9/10 NPS to 4/10 NPS. Reevaluation found full cervical range of motion.

Despite pain reductions, the patient continued to endorse nausea, dizziness, fatigue, and tinnitus. Acupuncture was performed with DBC Spring Ten stainless steel needles (0.20×30 , 0.20×15) in addition to chiropractic treatments for her continued symptoms during the subsequent 2 visits. Points included cervical huatuojiayi points (perpendicular insertion with slight angle towards the spine; depth, 0.5-1.0 cun), Fengchi (GB-20, angled towards the nose tip; depth, 0.5 cun), Jianjing (GB-21, perpendicular insertion with slight medial angling; depth, 0.3 cun), and Houxi (SI-3, perpendicular insertion; depth, 0.5 cun) locally and Kunlun (BL-60, perpendicular insertion; depth, 0.5 cun) distally. These points were chosen because of the channels' course passing through the affected area and their traditionally accepted therapeutic effects. The needles were inserted with a dispersing technique (including scraping and thrusting) and retained for 20 minutes each visit. Auricular endocrine, thalamus, omega 2, Point Zero, and shenmen were needled bilaterally at a depth of 0.1-0.2 cun perpendicularly for pain relief and were retained for 20 minutes.

After 2 treatments combining chiropractic manipulation and acupuncture over 3 days, the patient reported neck and headache pain levels to be 0/10 NPS and a complete resolution of the physical and cognitive mTBI symptoms. In addition, she was able to participate in all of her former activities without symptoms returning. She continued with follow-up care for an additional 4 weeks at 1-2 visits per week without relapse or regression of her symptoms. Examination, treatment, and follow-up

evaluation were performed by the author. The patient gave her written consent for use of her personal health information as it related to this case.

Discussion

This case report outlined the care of an adult female patient with a fall-induced mTBI with both physical and cognitive symptoms. The management and treatment of patients with mTBI have been a popular topic within the literature, as the reported incidence has increased rapidly over the last decade.² Currently, there is no one standard of care for mTBI outside of rest and avoidance of reinjury.^{5,9-13} This treatment method has been shown to decrease symptoms with time and reduce the risk of reinjury in athletes.^{1,10-12,16} In this case, the patient's mTBI symptoms interfered with her work duties and daily activities. Often, her daily tasks also increased the intensity and frequency of her symptoms, but she was unable to discontinue aggravating activities due to being unable to be away from work. This made rest an ineffective treatment due to her inability to maintain compliance with provider recommendations. In addition, the patient did not experience relief with medications aimed at reducing her symptoms.

In this patient's case, manual chiropractic manipulation of the cervical spine, soft tissue therapies, and acupuncture treatments expedited the reduction in her symptom frequency and intensity after a fall resulting in an mTBI. Notably, her pain levels decreased, range of motion improved to normal limits, and she reported a complete resolution of dizziness, nausea, tinnitus, and fatigue after 8 visits over the course of 2.5 weeks. This rate of improvement and return to regular activities was much quicker than the reported rate for rest alone.⁷ She did not relapse over the following 4 weeks of follow-up care, marking the treatment method as potentially long-lasting in nature.

Limitations

It is possible that the patient's mTBI symptoms improved regardless of care because 5 weeks had passed from the initial injury to when she reported to be symptom-free. The patient also may have engaged in other therapeutic activities that she did not report during the 2 weeks of treatment for active symptoms. It is also difficult to gauge the level of disruption to the

patient's daily activities because of the information being collected verbally rather than through the use of a validated questionnaire, such as the activity of daily living questionnaire, SCAT3, or Neck Disability Index. The findings of this case report may not necessarily be applicable to other patients with similar symptoms. More participants in a more distributed population are needed to test the results of these treatment methods with more objective evaluation measures.

Conclusion

The treatment outlined in this case report describes an active approach to mTBI treatment rather than rest and patient education alone. The use of gentle spinal manipulation, soft tissue therapies, and acupuncture reduced the patient's symptoms, improved her range of motion, and increased her quality of life by achieving full return to daily activities in a treatment period of 2 weeks.

Funding Sources and Conflicts of Interest

No funding sources or conflicts of interest were reported for this study.

References

- Johnson C, Green B, Nelson R, Moreau B, Nabhan D. Chiropractic and concussion in sport: a narrative review of the literature. *J Chiropr Med* 2013;12:216–29.
- Centers of Disease Control and Prevention. Injury prevention & control: traumatic brain injury. Atlanta, GA: Centers for Disease Control and Prevention; 2014 [Available from: www.cdc.gov/traumaticbraininjury/data/rates.html. Accessed 2015 June 02].
- Cantu RC, Aubry M, Dvorak J, et al. Overview of concussion consensus statements since 2000. *Neurosurg Focus* 2000;2006(21):E3.
- Cassidy JD, Carroll LJ, Peloso PM, et al. Incidence, risk factors and prevention of mild traumatic brain injury: results of the WHO collaborating centre task force on mild traumatic brain injury. *J Rehabil Med* 2004;36(Suppl 43):28–60.
- Patterson ZR, Holahan MR. Understanding the neuroinflammatory response following concussion to develop treatment strategies. *Front Cell Neurosci* 2012;6(58):1–10.
- Marshall S, Bayley M, McCullagh S, Velikonja D, Berrigan L. Clinical practice guidelines for mild traumatic brain injury and persistent symptoms. *Can Fam Physician* 2012;58:257–67.
- Donovan J, Cancelliere C, Cassidy JD. Summary of the findings of the International Collaboration on Mild Traumatic Brain Injury Prognosis. *Chiropr. Man Ther* 2014;22:38.
- Centers for Disease Control and Prevention. Heads up. Facts for physicians about mild traumatic brain injury (MTBI). Atlanta, GA: Centers for Disease Control and Prevention; 2007. Available from: www.cdc.gov/ncipc/pub-res/tbi_toolkit/physicians/mtbi/mtbi.pdf. Accessed 2015 March 01.
- Cushman JG, Agarwal N, Fabian TC, et al. Practice management guidelines for the management of mild traumatic brain injury: the EAST practice management guidelines work group. *J Trauma* 2001;51:1016–26.
- McCrary P, Meeuwisse W, Johnston K, et al. Consensus statement on concussion in sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008. *Br J Sports Med* 2009;43(Suppl 1):176–90.
- Standaert CJ, Herring SA, Cantu RC. Expert opinion and controversies in sports and musculoskeletal medicine: concussion in the young athlete. *Arch Phys Med Rehabil* 2007;88:1077–9.
- Willer B, Leddy JJ. Management of concussion and post-concussion syndrome. *Curr Treat Options Neurol* 2006;8:415–26.
- Gravel J, D'Angelo A, Carriere B, et al. Interventions provided in the acute phase for mild traumatic brain injury: a systematic review. *Syst Rev* 2013;2:63–75.
- Department of Defense. VA/DoD clinical practitioner guideline for management of concussion/mild traumatic brain injury. Washington, DC: Department of Veterans Affairs, Department of Defense; 2009. Available at http://www.healthquality.va.gov/guidelines/Rehab/mtbi/concussion_mtbi_full_L0.pdf; accessed 2015 March, 01.
- Moreau W, Nabhan D. Development of the 2012 American Chiropractic Board of Sports Physicians position statement on concussion in athletics. *J Chiropr Med* 2013;12:269–73.
- Clay M, Glover K, Lowe D. Epidemiology of concussion in sport: a literature review. *J Chiropr Med* 2013;12:230–51.
- Cassidy JD, Boyle E, Carroll LJ. Population-based, inception cohort study of the incidence, course, and prognosis of mild traumatic brain injury after motor vehicle collisions. *Arch Phys Med Rehabil* 2014;95(3 Suppl):S278–85.
- Carroll LJ, Cassidy JD, Côté P. The role of pain coping strategies in prognosis after whiplash injury: passive coping predicts slowed recovery. *Pain* 2006;124(1–2):18–26.
- Donovan J, Cassidy JD, Cancelliere C, et al. Beyond the spine: a new clinical research priority. *J Can Chiropr Assoc* 2015;59(1):6–12.
- Clar C, Tsertsvadze A, Court R, Hundt GL, Clarke A, Sutcliffe P. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: systematic review and update of UK evidence report. *Chiropr Man Ther* 2014;22:12.
- Zhao L, Guo Y, Wang W, Yan LJ. Systematic review on randomized controlled clinical trials of acupuncture therapy for neurovascular headache. *Chin J Integr Med* 2011;17(8):580–6.
- Linde K, Allais G, Brinkhaus B, Manheimer E, Vickers A, White AR. Acupuncture for migraine prophylaxis. *Cochrane Database Syst Rev* 2009;1:DC0012180.
- Watanabe TK, Bell KR, Walker WC, Schomer K. Systematic review of interventions for post-traumatic headache. *PM R* 2012;4(2):129–40.
- Khusid M. Clinical indications for acupuncture in chronic post-traumatic headache management. *Mil Med* 2015;2:132–6.

25. Jodorkovsky A, Rotenberg T. The acupuncture treatment of sensory loss following head trauma: a case study. *J Chin Med* 2013;103:34–6.
26. Sandberg M, Lundeberg T, Lindberg L, Gerdle B. Effects of acupuncture on skin and muscle blood flow in healthy subjects. *Eur J Appl Physiol* 2003;90(1-2):114–9.
27. Han J. Acupuncture and endorphins. *Neurosci Lett* 2004;361(1-3):258–61.
28. Vickers AJ, Cronin AM, Maschino AC, et al. Acupuncture for chronic pain: individual patient data meta-analysis. *Arch Intern Med* 2012;172(19):1444–53.
29. Lee C, Crawford C, Wallerstedt D, et al. The effectiveness of acupuncture research across components of the trauma spectrum response (TSR): a systematic review of reviews. *Syst Rev* 2012;1:46.
30. Lee A, Fan LT. Stimulation of the wrist acupuncture point P6 for preventing postoperative nausea and vomiting. *Cochrane Database Syst Rev* 2009;2:CD003281.
31. Linde K, Allais G, Brinkhaus B, Manheimer E, Vickers A, White A. Acupuncture for tension-type headache. *Cochrane Database Syst Rev* 2011;1:CD007587, <http://dx.doi.org/10.1002/14651858.CD007587>.