Published in final edited form as:

Angew Chem Int Ed Engl. 2015 September 28; 54(40): 11800–11803. doi:10.1002/anie.201505161.

Palladium-Catalyzed Decarbonylative Dehydration for the Synthesis of α -Vinyl Carbonyl Compounds and Total Synthesis of (–)-Aspewentin A, B, and C

Yiyang Liu^a, Scott C. Virgil^a, Robert H. Grubbs^{a,*}, and Brian M. Stoltz^{a,*}

^aThe Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 E. California Blvd, MC 101-20, Pasadena, CA 91125 (USA)

Abstract

The direct α -vinylation of carbonyl compounds that forms a quaternary stereocenter is a challenging transformation. We discovered that δ -oxocarboxylic acids can serve as masked vinyl compounds and be unveiled by palladium-catalyzed decarbonylative dehydration. The carboxylic acids are readily available through enantioselective acrylate addition or asymmetric allylic alkylation. A variety of α -vinyl quaternary carbonyl compounds are obtained in good yields, and an application in the first enantioselective total synthesis of (–)-aspewentin A, B, and C is demonstrated.

Keywords

α -vinyl carbonyl compounds; decarbonylative dehydration; palladium catalysis; quaternary stereocenters

An all-carbon quaternary center bearing an ethylene substituent is a common structural motif in many natural products (see Figure 1).^[1] An important approach to the construction of this unit is the α -vinylation of carbonyl compounds, and two general methods have been developed. One is the direct coupling of an enolate nucleophile with a vinyl electrophile such as an alkenyl ether,^[2] vinyl bromide,^[3] or acetylene itself.^[4,5] Although this approach can be extended to alkenylations as well, and asymmetric versions are known,^[3a,3c,4c] the scope of the enolate nucleophile is generally limited to 1,3-dicarbonyl compounds^[4] or those with only one enolizable position.^[2,3,5b] A second tactic involves addition of the enolate nucleophile to a vinyl surrogate such as vinyl sulfoxide,^[6] (phenylseleno)acetaldehyde,^[7] or ethylene oxide,^[8] followed by elimination. However, there are few reports of stereoselective additions that form the quaternary stereocenter,^[9] and none are catalytic or enantioselective. Due to these constraints, even the simplest 2-methyl-2-vinylcyclohexanone (**7**) is not known as a single enantiomer in the literature.

We envisioned an alternative approach to access α -vinyl carbonyl compounds, by employing a decarboxylative elimination of δ -oxocarboxylic acids (Scheme 1). These acids

may be prepared by numerous methods, including addition of an enolate nucleophile to an acrylate acceptor^[10] and palladium-catalyzed allylic alkylation.^[11] Importantly, these two methods allow for the enantioselective construction of the requisite quaternary stereocenter.

Recently, we reported on the palladium-catalyzed decarbonylative dehydration of fatty acids to form terminal olefins.^[12] Due to the importance of α -vinyl carbonyl compounds and the challenges in their preparation, we became interested in applying our decarbonylative dehydration chemistry as an alternative strategy to α -vinylation. Since the carboxylic acid, which bears a quaternary center two atoms away from the reactive carboxyl group, is more hindered than a simple fatty acid, we expected that the reaction conditions would need to be tuned for this particular class of substrates. Additionally, from a practical standpoint, our previous studies were typically conducted on ~5 g fatty acid substrate without solvent, under vacuum distillation conditions. Thus, a smaller scale alternative for implementation on laboratory scale and in the context of multistep organic synthesis would need to be developed.

At the outset of our investigation, we prepared carboxylic acid **8a** and subjected it to modified palladium-catalyzed decarbonylative dehydration conditions, when slightly higher loadings of catalyst, ligand, and additive were employed (Scheme 2). We were pleased to isolate vinyl cyclopentanone **9a** in 67% yield.^[13] This result demonstrated that steric bulk at the quaternary center does not significantly retard the reaction, but proximal functionality (e.g. the ketone) could alter the reaction pathway.

With this exciting initial result in hand, we proceeded to investigate the scope of the reaction (Table 1). First, we synthesized (*R*)-3-(1-methyl-2-oxocyclohexyl)propanoic acid (**8b**) by enantioselective d'Angelo Michael addition,^[10] and subjected it to decarbonylative dehydration (entry 2). We were delighted to obtain the desired product, (*R*)-2-methyl-2-vinylcyclohexanone (*ent*-**7**), in 60% yield and 92% ee. Likewise, 2-ethyl-2-vinylcyclohexanone (**9c**) was prepared in a similar fashion. Carboxylic acids bearing allyl or 2-methallyl substituents, which can be prepared via palladium-catalyzed allylic alkylation,^[11] also underwent decarbonylative dehydration smoothly to provide the corresponding 2-allyl-2-vinyl-substituted cyclohexanones **9d** and **9e** (entries 4 and 5), the latter in 92% ee from enantioenriched acid **8e**. It is worth noting that double bond isomerization in the allyl moiety is negligible for **9d** and does not occur at all for **9e**. Acyclic keto acid **8f** is converted to acyclic ketone **9f** in good yield (entry 6). Aside from keto acid substrates, we examined acids bearing other types of carbonyl functionalities (entries 7–10), and found that α -vinyl ester **9g**, lactam **9h**, and aldehyde **9i** can all be prepared in good yields. More complex scaffolds such as acid **8j**, obtained by oxidative cleavage of testosterone,^[14] also undergo the reaction to provide vinylated tricycle **9j** (entry 11). While the reaction can be carried out in the absence of a solvent at a fairly large scale (5 mmol, entries 1–7), we found that for smaller scale synthesis it is more convenient to use *N*-methylpyrrolidinone (NMP) as solvent along with slightly modified conditions (entries 8–11).^[15]

To further demonstrate the utility of our decarbonylative dehydration approach to vinylation, we embarked on a total synthesis of aspewentin B (**1**, Figure 1), a norditerpene natural

product isolated from *Aspergillus wentii*.^[16, 17] This terpenoid contains an α -vinyl quaternary cyclohexanone scaffold, and is therefore ideally suited for our chemistry. Retrosynthetically, we envisioned that the vinyl group could be formed by decarbonylative dehydration of δ -oxocarboxylic acid **11**, which might be obtained by elaboration of allyl ketone **12**.^[18] The quaternary stereocenter would be set by palladium-catalyzed enantioselective allylic alkylation of tricyclic ketone **13**, which could be constructed from known aryl bromide **14**.^[19]

We commenced our total synthesis by copper-catalyzed coupling of a Grignard reagent derived from aryl bromide **14** with ethyl 4-iodobutyrate (Scheme 4).^[20] α -Methylation of the coupled product produces ester **15**, which was saponified and then cyclized to form tricyclic ketone **13**. The ketone was converted to the corresponding allyl enol carbonate (**16**), and subjected to palladium-catalyzed enantioselective decarboxylative allylic alkylation to afford allyl ketone **12** in nearly quantitative yield and 94% ee. Interestingly, the allylic alkylation reaction proceeds efficiently with low palladium catalyst loading, employing a new catalytic protocol recently developed by our group.^[21] Only 2.7 mg Pd(OAc)₂ is needed for a reaction of 1.42 g starting material (**16**) to deliver 1.25 g allyl ketone product (**12**). Hydroboration/oxidation of the terminal olefin of **12**, followed by further oxidation, delivers carboxylic acid **11** in 73% yield.^[18] Gratifyingly, palladium-catalyzed decarbonylative dehydration^[22] furnishes α -vinyl ketone **17** in 93% yield.^[23] Removal of the *O*-methyl group provides (–)-aspewentin B (**1**) in 78% yield.^[24] Thus, (–)-aspewentin B (**1**) was synthesized in 9 steps and 25% overall yield from known starting materials. With **1** in hand, two other natural products of the aspewentin family can also be accessed. Reduction of the ketone moiety of **1** furnishes (–)-aspewentin A (**18**) in 85% yield.^[25] Oxidation of the phenol in **18** affords (–)-aspewentin C (**19**) in 13% yield and (+)-10-*epi*-aspewentin C (**20**) in 7% yield, over 2 steps.^[26]

In summary, we have developed a new approach to access α -vinyl quaternary carbonyl compounds via palladium-catalyzed decarbonylative dehydration of δ -oxocarboxylic acids. A variety of acids with different scaffolds and functional groups can be transformed into the corresponding α -vinyl carbonyl compounds without erosion of the stereointegrity of the neighboring quaternary center. We have also applied the method to the first enantioselective total synthesis of (–)-aspewentin A, B, and C. Further applications of this transformation in natural product synthesis are currently ongoing in our lab and will be reported in due course.

Experimental Section

See Supporting Information for experimental details.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors wish to thank the Resnick Sustainability Institute at Caltech (graduate fellowship to Y.L.), NIH (R01GM080269 to B.M.S., 5R01GM031332-27 to R.H.G.), the Gordon and Betty Moore Foundation, the Caltech

Center for Catalysis and Chemical Synthesis, and Caltech for financial support. Dr. David VanderVelde is acknowledged for NMR spectroscopy assistance. Dr. Mona Shahgholi and Naseem Torian are acknowledged for high-resolution mass spectrometry assistance. We would like to thank Prof. Richmond Sarpong of UC Berkeley for helpful discussions.

References

1. a) Bernauer K, Englert G, Vetter W, Weiss E. *Helv. Chim. Acta*. 1969; 52:1886–1905. b) Moore RE, Cheuk C, Patterson GML. *J Am. Chem. Soc.* 1984; 106:6456–6457. c) Masuda T, Masuda K, Shiragami S, Jitoe A, Nakatani N. *Tetrahedron*. 1992; 48:6787–6792. d) Rubal JJ, Moreno-Dorado FJ, Guerra FM, Jorge ZD, Saouf A, Akssira M, Mellouki F, Romero-Garrido R, Massanet GM. *Phytochemistry*. 2007; 68:2480–2486. [PubMed: 17669447]
2. Nishimoto Y, Ueda H, Yasuda M, Baba A. *Angew. Chem. Int. Ed.* 2012; 51:8073–8076.
3. a) Chieffi A, Kamikawa K, Åhman J, Fox JM, Buchwald SL. *Org. Lett.* 2001; 3:1897–1900. [PubMed: 11405739] b) Huang J, Bunel E, Faul MM. *Org. Lett.* 2007; 9:4343–4346. [PubMed: 17887766] c) Taylor AM, Altman RA, Buchwald SL. *J Am. Chem. Soc.* 2009; 131:9900–9901. [PubMed: 19580273]
4. a) Nakamura M, Endo K, Nakamura E. *Org. Lett.* 2005; 7:3279–3281. [PubMed: 16018640] b) Endo K, Hatakeyama T, Nakamura M, Nakamura E. *J Am. Chem. Soc.* 2007; 129:5264–5271. [PubMed: 17388598] c) Fujimoto T, Endo K, Tsuji H, Nakamura M, Nakamura E. *J Am. Chem. Soc.* 2008; 130:4492–4496. [PubMed: 18331035]
5. a) Yamaguchi M, Tsukagoshi T, Arisawa M. *J Am. Chem. Soc.* 1999; 121:4074–4075. b) Arisawa M, Miyagawa C, Yoshimura S, Kido Y, Yamaguchi M. *Chem. Lett.* 2001:1080–1081.
6. Tanikaga R, Sugihara H, Tanaka K, Kaji A. *Synthesis*. 1977:299–301.
7. a) Kowalski CJ, Dung J-S. *J Am. Chem. Soc.* 1980; 102:7950–7951. b) Clive DLJ, Russell CG, Suri SC. *J Org. Chem.* 1982; 41:1632–1641.
8. Nozoye T, Shibamura Y, Nakai T, Hatori Y. *Chem. Pharm. Bull.* 1988; 36:4980–4985.
9. a) Meyers AI, Hanreich R, Wanner KT. *J Am. Chem. Soc.* 1985; 107:7776–7778. b) Kawashima H, Kaneko Y, Sakai M, Kobayashi Y. *Chem. – Eur. J.* 2014; 20:272–278. [PubMed: 24273103]
10. Pfau M, Revial G, Guingant A, d'Angelo J. *J Am. Chem. Soc.* 1985; 107:273–274.
11. Mohr JT, Behenna DC, Harned AM, Stoltz BM. *Angew. Chem. Int. Ed.* 2005; 44:6924–6927.
12. Liu Y, Kim KE, Herbert MB, Fedorov A, Grubbs RH, Stoltz BM. *Adv. Synth. Catal.* 2014; 356:130–136. [PubMed: 24772061]
13. Cyclic lactone **10** was also isolated in 20% yield as the major byproduct. Byproducts of this type are observed in cases where cyclic ketones possess an enolizable position.

14. de Avellar IGJ, Vierhapper FW. *Tetrahedron*. 2000; 56:9957–9965.
15. The small-scale reactions were performed under 1 atm N₂ (no distillation). Acetic anhydride was found to be unsuitable under such conditions, while benzoic anhydride proved to be optimal.
16. Miao F-P, Liang X-R, Liu X-H, Ji N-Y. *J Nat. Prod.* 2014; 77:429–432. [PubMed: 24499164]
17. While the natural product has (+)-stereochemistry, we synthesized the (–)-enantiomer due to the ready availability of the (S)-*t*-BuPHOX ligand.
18. The enantioselective Michael addition does not work for 1-tetralone substrates. For details, see: d'Angelo J, Desmaële D, Dumas F, Guingant A. *Tetrahedron Asymmetry*. 1992; 3:459–505.
19. Allergan, Inc. Patent. WO2005/58301 A1. 2005.

20. Brown ML, Eidam HA, Paige M, Jones PJ, Patel MK. *Bioorg. Med. Chem.* 2009; 17:7056–7063. [PubMed: 19747831]
21. Stoltz BM, et al. Manuscript accepted by *Adv. Synth. Catal.*
22. Decarbonylative dehydration of **11** using Ac₂O under standard vacuum distillation conditions results in no conversion; the reaction mixture solidifies upon distillation of AcOH.
23. An alternative route from allyl ketone **12** to vinyl ketone **17** involves isomerization of the double bond to the internal position followed by ethenolysis. While the isomerization (i.e. **12** → **SI-17**) proceeded smoothly using the protocol developed by Nishida (Arisawa M, Terada Y, Nakagawa M, Nishida A. *Angew. Chem. Int. Ed.* 2002; 41:4732–4734.) to afford internal olefin **SI-17**, attempts at ethenolysis of the resulting internal olefin were unsuccessful, likely due to steric bulk at the adjacent quaternary center. See Supporting Information for details.

24. Spectroscopic data (¹H and ¹³C NMR) and exact mass of the synthetic compound matches those of natural (+)-aspewentin B. Optical rotation, however, is significantly different (synthetic compound: [α]_D²⁵ −90.5 (c 0.20, MeOH, 98% ee); natural product: [α]_D²⁰ +23.3 (c 0.20, MeOH)). HPLC analysis showed the synthetic compound to be of 98% ee. Based on the stereoselectivity of the Pd-PHOX catalyst in previous examples, we reasoned that the absolute configuration of our synthetic compound is opposite to that of the natural product, and thus the sign of optical rotation is correct. The difference in magnitude may arise from the possibility that the natural product contains an impurity, or that the natural product is a scalemic mixture (i.e. not enantiopure). For a review on non-enantiopure natural products, see: Finefield JM, Sherman DH, Kreitman M, Williams RM. *Angew. Chem. Int. Ed.* 2012; 51:4802–4836. Another possibility is that the isolated natural product contained a small amount of impurity that had a large effect on optical rotation.
25. Quandt G, Höfner G, Wanner KT. *Bioorg. Med. Chem.* 2013; 21:3363–3378. [PubMed: 23598250]
26. Carlsen PN, Mann TJ, Hoveyda AH, Frontier AJ. *Angew. Chem. Int. Ed.* 2014; 53:9334–9338.

Figure 1.
Natural products with ethylene substituents.

Scheme 1.
Decarboxylative approach to vinylation.

Scheme 2.
Decarbonylative dehydration of δ -oxocarboxylic acid **8a**.

Scheme 3.
Retrosynthetic analysis of (-)-aspewentin B.

Scheme 4.

Total synthesis of (-)-aspewentin A, B, and C.

Table 1

Decarbonylative dehydration of δ -oxocarboxylic acids.

$ \begin{array}{c} \text{PdCl}_2(\text{PPh}_3)_2 \text{ (0.2 mol\%)} \\ \text{Xantphos (0.24 mol\%)} \\ \text{(t-Bu)}_3\text{biphenol (1 mol\%)} \\ \text{Ac}_2\text{O (1.2+0.5+0.3+0.2 equiv)} \\ \text{neat, 132 } ^\circ\text{C, 2 h} \\ \text{vacuum distillation} \end{array} \rightarrow $			
entry ^a	δ -oxocarboxylic acid	product	yield
1			67%
2			60% 92% ee
3			66%
4			54% ^b
5			69% 92% ee
6			75%
7			50% 51%
9 ^c			57%
10 ^c			77%
11 ^c			41%

^[a] 5 mmol scale.^[b] Isolated as a 95:5 mixture of desired product and internal olefin isomer.

^[c]Condition: 0.5 mmol substrate (1 equiv), benzoic anhydride (1.2 equiv), PdCl₂(nbd) (1 mol%), Xantphos (1.2 mol%), NMP (0.25 mL), 1 atm N₂, 132 °C, 3 h.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript