Time course of facial emotion processing in women with borderline personality disorder: an ERP study

Natalie A. Izurieta Hidalgo, MSc; Rieke Oelkers-Ax, MD; Krisztina Nagy, PhD;
Falk Mancke, MD; Martin Bohus, MD; Sabine C. Herpertz, MD; Katja Bertsch, PhD

Background: Borderline personality disorder (BPD) is characterized by a negative perception of others. Previous studies have revealed deficits and biases in facial emotion recognition. This study investigates the behavioural and electrophysiological correlates underlying facial emotion processing in individuals with BPD. **Methods:** The present study was conducted between July 2012 and May 2014. In an emotion classification task, unmedicated female patients with BPD as well as healthy women had to classify faces displaying blends of anger and happiness while the electroencephalogram was recorded. We analyzed visual event-related potentials (ERPs) reflecting early (P100), structural (N170) and categorical (P300) facial processing in addition to behavioural responses. **Results:** We included 36 women with BPD and 29 controls in our analysis. Patients with BPD were more likely than controls to classify predominantly happy faces as angry. Independent of facial emotion, women with BPD showed enhanced early occipital P100 amplitudes. Additionally, temporo-occipital N170 amplitudes were reduced at right hemispherical electrode sites. Centroparietal P300 amplitudes were reduced particularly for predominantly happy faces and increased for highly angry faces in women with BPD, whereas in healthy volunteers this component was modulated by both angry and happy facial affect. **Limitations:** Our sample included only women, and no clinical control group was investigated. **Conclusion:** Our findings suggest reduced thresholds for facial anger and deficits in the discrimination of facial happiness in individuals with BPD. This biased perception is associated with alterations in very early visual as well as deficient structural and categorical processing of faces. The current data could help to explain the negative perception of others that may be related to the patients' impairments in interpersonal functioning.

Introduction

Patients with borderline personality disorder (BPD) have a biased and more negative perception of others. According to the DSM-5 model,¹ this may impair their empathy and thus contribute to their interpersonal dysfunctioning (e.g., by triggering reactive, aggressive responses).² There is increasing evidence for deficient or biased emotional face perception in individuals with BPD,³ which may contribute to the patients' exaggerated anticipation of social threat and difficulties in social interactions.⁴ Patients with BPD are, for instance, more likely to judge others as being more negative and aggressive;^{5,6} to make more errors in facial emotion classification tasks, particularly regarding negative emotional expressions;^{7–9} and to ascribe negative emotions to neutral faces.^{5,10} According to other findings, patients with BPD may be regarded as emotionally hypersensitive for subtle facial cues,

reflected by lower thresholds for emotion recognition in ambiguous or mixed emotional faces.^{11–13}

Daros and colleagues³ have proposed a model according to which the increased arousal of patients with BPD may either lead to an enhanced detection of subtle facial threat or hinder the classification of intense facial emotions in binding attentional resources by salient social cues. This model may explain seemingly inconsistent results from various emotion recognition tasks. It is also congruent with neuroimaging studies that have revealed enhanced and prolonged amygdala activations,^{14,15} a decreased prefrontal inhibition of the amygdala and altered insular and anterior cingulate activations^{16–18} for emotional stimuli in patients with BPD. In combining eye tracking with functional MRI (fMRI), we recently found more and faster initial eye movements toward the eyes, and thus the most threatening part of angry faces, associated with increased amygdala activation in patients with

Correspondence to: N.A. Izurieta Hidalgo, Department of General Psychiatry, Center of Psychosocial Medicine, University of Heidelberg, Voßstraße 4, Center of Psychosocial Medicine, 69115 Heidelberg, Germany; natalie.izurieta@med.uni-heidelberg.de

Submitted Aug. 6, 2014; Revised Feb. 23, 2015; Accepted Apr. 4, 2015; Early-released Aug. 18, 2015

DOI: 10.1503/jpn.140215

©2016 8872147 Canada Inc.

BPD.¹⁴ In line with the assumption of an initial hypersensitivity for acute social threat, this may indicate alterations in very early stages of facial emotion processing. Nevertheless, little is known about the time course of facial emotion processing and its emotion-specific differences in individuals with BPD.

The electroencephalography-based event-related potential (ERP) technique provides a high temporal resolution allowing a noninvasive online registration of cortical processes.¹⁹ Emotional faces elicit a specific sequence of ERPs over time,²⁰ starting with a very early positive wave, the P100, a component that is recorded at occipital sites about 100 ms after the presentation of any visual stimulus. The P100 is thus thought to reflect general primary visual analyses in the striate and extrastriate cortex that are not specific to (emotional) faces.²¹ This is followed by the N170, a negative wave recorded at temporo-occipital sites around 170 ms, which is thought to reflect the activation of face-specific neurons primarily in the fusiform gyrus or superior temporal sulcus with additional activation of a wider network, including the occipital cortex.^{22,23} The N170 may reflect the structural encoding of faces leading to the generation of a complete facial representation.²⁴ Although these early components primarily reflect initial, visual (P100) and structural face (N170) processing that are mostly independent of the semantic stimulus content,²⁵ there are also indications for modulatory influences of emotions that may be linked to stimulus saliency in terms of survival, reproduction and procreation arising from limbic input to the visual cortex.^{26–29} A later component, the P300, a positive wave that is recorded about 300–600 ms after stimulus onset at centroparietal sites, is particularly sensitive to psychological influences, such as anxiety, depression, alexithymia and aggression.^{30–33} The P300 is possibly generated in the extrastriate occipital inferior temporal and medial parietal cortices³⁴ and has been linked to subsequent higher stages of emotional face processing and decisional processes implicated in working memory updating or conscious detection of change, cognitive closure or emotion categorization.

There is limited literature on the electrophysiology underlying facial emotion processing in individuals with BPD. A magnetoencephalographic (MEG) study³⁵ revealed reduced M170 amplitudes, an N170 equivalent component, for faces displaying basic emotions, but also for neutral faces, houses and animals in patients with BPD compared with healthy individuals. This stimulus-independent reduction may reflect broad deficits in information processing at a relatively early perceptual level in individuals with BPD.³⁵ Furthermore, several studies revealed reductions (for exceptions, see studies by Meares and colleagues^{36,37}) in the P300 amplitude in patients with BPD compared with healthy volunteers (although not during emotion classification) in auditory oddball,^{38–40} go/no-go⁴¹ or Stroop tasks⁴² and after late feedback in a 2-choice reaction task.⁴³ Taken together, previous results suggest alterations in early and later ERP components in individuals with BPD, but investigations of the time course of these alterations during an emotion classification task are missing.

Thus, the aim of the study was to investigate the behavioural and electrophysiological correlates underlying emotional face processing in individuals with BPD with a particular focus on the patients' sensitivity for facial anger. We used a validated task^{12,44–47} with facial stimuli depicting blends of an-

gry and happy expressions, thus containing emotionally ambiguous information. This design bears the potential to unravel interpretative biases evident in an increased proportion of anger judgments of such stimuli. In fact, we have previously found a specific interpretative bias or hypersensitivity for facial anger in patients with BPD in a task with faces displaying blends of different emotional expressions.¹² In the present experiment, unmedicated female patients with BPD and healthy women had to classify facial emotions in a forced choice task while the electroencephalogram was recorded. The faces were preceded by task-irrelevant happily or angrily intoned sentences to increase the ecological validity of the task.⁴⁶ We expected that, compared with healthy volunteers, patients would show reduced thresholds for anger recognition but also more misclassifications and prolonged response latencies for predominantly angry faces. As for the emotionally intoned auditory stimuli, we expected an increased influence of the angry intonation in patients, leading to an increased likelihood of classifying ambiguous faces as angry. We explored the ERPs underlying structural and categorical facial emotion processing, expecting a hyperresponsiveness for facial anger in patients with BPD compared with healthy volunteers.

Methods

Participants

We studied unmedicated women with a current DSM-IV diagnosis of BPD and healthy women who had never received a psychiatric diagnosis or undergone any psychological or psychiatric treatment. Participants were recruited through the resident's registration office, advertisements and clinical referral from in- and outpatient units. Behavioural data were excluded in cases of positive toxicology screenings, brain damage or technical malfunctioning; EEG data were excluded if there was an insufficient number of trials owing to uncorrectable artifacts.

We verified handedness using the Edinburgh Handedness Inventory,⁴⁸ and participants were required to have normal or corrected-to-normal visual acuity. General exclusion criteria were neurologic disorders, current alcohol/drug abuse (urine toxicology screenings) or alcohol/drug abuse in the last 2 months (interview), severe medical illness and use of psychotropic medication for at least 2 weeks before participation in the study. Additional exclusion criteria for patients were lifetime diagnosis of schizophrenia, schizoaffective or bipolar disorder and reported alcohol/drug dependence in the last 12 months. Only patients who currently fulfilled at least 5 DSM-IV criteria for BPD were included in the study.

The Ethics Committee of the Medical Faculty of the University of Heidelberg approved the study. All participants provided written informed consent after the study procedures were fully explained, and they were paid for their participation.

Measures

For each patient with BPD and control, the diagnostic process comprised an extensive telephone interview (about 45 min) to screen for inclusion/exclusion criteria followed

by an onsite diagnostic appointment (about 3 h) involving structured clinical interviews (SCID-I;⁴⁹ International Personality Disorder Examination⁵⁰) to assess borderline diagnosis and axis I and II comorbidities as well as the Raven's progressive matrices⁵¹ as an estimate for intelligence. All interviews were performed by experienced diagnosticians (masters in psychology or doctorate in medicine with several years of clinical training) who underwent standardized diagnostic training before the study. In addition, we assessed borderline symptom severity using the Borderline Symptom List (BSL⁵²), state and trait anger using the State-Trait Anger Expression Inventory (STAXI,⁵³ state, trait, reaction, anger-in, anger-out, anger-control subscales), emotional dysregulation using the Difficulties in Emotion Regulation Scale (DERS,⁵⁴ acceptance, awareness, impulse, strategy, clarity, goals subscales) and depressiveness using the Beck Depression Inventory (BDI-II⁵⁵).

Experimental protocol

After arrival in the laboratory, participants provided a urine sample for toxicology screening and filled in questionnaires. To perform the experimental task, they were seated in a sound-attenuated, dimly lit room with their heads restrained in a chin rest 1 m from a 21-inch computer screen and equipped with in-ear headphones. The electrode sites on the participants' scalps and faces were cleaned and gently abraded before electrode placement. After a short training, participants completed 3 experimental runs of 15 min each (total participation time, including EEG preparation, training and short breaks between the runs was about 90 min).

Emotion classification paradigm

The emotion classification paradigm consisted of a previously validated^{12,45–47} forced-choice task in which participants had to classify faces depicting blends of angry and happy expressions by pressing a corresponding key for "angry" or "happy." This task has been deemed suitable to detect an interpretative bias or hypersensitivity for facial anger in patients with BPD.¹² Half of the faces within each emotional blend were preceded by a happily intoned sentence and half by an angrily intoned sentence with neutral content. These sentences were read by professional female and male actors and were validated in an independent sample before the experiment. Facial stimuli consisted of 2 male and 2 female faces⁵⁶ that were morphed in 10% steps, resulting in a series of 7 pictures showing blends of the angry and happy expressions: predominantly happy faces (80:20%, 70:30%, 60:40% happiness:anger), maximally ambiguous faces (50:50% happiness:anger) and predominantly angry faces (40:60%, 30:70%, 20:80% happiness:anger). In total, the experiment comprised 840 trials presented in 3 experimental runs of 280 trials each in pseudorandomized order with no more than 3 presentations of the same emotional condition in a row. Each trial started with the presentation of a fixation cross and auditory presentation of an angrily or happily intoned sentence (2000 ms), a fixation cross without auditory

stimulation (100 ms) followed by a facial stimulus (response-locked presentation, 1100 ms) and another fixation cross (450–650 ms).

Data acquisition and processing

The EEG was recorded with Ag/AgCl electrodes from 60 sites (equidistant reference system, Easy-Cap GmbH) using an average reference. Additionally, vertical and horizontal electro-oculograms (EOGs) were recorded from the epicanthus of each eye and from the supra- and infraorbital positions of the left eye. Electrode impedances were kept below 10 k Ω and signals were amplified with a 72-Kanal Quick-Amp amplifier (Brain Products, GmbH). The pass-band was set to 0.01–200 Hz; the signals were digitized at 1000 Hz and stored to hard disk for later analysis. Data processing was conducted using Brain Vision Analyzer 2.0 (Brain Products, GmbH) and included renaming of EEG channels to a 10/20 system⁵⁷ (small position deviations are marked with "''"), digital filtering (0.1–40 Hz, 24dB/octave roll-off), correction of eye-movement artifacts using an algorithm provided by Gratton and colleagues,⁵⁸ semiautomatic rejection of trials with nonphysiologic artifacts, segmentation into 1300 ms periods (–100 ms to 1200 ms from stimulus onset) and baseline correction (reference: –100 ms to 0 ms from stimulus onset). Separate averages were computed for each electrode and combination of facial and auditory emotion at individual and group levels. Statistical analyses were conducted on ERP peaks for the P100 (maximum positive peak 80–130 ms at O1', Oz, O2') and N170 (maximum negative peak 130–230 ms at TP7', TP8', TP9', TP10', P7', P8', P9', P10', O1', O2', O9', O10') components and mean amplitude for the P300 (350–600 ms at Cz, CP1', CP2', CPz, CP3', CP4', P1', Pz, P2').

Statistical analysis

Behavioural data (proportion of anger classifications and response latencies) were subjected to analyses of variance (ANOVAs) using group (BPD, control) as a between-subjects factor and facial emotion (7 levels of anger-happiness intensity) and voice intonation (angry, happy) as within-subjects factors. We also ran analyses using multivariate ANOVA, which yielded similar results for the behavioural and ERP data in that all reported effects were significant using multivariate analyses except for the group $\times$ emotion interaction in the behavioural data, which reached only trend-level significance ($p = 0.08$). Hypothesis-driven univariate ANOVAs, however, revealed significant group differences in the proportion of anger classification for faces with 20% ($p = 0.015$) and 30% anger ($p = 0.011$) and in response latencies for faces with 20% anger ($p = 0.028$). Behavioural results remained stable after exclusion of 2 participants without reliable EEG data. We averaged across the sex of the faces because preliminary behavioural and ERP data analyses did not reveal any effect of sex (all $p > 0.10$). For the ERP data, separate ANOVAs had to be performed for the P100, N170 and P300 owing to the different localizations of these ERP components

(see the data acquisition and processing section for the different sets of electrodes). Besides the factors group, facial emotion and voice intonation, analyses of P100 and N170 additionally comprised the within-subjects factor electrode site (P100: left, middle, right) and hemisphere (N170: left, right), respectively. Since response latencies were skewed, they were log-transformed before analysis. Where appropriate, we applied the Huynh–Feldt⁵⁹ procedure to correct for potential violations of the sphericity assumption. All statistical analyses were 2-tailed, and we considered results to be significant at $p < 0.05$. In cases of significant effects, we used Dunn's multiple comparison as post hoc tests, which include a Bonferroni correction for multiple testing. We additionally performed Pearson correlations to explore trade-off between accuracy (proportion of anger classification) and speed (response latencies) in the behavioural data and associations between behavioural and ERP data as well as self-reported traits. The results of these exploratory correlational analyses may be interpreted only descriptively owing to the limited power and the cumulation of type I errors.

Results

Participants

We recruited 36 unmedicated women with BPD (mean age 26.7 ± 6.0 [range 18–40] yr) and 29 healthy women (mean age 26.2 ± 6.0 [range 19–48] yr) for participation in the study. Behavioural data were excluded for 3 patients and 1 control (2 owing to positive toxicology screenings, 1 owing to organic brain damage and 1 owing technical malfunctioning), and EEG data were excluded for 1 patient and 1 control owing to uncorrectable artifacts. Thus, the results are based on the behavioural data of 33 patients and 28 controls and the EEG data of 32 patients and 27 controls. All participants were right-handed. Detailed demographic and psychometric information and group comparisons are provided in Table 1, and comorbid disorders of the patient group are presented in Table 2.

Table 1: Demographic and psychometric data

Category	Group; mean $\pm$ SD		p value
	BPD ($n = 33$)	Control ($n = 28$)	
Age, yr	26.7 ± 6.0	26.2 ± 6.0	0.55
IQ*	109.7 (11.4)	111.7 (11.3)	0.96
No. of DSM-IV BPD criteria (range)	6.58 ± 1.25 (5–9)	0 ± 0.2 (0–1)	< 0.001
Borderline symptomatology†	2.1 ± 0.7	0.2 ± 0.2	< 0.001
Anger‡	24.5 ± 6.7	13.5 ± 2.7	< 0.001
Depressiveness§	26.2 ± 11.1	3.4 ± 3.1	< 0.001
Difficulties in emotion regulation¶	118.1 ± 17.3	60.5 ± 13.2	< 0.001

BPD = borderline personality disorder; SD = standard deviation.

*Raven Progressive Matrices.

†Borderline Symptom Checklist.

‡State-Trait Anger Inventory.

§Beck Depression Inventory.

¶Difficulties in Emotion Regulation Scale.

Behavioural results

Proportion of anger classification

Across diagnostic groups, participants generally classified predominantly angry faces (60%–80% anger) more often as angry and predominantly happy faces (20%–40% anger) as happy, while there was no general tendency for maximally ambiguous faces (main effect of facial emotion $F_{6,354} = 973.55$, $p < 0.001$, $\eta^2 = 0.94$). Proportion of anger classification was also influenced by the auditory stimuli (main effect of voice intonation $F_{1,59} = 27.86$, $p < 0.001$, $\eta^2 = 0.32$; interaction of voice intonation and facial emotion $F_{6,354} = 5.68$, $p < 0.001$, $\eta^2 = 0.09$). Post hoc analyses showed that the voice intonation particularly influenced the rating of faces with 40%–60% anger (all $p < 0.01$). For instance, when an angrily intoned sentence preceded such a face, participants rated this face more often as angry.

The ANOVA also revealed a significant group $\times$ facial emotion interaction ($F_{6,354} = 2.89$, $p = 0.049$, $\eta^2 = 0.05$). According to post hoc tests, patients with BPD classified predominantly happy faces (20%–40% anger) more often as angry than controls ($p < 0.01$) and tended to misclassify predominantly angry faces (60%–80% anger) as happy ($p > 0.05$, Fig. 1A). There was no significant group $\times$ voice intonation interaction ($p = 0.21$).

Response latency

Overall, participants responded faster for the less ambiguous faces and slower as faces turned ambiguous (main effect of facial emotion $F_{6,354} = 55.78$, $p < 0.001$, $\eta^2 = 0.49$). The emotional voice intonation also had a significant effect: across groups, participants showed faster responses to predominantly happy faces (20%–40% anger) that were preceded by a happily intoned voice and to predominantly angry faces (60%–80% anger) preceded by an angry voice compared with predominantly happy faces (20%–40% anger) preceded by angry voices and predominantly angry faces (60%–80% anger) preceded by happy voices (interaction of voice intonation $\times$ facial emotion $F_{6,354} = 2.58$, $p = 0.023$, $\eta^2 = 0.04$).

According to a significant group $\times$ facial emotion interaction ($F_{6,354} = 3.00$, $p = 0.047$, $\eta^2 = 0.05$, Fig. 1B), patients with BPD responded slower than healthy volunteers, especially for

Table 2: Comorbidities of patients with borderline personality disorder

Comorbidity	Period; no. (%)	
	Current	Lifetime
Major depression	8 (22)	20 (56)
Dysthymia	1 (3)	0
Anxiety disorders	15 (42)	5 (14)
Posttraumatic stress disorder	12 (33)	2 (6)
Eating disorders	10 (28)	8 (22)
Substance use disorders*	2 (6)	9 (25)
Somatoform disorders	6 (17)	0
Avoidant personality disorder	10 (28)	1 (3)
Antisocial personality disorder	1 (3)	0

*Two current nicotine abusers and 9 participants with remitted substance use disorders.

predominantly happy faces (20%–40% anger, $p < 0.01$). There was no significant group $\times$ voice intonation interaction ($p = 0.64$).

As there were no significant interactions of group $\times$ voice intonation in the behavioural data (all $p > 0.20$) and no indications of such an effect in the ERP data (all $p > 0.10$ for mean amplitudes), the factor voice intonation was dropped from the analyses of the ERP data.

ERP results

P100

Across groups, P100 peak amplitudes were generally larger at lateral (O1' and O2') than midline (Oz) electrode sites (main effect of electrode $F_{2,114} = 40.89$, $p < 0.001$, $\eta^2 = 0.42$, $\epsilon = 0.93$).

Fig. 1. (A) Mean proportion of anger classification in percent (± 1 standard error). (B) Mean response latencies in milliseconds (± 1 standard error) of patients with borderline personality disorder (BPD) and healthy controls (HC) averaged across auditory stimulus conditions. The X axis shows percentage of facial anger. ** $p < 0.01$, * $p < 0.05$.

In addition, the ANOVA revealed a trend-level significant main effect of group ($F_{1,57} = 2.83, p = 0.10, \eta^2 = 0.05$) and a significant group $\times$ facial emotion $\times$ electrode interaction ($F_{12,684} = 1.91, p = 0.042, \eta^2 = 0.03, \epsilon = 0.83$; Fig. 2A and B). Post hoc tests showed that the P100 was lateralized in healthy volunteers, with enhanced left hemispheric amplitudes (O1' compared with Oz and O2'; anger intensities all $p < 0.01$), while patients with BPD did not show this lateralization effect (all $p > 0.05$ except for 50% anger: $p < 0.01$; Fig. 2C). Moreover, in healthy volunteers, the P100 amplitude was modulated by facial emotion. They had increased peaks for faces displaying 80% anger compared with faces displaying 20% anger ($p < 0.01$ for all electrodes), while patients with BPD showed enhanced P100 amplitudes independent of facial emotion (all $p > 0.10$).

Across groups, P100 peak latency tended to be increased for maximally ambiguous faces (main effect of facial emotion $F_{6,342} = 2.17, p = 0.06, \eta^2 = 0.04, \epsilon = 0.83$) and was delayed at left compared with midline and right hemispheric electrodes (main effect of electrode $F_{2,114} = 8.92, p < 0.001, \eta^2 = 0.14, \epsilon = 1.0$).

According to a significant main effect of group ($F_{1,57} = 4.17, p = 0.046, \eta^2 = 0.07$), P100 peaks were delayed in patients with BPD compared with healthy volunteers (100.4 ms v. 95.3 ms); however, neither facial emotion nor electrode significantly interacted with group (all $p < 0.10$).

N170

Across groups, N170 peak amplitudes were more pronounced at right hemispheric electrodes (main effect of hemisphere $F_{1,57} = 7.89, p = 0.007, \eta^2 = 0.12, \epsilon = 1.0$). N170 amplitudes were also modulated by facial emotion (main effect of facial emotion $F_{6,342} = 4.48, p < 0.001, \eta^2 = 0.07, \epsilon = 1.0$), with larger amplitudes for faces with 80% anger than faces displaying lower anger intensity (20%–70% anger, all $p < 0.01$).

In addition, there was a significant group $\times$ hemisphere interaction ($F_{1,57} = 13.62, p = 0.001, \eta^2 = 0.19, \epsilon = 1.0$) according to which the N170 lateralization was present only in healthy volunteers ($p < 0.01$) but not in patients with BPD ($p > 0.05$). Thus, compared with healthy volunteers, patients with BPD

Fig. 2. (A) Mean event-related potential (ERP) at Oz electrode position in patients with borderline personality disorder (BPD) and healthy controls (HC) averaged across auditory stimulus conditions. (B) Head views of P100 and N170. (C) Peak amplitude of P100. (D) Peak amplitude of N170. ** $p < 0.01$, * $p < 0.05$.

had reduced N170 peak amplitudes at right hemispheric electrodes ($p < 0.01$, Fig. 2D). There was no significant interaction of group $\times$ facial emotion ($p = 0.20$) or group $\times$ facial emotion $\times$ hemisphere ($p = 0.64$).

There were no significant main or interaction effects for N170 peak latency (all $p > 0.10$).

P300

P300 amplitudes were largest for predominantly angry (60%–80% anger) or happy (20%–40% anger) faces and smallest for maximally ambiguous faces (50% anger, main effect of facial emotion $F_{6,342} = 24.86$, $p < 0.001$, $\eta^2 = 0.30$, $\epsilon = 0.98$).

According to a significant interaction of group $\times$ facial emotion ($F_{6,342} = 3.72$, $p = 0.002$, $\eta^2 = 0.06$, $\epsilon = 0.98$), P300 amplitudes were generally reduced in patients with BPD compared with healthy volunteers (all $p < 0.01$), with the strongest group differences for predominantly happy faces (20%–40% anger; Fig. 3). A closer inspection revealed that patients showed increased P300 amplitudes only for the most angry faces (70%–80% anger) compared with maximally ambiguous faces (50% anger, $p < 0.01$) as well as the most happy

faces (20%–30% anger, $p < 0.01$). This strongly differed from the pattern found in healthy volunteers, whose P300 amplitudes were larger for both the most angry and happy faces compared with maximal ambiguous faces (all $p < 0.01$).

Correlations

In both groups, response latencies and proportion of anger classification were positively correlated for predominantly happy faces (BPD: $r_{31} \geq 0.61$, $p \leq 0.001$; control: $r_{27} \geq 0.40$, $p \leq 0.040$); participants who responded slowest also classified predominantly happy faces more often as angry. In patients with BPD, anger classification and response latencies for predominantly happy faces were positively associated with N170 amplitudes (20%–40% anger: $r_{30} \geq 0.32$, $p \leq 0.08$), and response latencies for ambiguous and predominantly angry faces were negatively associated with P300 amplitudes (50%–80% anger: $r_{30} \geq -0.37$, $p \leq 0.039$). In controls, there were no correlations between behavioural and ERP data ($r_{26} \leq 0.30$, $p \geq 0.10$). In both groups, P100 and N170 amplitudes were positively associated (BPD: right hemispheric electrode for 20%–50% anger $r_{30} \geq 0.33$, $p \leq 0.06$;

Fig. 3. (A) Mean event-related potential (ERP) at the Pz electrode position of patients with borderline personality disorder (BPD) and healthy controls (HC) averaged across auditory stimulus conditions. (B) Head views and (C) mean amplitude of P300. * $p < 0.05$; ** $p < 0.01$.

control: right and left hemispheric electrodes for 20%–80% anger $r_{26} \geq 0.40$, $p \leq 0.038$), with smaller (less negative) N170 amplitudes in those participants with greater (more positive) P100 amplitudes. In healthy volunteers, N170 and P300 amplitudes were positively correlated for predominantly happy and ambiguous faces ($r_{26} \geq 0.33$, $p \leq 0.038$). No correlations were found between N170 and P300 amplitudes in patients with BPD and between P100 and P300 amplitudes in both groups ($r < 0.30$, all $p > 0.10$).

We additionally found positive correlations between proportion of anger classification and the DERS emotional awareness subscale (30%–70% anger: $r_{31} \geq 0.34$, $p \leq 0.06$), BSL borderline symptom severity (20%–60% anger: $r_{30} \geq 0.31$, $p \leq 0.09$) and BDI depressiveness (40%–60% anger: $r_{30} \geq 0.32$, $p \leq 0.09$) in patients with BPD but not in healthy volunteers ($r_{27} \leq 0.30$, $p \geq 0.10$). In patients with BPD, P100 amplitudes were positively correlated with DERS emotional awareness (20%–80% anger: $r_{30} \geq 0.33$, $p \leq 0.07$; no associations in controls: $r_{27} \leq 0.30$, $p \geq 0.10$), while P300 amplitudes were positively related with anger suppression (STAXI anger-in; 20%–80% anger: $r_{30} \geq 0.40$, $p \leq 0.024$; no associations in controls: $r_{27} \leq 0.30$, $p \geq 0.10$).

Comorbid psychiatric disorders

As influences of depression, posttraumatic stress disorder (PTSD) and social anxiety on emotional faces processing have been previously reported, we compared patients with BPD with and without depressive disorder (current diagnosis of major depression or dysthymia), current PTSD and current social anxiety disorders (current diagnosis of social phobia or avoidant personality disorder) in additional group $\times$ emotional facial expression ANOVAs. These analyses revealed no significant main effect of group or group $\times$ facial expression interaction for depression ($n = 10$ with and $n = 32$ without current depressive disorder), PTSD ($n = 12$ with and $n = 21$ without current PTSD) and social anxiety ($n = 14$ with and $n = 18$ without current social anxiety disorder) for behavioural and ERP data (all $p > 0.05$).

Discussion

To our knowledge, this is the first study to investigate temporal dynamics of facial emotion processing in patients with BPD using an EEG ERP technique with a particular focus on the patients' sensitivity to facial anger. The results confirm alterations or biases in very early stages of face processing, as has been hypothesized on the basis of eye-tracking and neuroimaging studies.^{14–16} In addition to these early alterations, patients with BPD showed deficiencies in further stages of structural and categorical face processing that may be related to previously reported deficits in emotion classification, supporting a recent model on altered facial classification.³

The present study revealed an enhanced likelihood to misclassify predominantly happy faces (20%–40% anger) as angry in patients with BPD. This is consistent to previous results revealing a specific bias for anger in a similar task with faces displaying blends of all 6 basic emotions¹² and may thus indicate an increased sensitivity or lower threshold for subtle cues of social threat. This is in line with increased reflexive gaze orientation toward threatening facial cues.¹⁴ Further in-

dications for such a hypersensitivity were also found in neuroimaging studies that revealed enhanced and prolonged amygdala activations and a decreased prefrontal inhibition of amygdala for emotional faces and socially threatening stimuli in patients with BPD.^{14–16} In the present study, a higher proportion of anger classification was found in patients with increased levels of self-reported unawareness of their own emotions. In addition, misclassification of predominantly happy faces (20%–40% anger) as angry was related to the severity of borderline symptomatology, whereas classification of more ambiguous faces (40%–60% anger) as angry was related to depressiveness. These exploratory correlational results support the alternative DSM-5 model of BPD claiming a more negative perception of others that might underlie the patients' impairments in interpersonal functioning.¹ Importantly, the present results may also indicate deficits in the classification of positive facial emotions, such as happiness, in patients with BPD. This is consistent with the results of a recent behavioural study⁶⁰ that revealed both an earlier detection for anger and a slower recognition of facial happiness in patients with BPD using different intensities of happy, angry and fearful facial expressions. In line with the previously reported behavioural deficiencies in classifying faces displaying high intensities of negative and particularly angry expressions,³ patients in the present study also tended to classify predominantly angry faces (60%–80% anger) less accurately than controls. Based on the model proposed by Daros and colleagues³ in which misclassification of intense negative facial emotions are partially explained by a depletion of cognitive resources required to disengage attention from highly salient emotional stimuli, we expected longer response latencies specifically for the most angry faces (i.e., 80% anger). However, patients with BPD responded generally slower than healthy volunteers (20%–80% anger), and this effect was most pronounced for predominantly happy faces (20%–40% anger). The positive correlations between proportion of anger classification and response latencies suggests that the patients' misclassifications did not result from a trade-off between accuracy and speed and thus do not reflect an impulsive response style. Importantly, subgroups of patients with BPD with and without a current diagnosis of depressive disorders, PTSD and social anxiety disorders, which are highly prevalent in individuals with BPD and have been previously associated with a hypervigilance for social threat cues^{61,62} and a reduced facial happiness recognition, did not significantly differ in the proportion of anger classification, response latencies and ERP data.

By adding emotionally intonated auditory stimuli, we could investigate the sensitivity for facial anger in a more ecologically valid setting.⁴⁶ Contrary to our hypotheses and the previously reported deficits in multisensory integration of emotional information in patients with BPD,⁶³ the emotionally intonated auditory stimuli affected the proportion of facial anger classification similarly in patients and healthy volunteers. Both groups showed faster responses in congruent trials of voice and facial emotion than in incongruent trials, and the voice had strongest effects on the classification of faces with 40%–60% anger. This result means that patients with BPD — comparable to healthy

controls — are able to benefit from a second sensory channel in case the information is synergistic. Further research is required to investigate auditory/bimodal emotion processing in individuals with BPD.

Besides behavioural alterations, the present study revealed a pattern of changes in very early and intermediate stages of facial emotion processing in patients with BPD. Irrespective of facial emotion, patients showed increased P100 amplitudes, a component that reflects primary visual analyses in the striate and extrastriate cortex.²¹ Contrary to this, P100 amplitudes were modulated by facial emotion in healthy volunteers, with significantly greater amplitudes for highly angry (80% anger) faces than faces with 20%–70% anger. This pattern relates to previous reports of enhanced P100 amplitudes for negative compared with happy⁶⁴ or neutral faces⁶⁵ in healthy volunteers and may reflect increased limbic input to the visual cortex in the presence of social threat.²⁶ Interestingly, enhanced P100 amplitudes for various emotional faces were previously reported in students with high social anxiety⁶⁶ as well as in healthy volunteers after an experimental provocation of reactive aggression,⁶⁷ indicating a trait- (social anxiety, BPD) and state-dependent (provocation) early hypervigilance in the visual cortex, which may increase the attention for incoming social information. The enhanced P100 amplitudes of patients with BPD in the present study could thus reflect the patients' general hyperresponsiveness for (potentially threatening) information, which, according to the alternative DSM-5¹ model of BPD, may, in association with a more negative perception of others, lead to impaired interpersonal functioning.

In addition to this early hyperresponsiveness, patients with BPD showed deficits in subsequent stages of emotional face processing, as reflected by reduced temporo-occipital N170 and centroparietal P300 amplitudes. The N170, which reflects structural analyses of facial information, including the structural encoding of faces, in the fusiform face area and the superior temporal sulcus²³ was lateralized in healthy volunteers. This lateralization is in agreement with the hypothesis of a preferential involvement of the right hemisphere in configurational facial analysis that has been confirmed by increased right posterior–inferior temporal N170 amplitudes for facial stimuli in previous ERP studies.^{25,68} Patients with BPD had significantly reduced right hemispheric N170 amplitudes and thus no right-lateralization. Comparably, Merkl and colleagues³⁵ reported decreased temporo-occipital M170 amplitudes for facial stimuli in patients with BPD compared with healthy volunteers. Positive correlations between P100 and N170 amplitudes suggest a cascade effect with strongest deficits in structural processing and encoding of faces in patients with the largest initial visual hyperresponsiveness, which might explain the patients' face classification impairments.

Patients with BPD also had reduced centroparietal P300 amplitudes. This component is thought to reflect higher stages of emotional face processing, including emotional categorization, working memory updating and cognitive closure processes that are highly sensitive to psychological influences. For instance, reduced P300 amplitudes in impulse control tasks have been reported in individuals with BPD^{41–43} and impulse control disorders or reactive aggression (for a review,

see Patrick⁶⁹) and have been interpreted as a correlate of the externalizing factor that these disorders have in common.⁶⁹ Although the results of these tasks are difficult to compare with the present facial processing results, the general reductions in P300 amplitudes may partly reflect enhanced externalizing behaviour of patients with BPD. Supporting this assumption, we found the largest P300 amplitudes in patients who reported greatest withholding or suppression of anger. Additionally, the pattern of P300 alterations of patients also indicates deficits in attentive categorization of the emotional intensity. In line with the previously described P300 modulation by both pleasant and unpleasant stimuli,²⁰ healthy participants showed increased amplitudes for both predominantly happy (20%–40% anger) and angry (60%–80% anger) compared with maximally ambiguous facial expressions (50% anger), while patients with BPD showed poorer discrimination between faces displaying 20%–60% anger and only enlarged P300 amplitudes for faces displaying 70%–80% anger. Similarly reduced P300 amplitudes for predominantly happy faces have been previously reported in depressive individuals,⁷⁰ which may suggest an association with the patients' negative perception of others and emotional negativity. Thus, the results indicate deficits in the processing and discrimination of facial happiness and a hypervigilance to full-blown expressions of anger, which could at least partially explain the deficits observed in the behavioural task.

Limitations

To our knowledge, this is the first study to examine the temporal dynamics of facial emotion processing in a large group of patients with BPD compared with age- and IQ-matched healthy women using an EEG-based ERP technique in an ecologically more valid setting. However, several limitations of the present study should be noted. First, we included only female patients with a number of comorbidities. Although this is in line with most of the previously published studies on emotional face processing in patients with BPD³ and reflects a typical pattern of comorbid psychiatric disorders,⁷¹ borderline-specific conclusions should be drawn with caution. We would therefore encourage studies that include clinical control groups, such as patients with major depression, social anxiety disorder or PTSD, as well as male participants. Second, further studies need to include self-reports and physiologic measures to assess arousal, alexithymia or empathy, which have been hypothesized to be important factors in explaining the alterations in emotional face processing in patients with BPD.^{3,72} Third, the temporal restrictions and the need of a large amount of trials owing to the small signal-to-noise ratio of the ERPs made us focus on facial anger and happiness and to not include a silent control condition. Therefore, we cannot rule out a general effect of auditory stimulation on information processing in patients with BPD. It remains unclear why the auditory stimulation did not have strong effects on the facial emotion processing in the present study, which is contrary to auditory modulation of early stages of face processing found in healthy individuals.⁷³ It may be noted that auditory stimuli were

presented before facial expression and were task-irrelevant, which may have reduced their impact on facial emotion processing.⁴⁶ The present results therefore need to be regarded as a first attempt to investigate the time course of alterations in facial information processing in patients with BPD that raises further questions for future studies, including a more precise investigation of interactions between auditory and visual stimulation. Moreover, we cannot disentangle a specific bias for facial anger from an inability to recognize and discriminate happiness or a general tendency to misclassify emotions. Yet, our results are highly consistent to a previously reported lower threshold for facial anger in a study using various combinations of all 6 basic emotions without an auditory stimulation.¹² Further studies are, however, needed to disentangle deficits in the processing and recognition of facial happiness from an anger hypersensitivity.

Conclusion

Our data indicate a very early hyperresponsiveness for emotional faces independent of the emotional intensity or its valence in patients with BPD and subsequent deficits in both structural and categorical emotional face processing (e.g., poorer discrimination of happy and emotionally ambiguous faces, but a hyperresponsiveness for highly angry faces). These electrophysiological findings may underline the behavioural findings of enhanced sensitivity for subtle cues of facial threat, deficits in the discrimination of facial happiness as well as deficits in the classification of highly socially threatening facial stimuli. Electrophysiological and behavioural alterations were strongest in patients with most negative affectivity and unawareness of own emotions. The ability to decode emotional information from facial expressions is important for the regulation of emotion and the fine graining of social responses. Different mechanisms, such as an enhanced sensitivity for facial anger or deficits in the recognition of positive facial information, may contribute to findings of a more negative perception of others in patients with BPD and may be a prerequisite for the patients' impaired interpersonal functioning. The results can thus be regarded as a further step to understand the psychological and neurobiological underpinnings of BPD, which may help to improve psychotherapeutic strategies for the patients' impairments in interpersonal functioning.

Acknowledgments: The authors thank K. Herwig for supporting the EEG measurement and analysis, L. Kramer, L. Müller, A. Gäbel and D. Gescher for their help with data collection, the Informatics team for programming, and the team of the Clinical Research Group on Mechanisms of Disturbed Emotion Processing in Borderline Personality Disorder (KFO 256) for participant recruitment and organization. The study was supported by grants from the German Research Foundation (DFG) awarded to S.C. Herpertz (He 2660/12-1; He 2660/7-2). Preliminary data of this study were presented at the DGPPN conference 2013 in Berlin and at the EPA Conference 2014 in Munich, Germany.

Affiliations: From the Department for General Psychiatry, Center of Psychosocial Medicine, University of Heidelberg, Germany (Izurieta Hidalgo, Nagy, Mancke, Herpertz, Bertsch); the Department for Child and Adolescent Psychiatry, Center of Psychosocial Medicine,

University of Heidelberg, Heidelberg, Germany (Oelkers-Ax); and the Department of Psychosomatic Medicine, Central Institute of Mental Health Mannheim, Medical Faculty Mannheim/Heidelberg University, Heidelberg, Germany (Bohus).

Competing interests: None declared.

Contributors: N. Izureta, R. Oelkers-Ax, M. Bohus, S. Herpertz and K. Bertsch designed the study. N. Izureta, K. Nagy, F. Mancke, M. Bohus and K. Bertsch acquired the data, which N. Izureta, R. Oelkers-Ax, S. Herpertz and K. Bertsch analyzed. N. Izureta and K. Bertsch wrote the article, which all authors reviewed and approved for publication.

References

1. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*, 5th ed. Arlington (VA): American Psychiatric Publishing; 2013.
2. Mancke F, Herpertz SC, Bertsch K. Aggression in borderline personality disorder — a multidimensional model. *Personal Disord* 2015; 6:278-91.
3. Daros AR, Zakzanis KK, Ruocco AC. Facial emotion recognition in borderline personality disorder. *Psychol Med* 2013;43:1953-63.
4. Domes G, Schulze L, Herpertz SC. Emotion recognition in borderline personality disorder — a review of the literature. *J Pers Disord* 2009; 23:6-19.
5. Arntz A, Veen G. Evaluations of others by borderline patients. *J Nerv Ment Dis* 2001;189:513-21.
6. Barnow S, Stopsack M, Grabe HJ, et al. Interpersonal evaluation bias in borderline personality disorder. *Behav Res Ther* 2009;47:359-65.
7. Levine D, Marziali E, Hood J. Emotion processing in borderline personality disorder. *J Nerv Ment Dis* 1997;185:240-6.
8. Bland AR, Williams CA, Scherer K, et al. Emotion processing in borderline personality disorders. *Issues Ment Health Nurs* 2004;25:655-72.
9. Unoka Z, Fogd D, Füzy M, et al. Misreading the facial signs: specific impairments and error patterns in recognition of facial emotions with negative valence in borderline personality disorder. *Psychiatr Res* 2011;189:419-25.
10. Wagner AW, Linehan MM. Facial expression recognition ability among women with borderline personality disorder: implications for emotion regulation? *J Pers Disord* 1999;13:329-44.
11. Daros AR, Uliaszek AA, Ruocco AC. Perceptual biases in facial emotion recognition in borderline personality disorder. *Personal Disord* 2014;5:79-87.
12. Domes G, Czeschnek D, Weidler F, et al. Recognition of facial affect in borderline personality disorder. *J Pers Disord* 2008;22:135-47.
13. Lynch TR, Rosenthal MZ, Kosson DS, et al. Heightened sensitivity to facial expressions of emotion in borderline personality disorder. *Emotion* 2006;6:647-55.
14. Bertsch K, Gamer M, Schmidt B, et al. Oxytocin and reduction of social threat hypersensitivity in women with borderline personality disorder. *Am J Psychiatry* 2013;170:1169-77.
15. Hazlett EA, Zhang J, New AS, et al. Potentiated amygdala response to repeated emotional pictures in borderline personality disorder. *Biol Psychiatry* 2012;72:448-56.
16. Ruocco AC, Amirthavasagam S, Choi-Kain LW, et al. Neural correlates of negative emotionality in borderline personality disorder: an activation-likelihood-estimation meta-analysis. *Biol Psychiatry* 2013;73:153-60.
17. Koenigsberg HW, Denny BT, Fan J, et al. The neural correlates of anomalous habituation to negative emotional pictures in borderline and avoidant personality disorder patients. *Am J Psychiatry* 2014; 171:82-90.
18. Schulze L, Domes G, Krüger A, et al. Neuronal correlates of cognitive reappraisal in borderline patients with affective instability. *Biol Psychiatry* 2011;69:564-73.
19. Hillyard SA, Kutas M. Electrophysiology of cognitive processing. *Annu Rev Psychol* 1983;34:33-61.
20. Eimer M, Holmes A. Event-related brain potential correlates of emotional face processing. *Neuropsychologia* 2007;45:15-31.
21. Gomez Gonzalez CM, Clark VP, Luck SJ, et al. Sources of attention-sensitive visual event-related potentials. *Brain Topogr* 1994;7:41-51.

22. Herrmann MJ, Ehlis AC, Muehlberger A, et al. Source localization of early stages of face processing. *Brain Topogr* 2005;18:77-85.
23. Eimer M, Kiss M, Nicholas S. Response profile of the face-sensitive N170 component: a rapid adaptation study. *Cereb Cortex* 2010;20:2442-52.
24. Jeffreys DA. Evoked potentials studies of face and object processing. *Vis Cogn* 1996;3:1-38.
25. Balconi M, Lucchiari C. Event-related potentials related to normal and morphed emotional faces. *J Psychol* 2005;139:176-92.
26. Vuilleumier P, Driver J. Modulation of visual processing by attention and emotion: windows on causal interactions between human brain regions. *Philos Trans R Soc Lond B Biol Sci* 2007;362:837-55.
27. Delle-Vigne D, Kornreich C, Verbanck P, et al. Subclinical alexithymia modulates early audio-visual perceptive and attentional event-related potentials. *Front Hum Neurosci* 2014;8:106.
28. Blau VC, Maurer U, Tottenham N, et al. The face-specific N170 component is modulated by emotional facial expression. *Behav Brain Funct* 2007;3:7.
29. Schupp HT, Stockburger J, Bublatzky F, et al. Explicit attention interferes with selective emotion processing in human extrastriate cortex. *BMC Neurosci* 2007;8:16.
30. Cuthbert BN, Schupp H, Bradley M, et al. Brain potentials in affective picture processing: covariation with autonomic arousal and affective report. *Biol Psychol* 2000;52:95-111.
31. Campanella S, Falbo L, Rossignol M, et al. Sex differences on emotional processing are modulated by subclinical levels of alexithymia and depression: a preliminary assessment using event-related potentials. *Psychiatry Res* 2012;197:145-53.
32. Rossignol M, Anselme C, Vermeulen N, et al. Categorical perception of anger and disgust facial expression is affected by non-clinical social anxiety. *Brain Res* 2007;1132:166-76.
33. Bertsch K, Böhnke R, Kruk MR, et al. Influence of aggression on information processing in the emotional stroop task — an event-related potential study. *Front Behav Neurosci* 2009;3:28.
34. Sabatinelli D, Lang PJ, Keil A, et al. Emotional perception: correlation of functional MRI and event-related potentials. *Cereb Cortex* 2007;17:1085-91.
35. Merkl A, Ammelburg N, Aust S, et al. Processing of visual stimuli in borderline personality disorder: a combined behavioural and magnetoencephalographic study. *Int J Psychophysiol* 2010;78:257-64.
36. Meares R, Melkonian D, Gordon E, et al. Distinct pattern of P3a event-related potential in borderline personality disorder. *Neuroreport* 2005;16:289-93.
37. Meares R, Schore A, Melkonian D. Is borderline personality a particularly right hemispheric disorder? A study of P3a using single trial analysis. *Aust N Z J Psychiatry* 2011;45:131-9.
38. Blackwood DH, St Clair DM, Kutcher SP. P300 event-related potential abnormalities in borderline personality disorder. *Biol Psychiatry* 1986;21:560-4.
39. Kutcher SP, Blackwood DH, St Clair D, et al. Auditory P300 in borderline personality disorder and schizophrenia. *Arch Gen Psychiatry* 1987;44:645-50.
40. Drake ME Jr, Phillips BB, Pakalnis A. Auditory evoked potentials in borderline personality disorder. *Clin Electroencephalogr* 1991;22:188-92.
41. Ruchow M, Groen G, Kiefer M, et al. Response inhibition in borderline personality disorder: event-related potentials in a Go/Nogo task. *J Neural Transm* 2008;115:127-33.
42. Houston RJ, Bauer LO, Hesselbrock VM. Effects of borderline personality disorder features and a family history of alcohol or drug dependence on P300 in adolescents. *Int J Psychophysiol* 2004;53:57-70.
43. De Bruijn ER, Grootens KP, Verkes RJ, et al. Neural correlates of impulsive responding in borderline personality disorder: ERP evidence for reduced action monitoring. *J Psychiat Res* 2006;40:428-37.
44. Calder A, Young D, Rowland D, et al. Facial emotion recognition after bilateral amygdala damage: differentially severe impairment of fear. *Cognitive Neuropsych* 1996;13:699-745.
45. de Gelder B, Böcker KB, Tuominen J, et al. The combined perception of emotion from voice and face: early interaction revealed by human electric brain responses. *Neurosci Lett* 1999;260:133-6.
46. DeGelder B, Vroomen J. The perception of emotions by ear and by eye. *Cognition and Emotion* 2000;14:289-311.
47. Jusyte A, Schönenberg M. Threat processing in generalized social phobia: an investigation of interpretation biases in ambiguous facial affect. *Psychiatry Res* 2014;217:100-6.
48. Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 1971;9:97-113.
49. Wittchen HU, Wunderlich U, Gruschwitz S, et al. Strukturiertes Klinisches Interview für DSM-IV (SKID-I), Achse I: Psychische Störungen; Germany, Ed. Hogrefe; 1997.
50. Loranger AW, Sartorius N, Andreoli A, et al. The International Personality Disorder Examination. The World Health Organization/Alcohol, Drug Abuse, and Mental Health Administration international pilot study of personality disorders. *Arch Gen Psychiatry* 1994;51:215-24.
51. Heller KA, Kratzmeier H, Lengfelder A. *Matrizen-Test-Manual. Ein Handbuch zu den Advanced Progressive Matrices von Raven*. Göttingen: Beltz-Testgesellschaft; 1998.
52. Bohus M, Limberger M, Frank U et al. Psychometric properties of the Borderline Symptom List (BSL). *Psychopathology* 2007;40:126-132.
53. Schwenkmezger P, Hodapp V. A Questionnaire for assessing anger and expression of anger. *Z Klin Psychol Psychopathol Psychother* 1991;39:63-8.
54. Gratz KL, Roemer L. Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *J Psychopathol Behav* 2004;26:41-54.
55. Beck AT, Ward CH, Mendelson M, et al. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561-71.
56. Ekman P, Friesen WV. Constants across cultures in the face and emotion. *J Pers Soc Psychol* 1971;17:124-9.
57. Jasper HH. The ten-twenty electrode system of the International Federation. *Electroencephalogr Clin Neurophysiol* 1958;10:370-5.
58. Gratton G, Coles MG, Donchin E. A new method for off-line removal of ocular artifact. *Electroencephalogr Clin Neurophysiol* 1983;55:468-84.
59. Huynh H, Feldt LS. Estimation of the box correction for degrees of freedom from sample data in randomized block and split-plot designs. *J Educ Stat* 1976;1:69-82.
60. Veague HB, Hooley JM. Enhanced sensitivity and response bias for male anger in women with borderline personality disorder. *Psychiatry Res* 2014;215:687-93.
61. Bar-Haim Y, Lamy D, Pergamin L, et al. Threat-related attentional bias in anxious and nonanxious individuals: a meta-analytic study. *Psychol Bull* 2007;133:1-24.
62. Joormann J, Talbot L, Gotlib IH. Biased processing of emotional information in girls at risk for depression. *J Abnorm Psychol* 2007;116:135-43.
63. Minzenberg MJ, Poole JH, Vinogradov S. Social-emotion recognition in borderline personality disorder. *Compr Psychiatry* 2006;47:468-74.
64. Halgren E, Raj T, Marinkovic K, et al. Cognitive response profile of the human fusiform face area as determined by MEG. *Cereb Cortex* 2000;10:69-81.
65. Holmes A, Nielsen MK, Green S. Effects of anxiety on the processing of fearful and happy faces: An event-related potential study. *Biol Psychol* 2008;77:159-73.
66. Rossignol M, Campanella S, Maurage P, et al. Enhanced perceptual responses during visual processing of facial stimuli in young socially anxious individuals. *Neurosci Lett* 2012;526:68-73.
67. Bertsch K, Böhnke R, Kruk MR, et al. Exogenous cortisol facilitates responses to social threat under high provocation. *Horm Behav* 2011;59:428-34.
68. Campanella S, Hanoteau C, Dépy D, et al. Right N170 modulation in a face discrimination task: an account for categorical perception of familiar faces. *Psychophysiology* 2000;37:796-806.
69. Patrick CJ. Psychophysiological correlates of aggression and violence: an integrative review. *Philos Trans R Soc Lond B Biol Sci* 2008;363:2543-55.
70. Cavanagh J, Geisler MW. Mood effects on the ERP processing of emotional intensity in faces: a P3 investigation with depressed students. *Int J Psychophysiol* 2006;60:27-33.
71. Zanarini MC, Frankenburg FR, Hennen J, et al. Axis I comorbidity in patients with borderline personality disorder: 6-year follow-up and prediction of time to remission. *Am J Psychiatry* 2004;161:2108-14.
72. New AS, Aarseth Rot M, Ripoll LH, et al. Empathy and alexithymia in borderline personality disorder: clinical and laboratory measures. *J Pers Disord* 2012;26:660-75.
73. Joassin F, Maurage P, Bruyer R, et al. When audition alters vision: an event-related potential study of the cross-modal interactions between faces and voices. *Neurosci Lett* 2004;369:132-7.