Published in final edited form as:

Curr Opin Struct Biol. 2015 December ; 35: 1–6. doi:10.1016/j.sbi.2015.07.004.

Structural insights into the regulation of aromatic amino acid hydroxylation

Paul F. Fitzpatrick

Department of Biochemistry, University of Texas Health Science Center, San Antonio, TX 78229, fitzpatrickp@uthscsa.edu

Abstract

The aromatic amino acid hydroxylases phenylalanine hydroxylase, tyrosine hydroxylase, and tryptophan hydroxylase are homotetramers, with each subunit containing a homologous catalytic domain and a divergent regulatory domain. The solution structure of the regulatory domain of tyrosine hydroxylase establishes that it contains a core ACT domain similar to that in phenylalanine hydroxylase. The isolated regulatory domain of tyrosine hydroxylase forms a stable dimer, while that of phenylalanine hydroxylases undergoes a monomer-dimer equilibrium, with phenylalanine stabilizing the dimer. These solution properties are consistent with the regulatory mechanisms of the two enzymes, in that phenylalanine hydroxylase is activated by phenylalanine binding to an allosteric site, while tyrosine hydroxylase is regulated by binding of catecholamines in the active site.

Graphical abstract

Introduction

The reactions catalyzed by the pterin-dependent aromatic amino acid hydroxylases are shown in Figure 1 [1]. Phenylalanine hydroxylase (PheH) in the liver catalyzes the hydroxylation of excess phenylalanine to form tyrosine; a deficiency in the enzyme results in the disease phenylketonuria. Tyrosine hydroxylase (TyrH) in the nervous system and the adrenal gland catalyzes the hydroxylation of tyrosine to form 3,4-dihydroxyphenylalanine in the key regulatory step in the biosynthesis of the catecholamine neurotransmitters. Tryptophan hydroxylase (TrpH) in the brain and gastric system catalyzes the hydroxylation

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

of tryptophan to form 5-hydroxytryptophan during serotonin biosynthesis. The physiological pterin substrate is typically tetrahydrobiopterin. The hydroxylation reaction results in hydroxylation of the tetrahydropterin substrate at the 4a position; this 4a-hydroxypterin is formally at the dihydropterin oxidation state and rapidly eliminates water to generate the dihydropterin. While TyrH and TrpH have only been reported for eukaryotic organisms, PheH is also found in many bacteria. The microbial enzymes catalyze the same reaction, although there is one report of a bacterial PheH that catalyzes the formation of 3-hydroxyphenylalanine [2]. This review will focus on recent advances in our understanding of the structural basis for the regulation of PheH and TyrH, since little is known about the regulation of TrpH.

Overall structures

Even before high-resolution structures of the aromatic amino acid hydroxylases became available, three functional domains had been identified in the eukaryotic enzymes based on their sequences: a homologous central catalytic domain of ~290 residues, a divergent regulatory N-terminal domain, and a C-terminal tetramerization domain containing an extended helix that is responsible for the homotetrameric quaternary structures [1]. A sequence alignment of the regulatory domains of the human enzymes is shown in Figure 2A, illustrating their diversity in size and sequence. Figure 2B shows the similar structures of the catalytic domains of the three eukaryotic enzymes. Both PheH and TyrH are single-copy genes, but alternate splicing of the mRNA for human TyrH results in four different isoforms differing in the length of an insert after Met30 in the regulatory domain [3]. The functional differences among the isoforms have not been established. The kinetic properties are very similar, although increasing the length of the regulatory domains somewhat attenuates catecholamine inhibition [4], and the levels of expression vary within different areas of the brain [3]. Ser31 of human isoform 1 is a phosphorylation site; the insertion may change the kinase responsible for phosphorylating this residue [5]. There are two genes for human TrpH coding for proteins that differ in the size of their regulatory domains; the more recently discovered TrpH2 is expressed in neuronal cells, while TrpH1 is expressed in peripheral tissues [6,7]. The former is also 46 residues longer, with the additional sequence at the N-terminus. The bacterial PheHs are monomeric proteins homologous to the catalytic domains of the eukaryotic enzymes and lack the regulatory and tetramerization domains.

The seminal structures of the aromatic amino acid hydroxylases were of tetrameric rat TyrH lacking the regulatory domain and a dimeric form of rat PheH lacking the C-terminal helix (Figure 3A) [8,9]; these allowed identification of residues in the active site and of the relative arrangement of the individual domains. Additional structures of substrate and inhibitor complexes of all three eukaryotic enzymes and of several bacterial PheHs have confirmed the very similar structures of the catalytic domains and provided additional details relevant to catalysis; all published structures are of enzymes lacking either or both the N-terminal regulatory domain and the C-terminal tetramerization domain. It is possible to build a model for the intact PheH tetramer using the structures of the dimeric enzyme lacking the tetramerization domain and of the tetrameric enzyme lacking the regulatory domain (Figure 3B). This shows that the protein is a dimer of dimers in which the individual regulatory domains do not interact with one another. One interface between the dimers must

be weaker in PheH than in the other two enzymes, since PheH is reported to form a mixture of dimers and tetramers in solution [10]. Recently, the solution structure of the isolated regulatory domain of rat TyrH was determined [11], so that it is now possible to build a model of intact TyrH, as is described below. All of the available structures of TrpH lack both the regulatory domain and the tetramerization domain, but the sequences of PheH and TrpH are similar enough to suggest that these two proteins have similar structures.

Regulation of phenylalanine hydroxylase

Liver PheH is an allosteric enzyme [12], exhibiting maximal activity only when activated by phenylalanine. The current model for the regulation of PheH (Figure 4) is based in large part on results with the rat enzyme [13]. This model proposes that the resting form has little or no activity, and that binding of phenylalanine at an activating site converts the enzyme to an active form. In addition, tetrahydrobiopterin potentiates the allosteric affect by binding to the low activity form of the enzyme. Human PheH is reported to have significant activity even in the absence of activation by phenylalanine, but otherwise the regulatory mechanism is assumed to be the same [14]. Mutant forms of rat or human PheH lacking the N-terminal ~117 residue regulatory domain are fully active [12,15], so that the regulatory domain is required to form the low activity resting form of the enzyme. The activated enzyme exhibits positive cooperativity in its kinetics with phenylalanine as a substrate [16], so that the regulatory mechanism may be more complex than indicated by the model in Figure 4.

The structure of rat PheH containing the regulatory and catalytic domain provided a structural explanation for the enzyme's allosteric behavior [9], in that this structure (Figure 3A) shows that the N-terminus of the regulatory domain partially occludes the active site, especially the binding pocket for the tetrahydropterin. Phenylalanine activation would cause a conformational change that moved this inhibitory peptide out of the active site. Consistent with such a model, deletion of the N-terminal 29 residues of rat PheH yields enzyme that is fully active without prior treatment with phenylalanine [17]. A recent analysis using catecholamine binding to the iron as a probe of active site accessibility confirmed that the regulatory domain at least partly blocks the active site and that phenylalanine binding to the allosteric site reverses this [18]. Structures of the catalytic domain of PheH with both tetrahydrobiopterin and an amino acid bound show that several surface loops close down over the active site when both substrates are bound [19]; the positions of several residues in these loops would conflict with residues in the N-terminal regulatory domain.

The location of the site at which phenylalanine binds to activate PheH has been controversial. Both the active site and a separate site in the regulatory domain have been proposed [12,20], but recent results provide support for a site in the regulatory domain. Binding of phenylalanine to the isolated regulatory domain of PheH alters the NMR spectrum of the protein and results in it eluting earlier from a gel filtration column [21], establishing that there is a binding site for phenylalanine in the regulatory domain. Activation of PheH by phenylalanine correlates with a change in the fluorescence of the protein [22], and elimination of phenylalanine binding in the active site does not eliminate this fluorescence change [18], leaving the site in the regulatory domain as the likely allosteric site. The effect of phenylalanine of the oligomerization state of the regulatory

domain of PheH (see below) is consistent with one phenylalanine site per regulatory domain [23]. Still, the location of the phenylalanine site within the regulatory domain has not been established.

PheH is also regulated by phosphorylation of Ser16 in the N-terminal regulatory domain. This is reported to decrease the concentration of phenylalanine needed for activation [24]; such an effect could arise from either a change in the intrinsic binding constant for phenylalanine or a shift in the equilibrium between E_i and E_a in Scheme 4. The structure of phosphorylated rat PheH lacking the C-terminal tetramerization domain has been determined [9]; surprisingly, this structure is identical to that of the unphosphorylated form of the same protein. Unfortunately, the N-terminal 18 residues that contain the phosphorylation site were not seen in either crystal structure; this suggests that the structural change upon phosphorylation is localized. HXMS analyses of rat PheH show that the conformational change upon phosphorylation is more localized than that associated with phenylalanine activation [25,26].

Regulation of tyrosine hydroxylase

TyrH is regulated by a combination of phosphorylation and feed back inhibition (Figure 4) [27]. The active-site iron in the pterin-dependent hydroxylases must be ferrous for catalysis, but it is readily oxidized to Fe(III) by oxygen. Catecholamines bind to the ferric form of TyrH with K_d values ~ 5 nM, effectively inactivating the enzyme. Phosphorylation by protein kinase A of a serine residue in the regulatory domain (Ser40 in rat TyrH and human isoform 1) reverses the inhibition by increasing the rate constants for dissociation of catecholamines from the iron by 2-3 orders of magnitude [28]. In the absence of a structure of TyrH containing both the catalytic and regulatory domains, the structure of PheH has served as a model despite the differences in sequence and size of the two regulatory domains. The N-terminus of TyrH is proposed to act as a flexible peptide that can extend across the active site; a catecholamine bound in the active site would stabilize this inhibitory conformation, while phosphorylation would destabilize it [29,30]. Consistent with such a model, the NMR spectrum of the isolated regulatory domain of TyrH showed that the N-terminal ~ 70 residues are conformationally flexible and that they can be removed without disrupting the core domain structure [11]. The core of the TyrH regulatory domain has the same structure as the core of the regulatory domain of PheH (Figure 2C). This fold, known as an ACT domain, has been seen in a number of bacterial proteins where it acts as an allosteric regulatory domain [31].

Role of regulatory domain dimerization in regulation

While the model for the structure of intact PheH shown in Figure 3B shows no interactions between the regulatory domains in PheH, the isolated regulatory domain of TyrH forms a stable dimer similar to those of other ACT domains (Figure 2D). This dimeric structure can be combined with the known structure of the tetrameric catalytic and tetramerization domains of TyrH to build a model for the intact protein (Figure 3C). Recent studies of the isolated regulatory domain of PheH showed that, rather than being a stable monomer as predicted by the X-ray structure or a dimer as predicted by the solution structure of the

regulatory domain of TyrH, it undergoes a monomer-dimer equilibrium [23]. Moreover, addition of phenylalanine to the monomer converts the protein to a dimer. Jaffe et al. [10] have proposed that the conformational change associated with activation of PheH involves a rotation of the regulatory domain with respect to the catalytic domain, allowing the regulatory domain to form an ACT dimer. The solution behavior of the isolated regulatory domain of PheH supports this model, and the structure of intact TyrH in Figure 4C provides a structural model for activated PheH involving a regulatory domain dimer.

Thus, both TyrH and PheH are proposed to alternate between active and inactive conformations that differ in the accessibility of the active site. The observation that the regulatory domain of TyrH forms a stable dimer is consistent with a model in which the conformational change in that enzyme is limited to relatively unstructured residues at the N-terminus of the regulatory domain, with no significant change in the structure of its ACT core. In contrast, movement of the autoinhibitory N-terminus of the regulatory domain of PheH away from the active site would result from phenylalanine binding in the allosteric site being linked to a large conformational change in which the regulatory domain rotates to form an ACT dimer. This proposal is supported by HXMS studies of the two enzymes. Phosphorylation and dopamine binding to TyrH result in changes in the protein dynamics that are limited to the flexible N-terminus of the regulatory domain outside of the ACT domain and to a peptide in the catalytic domain at the edge of the active site [30]. In contrast, activation of PheH by phenylalanine results in altered dynamics of peptides throughout the interface between the regulatory and catalytic domains and within the ACT domain. [26]. Thus, similar overall structures and similar changes in accessibility of the active site associated with activation result from different molecular mechanisms.

Regulation of TrpH

Problems in obtaining intact recombinant TrpH have limited the understanding of the regulation and structure of that enzyme. The ACT domain of TrpH is much more similar in sequence to that of PheH than is the ACT domain of TyrH (Figure 2A), so that the structures of the cores of the regulatory domains of all three hydroxylases are likely to be similar. The sequence of the regulatory domain of human TrpH1 is significantly shorter than that of TrpH2, suggesting that the latter has an extended unstructured N-terminus that is lacking in the former. However, there are no reports of activation of TrpH2 by tryptophan, so that the regulatory mechanism of PheH, including substrate-induced dimerization of the regulatory domain, may not be applicable. There is also no direct evidence that the activity of TrpH is regulated at the post-translational level. While TrpH2 is phosphorylated on Ser19, this is reported to affect stability rather than activity [32].

Binding of 14-3-3 proteins

14-3-3 proteins, a family of homodimeric proteins that bind phosphorylated proteins [33], will bind to both TyrH and TrpH, suggesting a higher level of regulation. Tight binding of TyrH1 to a 14-3-3 protein requires that Ser19 be phosphorylated, and the binding involves the flexible N-terminus of the regulatory domain [34,35]. The stoichiometry of binding is consistent with one 14-3-3 dimer binding one TyrH regulatory dimer in which both Ser19s

are phosphorylated. 14-3-3 proteins also bind human TrpH2 when Ser19 in that enzyme is phosphorylated [36]. In neither case is the functional role of the interaction understood.

Conclusion

Recent advances in understanding the structures and solution properties of the regulatory domains of PheH and TyrH have begun to give a more complete understanding of the structural changes involved in regulation of the two enzymes. All three of the eukaryotic aromatic amino acid hydroxylases can be described as containing homologous catalytic domains and ACT fold-containing regulatory domains. The regulatory domain of PheH appears to behave more like a classic ACT domain, in that interactions between discrete subunits are altered by binding of phenylalanine to an allosteric site, while the ACT domain of TyrH is unresponsive to the presence of the amino acid substrate.

Acknowledgments

The research described here from the author's laboratory was supported by grants from the National Institutes of Health (GM047291 and GM098140).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

* of special interest

**of outstanding interest

1. Fitzpatrick PF. Tetrahydropterin-dependent amino acid hydroxylases. *Ann. Rev. Biochem.* 1999; 68:355–381. [PubMed: 10872454]
2. Zhang W, Ames BD, Walsh CT. Identification of phenylalanine 3-hydroxylase for meta-tyrosine biosynthesis. *Biochemistry.* 2011; 50:5401–5403. [PubMed: 21615132]
3. Lewis DA, Melchitzky DS, Haycock JW. Four isoforms of tyrosine hydroxylase are expressed in human brain. *Neuroscience.* 1993; 54:477–492. [PubMed: 8101641]
4. Sura G, Daubner SC, Fitzpatrick PF. Effects of phosphorylation by protein kinase A on binding of catecholamines to the human tyrosine hydroxylase isoforms. *J. Neurochem.* 2004; 90:970–978. [PubMed: 15287903]
5. Le Bourdellès B, Horellou P, Le Caer J-P, Denèfle P, Latta M, Haavik J, Guibert B, Mayaux J-F, Mallet J. Phosphorylation of human recombinant tyrosine hydroxylase isoforms 1 and 2: An additional phosphorylated residue in isoform 2, generated through alternative splicing. *J. Biol. Chem.* 1991; 266:17124–17130. [PubMed: 1680128]
6. Amireault P, Sibon D, Côté F. Life without peripheral serotonin: Insights from tryptophan hydroxylase 1 knockout mice reveal the existence of paracrine/autocrine serotonergic networks. *ACS Chem. Neurosci.* 2012; 4:64–71. [PubMed: 23336045]
7. Mosienko V, Beis D, Pasqualetti M, Waider J, Matthes S, Qadri F, Bader M, Alenina N. Life without brain serotonin: Reevaluation of serotonin function with mice deficient in brain serotonin synthesis. *Behav. Brain Res.* 2015; 277:78–88. [PubMed: 24928769]
8. Goodwill KE, Sabatier C, Marks C, Raag R, Fitzpatrick PF, Stevens RC. Crystal structure of tyrosine hydroxylase at 2.3 Å and its implications for inherited neurodegenerative diseases. *Nat. Struct. Biol.* 1997; 4:578–585. [PubMed: 9228951]
9. Kobe B, Jennings IG, House CM, Michell BJ, Goodwill KE, Santarsiero BD, Stevens RC, Cotton RGH, Kemp BE. Structural basis of autoregulation of phenylalanine hydroxylase. *Nat. Struct. Biol.* 1999; 6:442–448. [PubMed: 10331871]

- **10. Jaffe EK, Stith L, Lawrence SH, Andrake M, Dunbrack RL Jr. A new model for allosteric regulation of phenylalanine hydroxylase: Implications for disease and therapeutics. *Arch. Biochem. Biophys.* 2013; 530:73–82. This paper proposes that the structural change accompanying activation of phenylalanine hydroxylase by phenylalanine involves dimerization of the regulatory domains and describes a possible pathway for the conformational change. [PubMed: 23296088]
- **11. Zhang S, Huang T, Ilangovan U, Hinck AP, Fitzpatrick PF. The solution structure of the regulatory domain of tyrosine hydroxylase. *J. Mol. Biol.* 2014; 426:1483–1497. This paper describes the solution structure of the regulatory domain of the enzyme and its formation of a dimer. [PubMed: 24361276]
12. Fitzpatrick PF. Allosteric regulation of phenylalanine hydroxylase. *Arch. Biochem. Biophys.* 2012; 519:194–201. [PubMed: 22005392]
13. Mitnaul LJ, Shiman R. Coordinate regulation of tetrahydrobiopterin turnover and phenylalanine hydroxylase activity in rat liver cells. *Proc. Natl. Acad. Sci. USA.* 1995; 92:885–889. [PubMed: 7846072]
14. Kowlessur D, Citron BA, Kaufman S. Recombinant human phenylalanine hydroxylase: Novel regulatory and structural properties. *Arch. Biochem. Biophys.* 1996; 333:85–95. [PubMed: 8806757]
15. Roberts KM, Pavon JA, Fitzpatrick PF. Kinetic mechanism of phenylalanine hydroxylase: intrinsic binding and rate constants from single-turnover experiments. *Biochemistry.* 2013; 52:1062–1073. [PubMed: 23327364]
16. Flydal MI, Martinez A. Phenylalanine hydroxylase: Function, structure, and regulation. *IUBMB Life.* 2013; 65:341–349. [PubMed: 23457044]
17. Jennings IG, Teh T, Kobe B. Essential role of the N-terminal autoregulatory sequence in the regulation of phenylalanine hydroxylase. *FEBS Lett.* 2001; 488:196–200. [PubMed: 11163771]
- *18. Roberts KM, Khan CA, Hinck CS, Fitzpatrick PF. Activation of phenylalanine hydroxylase by phenylalanine does not require binding in the active site. *Biochemistry.* 2014; 53:7846–7853. The results in this paper establish that elimination of phenylalanine binding in the active site of PheH does not prevent the conformational change that accompanies phenylalanine activation. [PubMed: 25453233]
19. Andersen OA, Stokka AJ, Flatmark T, Hough E. 2.0 Å Resolution crystal structures of the ternary complexes of human phenylalanine hydroxylase catalytic domain with tetrahydrobiopterin and 3-(2-thienyl)-L-alanine or L-norleucine: substrate specificity and molecular motions related to substrate binding. *J. Mol. Biol.* 2003; 333:747–757. [PubMed: 14568534]
20. Thorolfsson M, Ibarra-Molero B, Fojan P, Petersen SB, Sanchez-Ruiz JM, Martinez A. L-Phenylalanine binding and domain organization in human phenylalanine hydroxylase: a differential scanning calorimetry study. *Biochemistry.* 2002; 41:7573–7585. [PubMed: 12056888]
21. Li J, Ilangovan U, Daubner SC, Hinck AP, Fitzpatrick PF. Direct evidence for a phenylalanine site in the regulatory domain of phenylalanine hydroxylase. *Arch. Biochem. Biophys.* 2011; 505:250–255. [PubMed: 20951114]
22. Phillips RS, Parniak MA, Kaufman S. Spectroscopic investigation of ligand interaction with hepatic phenylalanine hydroxylase: Evidence for a conformational change associated with activation. *Biochemistry.* 1984; 23:3836–3842. [PubMed: 6487579]
- **23. Zhang S, Roberts KM, Fitzpatrick PF. Phenylalanine binding is linked to dimerization of the regulatory domain of phenylalanine hydroxylase. *Biochemistry.* 2014; 53:6625–6627. This paper describes the use of analytical ultracentrifugation to demonstrate that phenylalanine binding and dimerization of the regulatory domain of PheH are linked. [PubMed: 25299136]
24. Doskeland AP, Martinez A, Knappskog PM, Flatmark T. Phosphorylation of recombinant human phenylalanine hydroxylase: Effect on catalytic activity, substrate activation and protection against non-specific cleavage of the fusion protein by restriction protease. *Biochem. J.* 1996; 313:409–414. [PubMed: 8573072]
25. Li J, Fitzpatrick PF. Regulation of phenylalanine hydroxylase: Conformational changes upon phosphorylation detected by H/D exchange and mass spectrometry. *Arch. Biochem. Biophys.* 2013; 535:115–119. [PubMed: 23537590]

26. Li J, Dangott LJ, Fitzpatrick PF. Regulation of phenylalanine hydroxylase: Conformational changes upon phenylalanine binding detected by hydrogen/deuterium exchange and mass spectrometry. *Biochemistry*. 2010; 49:3327–3335. [PubMed: 20307070]
27. Daubner SC, Le T, Wang S. Tyrosine hydroxylase and regulation of dopamine synthesis. *Arch. Biochem. Biophys.* 2011; 508:1–12. [PubMed: 21176768]
28. Ramsey AJ, Fitzpatrick PF. Effects of phosphorylation of serine 40 of tyrosine hydroxylase on binding of catecholamines: Evidence for a novel regulatory mechanism. *Biochemistry*. 1998; 37:8980–8986. [PubMed: 9636040]
29. Wang S, Lasagna M, Daubner SC, Reinhart GD, Fitzpatrick PF. Fluorescence spectroscopy as a probe of the effect of phosphorylation at Ser40 of tyrosine hydroxylase on the conformation of its regulatory domain. *Biochemistry*. 2011; 50:2364–2370. [PubMed: 21302933]
30. Wang S, Sura GR, Dangott LJ, Fitzpatrick PF. Identification by hydrogen/deuterium exchange of structural changes in tyrosine hydroxylase associated with regulation. *Biochemistry*. 2009; 48:4972–4979. [PubMed: 19371093]
31. Lang EJM, Cross PJ, Mittelstädt G, Jameson GB, Parker EJ. Allosteric ACTION: the varied ACT domains regulating enzymes of amino-acid metabolism. *Curr. Opin. Struct. Biol.* 2014; 29:102–111. [PubMed: 25543886]
32. Murphy KL, Zhang X, Gainetdinov RR, Beaulieu JM, Caron MG. A regulatory domain in the N terminus of tryptophan hydroxylase 2 controls enzyme expression. *J. Biol. Chem.* 2008; 283:13216–13224. [PubMed: 18339632]
33. van Heusden GPH. 14-3-3 Proteins: Regulators of numerous eukaryotic proteins. *IUBMB Life*. 2005; 57:623–629. [PubMed: 16203681]
- *34. Kleppe R, Rosati S, Jorge-Finnigan A, Alvira S, Ghorbani S, Haavik J, Valpuesta JM, Heck AJ, Martinez A. Phosphorylation dependence and stoichiometry of the complex formed by tyrosine hydroxylase and 14-3-3gamma. *Mol. Cell. Proteomics*. 2014; 13:2017–2030. This paper describes an analysis of the stoichiometry of binding of a 14-3-3 protein to TyrH and the effect of phosphorylation. [PubMed: 24947669]
35. Hritz J, Byeon I-Ja L, Krzysiak T, Martinez A, Sklenar V, Gronenborn Angela M. Dissection of binding between a phosphorylated tyrosine hydroxylase peptide and 14-3-3ζ: A complex story elucidated by NMR. *Biophys. J.* 2014; 107:2185–2194. [PubMed: 25418103]
36. Winge I, McKinney JA, Ying M, D'Santos CS, Kleppe R, Knappskog PM, Haavik J. Activation and stabilization of human tryptophan hydroxylase 2 by phosphorylation and 14-3-3 binding. *Biochem. J.* 2008; 410:195–204. [PubMed: 17973628]

Highlights

The aromatic amino acid hydroxylases contain catalytic and regulatory domains.

The regulatory domains of tyrosine and phenylalanine hydroxylase are ACT domains.

The regulatory domains of tyrosine and phenylalanine hydroxylase form dimers.

Dimerization of the ACT domain of phenylalanine hydroxylase is linked to binding of phenylalanine.

Figure 1.

The reactions catalyzed by the aromatic amino acid hydroxylases phenylalanine hydroxylase (PheH), tyrosine hydroxylase (TyrH), and tryptophan hydroxylase (TrpH).

Figure 2A

TyrH1 1 MPTPDATTPQAKGFRAVSELDKQAEAIM-----
TyrH2 1 MPTPDATTPQAKGFRAVSELDKQAEAIMVRGQ-----
TyrH3 1 MPTPDATTPQAKGFRAVSELDKQAEAIM---GAPGPSLTGSPWPGTAAPAAS
TyrH4 1 MPTPDATTPQAKGFRAVSELDKQAEAIMVRGQGAPGPSLTGSPWPGTAAPAAS

TyrH1 31 -----SPRFIGRRQSLIEDARKEREAAAAAAAAVPSEPGDPLEAVAFEKEGK
TyrH2 31 -----SPRFIGRRQSLIEDARKEREAAAAAAAAVPSEPGDPLEAVAFEKEGK
TyrH3 58 YTPTPRSPRFIGRRQSLIEDARKEREAAAAAAAAVPSEPGDPLEAVAFEKEGK
TyrH4 62 YTPTPRSPRFIGRRQSLIEDARKEREAAAAAAAAVPSEPGDPLEAVAFEKEGK
PheH 1 MSTAVLENPGLGRKLSDFGQETSYIEDNCNQ
TrpH1 1 MIEDNKENKDHSLER
TrpH2 1 MQPAMMMFSSKYWLDSAVPEEHQLGSSSTLNKPNSGKNDDKGNKGSSKREAAATES

TyrH 80 AVLNLLFSPRATKP---SALSRAVKVFETFEAKIHLETRPAQRPRAGGPH---L
PheH 33 GAISLIFSLK---EEVGALAKVLRLEENDVNLTHIESRPSR-----LKKDEY
TrpH1 16 GRASLIFSLK---NEVGGLIKALIKFQEKHVNLHIESRKSRR-----RNSSEF
TrpH2 56 GKTAVVFSLK---NEVGGLVKALRLFQEKRVNMVHIESRKSRR-----RSSEV

TyrH 129 EYFVRL---EVRRGDLAALLSGVR-QVSEDVRSF---AGF---KVP
PheH 78 EFFTLDKRS---LPALTNIILKLRHDIGATVHELSDKKKD---TVP
TrpH1 61 EIFVDCDINREQLNDFHLLKSHTNVLSVNLDPNFTLKEGDMETVP
TrpH2 101 EIFVDCCEGKTEFNELIQLLKFTTIVTLNFPENIWTEEELEDVP

Figure 2B

Figure 2C

Figure 2D

Figure 2.

A, sequence alignments of the regulatory domains of human PheH, TyrH, and TrpH. The alignment of PheH and TyrH is based on the structures while the alignments of TrpH1 and TrpH2 are sequence-based. Residues conserved in all are in blue and residues conserved only in PheH, TrpH1, and TrpH2 are in red; B, overlay of the structures of the catalytic domains of human PheH (PDB file 1pah, red), TyrH (PDB file 2xsn, green) and TrpH1 (PDB file 1mlw, blue); C, overlay of the structures of the regulatory domains of rat PheH (blue, PDB file 2phm, residues 32-117) and rat TyrH (orange, PDB file 2mda, residues 75-159); D, overlay of the structures of the regulatory domain dimer of rat TyrH (blue, PDB file 2mda) and the ACT regulatory domain dimer of *E. coli* glyceraldehyde-3-phosphate dehydrogenase (green, PDB file 1psd).

Figure 3A

Figure 3B

Figure 3C

Figure 3.

A, structure of the combined regulatory (red) and catalytic (cyan) domains of rat PheH (PDB file 2phm); B, model of the intact PheH tetramer, constructed from PDB files 2phm and 2pah; C, model of the structure of intact rat TyrH constructed from the separate structures of the catalytic and regulatory domains (PDB files 1toh and 2mda). In each, the regulatory domains are in red, the catalytic domains in cyan, and the tetramerization domains in blue.

PheH: **TyrH:** **Figure 4.**

The regulatory mechanisms of phenylalanine hydroxylase and tyrosine hydroxylase. For both mechanisms E_a is active enzyme and E_i is inactive enzyme.