Published in final edited form as:

Angew Chem Int Ed Engl. 2015 September 14; 54(38): 11157–11161. doi:10.1002/anie.201505416.

The energetics of a three-state protein folding system probed by high-pressure relaxation dispersion NMR from 1 to 2500 bar

Vitali Tugarinov, David S. Libich, Virginia Meyer, Julien Roche, and G. Marius Clore

Laboratory of Chemical Physics, Building 5, National Institutes of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892-0520, U.S.A.

Vitali Tugarinov: vitali.tugarinov@nih.gov; G. Marius Clore: mariusc@mail.nih.gov

Abstract

The energetic and volumetric properties of a three-state protein folding system, comprising a metastable triple mutant of the Fyn SH3 domain, have been investigated using pressure-dependent ^{15}N -relaxation dispersion NMR from 1 to 2500 bar. Changes in partial molar volumes (ΔV) and isothermal compressibilities ($\Delta \kappa_T$) between all the states along the folding pathway have been determined to reasonable accuracy. The partial volume and isothermal compressibility of the folded state are $\sim 100 \text{ ml}\cdot\text{mol}^{-1}$ and $\sim 40 \text{ }\mu\text{l}\cdot\text{mol}^{-1}\cdot\text{bar}^{-1}$, respectively, higher than those of the unfolded ensemble. Of particular interest are the findings related to the energetic and volumetric properties of the on-pathway folding intermediate. While the latter is energetically close to the unfolded state, its volumetric properties are similar to those of the folded protein. The compressibility of the intermediate is larger than that of the folded state reflecting the less rigid nature of the former relative to the latter.

Keywords

Protein folding kinetics; High pressure NMR; Relaxation dispersion; Invisible states

Pressure, like temperature, constitutes a fundamental thermodynamic variable that can be used to modulate the free energy landscape of proteins, thereby providing a probe to study conformational fluctuations and folding events.^[1] To date, high pressure protein NMR has largely been confined to examining equilibrium properties^[2] and hydrogen/deuterium exchange,^[3] as well as slow conformational transitions (on the order of minutes to hours) subsequent to the application of a pressure jump.^[4] Two relatively recent studies have made use of pressure variation and relaxation dispersion to analyse the kinetics of the folding pathways of several metastable proteins.^[5,6] The latter studies, however, were confined to a narrow range of pressures extending up to only 120 bar in one case^[5] and 250 bar in the other.^[6] While relaxation dispersion experiments are sensitive to small changes in rate constants, the changes observed over this narrow pressure range are too small to fully probe the effects of pressure on kinetic parameters and reliably analyse higher order effects such as compressibility.

To gain detailed insight into the energetic and volumetric properties of a multi-step folding pathway we have carried out ^{15}N Carr-Purcell-Meinboom-Gill (CPMG) relaxation dispersion experiments^[7] on a metastable variant of the Fyn SH3 domain^[8] over a wide range of pressures from 1 to 2500 bar (0.1 to 250 MPa). The Fyn SH3 variant employed contains three destabilizing mutations, A39V, N53P, V55L (referred to hereafter as Fyn SH3-VPL), and has been shown to fold by a three-state pathway involving rapid exchange between the major folded species **F**, a folding intermediate **I** and the unfolded state **U**.^[8a] At 35°C, the temperature used in the current work, the populations of **I** and **U** at atmospheric pressure are ~2 and 0.7%, respectively.^[8a] The structure of the **I** state has been solved based on backbone chemical shifts and residual dipolar couplings derived from the relaxation dispersion data and reveals partial unfolding at the N- and C-termini, thereby exposing the side chains of Phe4 and Ala56 and increasing the size of a surface hydrophobic patch relative to the folded state.^[8b]

Figure 1A shows an overlay of a region of the ^1H - ^{15}N correlation spectrum of U- ^{15}N -labeled Fyn SH3-VPL acquired at six different pressures ranging from 1 to 2500 bar. Significant changes in $^1\text{H}_\text{N}/^{15}\text{N}$ chemical shifts of the folded state **F** are observed as a function of pressure for the majority of residues. Substantial effects of pressure are also evident in the ^{15}N -CPMG relaxation dispersion profiles, as illustrated, for example, by the data for L55 and Y10 shown in Figures 1B and C, respectively, where increasing pressure results in larger dispersion for the former but smaller for the latter. This is a direct consequence of the interplay between the differences in chemical shifts between the various states (ω) and the rates of exchange between them (k). Qualitatively, as all the interconversion rates between **F**, **I** and **U** decrease as the pressure is increased (*vide infra*), residues with larger ω values (e.g. L55) are driven to the slower exchange regime ($k < \omega$) leading to reduced relaxation dispersion, while residues with smaller ω values (e.g. Y10) are driven closer to the intermediate exchange regime ($k \approx \omega$) resulting in larger relaxation dispersion.

The pressure dependence of the interconversion rate constants k_{nm} , pertinent to the three-state folding/unfolding pathway shown in Figure 2A, follows the relationship:^[3,5,9]

$$k_{nm} = (k_B \kappa T / h) \exp(-\Delta G_{\text{TS}_{mn-n}}(P) / RT) \quad [1]$$

where the difference in free energy $\Delta G_{\text{TS}_{mn-n}}(P)$ between the transition states connecting states m and n (TS_{mn}) and state n as a function of the pressure P is given by:

$$\Delta G_{\text{TS}_{mn-n}}(P) = \Delta G_{\text{TS}_{mn-n}}^0 + \Delta V_{\text{TS}_{mn-n}}(P - P_0) - \frac{1}{2} \Delta \kappa_{\text{T,TS}_{mn-n}}(P - P_0)^2 \quad [2]$$

The subscripts $n, m \in \{\text{F}, \text{I}, \text{U}\}$; $\Delta G_{\text{TS}_{mn-n}}^0(P)$ is the free energy difference at $P_0 = 1$ bar (0.1 MPa); $\Delta V_{\text{TS}_{mn-n}}$ and $\Delta \kappa_{\text{T,TS}_{mn-n}}$ are the corresponding changes in partial volume V and isothermal compressibility κ_{T} (defined as dV/dP); k_B is the Boltzmann constant, h Planck's constant, T the temperature (in Kelvin), R the universal gas constant, and κ the transmission coefficient. A value of $\kappa = 1.6 \times 10^{-7}$ is commonly used for protein folding,^[10] corresponding to $(k_B \kappa / h) = 3000 \text{ s}^{-1} \text{K}^{-1}$.

Global fitting of all relaxation dispersion profiles at all pressures simultaneously to the three-state model shown in Figure 2A with the pressure dependence of the rate constants governed by Eqs. [1–2] was carried out as described in detail in the Supporting Information. The values of $V_{TS_{mn}-n}$, $\kappa_{T,TS_{mn}-n}$ and k_{nm}^0 (the rate constants connecting states n and m at 1 bar) obtained from the global best-fits are provided in Table 1. To ensure smooth changes in chemical shift differences as a function of pressure, the values of ϖ between states **F** and **I** (ϖ_{FI}) and **F** and **U** (ϖ_{FU}) were constrained to a quadratic polynomial function of pressure (see Supporting Information Eq. [S7]). The pressure dependence of the four rate constants in Figure 2A was modeled using either a simple linear relationship assuming $\kappa_{T,TS_{mn}-n} = 0$ (referred to hereafter as ‘model 1’),

$$k_{nm}(P) = k_{nm}^0 \exp\{-\Delta V_{TS_{mn}-n}(P - P_0)/RT\} \quad [3]$$

or the full quadratic relationship (referred to as ‘model 2’),

$$k_{nm}(P) = k_{nm}^0 \exp\{[-\Delta V_{TS_{mn}-n}(P - P_0) + \frac{1}{2}\Delta\kappa_{T,TS_{mn}-n}(P - P_0)^2]/RT\} \quad [4]$$

Excellent fits were obtained with $\chi^2 = 1.13$ and 1.09 for models 1 and 2, respectively. Although the small decrease in χ^2 for model 2 relative to model 1 is statistically significant (partial F -test $p < 0.0001$), and the optimized values of κ_T are both well defined by the experimental data and lie within the expected range,^[11] we report and compare the results from both fits as there is a significant “cross-talk” in the fits between V and κ_T , thereby providing a better appreciation of the range of variability of the optimized parameters depending on the model employed. We also note that very similar results are obtained when the relaxation dispersion data are fit separately at each pressure (*i.e.* without constraining the rate constants to a particular pressure dependence model). Simultaneous global fitting of the data at all pressures, however, is more robust and ensures smooth continuity of changes in rate constants and ϖ values as a function of pressure (see Supporting Information for discussion of the changes in ϖ_{FI} and ϖ_{FU} as a function of pressure).

The pressure dependence of the species populations and rate constants are shown in Figures 2B and 2C–F, respectively. The primary effect of pressure is to increase the population (p_U) of the unfolded state **U** from ~0.7 % at 1 bar to ~5 % at 2500 bar; the population (p_I) of the folding intermediate **I** is virtually pressure-independent (Figure 2B). This is a direct consequence of the differential effects of pressure on the interconversion rate constants between the various states. While k_{FI} and k_{IF} decrease in concert by less than two-fold from 1 to 2500 bar (Figures 2C and D, respectively), k_{IU} (Figure 2E) and k_{UI} (Figure 2F) are affected very differently by pressure, with by far the largest effect exerted on k_{UI} which decreases by almost an order of magnitude from 1 to 2500 bar. These observations are valid irrespective of whether model 1 or 2 is used in the fit, even though significant deviations from exponentiality are seen for the pressure dependence of k_{FI} and k_{IU} using model 2 (red symbols in Figures 2C and E).

The changes in partial molar volumes and isothermal compressibilities obtained from the global fits are referenced to the unfolded state **U** in Figures 3A and 3B, respectively. Not

surprisingly, the volume and compressibility of the folded state **F** are higher than those of the unfolded state **U** by $\sim 100 \text{ ml}\cdot\text{mol}^{-1}$ and $\sim 40 \text{ }\mu\text{l}\cdot\text{mol}^{-1}\cdot\text{bar}^{-1}$, respectively. The largest volume change occurs between state **U** and the transition state TS_{UI} between states **I** and **U**. As expected from the close similarity in the structures of **F** and **I**,^[8b] the volume of **I** is close to that of **F** irrespective of the model employed, with V_{I} larger than V_{F} by only 8 and 14 $\text{ml}\cdot\text{mol}^{-1}$ for models 1 and 2, respectively (Figure 3A; Table 1). As the differences between the structures of **F** and **I** are minor,^[8b] it is conceivable that the somewhat higher partial volumes of **I** result primarily from changes in hydration occurring upon partial unfolding of the N- and C-termini. Of note, the compressibility of **I** is also higher than that of **F** (Figure 3B; Table 1) - likely a consequence of the presence of some non-native interactions and solvent exposure of hydrophobic side-chains in **I**.

The pressure dependence of the activation free energies $G_{\text{TS}_{mn-n}}$ is shown in Figure 4A. All the values of $G_{\text{TS}_{mn-n}}$ increase with pressure irrespective of the fitting model, with the largest free energy change occurring between **U** and the corresponding transition state TS_{UI} (cf. the pressure dependence of k_{UI} shown in Figure 2F).

The differences in free energies between states m and n , G_{m-n} ($n, m \in \{\text{F}, \text{I}, \text{U}\}$), at each pressure can be calculated from the corresponding equilibrium constants K_{nm} ($= k_{nm}/k_{mn}$),

$$\Delta G_{m-n} = -RT \ln K_{nm} \quad [5]$$

The pressure dependence of the G values between the **F**, **I** and **U** states is shown in Figure 4B. The free energy of **I** is close to that of **U** and becomes equi-energetic with the latter at a pressure of $\sim 1000 \text{ bar}$ (the intersection of $G_{\text{I-F}}$ and $G_{\text{U-F}}$ in Figure 4B) when states **U** and **I** are equally populated (cf. Figure 2B). This implies an approximately equal destabilization of states **I** and **F** by pressure, while the unfolded state **U** remains largely unaffected (see inset in Figure 4B).

To the best of our knowledge, the kinetics of only one other three-state protein folding system as a function of pressure has been investigated to date. In the pressure-dependent relaxation dispersion study of the G48M mutant of the Fyn SH3 domain, Kay and co-workers used a narrow range of pressures from 1 to 120 bar to derive the volumetric properties of the system using the linear model of pressure dependence (i.e. $\kappa_{\text{T}} = 0$).^[5] Although the reported volume change, $V_{\text{F-U}}$, between the states **F** and **U**, $V_{\text{F-U}}$, of $70 \pm 3 \text{ ml}\cdot\text{mol}^{-1}$ is comparable to that found here for Fyn SH3-VPL, some important differences emerge in relation to the volume of the intermediate state **I**. In particular, in the G48M mutant, the volume of the **I** state lies *in between* those of **U** and **F**,^[5] while in Fyn SH3-VPL (at 35 °C) the volume of **I** is slightly higher than that of **F** irrespective of the fitting model employed (Figure 3A), indicating differences in structure and/or hydration between the folding intermediates of the Fyn SH3-G48M and VPL mutants. In addition, the much wider range of pressures (1 to 2500 bar) used in the present work permits reliable estimates of κ_{T} for all the states on the Fyn SH3-VPL folding pathway (Figure 3B). Notably, the *positive* values of $\kappa_{\text{T,F-U}}$ obtained for Fyn SH3-VPL (Table 1) are consistent with expectations based on prior relaxation dispersion and volumetric measurements,^[6] as well as theoretical predictions.^[11a]

In summary, we have used ^{15}N CPMG relaxation dispersion NMR to measure the pressure-dependent folding/unfolding kinetics of a triple metastable mutant of the Fyn SH3 domain, which is known to fold through a sparsely populated on-pathway intermediate.^[8] The exquisite sensitivity of NMR relaxation dispersion to even small changes in rate constants and the wide range of pressures used (up to 2500 bar) enabled us to reliably quantify the changes in partial molar volumes and isothermal compressibilities along the folding trajectory. The insights gained into the energetic and volumetric properties of the on-pathway intermediate are of particular interest: while the intermediate state is energetically close to the unfolded state, its volumetric properties are similar to those of the native folded protein. Further, the compressibility of the intermediate state is larger than that of the folded state, indicating the structure of the intermediate is less rigid than that of the folded state. Pressure-dependent relaxation-based NMR methods represent a powerful complement to conventional approaches for measuring volumetric and energetic properties associated with protein folding where pH, denaturants or temperature are used to perturb the folding equilibria – all the more so in cases of complex, multi-step folding pathways.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the Intramural Program of NIDDK, NIH and by the AIDS Targeted Antiviral Program of the Office of the Director of the NIH (to G.M.C.)

References

1. a) Jonas J. *Science*. 1982; 216:1179–1184. [PubMed: 17830561] b) Akasaka K. *Chem. Rev.* 2006; 106:1814–1835. [PubMed: 16683756] c) Akasaka K, Kitahara R, Kamatari YO. *Arch. Biochem. Biophys.* 2013; 531:110–115. [PubMed: 23246376] d) Hirata F, Akasaka K. *J. Chem. Phys.* 2015; 142:044110. [PubMed: 25637972] e) de Oliveira GAP, Silva JL. *Proc. Natl. Acad. Sci. U. S. A.* 2015; 112:E2775–E2784. [PubMed: 25964355]
2. a) Rouget J-B, Aksel T, Roche J, Saldana JL, Garcia AE, Barrick D, Royer CA. *J. Am. Chem. Soc.* 2011; 133:6020–6027. [PubMed: 21446709] b) Fu Y, Kasinath V, Moorman VR, Nucci NV, Hilser VJ, Wand AJ. *J. Am. Chem. Soc.* 2012; 134:8543–8550. [PubMed: 22452540] c) Roche J, Caro JA, Norberto DR, Barthe P, Roumestand C, Schlessman JL, Garcia AE, Garcia-Moreno BE, Royer CA. *Proc. Natl. Acad. Sci. U. S. A.* 2012; 109:6945–6950. [PubMed: 22496593] d) Vakpai N, Nisius L, Wiktor M, Grzesiek S. *Proc. Natl. Acad. Sci. U. S. A.* 2013; 110:E368–E376. [PubMed: 23284170] e) Kalbitzer HR, Rosnizeck IC, Muent CE, Narayanan SP, Kropf V, Spoener M. *Angew. Chem. Int. Ed.* 2013; 52:14242–14246. f) Erlach MB, Koehler J, Moeser B, Horinek D, Kremer W, Kalbitzer HR. *J. Phys. Chem. B.* 2014; 118:5681–5690. [PubMed: 24798035]
3. Fuentes EJ, Wand AJ. *Biochemistry*. 1998; 37:9877–9883. [PubMed: 9665691]
4. a) Kremer W, Arnold M, Munte CE, Hartl R, Erlach MB, Koehler J, Meier A, Kalbitzer HR. *J. Am. Chem. Soc.* 2011; 133:13646–13651. [PubMed: 21774550] b) Roche J, Dellarole M, Caro JA, Norberto DR, Garcia AE, Garcia-Moreno B, Roumestand C, Royer CA. *J. Am. Chem. Soc.* 2013; 135:14610–14618. [PubMed: 23987660]
5. Bezsonova I, Korzhnev DM, Prosser RS, Forman-Kay JD, Kay LE. *Biochemistry*. 2006; 45:4711–4719. [PubMed: 16605239]
6. Korzhnev DM, Bezsonova I, Evanics F, Taulier N, Zhou Z, Bai Y, Chalikian TV, Prosser RS, Kay LE. *J. Am. Chem. Soc.* 2006; 128:5262–5269. [PubMed: 16608362]

7. a) Hansen DF, Vallurupalli P, Kay LE. *J. Phys. Chem. B.* 2008; 112:5898–5904. [PubMed: 18001083] b) Palmer AG 3rd. *J. Magn. Reson.* 2014; 241:3–17. [PubMed: 24656076]
8. a) Neudecker P, Zarrine-Afsar A, Choy WY, Muhandiram DR, Davidson AR, Kay LE. *J. Mol. Biol.* 2006; 363:958–976. [PubMed: 16989862] b) Neudecker P, Robustelli P, Cavalli A, Walsh P, Lundstrom P, Zarrine-Afsar A, Sharpe S, Vendruscolo M, Kay LE. *Science.* 2012; 336:362–366. [PubMed: 22517863]
9. Prehoda KE, Mooberry ES, Markley JL. *Biochemistry.* 1998; 37:5785–5790. [PubMed: 9558311]
10. Hagen SJ, Hofrichter J, Szabo A, Eaton WA. *Proc. Natl. Acad. Sci. U. S. A.* 1996; 93:11615–11620. [PubMed: 8876184]
11. a) Taulier N, Chalikian TV. *Biochim. Biophys. Acta.* 2002; 1595:48–70. [PubMed: 11983386] b) Chalikian TV. *Annu. Rev. Biophys. Biomol. Struct.* 2003; 32:207–235. [PubMed: 12543706] c) Silva JL, Weber G. *Annu. Rev. Phys. Chem.* 1993; 44:89–113. [PubMed: 8257561]

Figure 1.

Pressure-dependent NMR of Fyn SH3-VPL. (A) ^1H - ^{15}N correlation spectra acquired at six different pressures ranging from 1 to 2500 bar (800 MHz, 35 °C). The cross-peaks corresponding to the different pressure values are color-coded as indicated on the plot. Selected cross-peaks are labeled. The direction of chemical shift changes with pressure is indicated with an arrow for the peak of T47. (B) ^{15}N CPMG relaxation dispersion^[7a] profiles for L55 (left) and Y10 (right) acquired at six pressures and two spectrometer fields (500 and 800 MHz) plotted using the same color code as in panel (A). The continuous lines

connecting the experimental points represent the results of the global best-fit at all pressures simultaneously to model 2 (see text). Chemical shift differences between states **F** and **I** (ω_{FI}) and **F** and **U** (ω_{FU}) for L55 and Y10 are: $\omega_{FI} = 3.1$ and 2.8 ppm, respectively, and $\omega_{FU} = 5.4$ and -1.2 ppm, respectively.^[8a] Additional examples of ^{15}N CPMG relaxation dispersion fits, and experimental details related to sample preparation and acquisition of the ^{15}N CPMG relaxation dispersion data (that measure the effective relaxation rates of in-phase ^{15}N coherences^[7a]) are provided in the Supporting Information.

Figure 2.

Pressure dependence of folding/unfolding kinetics of Fyn SH3-VPL. **(A)** Three-state model of protein (un)folding used for all the fits to the relaxation dispersion data in this work. k_{nm} are the interconversion rate constants and p_n the population of each state. **(B)** Change of p_I and p_U (%) with pressure. Pressure dependencies of **(C)** k_{FI} , **(D)** k_{IF} , **(E)** k_{IU} and **(F)** k_{UI} rate constants. In each of the plots (B–F), the blue and red filled-in circles represent the results of the global fit to models 1 ($\kappa_T = 0$; cf. Eq. [3]) and 2 (cf. Eq. [4]), respectively. The lines

connecting the values fitted for each pressure are drawn to guide the eye and do not represent the results of a fit.

Figure 3.

Partial volumes and isothermal compressibilities for the Fyn SH3-VPL three-state protein folding system. **(A)** Partial molar volumes of the Fyn SH3-VPL system referenced to the volume of the unfolded state ($V_U = 0$) obtained from global fitting of the relaxation dispersion data at all pressures to either model 1 (blue; $\kappa_T = 0$; cf. Eq. [3]) or model 2 (red; cf. Eq. [4]). **(B)** Isothermal compressibilities referenced to the compressibility of the unfolded state ($\kappa_{T,U} = 0$). The values of the partial volumes and compressibilities of each equilibrium (F, I, U) and transition (TS) state are indicated below the solid horizontal bars.

Figure 4.

Pressure dependence of activation free energies and free energies for the Fyn SH3-VPL three-state protein folding system. **(A)** Activation free energies (Eq. [2]) and **(B)** free energy differences (Eq. [5]) between the states derived from the results of the global fit of the relaxation dispersion data at all pressures to either model 1 ($\kappa_T = 0$; blue circles; cf. Eq. [3]) or model 2 (red circles; cf. [Eq. [4]). Note that $G_{U-F} = G_{I-F} + G_{U-I}$. Free energy level diagrams of the three-state folding equilibrium are shown in the insets of both panels.

The arrows in the inset of panel (B) indicate the direction of the free energy changes of states **F** and **I** with pressure.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Values of rate constants (k), changes in partial volumes (ΔV), and changes in isothermal compressibility ($\Delta \kappa_T$) obtained from the global fit of the Fyn SH3-VPL relaxation dispersion data at six pressures ranging from 1 to 2500 bars (35 °C).^[a]

	k_0 (s ⁻¹) ^[b] (at 1 bar)	ΔV (ml/mol)	$\Delta \kappa_T$ (μl/mol/bar)
TS _{IF} -F (k_{FI})	66±4 (80±4)	43±10 (33±3)	22±7
TS _{IF} -I (k_{IF})	3581±100 (3723±21)	30±4 (25±1)	7±3
TS _{UI} -I (k_{IU})	382±23 (361±16)	-4±9 (13±3)	-20±7
TS _{UI} -U (k_{UI})	904±17 (867±12)	132±3 (97±1)	33±3
F-U	-	122±15 (76±5) ^[c]	38±11 ^[c]
F-I	-	-14±18 (-8±6) ^[c]	-15±14 ^[c]

^[a] All data are reported with ± 1 standard deviation as obtained from the variance-covariance matrix of the non-linear fit. The values fitted using the linear model of pressure dependence (i.e., $\Delta \kappa_T = 0$, model 1; cf. Eq. [3]) are listed in parentheses.

^[b] Note that values of k_{FI} , k_{IF} , k_{IU} and k_{UI} , reported previously by Neudecker et al.^[8a] at 35°C and atmospheric pressure (1 bar) are 51±1.6, 2951±55, 389±9, and 924±15 s⁻¹, respectively, corresponding to p_I and p_U values of 1.69 and 0.71 %, respectively.

^[c] The values of ΔV and $\Delta \kappa_T$ between **F** and **U**, and **F** and **I**, are calculated from the corresponding fitted values for the transition states (first four rows).