Spitz Nevus on the Perianal Area of a Child: An Unusual Location

Dong Yeup Lee, Jong Heon Baek, Dong Joo Kim, Soo Kyung Lee, Myoung Shin Kim, Un Ha Lee, Jeong Hee Hahm

Department of Dermatology, Sanggye Paik Hospital, Inje University College of Medicine, Seoul, Korea

Dear Editor:

Spitz nevus is uncommon melanocytic nevus, and the head and neck are known to be the most common involved sites. To our knowledge, this is the first report of Spitz nevus arising on the perianal area.

A 3-year-old boy presented with a pedunculated nodule on his perianal area (Fig. 1). The nodule was about 1 cm in diameter, with erythema and tiny black-pigmented spots anterior to the anal orifice. It had developed 1 year previously and had been gradually increasing in size. Shave excision was done for the diagnosis and treatment. Histopathological examination revealed numerous nests of cells in the entire dermis (Fig. 2A). In the epidermis, there were sparse basal pigmentations along the dermoepidermal junction and some nevus cells with large nuclei on the tip of the rete ridges (Fig. 2B). In the dermis, tumor cells were composed of large epithelioid nevocytes with abundant eosinophilic cytoplasm (Fig. 2C, D). Many tumor cells were uniformly atypical large epithelioid cells especially in the superficial dermis; however, there was no mitosis or Kamino body. Immunohistochemical staining revealed that the nest cells were superficially positive for HMB-45 and Ki-67. Because the lower margin contained tumor cells, we could not identify whether there was progressive dispersion of melanocytes to smaller aggregates or to single melanocytes in the deepest part of the

lesion; therefore, we concluded that further excision with a clear margin or close follow-up for recurrence is needed. Since shave excision was accomplished, there had been no new lesions for about 1 year; however, he has not made a follow-up visit thereafter.

Traditionally, the sites of predilection of Spitz nevus are known to be the face, head, and neck in children, whereas in adults, the lower limbs have been recognized as the most common involved sites. However, in a study of 349 patients with Spitz nevus¹, the most common location was the lower extremities in all age groups, except in the group older than 45 years, in whom the trunk was the most common site. There have been only three reports of penile solitary Spitz nevi in young adults²⁻⁴ and one report of Spitz nevus on the inner surface of the labium major⁵. There was no characteristic and common feature between all of these genital Spitz nevi. Although there are few re-

Fig. 1. Spitz nevus on the perianal area: an erythematous pedunculated nodule with tiny black pigmentation anterior to the anal orifice.

Received August 11, 2014, Revised January 10, 2015, Accepted for publication February 26, 2015

Corresponding author: Myoung Shin Kim, Department of Dermatology, Sanggye Paik Hospital, 1342 Dongil-ro, Nowon-gu, Seoul 01757, Korea. Tel: 82-2-950-1131, Fax: 82-2-931-8720, E-mail: etihwevol@naver.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fig. 2. Histological examination showed (A) numerous nests of cells in the entire dermis (H&E, ×40). On the tip of the rete ridges (B), some proliferation of nevus cells and melanocytes are visible (H&E, ×40). In the superficial dermis (C) and deep dermis (D), the cell nests are composed of epithelioid nevocytes that have a large cell size, abundant cytoplasm, cleft around cells, and prominent nucleoli (H&E, ×200). There was no atypical mitosis or pagetoid spreading.

ports of Spitz nevi of the genital area, to the best of our knowledge, this is the first reported case of Spitz nevus on the perianal area.

Unlike the typical Spitz nevus, the histopathologic findings of this case included epithelioid cells of large and pleomorphic morphology without a Kamino body. However, histologic findings that suggest the atypical Spitz tumor, such as higher cellular density and considerable mitotic rate, were not observed in this case.

In conclusion, this is the first case of Spitz nevus arising from the perianal area, and we suggest that Spitz nevus should be considered as a differential diagnosis in an unusual papular/nodular lesion on the perianal area in children.

REFERENCES

1. Requena C, Requena L, Kutzner H, Sánchez Yus E. Spitz nevus: a clinicopathological study of 349 cases. *Am J Dermatopathol* 2009;31:107-116.
2. Aoyagi S, Sato-Matsumura KC, Akiyama M, Tanimura S, Shibaki H, Shimizu H. Spitz naevus of the glans penis: an unusual location. *Acta Derm Venereol* 2004;84:324-325.
3. Filippov SV, Kniaz'kin IV, Anichkov NM, Zeziulin PN, Shinkarenko AV, Bykov NM. Spitz nevus (juvenile nevus) of the penile skin. *Arkh Patol* 2002;64:46-48.
4. Cho SB, Kim HS, Jang H. A pedunculated hyalinizing Spitz nevus on the penile shaft. *Int J Dermatol* 2009;48:1134-1136.
5. Polat M, Topcuoglu MA, Tahtaci Y, Hapa A, Yilmaz F. Spitz nevus of the genital mucosa. *Indian J Dermatol Venereol Leprol* 2009;75:167-169.