Published in final edited form as:

Org Lett. 2016 January 4; 18(1): 36–39. doi:10.1021/acs.orglett.5b03127.

Asymmetric Organocatalytic Reductive Coupling Reactions between Benzyldiene Pyruvates and Aldehydes

Matthew A. Horwitz, Blane P. Zavesky, Jesus I. Martinez-Alvarado, and Jeffrey S. Johnson*

Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, United States

Abstract

An organocatalytic three-component reductive coupling reaction between dimethyl phosphite, benzyldiene pyruvates, and aldehydes is reported. A chiral triaryliminophosphorane catalyst promotes Pudovik addition, which is followed by phospho-Brook rearrangement to transiently generate enolates that are trapped stereoselectively by aldehydes. This reductive coupling provides vicinal polyfunctionalized stereocenters from readily available prochiral starting materials with excellent diastereoselectivity, enantioselectivity, and yield.

The reductive union of two prochiral starting materials into products bearing vicinal stereogenic centers builds molecular complexity and as such is an actively sought transformation in chemical synthesis. Pinacol-type reductive coupling reactions deliver

*Corresponding Author. jsj@unc.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b03127.

Crystallographic data for **2a** (CIF)

Experimental procedures, characterization, spectral data for all compounds (PDF)

The authors declare no competing financial interest.

vicinal diols,^{1–5} but drawbacks remain. Commonly used single-electron transfer methods rely on stoichiometric amounts of low-valent metal,^{3n–r} and stereocontrol can be challenging.^{5b,c} In the methodologies that have successfully achieved selectivity in the reductive coupling reaction, there is still the issue of making the vicinal alcohols react orthogonally in downstream transformations.^{3g} Some of these issues were recently addressed through the use of a base-catalyzed, phosphite-mediated asymmetric reductive coupling of two different carbonyls.^{6,7} In this mechanistic manifold, a Pudovik addition of a dialkylphosphite to an isatin triggers phospho-Brook rearrangement and subsequent catalyst controlled trapping of the resultant enolate with an aldehyde.^{6–9} Herein, we extend this reaction framework and report a highly stereoselective phosphite-mediated reductive coupling reaction between benzylidene pyruvates and aryl aldehydes (Scheme 1).

Our goal of introducing a higher level of functionality into the product carries with it challenges not faced in our prior work (Scheme 2). In order to achieve a stereoselective cross-coupled product from ambident benzylidene pyruvates and aryl aldehydes, it is necessary to be able to control (a) the chemoselectivity of the phosphite addition (pyruvate vs aldehyde),¹⁰ (b) the regioselectivity of the phosphite addition (1,2- vs 1,4-addition),¹¹ (c) the nucleophilicity of the nascent enolate (α - vs γ -trapping),¹² (d) the chemoselectivity of the enolate trap (pyruvate vs aldehyde),^{12,13} and (e) the stereoselectivity of the enolate addition into the aryl aldehyde. Fortunately, the relative electron deficiency of the benzylidene pyruvates made the chemo- and regioselectivity issues manageable. Furthermore, the chiral triaryliminophosphoranes developed by Dixon and co-workers¹⁴ guided the stereodefining C–C bond construction with excellent levels of diastereo- and enantiocontrol.

Initially, we studied the dimethyl phosphite mediated reductive coupling of benzylidene pyruvate **1a** with *para*-bromobenzaldehyde (Table 1). Using 10 mol % KO^tBu at 0 °C, the reaction was complete in minutes and hydroxy phosphate **2a** was formed exclusively (dr 1.2:1). Having found that the enolate formed by the Pudovik–phospho–Brook sequence was both nucleophilic at the correct position and capable of being trapped by aryl aldehydes, we turned our attention to the development of the asymmetric variant. In our previous experience with this type of reductive coupling reaction, we demonstrated through crossover experiments that a stereoablative retro aldol process becomes possible somewhere in the cryogenic range;⁷ therefore, we sought to carry out the reactions at as low a temperature as possible. We observed that cinchona alkaloid-derived thiourea catalysts were not basic enough to permit the reaction to proceed at cryogenic temperatures, which caused us to move toward other catalyst families. The evaluation of chiral triaryliminophosphorane **C1** revealed that, after 48 h at –60 °C, the starting material was completely consumed and a 6:1 ratio of products was obtained arising from aldehyde trapping (**2a**) relative to proton trapping (**3a**), the former with a diastereomer ratio of 13:1. This encouraging result led us to synthesize and evaluate catalyst **C2**, which gave a >20:1 ratio of **2a:3a**, with >20:1 dr and 97:3 er.

The application of catalyst **C2** to a broader range of reaction partners was then undertaken (Scheme 3). The reaction proceeds with electron-withdrawing groups on the benzylidene pyruvate; placing the electron-withdrawing group on the *ortho* (**2f**, **2g**) or *para* (**2b–2d**)

positions on the benzylidene pyruvate led to comparable yields and stereoselectivities to the unsubstituted case, but we found that using a *meta*-bromo benzylidene pyruvate gave only 5.4:1 dr. Additionally, while substrates with *meta* and *para* electron-withdrawing groups gave upward of 96:4 er, we observed enantioselectivities of 90.5:9.5 for **2f** and 93:7 for **2g** (*o*-bromo and *o*-fluoro, respectively). Using a 4-methyl substituted benzylidene pyruvate we observed that the reaction was complete in 24 h (**2h**), though, with stronger electron-donating groups on the ring, the reaction is slower likely due to depressed rate of Pudovik addition (**2i–2j**). Using the 2-thienylidene pyruvate gave **2k** in >20:1 dr, with 87% yield and 92:8 er, but extending the conjugation of the starting material as in **2l** gave 14:1 dr and 92:8 er, with a 74% yield. The reaction was found to proceed with other electron-deficient aryl aldehydes as well (**2m–2r**), either in the *para* or *meta* position, although there was a noticeable drop in stereoselectivity with *para*-nitrobenzaldehyde. We attempted to use benzaldehyde as a coupling partner, but observed that the major product formed in that reaction was **3**.¹⁵

The asymmetric reductive coupling reaction on gram scale works comparably to those reactions conducted on smaller scale. Figure 2 illustrates the conversion of 1 g of **1a** to 1.88 g of the derived coupled product **2a** with >20:1 dr and 97.5:2.5 er after a single recrystallization. An X-ray diffraction study of this material revealed the absolute configuration of the coupled product to be (*1R,2R*) (Scheme 4).¹⁶

The work described here expands on organic reductant-based organocatalytic reductive coupling. The title process exhibits high levels of chemo- and stereoselectivity in the face of multiple potential reaction pathways. Specifically, this work presents new possibilities for the coupling partners that can participate in this reaction. Research into potential applications of these new motifs is currently underway in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

Financial support was provided by Award R01 GM103855 from the National Institute of General Medical Sciences.

REFERENCES

1. Reviews: Terra BS, Macedo F Jr. ARKIVOC. 2012:134–151. Chatterjee A, Joshi NN. Tetrahedron. 2006; 62:12137–12158.
2. Seminal papers on cross pinacol reactions with low-valent metal reagents: McMurry JE, Krepski LR. J. Org. Chem. 1976; 41:3929–3930. McMurry JE. Chem. Rev. 1989; 89:1513–1524.
3. Developments toward using aldehydes as coupling partners in stereoselective reductive coupling reactions: Takahara PM, Freudenberger JH, Konradi AW, Pedersen SF. Tetrahedron Lett. 1989; 30:7177–7180. Freudenberger JH, Konradi AW, Pedersen SF. J. Am. Chem. Soc. 1989; 111:8014–8016. Annunziata R, Cinquini M, Cozzi F, Giaroni PF. Tetrahedron: Asymmetry. 1990; 1:355–358. Annunziata R, Cinquini M, Cozzi F, Glaroni P, Benaglia MF. Tetrahedron. 1991; 47:5737–5758. Annunziata R, Benaglia M, Cinquini M, Cozzi F, Giaroni PF. J. Org. Chem. 1992; 57:782–784. Konradi AW, Pedersen SF. J. Org. Chem. 1990; 55:4506–4508. Konradi AW, Kemp SJ, Pedersen SF. J. Am. Chem. Soc. 1994; 116:1316–1323. Park J, Pedersen SF. J. Org. Chem. 1990; 55:5924–

5926. Askham FR, Carroll KM. *J. Org. Chem.* 1993; 58:7328–7329. Yoda H, Matsuda K, Nomura H, Takabe KF. *Tetrahedron Lett.* 2000; 41:1775–1779. Takai K, Nitta K, Utimoto KF. *Tetrahedron Lett.* 1988; 29:5263–5266. Takai K, Morita R, Toratsu C. *Angew. Chem. Int. Ed.* 2001; 40:1116–1119. Takai K, Morita R, Matsushita H, Toratsu CF. *Chirality.* 2003; 15:17–23. [PubMed: 12467037] Boeckman RK Jr, Hudack RA Jr. *J. Org. Chem.* 1998; 63:3524–3525. Jung M, Groth UF. *Synlett.* 2002:2015–2018. Groth U, Jung M, Vogel TF. *Chem. - Eur. J.* 2005; 11:3127–3135. [PubMed: 15776493] Fischer S, Groth U, Jung M, Lindenmaier M, Vogel TF. *Tetrahedron Lett.* 2005; 46:6679–6682. Miyoshi N, Fukuma T, Wada MF. *Chem. Lett.* 1995; 24:999–1000. Maekawa H, Yamamoto Y, Shimada H, Yonemura K, Nishiguchi IF. *Tetrahedron Lett.* 2004; 45:3869–3872. Nair V, Ros S, Jayan CN, Rath NP. *Tetrahedron Lett.* 2002; 43:8967–8969. Yang Y-S, Shen Z-L, Loh T-P. *Org. Lett.* 2009; 11:2213–2215. [PubMed: 19385662] Kise N, Shiozawa Y, Ueda NF. *Tetrahedron.* 2007; 63:5415–5426. Takenaka N, Xia G, Yamamoto HF. *J. Am. Chem. Soc.* 2004; 126:13198–13199. [PubMed: 15479054] Duan X-F, Feng J-X, Zi G-F, Zhang Z-B. *Synthesis.* 2009; 2009:277–282.
4. Fittig RF. *Justus Liebigs Ann. Chem.* 1859; 110:23–45.
5. Recent developments in using ketones in pinacol reactions: Matsukawa S, Hinakubo YF. *Org. Lett.* 2003; 5:1221–1223. [PubMed: 12688724] Shi L, Fan C-A, Tu Y-Q, Wang M, Zhang F-M. *Tetrahedron.* 2004; 60:2851–2855. Aspinall HC, Greeves N, Valla CF. *Org. Lett.* 2005; 7:1919–1922. [PubMed: 15876019]
6. Phosphonate–phosphate rearrangements starting from α -hydroxy trialkyl phosphonoacetates: Corbett MT, Uraguchi D, Ooi T, Johnson JS. *Angew. Chem. Int. Ed.* 2012; 51:4685–4689.
7. Asymmetric reductive coupling of isatins and aldehydes: Horwitz MA, Tanaka N, Yokosaka T, Uraguchi D, Johnson JS, Ooi TF. *Chem. Sci.* 2015; 6:6086–6090. [PubMed: 26508995]
8. Addition of phosphite to α -keto amides followed by phosphonate–phosphate rearrangement and intramolecular trapping: Kondoh A, Aoki T, Terada MF. *Org. Lett.* 2014; 16:3528–3531. [PubMed: 24955761]
9. Developments in phosphonate–phosphate rearrangement using acyl phosphonates as starting materials: Bausch CC, Johnson JS. *Adv. Synth. Catal.* 2005; 347:1207–1211. Demir AS, Reis Ö, dir AÇ, Esiringü , Eymur SF. *J. Org. Chem.* 2005; 70:10584–10587. [PubMed: 16323878]
10. Phosphite additions to aldehydes catalyzed by iminophosphoranes: Uraguchi D, Ito T, Ooi TF. *J. Am. Chem. Soc.* 2009; 131(3836)
11. Phospha-Michael additions to enones and ynones with dialkylphosphite nucleophiles: Li G, Wang L, Yao Z, Xu FF. *Tetrahedron: Asymmetry.* 2014; 25:989–996. Lin Y-D, Kao J-Q, Chen C-T. *Org. Lett.* 2007; 9:5195–5198. [PubMed: 17999511] Zhang A, Cai L, Yao Z, Xu F, Shen QF. *Heteroat. Chem.* 2013; 24:345–354. Yavari I, Hossaini Z, Alizadeh AF. *Monatsh. Chem.* 2006; 137:1083–1088. Zhao D, Yuan Y, Chan A, Wang RF. *Chem. - Eur. J.* 2009; 15:2738–2741. [PubMed: 19212992] Wang F, Wang S, Zhu X, Zhou S, Miao H, Gu X, Wei Y, Yuan QF. *Organometallics.* 2013; 32:3920–3931. Oehler E, Zbiral EF. *Liebigs Ann. Chem.* 1991; 1991:229–236. Touil S, Zantour HF. *Phosphorus, Sulfur Silicon Relat. Elem.* 1997; 131:183–189. Galkin VI, Khabibullina AB, Bakhtiyarova IV, Cherkasov RA, Pudovik AN. *Zh. Obshch. Khim.* 1988; 58:1002–1011. Galkin VI, Galkina IV, Khabibullina AB, Kurdi KA, Cherkasov RA, Pudovik AN. *Dokl. Akad. Nauk SSSR Ser. Khim.* 1990; 314:1408–1411. Vysotskii VI, Prikhod'ko YV, Pavel' GV, Tilichenko MN. *Zh. Obshch. Khim.* 1983; 53:2675–2680. Khidre MD, Abou-Yousef HM, Mahran MRH. *Phosphorus, Sulfur Silicon Relat. Elem.* 2000; 160:181–194.
12. Radosevich and Wang have demonstrated that similar enolates are α -nucleophilic instead of γ -nucleophilic and are also capable of homodimerization: Wang SR, Radosevich AT. *Org. Lett.* 2013; 15:1926–1929. [PubMed: 23578192]
13. Nakamura and Hayashi have shown that related enolates are subject to enantioselective protonation: Hayashi M, Nakamura SF. *Angew. Chem. Int. Ed.* 2011; 50:2249–2252.
14. Núñez MG, Farley AJM, Dixon DJ. *J. Am. Chem. Soc.* 2013; 135:16348–16351. [PubMed: 24107070]
15. Alkylidene pyruvates also undergo the reaction, but the products are not stable to column chromatography.

16. CCDC1432958 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Centre via. www.ccdc.cam.ac.uk/data_request/cif.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Scheme 1.
Asymmetric Reductive Multicomponent Coupling Reactions

Scheme 2.
Chemoselectivity Issues

Scheme 3.**Asymmetric Reductive Coupling Reactions^{a,b}**

^aAll reactions were conducted on 0.1 mmol scale, using 1.1 equiv of dimethyl phosphite and 5.0 equiv of ArCHO. % yields refer to isolated yields. All dr, er, and % yields are the averages of two trials. ^bReaction time = 48 h.

Scheme 4.

Asymmetric Reductive Coupling Reaction on Gram-Scale and X-ray Diffraction Study of **2a**

^aThe reaction was conducted using 1.1 equiv of dimethyl phosphite and 5.0 equiv of ArCHO. % yield refers to isolated yield. Reaction was run for 24 h.

Table 1

Reaction Optimization

1a	+		+
<p>C1 R¹ = Ph, R² = H C2 R¹ = H, R² = <i>t</i>Bu</p> <p>(10 mol %)</p> <p>3a</p> <p>THF (0.1 M)</p>									
entry	temp (°C)	catalyst	dr	er	2a:3a				
1 ^a	0	KO ^t Bu	1.2:1	50:50	100:0				
2 ^b	-60	C1	13:1	9:91	6:1				
3 ^b	-60	C2	17:1	97:3	>20:1				