Published in final edited form as:

Curr Opin Behav Sci. 2016 February ; 7: 82–90. doi:10.1016/j.cobeha.2015.12.002.

Early Caregiving and Human Biobehavioral Development: A Comparative Physiology Approach

Amie A. Hane and Nathan A. Fox

Abstract

A large and growing body of evidence demonstrates associations between quality of the early caregiving environment and risk for stress-related illness across the lifespan. The recent research examining associations between early caregiving environments and subsequent development is reviewed, with particular attention to early programming and subsequent malleability of systems underlying stress responsivity. A developmental comparative physiology model is suggested; one in which postnatal programming and phenotypic plasticity act in concert as mechanisms underlying the persisting effects of early care environments for biobehavioral outcomes.

The field of human development has been inundated with evidence documenting the toxicity of early life stress as a risk factor for lifelong physical and psychiatric disorders. Evidence drawn from large epidemiological studies shows that adults reporting higher levels of childhood adversity report poor health outcomes across the lifespan, including sleep disturbance¹; risk for autoimmune disease²; and risk for mental health disturbance^{3,4}. Adversity in the early rearing ecology appears to set into motion a cascade of stress and physiological responses that undermine long-term health. For instance, the accumulation of adverse events from early- to middle-childhood is associated with elevated biomarkers of inflammation in adolescence⁵. Chronicity of adversity over childhood magnifies the effects of each new stressor, leading to individuals who are less capable of adapting to later life stress given vulnerability imparted by earlier exposures³. Elucidating the mechanisms underlying the early programming of stress physiology is imperative to heading off the cascade of exacerbated risk for reduced health and quality of life that begins in the early care environment (ECE).

There are multiple sources of evidence for the biobehavioral consequences of extremely adverse ECEs on child and adult outcomes. These include exposure to painful procedures in the neonatal intensive care unit⁶, childhood maltreatment⁷ and the global deprivation of

Corresponding Author: Amie A. Hane, Department of Psychology, Williams College, 18 Hoxsey Street, Williamstown, MA 01267, ahane@williams.edu.

Secondary Institutional Affiliation of Corresponding Author:

Division of Developmental Neuroscience, New York State Psychiatric Institute, New York, NY

Co-Author:

Nathan A. Fox, Department of Human Development and Quantitative Methodology, University of Maryland, College Park, 3304 Benjamin Building, College Park, MD 20742

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

institutionalization⁸. Together this evidence supports the importance of the quality of the ECE for the development of stress physiology. Less extreme gradations in the quality of the ECE also show that low quality maternal caregiving behavior (MCB) is associated with suboptimal biobehavioral outcomes^{9–11}. Here we review research demonstrating that the ECE predicts biobehavioral development of the child, with particular attention to postnatal programming and phenotypic plasticity of systems underlying stress responding. We integrate both neuroscience and ecological research as sources of potential mechanisms underlying these effects. A developmental comparative physiology model¹² is suggested; one in which postnatal programming and phenotypic plasticity act in concert as mechanisms underlying the persisting effects of ECE for biobehavioral outcomes.

Early Maternal Care and Postnatal Programming

Maternal caregiving in early infancy sets the stage for biobehavioral adaptations that are to become the basis for stress responsivity and regulation to challenges in the environment. Central to Bowlby's attachment theory is that the infant is prepared to seek proximity to a primary caregiver who serves as a source of safety and comfort in the face of fatigue, distress or threat¹³. Hofer¹⁴ has provided psychobiological evidence to support this, showing that the acute response to maternal separation seen in infantile rodent offspring results from the loss of the individual components packaged within maternal care (i.e., warmth, satiety, tactile stimulation), each of which plays a critical role in supporting homeostasis in pups.

The regulatory capacity of maternal care depends upon the quality of caregiving behavior. Naturally-occurring individual differences in the rodent in the frequency of licking and grooming (LG) behavior are associated with alterations in central corticotropin-releasing factor and stress responsive systems¹⁵ and accompanying behavioral differences that persist into adulthood. Rat pups that experience low levels of LG display, as adults, a stress-reactive neuroendocrine profile¹⁶ and a corresponding behavioral profile of elevated stress reactivity^{17,18}. These effects are mediated by epigenetic alterations to stress-related genes, with high (relative to low) LG offspring showing decreased DNA methylation of the glucocorticoid receptor region of the hippocampus, which is associated with the reduction in stress responding documented in high LG offspring¹⁹.

Early Care and Postnatal Programming in Humans

Rodent models affirm the developmental relevance of early-occurring maternal care to biobehavioral development via postnatal programming that is accompanied by epigenetic alterations to systems underlying stress responding. Parallel examples exist in the human literature. Similar to the rodent model, ordinary variations in the quality of maternal care in full-term healthy infants is associated with phenotypic changes associated with increased stress and social difficulties¹⁰. Within the context of a large, longitudinal study of low-risk mothers and their 9-month-old infants, mothers and infants were observed in the home and rated for quality of maternal behavior for sensitive, non-intrusive interventions during the course of routine care in the home (feeding, application of lotion, and dressing). A quality of maternal caregiving behavior (MCB) variable was derived from ratings of maternal sensitivity. Relative to infants who received high quality MCB, those who experienced low

quality MCB displayed a stress-reactive biobehavioral profile, marked by more fearfulness during the presentation of novel stimuli, less positive joint attention to a shared object with an unfamiliar experimenter, and more negative affect during mother-child interaction. Infants who experienced low quality MCB also showed a pattern of resting relative right frontal EEG asymmetry, which previous research has shown to be associated with higher basal and stress-induced salivary cortisol concentrations in 6-month-olds²⁰ and withdrawal motivation across early development²¹. These findings suggest that ‘ordinary’ variations in the quality of the early caregiving environment yields contemporaneous phenotypic changes to the systems involved in regulation of stress—effects that closely parallel the rodent postnatal programming models.

Quality of maternal care may have important implications for the neurological substrates that influence subsequent parenting behavior. For instance, relative to mothers who retrospectively perceived high quality maternal care in childhood, those who retrospectively perceived experiencing low quality maternal care in childhood showed more activation of the hippocampus when exposed to infant crying²². Such hippocampal activation to infant crying may suggest that early maternal care influences maternal neural response to parenting-related stress in adulthood.²³ This profile of neural activation of stress in the context of parenting may be a function of the reactivation of stress that originated in the ECE. The adverse ECE may prime mothers for stress that is reactivated upon exposure to the context of parenting in the future via the mechanism of phenotypic plasticity.

Phenotypic Plasticity

Evolutionary biology has long documented the phenomenon of phenotypic plasticity, defined as the ability of a single genotype to produce different phenotypes based on the demands of the environment²⁴. Ecologists offer compelling evidence demonstrating that environmental factors predispose organisms to not only respond defensively, or adapt to their environment; but also to be primed for malleability to mount the same defense again in the future. Reactivation of a defensive adaptation is context-dependent, and occurs when the organism is confronted with the same ecology in the future. A compelling illustration comes from research on the intertidal snails that dwell in rocky tide pools on the coast of Maine. Trussell sampled intertidal snails from two natural habitat conditions—i.e., from tide pools where exposure to predatory green crabs was high; and from tide pools where threat of predation was low. When exposed experimentally to a predatory crab in the laboratory, the snails that were sampled from tide pools with high risk of predation showed morphological changes—exhibiting significantly greater shell density growth and smaller body size than snails sampled from tide pools in which the risk for predation was low²⁵. In essence, ecological adversity primed these organisms for change, increasing malleability, thereby insuring that a defensive response would be mounted more readily, or reactivated, when encountering threat in the future. Reactivation is also dose-dependent, as the amount of shell density growth was influenced by the intensity of the risk for predation²⁶. The defense is formidable, as the denser shell is also stronger and the reduced body size limits the crab’s accessibility to the snail’s soft tissue²⁶. The stress of predation is associated with a decrease in growth efficiency associated with less foraging behavior, thereby making the snail a less efficient source of energy for its predator²⁷. However, each exposure to predatory risk adds

the additional burden of carrying the weight of the thicker shell along with the accompanying reduced body mass.

Contemporary research provides evidence for epigenetic mediation of phenotypic adaptations²⁴. The intertidal snail shows transcriptional alterations in RNA in response to thermal stress and risk for predation, and the profile of transcriptional activity is distinct for each type of stressor²⁸. Recent evidence from another intertidal organism, the sea urchin, highlights the developmental nature of phenotypic plasticity. Sea Urchin larvae float freely in open waters before settling into intertidal habitats. The larvae are commonly confronted with food (i.e., algae) scarcity. Food-deprived larvae adapt with morphological changes, including development of longer arms and reabsorption of stomach tissue, leading to smaller rudiments. These morphological changes are associated with epigenetic differences—when experimentally compared with well-fed larvae, starved larvae show epigenetic alterations to the genes responsible for metabolic rate, growth, mitochondrial activity, regulation of homeostasis, and resistance to stress²⁹. Together these examples from the ecology literature highlight two points: First, when faced with adversity organisms adapt to their situation with significant morphological changes. And second, the mechanisms for these adaptations appear to be epigenetic in nature. Both of these processes may be present in the human response to adversity as well.

Early Care Environments and Phenotypic Plasticity

In the rodent model, it has been suggested that a general profile of *defensive reactivity* develops in low LG rodents because behavioral defensiveness and the associated release of stress-related hormones are adaptive, allowing for detection of threat and the mobilization of metabolic resources under suboptimal early care conditions³⁰. Much like the intertidal snail, this mounting of a defensive response may serve to prime the system to mount such a reaction in the future under similar environmental conditions, and to do so more readily than others of the same species reared under less aversive conditions.

Phenotypic plasticity is also applicable to humans, with infants showing both a behavioral and physiological defensive response during the experience of low-quality caregiving. Hane and Philbrook observed mothers as they undressed, bathed, and re-dressed their 4–8 week old full-term infants in the home and MCB was coded. Salivary cortisol was measured in infants prior to tub bathing and again 15 minutes following removal of the neonate from the bath water. Infants of mothers who provided lower quality MCB showed a significantly larger increase in cortisol (over basal levels) following this experience of routine care⁹. Hence, the experience of being cared for by an insensitive caregiver early in life was associated with elevated stress responding linked to the early care experience.

A growing body of literature demonstrates that ECEs are associated with epigenetic alterations in peripheral tissue sampling in humans^{31–35} and this research is consistent with the phenotypic plasticity modelⁱ. Table 1 highlights four studies that document that epigenetic changes are dynamically linked to the ECE, with evidence of change over time as

ⁱAn extensive review of epigenetics and ECEs for humans is beyond the scope of this review, readers are referred to Boyce and Kober⁵⁴

a function of prolonged exposure to adversity. For example, in preterm infants, the experience of the NICU itself is associated with epigenetic changes in methylation over time, varying from birth to postnatal day four³² to time of discharge from the NICU³¹. In institutionalized children, epigenetic alterations to telomere length is associated with duration of time spent in the institution³³, with a steady decrease in telomere length from ages 8–14 for children who remain institutionalized.³⁶ Hence, epigenetic markers are further altered as exposure to adversity increases. It may not be surprising then that widespread changes in methylation across the genome have been found for children with a history of earlier exposure to maltreatment³⁵ and in adolescents with exposure to high levels of parental stress³⁴. Dramatic epigenetic changes associated with a history of adverse ECEs by middle childhood and adolescence may be the result of continual epigenetic changes that accumulate across time as exposure to adversity or new stressors are encountered.

Context-Dependent Reactivation

There is also support for context-dependent reactivation of stress responding based on features of the ECE. Even a singular disruption to mother-infant interaction may lead to context-dependent reactivation. Six-month infants were brought to the lab and randomly assigned to either a mother-infant face-to-face interaction (control group) or the maternal still-face paradigm (SFP). The infants who were confronted with the maternal termination of contingent responsiveness in the SFP showed a significant increase in salivary cortisol following the SFP. When brought back to the laboratory the next day, placement of the infant in the same context, but not proceeding with the SFP resulted in reactivation of the cortisol response for infants in the SFP condition. No such effect was found for control infants on days 1 or 2 of the experiment.

Additional evidence comes from the rodent and human maternal care literatures. If the receipt of insensitive care in the early relational context primes the system for defensive responding in similar contexts, then social interactions with others may trigger reactivation of stress physiology. One might expect that offspring who experience stress during early care may have difficulties later on while interacting with peers. This too is supported by animal models. Juvenile male offspring who received low LG as pups engaged in more play fighting in multiple play partners housing than high LG males³⁷ and adult offspring receiving low LG as pups manifested more aggressive and defensive behavior during a resident-intruder test, as compared to adult offspring who received high LG as pups³⁸. The ECE of the human is also associated with a profile of defensive play behavior with peers. In a longitudinal follow-up of MCB and biobehavioral responding, we found relative to children who experienced high quality MCB as infants (high MCB children), those who received low quality MCB (low MCB children) continued to show increased stress reactivity on measures that parallel those used in our earlier report, including inhibited social behavior with adults and right frontal EEG asymmetry¹¹. As well, low MCB children manifested more aggression during play with a novel peer than low MCB children. Mothers reported that low MCB children tended to show more internalizing problems and more proneness to anger in social situations.

Additional support for defensive reactivation as a function of ECE is present for children exposed to more extreme adversity in the ECE. Longitudinal follow-up of a children born very preterm shows that higher exposure to procedural pain (routine blood draw) is associated with reduced cortisol responding to acute pain at age 4-months³⁹ and lower basal cortisol and less variable diurnal rhythms in cortisol at school age⁶. Children who were institutionalized in infancy also exhibit differences in stress responding later in childhood⁴⁰. In a longitudinal follow-up of children from the Bucharest Early Intervention Project⁴¹, children underwent the Trier Social Stress Test (TSST) at age 12 and neuroendocrine and autonomic stress responding were examined. Relative to children who were randomly assigned to foster care, continually institutionalized children showed blunted cortisol responding across the stress task. Autonomic responding was also blunted for the continually institutionalized group, in terms of heart rate, diastolic blood pressure, and pre-ejection period. Placement into foster care before ages 18–24 months was significantly associated with the increase in stress physiology, similar to that found in the never institutionalized community controls. Hence, there may be a sensitive period for ECE programming of stress physiology⁴⁰. Autonomic responding is also dampened upon reactivation in children with a history of abuse/neglect. Children with a history of maltreatment failed to show an increase in blood pressure in response to a social laboratory stressor; while non-maltreated children showed a significant increase to the laboratory stress paradigm.⁷ Hence, for instances of extremely adverse ECE's, defensive reactivation manifests as a less variable, or blunted physiological response to stress, indicative of physiological dysregulation that is specific to the nature of the ECE.

When does Reactivation Become Maladaptive?

Adverse rearing ecologies prime the system to adapt to the conditions within which development ensued. The infant of a low MCB mother benefits from the activation of stress physiology to meet the metabolic demand of discomfort. When remaining in the context of ecological adversity, increased reactivity and defensive adaptations may be advantageous for survival. For instance, in the rodent, female offspring of low- and mid- LG dams show play dominance and are particularly likely to engage in pinning behavior when female peers are in estrus, which may promote their social rank and reproductive advantage in competitive environments in which resources are scarce⁴². Female children of mothers who were rated as high in maternal harshness in early childhood were significantly more likely to reach menarche earlier, and early menarche was predictive of sexual risk-taking behavior⁴³. Hence, programming from the early rearing environment may result in activation of defensive responding that is adaptive for offspring who remain in harsh ecologies. Ellis has suggested that 'fast' life history strategies promote reproductive success at an earlier age for offspring whose ecology forecasts less longevity⁴⁴.

The low MCB child epigenetically programmed for heightened stress responding may manifest context-dependent reactivation of stress responding in each new social setting outside of the home. When the formative social experience offered by mother signals discomfort and stress in the presence of another, each new social context thereafter may grow increasingly more difficult to navigate, as the increase in stress physiology and defensive reactivation costs the individual, becoming a thicker shell and a heavier burden

over time. The physiological load of continually heightened stress in the social world is carried within the child, creating anxiety in early childhood^{11,45} that may influence cardiovascular health beginning in childhood⁴⁶ and inflammatory responses that compromise immune health into adulthood⁴⁷.

More troubling is that the system is primed for reactivation under extremely adverse ECE. Institutionalized infants adapt to multiple caregivers with indiscriminate sociability. This is adaptive in the institution, where indiscriminate sociability ensures receipt of care—the child who reaches out to more caregivers is more likely to receive care. However, under conditions of lower adversity, this behavior is maladaptive, diagnosed as Reactive Attachment Disorder, and interferes with the development of a healthy attachment to a singular primary caregiver⁴⁸. Maltreated children show greater ERP responding to angry (vs. sad or neutral) faces, which is adaptive when the child's sensitivity to the affective valence of the abusive caregiver signals impending danger. However, in the non-abusive context outside of the home of origin, this type of hypervigilance to mood interferes with attentional resources that need to be directed toward learning and healthy social functioning⁴⁹. The long-term health consequences of altered stress physiology as seen in low MCB children, NICU survivors, institutionalized infants, and maltreated children may result in lifelong risk for developmental psychopathology, as context-dependent reactivation continues to occur across the lifespan.

Context Dependent Reactivation in Adulthood

Phenotypic plasticity as a function of the ECE and the associated epigenetic alterations are not limited to childhood. There is evidence for reactivation of defensive responding from stress encountered in adulthood in rodent models. For instance, female adult offspring who experienced repeated postnatal maternal separation in the first two weeks of life (in comparison to non-handled controls), were assessed for anhedonic responding after exposure to stress introduced in adulthood. Postnatally maternally-separated rats showed a blunted anhedonic response to an initial acute exposure to social defeat (pinning by a male resident-intruder), but an elevated anhedonic response was found after 7 daily exposures of the same social defeat paradigm. Hence, the ECE predicted later reactivation characterized by an initial protection against the anhedonic response to a single social stressor, but a swift reactivation, with an increased anhedonic response, when confronted with the same social stressor repeatedly⁵⁰. Further, joint exposure of postnatal maternal separation (vs. handling or normal rearing) and adulthood stress (novel cage or social defeat) was associated with increased *slc6a4* mRNA expression in serotonergic neurons only for rats that experienced postnatal separation *and* social defeat in adulthood. Hence, subsequent social stress in adulthood further increased the epigenetic alterations associated with dysregulation of the serotonergic systems involved in depression and anxiety resulting from the ECE⁵¹. In the human, early life trauma, and particularly child abuse, is associated with PTSD in adulthood following exposure to re-victimization in adulthood, particularly when both the ECE and the later life trauma are contextually similar⁵². Depression is associated with inflammatory processes that are associated with early-life exposure to stress⁵³. In a study of depressed men with and without a history of early life trauma, only depressed men with a history of early adversity showed an increase in inflammatory response (increased proinflammatory

markers in plasma and DNA binding) to the TSST⁵⁴. Depressed women with a history of depression and a history of child abuse showed significantly higher neuroendocrine responding to the TSST (greater plasma ACTH and cortisol concentrations) relative to depressed women with no abuse history and community controls⁵⁵. Taken together, this evidence suggests that stressful life events in adulthood may reactivate programmed stress responding that originated in the ECE, particularly when the context of reactivation shares features of the ECE, i.e., social stressors in particular. Reactivation of stress physiology throughout the lifespan may also give rise to further increases in epigenetic alterations that originated in the ECE.

Summary and Conclusions

Animal models and studies in humans indicate that adverse early care environments program the system via epigenetic alterations to the systems responsible for biobehavioral stress responding^{32,56,57}. Here we suggest that developing stress physiology is phenotypically plastic. Recent advances in the field of ecology show that phenotypic plasticity is mediated by epigenetic changes induced by the ecology of early development²⁹. Context-dependent reactivation of postnatal programming as a function of the ECE may be the mechanism underlying the significant health footprint left in the wake of suboptimal care¹⁻⁴. As infants reared in adverse ECE's enter new social contexts, the burden of early adversity carries with them; setting into motion the cascade of context-dependent reactivation of defenses initially acquired in the ECE. These adaptations triggered by formative social experiences carry over to new social contexts, when they may no longer be of adaptive value, increasing the likelihood of reactivation in each new stressful context, which further contributes to more stress-related vulnerability. Context-dependent reactivation and the accompanying epigenetic changes remain ongoing processes across development that accumulates risk for the plethora of stress-related illnesses across the lifespan of individuals who experienced early adversity¹⁻⁴—be it nuanced differences in the quality of maternal care or extreme instances of adversity.

Acknowledgments

This research was supported by funds from the Groff Foundation to Amie A. Hane and a grant from the National Institutes of Mental Health (MH093349) to Nathan A. Fox. We would like to thank S. Wieman and M. Mitsuyama of Williams College for their assistance with this research. We extend our most sincere gratitude to the families who continue to participate in our research.

Glossary

Early Care Environment (ECE)

Refers to the characteristics of the rearing environment. Across species, this includes the elements of the caregiving environment that reduce or activate stress responding in offspring. In the human these include insensitive maternal caregiving behavior and exposures to extremely aversive conditions, such as parental separation and procedural pain of the Neonatal Intensive Care Unit; parental separation and global neglect of institutionalization; and parental maltreatment of the child, which includes neglect and abuse

Postnatal Programming

Drawn from rodent models of maternal care, refers to the processes through which quality of maternal caregiving behavior, defined in terms of high or low frequencies of licking/grooming behavior (usually occurring in the context of arch-backed nursing), in the first two weeks of life, induce changes in the behavioral and physiological regulation of stress that result in altered behavioral and physiological differences that persist across the lifespan. These effects are mediated by epigenetic changes, alterations in the expression of genes involved in regulation of stress and stress-related illness¹

Phenotypic Plasticity

Drawn from ecology and research in lower life forms, refers to the ability of a single genotype to produce more than one phenotype based on the demands of their environment². The differential expression of a phenotype depends upon these environmental demands, which elicit adaptive/defensive responses in organisms in response to adversity (e.g., food scarcity, climate change, and risk of predation) that results in morphological changes. Based on exposure to the challenges of the environment of origin, the organism then becomes phenotypically flexible, and will mount the same defense again in the future. These effects are also associated with transcriptional alterations in the candidate genes associated with the biochemical processes underlying the altered phenotype^{2,3}

Context Dependent Reactivation

A term extrapolated from the phenotypic plasticity literature, here we refer to the human response in which infants, children, or adults previously programmed for behavioral and physiological stress by the elements of an adverse ECE show reactivation of the stress response when encountering future stressors—particularly stressors that share contextual features with the stress-inducing features of the ECE (usually social stressors, as the context of early care represents the formative social experience)

Annotated Bibliography

1. Chapman DP, Wheaton AG, Anda RF, et al. Adverse childhood experiences and sleep disturbances in adults. *Sleep Medicine*. 2011; 12(8):773–779. [PubMed: 21704556]
2. Dube SR, Fairweather D, Pearson WS, Felitti VJ, Anda RF, Croft JB. Cumulative childhood stress and autoimmune diseases in adults. *Psychosomatic Medicine*. 2009; 71(2):243–250. [PubMed: 19188532]
3. Nurius PS, Green S, Logan-Greene P, Borja S. Life course pathways of adverse childhood experiences toward adult psychological well-being: A stress process analysis. *Child Abuse & Neglect*. 2015; 45:143–153. [PubMed: 25846195]

¹Weaver ICG, Cervoni N, Champagne FA, et al. Epigenetic programming by maternal behavior. *Nat. Neurosci.* Aug 2004;7(8):847–854.;

²Healy TM, Schulte PM. Phenotypic plasticity and divergence in gene expression. *Molecular Ecology*. Jul 2015;24(13):3220–3222.;

³Chu ND, Miller LP, Kaluziak ST, Trussell GC, Vollmer SV. Thermal stress and predation risk trigger distinct transcriptomic responses in the intertidal snail *Nucella lapillus*. *Molecular Ecology*. Dec 2014;23(24):6104–6113.

4. McCrory C, Dooley C, Layte R, Kenny RA. The lasting legacy of childhood adversity for disease risk in later life. *Health Psychology*. 2015; 34(7):687–696. [PubMed: 25150540]
5. Slopen N, Kubzansky LD, McLaughlin KA, Koenen KC. Childhood adversity and inflammatory processes in youth: a prospective study. *Psychoneuroendocrinology*. Feb; 2013 38(2):188–200. [PubMed: 22727478]
6. Brummelte S, Chau CMY, Cepeda IL, et al. Cortisol levels in former preterm children at school age are predicted by neonatal procedural pain-related stress. *Psychoneuroendocrinology*. 2015; 51:151–163. [PubMed: 25313535]
7. Leitzke BT, Hilt LM, Pollak SD. Maltreated youth display a blunted blood pressure response to an acute interpersonal stressor. *Journal of Clinical Child and Adolescent Psychology*. 2015; 44(2):305–313. [PubMed: 24175880]
8. Sigal JJ, Perry JC, Rossignol M, Ouimet MC. Unwanted infants: Psychological and physical consequences of inadequate orphanage care 50 years later. *American Journal of Orthopsychiatry*. 2003; 73(1):3–12. [PubMed: 12674514]
9. Hane AA, Philbrook LE. Beyond licking and grooming: Maternal regulation of infant stress in the context of routine care. *Parenting: Science and Practice*. 2012; 12(2–3):144–153.
10. Hane AA, Fox NA. Ordinary Variations in Maternal Caregiving Influence Human Infants' Stress Reactivity. *Psychological Science*. 2006; 17(6):550–556. [PubMed: 16771807]
11. Hane AA, Henderson HA, Reeb-Sutherland BC, Fox NA. Ordinary variations in human maternal caregiving in infancy and biobehavioral development in early childhood: A follow-up study. *Developmental Psychobiology*. 2010; 52(6):558–567. [PubMed: 20806328]
- 12*. Mueller CA, Eme J, Burggren WW, Roghair RD, Rundle SD. Challenges and opportunities in developmental integrative physiology. *Comp Biochem Physiol A-Mol Integr Physiol*. Jun.2015 184:113–124. This is a review paper of the research integrating the effects of early experience on epigenetic changes across animal taxa. During critical windows, periods of plasticity, animals and humans are vulnerable to maternal and environmental changes, but also receptive to recovery and intervention. [PubMed: 25711780]
13. Bowlby, J. A secure base: Parent-child attachment and healthy human development. New York, NY, US: Basic Books; 1988.
- 14*. Hofer MA. Psychobiological Roots of Early Attachment. *Current Directions in Psychological Science*. 2006; 15(2):84–88. A review of a series of elegant studies demonstrating the 'hidden regulators' packaged within maternal care. When separated from the dam in the postnatal period, the following effects were found: Loss of warmth provided by the dam was associated brain changes that result in slowed activity level; loss of satiety provided by maternal milk resulted in loss of sympathetic tone, resulting in a 40% reduction in heart rate; loss of tactile stimulation provided by mother resulted in behavioral reactivity to novelty.
15. Francis DD, Meaney MJ. Maternal care and the development of stress responses. *Current Opinion in Neurobiology*. Feb; 1999 9(1):128–134. [PubMed: 10072372]
16. Liu D, Diorio J, Tannenbaum B, et al. Maternal care, hippocampal glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress. *Science*. 1997; 277(5332):1659–1662. [PubMed: 9287218]
17. Caldji C, Tannenbaum B, Sharma S, Francis D, Plotsky PM, Meaney MJ. Maternal care during infancy regulates the development of neural systems mediating the expression of fearfulness in the rat. *Proceedings of the National Academy of Sciences of the United States of America*. Apr; 1998 95(9):5335–5340. [PubMed: 9560276]
18. Francis D, Diorio J, Liu D, Meaney MJ. Nongenomic transmission across generations of maternal behavior and stress responses in the rat. *Science*. Nov; 1999 286(5442):1155–1158. [PubMed: 10550053]
- 19*. Weaver ICG, Cervoni N, Champagne FA, et al. Epigenetic programming by maternal behavior. *Nat Neurosci*. Aug; 2004 7(8):847–854. Seminal paper demonstrating DNA methylation of the glucocorticoid receptor (GR) gene promoter in the hippocampus and acetylation of histones as a function of maternal care in the rat. Results show that relative to low LG rats, high LG rats experience changes in their hippocampal GR RNA levels and glucocorticoid negative feedback sensitivity. These changes mediate a more modest HPA axis stress response in high LG rats. [PubMed: 15220929]

20. Buss KA, Schumacher JRM, Dolski I, Kalin NH, Goldsmith HH, Davidson RJ. Right frontal brain activity, cortisol, and withdrawal behavior in 6-month-old infants. *Behav Neurosci.* 2003; 117(1): 11–20. [PubMed: 12619903]
21. Hane AA, Fox NA, Henderson HA, Marshall PJ. Behavioral reactivity and approach-withdrawal bias in infancy. *Developmental Psychology.* 2008; 44(5):1491–1496. [PubMed: 18793079]
22. Kim P, Leckman JF, Mayes LC, Newman M-A, Feldman R, Swain JE. Perceived quality of maternal care in childhood and structure and function of mothers' brain. *Developmental Science.* 2010; 13(4):662–673. [PubMed: 20590729]
23. Kim P, Strathearn L, Swain JE. The maternal brain and its plasticity in humans. *Hormones and Behavior.* 2015
24. Healy TM, Schulte PM. Phenotypic plasticity and divergence in gene expression. *Molecular Ecology.* 2015; 24(13):3220–3222. [PubMed: 26096949]
25. Trussell GC. Phenotypic plasticity in an intertidal snail: The role of a common crab predator. *Evolution.* Feb; 1996 50(1):448–454.
26. Trussell GC, Nicklin MO. Cue sensitivity, inducible defense, and trade-offs in a marine snail. *Ecology.* Jun; 2002 83(6):1635–1647.
27. Trussell GC, Ewanchuk PJ, Matassa CM. The fear of being eaten reduces energy transfer in a simple food chain. *Ecology.* Dec; 2006 87(12):2979–2984. [PubMed: 17249221]
28. Chu ND, Miller LP, Kaluziak ST, Trussell GC, Vollmer SV. Thermal stress and predation risk trigger distinct transcriptomic responses in the intertidal snail *Nucella lapillus*. *Molecular Ecology.* Dec; 2014 23(24):6104–6113. [PubMed: 25377436]
29. Carrier TJ, King BL, Coffman JA. Gene Expression Changes Associated With the Developmental Plasticity of Sea Urchin Larvae in Response to Food Availability. *Biol Bull.* Jun; 2015 228(3): 171–180. [PubMed: 26124444]
30. Parent C, Zhang TY, Caldji C, et al. Maternal care and individual differences in defensive responses. *Current Directions in Psychological Science.* Oct; 2005 14(5):229–233.
31. Provenzi L, Fumagalli M, Sirgiovanni I, et al. Pain-related stress during the Neonatal Intensive Care Unit stay and SLC6A4 methylation in very preterm infants. *Frontiers in Behavioral Neuroscience.* 2015:9. [PubMed: 25691862]
32. Kantake M, Yoshitake H, Ishikawa H, Araki Y, Shimizu T. Postnatal epigenetic modification of glucocorticoid receptor gene in preterm infants: a prospective cohort study. *BMJ Open.* 2014; 4(7):8.
33. Drury SS, Theall K, Gleason MM, et al. Telomere length and early severe social deprivation: linking early adversity and cellular aging. *Molecular psychiatry.* Jul; 2012 17(7):719–727. [PubMed: 21577215]
34. Essex MJ, Boyce WT, Hertzman C, et al. Epigenetic vestiges of early developmental adversity: childhood stress exposure and DNA methylation in adolescence. *Child Dev.* Jan-Feb; 2013 84(1): 58–75. [PubMed: 21883162]
35. Yang B-Z, Zhang H, Ge W, et al. Child abuse and epigenetic mechanisms of disease risk. *American Journal of Preventive Medicine.* 2013; 44(2):101–107. [PubMed: 23332324]
36. Melicher D, Buzas EI, Falus A. Genetic and epigenetic trends in telomere research: a novel way in immunoepigenetics. *Cellular and molecular life sciences : CMLS.* Nov; 2015 72(21):4095–4109. [PubMed: 26190020]
37. Parent CI, Meaney MJ. The Influence of Natural Variations in Maternal Care on Play Fighting in the Rat. *Developmental Psychobiology.* Dec; 2008 50(8):767–776. [PubMed: 18846499]
38. Menard JL, Hakvoort RM. Variations of maternal care alter offspring levels of behavioural defensiveness in adulthood: Evidence for a threshold model. *Behavioural Brain Research.* Jan; 2007 176(2):302–313. [PubMed: 17084914]
39. Morison SJ, Grunau RE, Oberlander TF, Whitfield MF. Relations between behavioral and cardiac autonomic reactivity to acute pain in preterm neonates. *The Clinical Journal of Pain.* 2001; 17(4): 350–358. [PubMed: 11783816]
40. McLaughlin KA, Sheridan MA, Tibu F, Fox NA, Zeanah CH, Nelson CA. Causal effects of the early caregiving environment on development of stress response systems in children. *Proceedings*

of the National Academy of Sciences of the United States of America. May; 2015 112(18):5637–5642. [PubMed: 25902515]

- 41*. Zeanah CH, Nelson CA, Fox NA, et al. Designing research to study the effects of institutionalization on brain and behavioral development: The Bucharest Early Intervention Project. *Development and Psychopathology*. Fall;2003 15(4):885–907. Foundational paper describing the methodology of the Bucharest Early Intervention Project, in which the developmental effects of institutionalization on young children are examined. The Intervention Project consisted of three conditions: Institutionalized (41), Foster Care (57), and Never Institutionalized (60). Children in the Institutionalized and Foster Care conditions were from orphanages in and around Bucharest. At the outset of the study, the Foster Care group did not differ from the Institutionalized group on key measures such as gender distribution, age, birth weight, or percentage of life spent in the institution. [PubMed: 14984131]
42. Parent CI, Del Corpo A, Cameron NM, Meaney MJ. Maternal Care Associates With Play Dominance Rank Among Adult Female Rats. *Developmental Psychobiology*. Nov; 2013 55(7):745–756. [PubMed: 22786820]
43. Belsky J, Steinberg L, Houts RM, Halpern-Felsher BL. The development of reproductive strategy in females: Early maternal harshness → earlier menarche → increased sexual risk taking. *Developmental Psychology*. 2010; 46(1):120–128. [PubMed: 20053011]
- 44*. Ellis BJ, Del Giudice M. Beyond allostatic load: Rethinking the role of stress in regulating human development. *Development and Psychopathology*. 2014; 26(1):1–20. Although necessary, stress responses can have negative long-term effects on development. This paper presents the life history theory that highlights the malleability of stress response systems to be adaptive in regulating alternative developmental pathway. Drawing on the adaptive calibration model, this theory proposes the need for consideration of biological fitness trade-offs in order to explain the relations between developmental exposures to stress, stress responsivity, behavioral strategies, and health. [PubMed: 24280315]
45. Mount KS, Crockenberg SC, J6 PSB, Wagar J-L. Maternal and child correlates of anxiety in 2½-year-old children. *Infant Behav Dev*. 2010; 33(4):567–578. [PubMed: 20723999]
46. Monk C, Kovelenco P, Ellman LM, et al. Enhanced stress reactivity in paediatric anxiety disorders: Implications for future cardiovascular health. *International Journal of Neuropsychopharmacology*. 2001; 4(2):199–206. [PubMed: 11468091]
47. Moons WG, Shields GS. Anxiety, Not Anger, Induces Inflammatory Activity: An Avoidance/Approach Model of Immune System Activation. *Emotion*. 2015
48. Zeanah CH, Smyke AT, Koga SF, Carlson E. Attachment in Institutionalized and Community Children in Romania. *Child Development*. 2005; 76(5):1015–1028. [PubMed: 16149999]
49. Pollak SD, Klorman R, Thatcher JE, Cicchetti D. P3b reflects maltreated children's reactions to facial displays of emotion. *Psychophysiology*. 2001; 38(2):267–274. [PubMed: 11347872]
- 50*. Der-Avakian A, Markou A. NEONATAL MATERNAL SEPARATION EXACERBATES THE REWARD-ENHANCING EFFECT OF ACUTE AMPHETAMINE ADMINISTRATION AND THE ANHEDONIC EFFECT OF REPEATED SOCIAL DEFEAT IN ADULT RATS. *Neuroscience*. Nov; 2010 170(4):1189–1198. During post-natal days 1–14, Long Evans rat pups were separated from their mothers for 3 hours each day in order to examine how neonatal maternal separation affects anhedonia under baseline conditions and in response to amphetamine administration. Anhedonic responding was measured via intracranial self-stimulation of the posterior lateral hypothalamus to assess neurological activity associated with activation of reward centers. Results show that anhedonic responding was jointly affected by postnatal separation and subsequent exposure to stress (social defeat). [PubMed: 20691770]
51. Gardner KL, Hale MW, Lightman SL, Plotsky PM, Lowry CA. Adverse early life experience and social stress during adulthood interact to increase serotonin transporter mRNA expression. *Brain Research*. Dec.2009 1305:47–63. [PubMed: 19781533]
52. Pratchett LC, Yehuda R. Foundations of posttraumatic stress disorder: Does early life trauma lead to adult posttraumatic stress disorder? *Development and Psychopathology*. 2011; 23(2):477–491. [PubMed: 23786690]

53. Cattaneo A, Macchi F, Plazzotta G, et al. Inflammation and neuronal plasticity: a link between childhood trauma and depression pathogenesis. *Frontiers in cellular neuroscience*. 2015; 9:40. [PubMed: 25873859]
54. Pace TWW, Mletzko TC, Alagbe O, et al. Increased stress-induced inflammatory responses in male patients with major depression and increased early life stress. *American Journal of Psychiatry*. Sep; 2006 163(9):1630–U1631. [PubMed: 16946190]
55. Heim C, Newport DJ, Heit S, et al. Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *Jama*. Aug 2; 2000 284(5):592–597. [PubMed: 10918705]
56. Provenzi L, Fumagalli M, Sirgiovanni I, et al. Pain-related stress during the Neonatal Intensive Care Unit stay and SLC6A4 methylation in very preterm infants. *Frontiers in Behavioral Neuroscience*. Apr.2015 9:9. [PubMed: 25691862]
57. Chau CMY, Ranger M, Sulistyoningrum D, Devlin AM, Oberlander TF, Grunau RE. Neonatal pain and COMT Val158Met genotype in relation to serotonin transporter (SLC6A4) promoter methylation in very preterm children at school age. *Frontiers in Behavioral Neuroscience*. Dec. 2014 8:12. [PubMed: 24478659]
- 54**. Boyce WT, Kobor MS. Development and the epigenome: the ‘synapse’ of gene-environment interplay. *Developmental Science*. Jan; 2015 18(1):1–23. A summary of research examining the interplay of genes and environment as mediated by epigenetics. Through chromatin reorganization, epigenetic processes alter the expression of a gene without changing the genetic code. Epigenetic changes are inheritable and could play a key role in developmental trajectory differences, sustained effects of early adversity, and varying vulnerability during critical periods. [PubMed: 25546559]

Highlights

- Reviews the research examining early caregiving and subsequent development.
- Programming and the malleability of stress-responding systems are emphasized.
- Our comparative physiology model integrates research in neuroscience and ecology.
- We present a model of phenotypic plasticity in human stress responding.

Figure. Quality of Early Caregiving Influences Biobehavioral Development in Humans: A Model of Phenotypic Plasticity

The quality of the early care environment (ECE) gives rise to differences in stress responding through *Postnatal Programming* (including epigenetic alterations) of the systems that regulate physiological stress responding. This programming primes the child by eliciting defensive and adaptive biobehavioral stress responding in the ECE. The child is also primed to respond with the same defensive biobehavioral responding upon exposure to future stressors across development via the process of *Context Dependent Reactivation*. Each subsequent exposure to new stress across development yields increased vulnerability for the child based on malleability to stress responding acquired in the ECE, as programmed defensive responses are reactivated. Reactivation 1) occurs when the child is confronted with similar stressors in the future; 2) can be considered adaptive in the ECE but maladaptive in less stressful contexts; and 3) may be associated with continued dynamic shifts in the epigenome that ultimately give rise to stress-related illness and difficulties in social functioning across the lifespan.

Table 1

Summary of Research Demonstrating Change Over Time in Epigenetic Alterations Linked to Early Care Environments.

Early Care Environment	Citation	Nature of Study Sample	Type of Peripheral Tissue and Age at DNA Sampling	Epigenetic Alteration	Functional Significance
Neonatal Intensive Care Unit	Provenzi et al., 2014 ¹	56 children born very preterm (24–28 weeks gestation) and 32 full-term controls. Very preterm infants were divided into high ($n=31$) vs. low ($n=25$) exposure to procedural pain (skin-breaking procedures) while in the NICU	Cord blood (at birth) and peripheral blood (in the very preterm group only at time of hospital discharge)	Methylation at 20 CpG sites within the promoter region of the <i>SLC6A4</i> gene were examined. No differences found at birth between very preterm and full term infants. Relative to very preterm infants who experienced lower procedural pain, those who experienced high procedural pain were found to have significantly increased methylation at CpG sites 5 and 6 at time of discharge.	Higher <i>SLC6A4</i> methylation is associated with intrasynaptic serotonin signaling, yielding increased serotonin concentrations consistent with biochemistry of depressed and anxious adults.
	Kantake et al., 2014 ²	40 infants (20 full term and 20 preterm)	Cord blood at birth and peripheral blood at day postnatal day 4	Methylation rates in the 1-F promoter region of the glucocorticoid receptor gene in 33 CpG sites were examined at birth and postnatal day 4. In preterm infants, methylation rates significantly increased at multiple sites from birth to postnatal day 4. Methylation rates remained stable in full-term infants.	Increased methylation of the glucocorticoid receptor gene is associated with a decrease in the sensitivity of immune cells to glucocorticoid hormones that are involved in terminating inflammatory processes and may be a mechanism underlying increased risk for adrenal insufficiency.
Institutionalization	Drury et al., 2012 ³	Longitudinal follow-up of 64 children enrolled in the Bucharest Early Intervention Project ⁴ including 28 children who remained in the institution and 36 children who were placed in foster care.	Buccal cells acquired once in children ranging in age for 6–10 years	Percent of time spent in institutional care was significantly associated with shorter telomere length. Association remained significant when controlling for study group (foster care or institutionalized), gender, ethnicity, low birth weight, and age at DNA sampling.	Decreased Telomere length is a marker for biological aging and is associated with cardiovascular disease, diabetes, obesity, and health risks related to early adversity.
	Humphreys et al., in press	Longitudinal follow-up of 79 children enrolled in the Bucharest Early Intervention Project ⁴ comparing 50 children with a history of institutionalization to 29 never institutionalized children.	Buccal cells acquired repeatedly, (2–4 times) from ages 8–14 years.	Telomere shortening was significantly accelerated over time in children with a history of institutionalization relative to never institutionalized children.	

¹ Provenzi L, Fumagalli M, Sirgiovanni I, et al. Pain-related stress during the Neonatal Intensive Care Unit stay and *SLC6A4* methylation in very preterm infants. *Frontiers in Behavioral Neuroscience*. 2015;9;.

² Kantake M, Yoshitake H, Ishikawa H, Araki Y, Shimizu T. Postnatal epigenetic modification of glucocorticoid receptor gene in preterm infants: a prospective cohort study. *BMJ Open*. 2014;4(7):8;.

³ Drury SS, Theall K, Gleason MM, et al. Telomere length and early severe social deprivation: linking early adversity and cellular aging. *Molecular psychiatry*. Jul 2012;17(7):719–727;.

⁴ Zeanah CH, Nelson CA, Fox NA, et al. Designing research to study the effects of institutionalization on brain and behavioral development: The Bucharest Early Intervention Project. *Development and Psychopathology*. Fall 2003 2003;15(4):885–907.

⁴Humphreys, K.L., Esteves, K., Zeanah, C.H., Fox, N.A., Nelson, C.A., Drury, S.S. Accelerated telomere shortening: Tracking the lasting impact of early institutional care at the cellular level (Under Review).