Published in final edited form as:

J Hosp Med. 2016 January ; 11(1): 48–51. doi:10.1002/jhm.2446.

Refractory Primary Medication Non-Adherence: Prevalence and Predictors After Pharmacist Counseling at Hospital Discharge

Kathleene Wooldridge, MD^{1,2}, Jeffrey L. Schnipper, MD, MPH^{3,4,5}, Kathryn Goggins, MPH^{6,7}, Robert S. Dittus, MD, MPH^{1,8,9}, and Sunil Kripalani, MD, MSc^{2,6,7}

¹Geriatric Research, Education, and Clinical Center, Veterans Affairs Tennessee Valley Healthcare System, Nashville, TN

²Section of Hospital Medicine, Division of General Internal Medicine and Public Health, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN

³Brigham and Women's Hospital Division of General Medicine and Primary Care, Boston, MA

⁴Brigham and Women's Hospital Hospitalist Service, Boston, MA

⁵Harvard Medical School, Boston, MA

⁶Center for Clinical Quality and Implementation Research, Vanderbilt University

⁷Center for Health Services Research, Vanderbilt University Medical Center, Nashville, TN

⁸Division of General Internal Medicine and Public Health, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN

⁹Institute for Medicine and Public Health, Vanderbilt University Medical Center, Nashville, TN

Abstract

Successful secondary prevention of cardiovascular disease relies on medication therapy, thus minimizing non-adherence is a focus for improving patient outcomes. Receipt of discharge medication counseling has been associated with improved drug knowledge and adherence. We evaluated the prevalence and predictors of post-discharge primary non-adherence (not filling new prescriptions) in patients who received discharge medication counseling by a pharmacist (i.e., refractory to intervention) as part of a randomized, controlled trial. Of 341 patients, 9.4% of patients did not fill all prescriptions after discharge. Patients who were living alone were more

Address for Correspondence: Kathleene Wooldridge, MD, Section of Hospital Medicine, Division of General Internal Medicine and Public Health, Department of Medicine, Vanderbilt University, 1215 21st Ave S, Suite 6000 Medical Center East, Nashville 37232, TN, USA. Phone: (615) 936-8219 Fax: (615) 936-1269, kathleene.t.wooldridge@vanderbilt.edu.

Prior Presentations: A preliminary version of this research was presented at the AcademyHealth Annual Research Meeting, June 16, 2015, Minneapolis, MN.

Potential Conflicts of Interest:

Kathleene Wooldridge: None

Jeffrey Schnipper: PI, investigator-initiated study funded by Sanofi-Aventis to develop, implement, and evaluate a multi-faceted intervention to improve transitions of care in patients with diabetes mellitus discharged on insulin

Kathryn Goggins: None

Robert Dittus: Passive co-owner, Medical Decision Modeling, Inc.; Bayer HealthCare. One day consultation and panelist on educational video for population health. (consultant fee); GlaxoSmithKline. One day consultant for population health, envisioning the future. (consultant fee)

Sunil Kripalani: Bioscape Digital, stock ownership

likely to not fill their medications compared to those who were married or cohabitating (odds ratio [OR] 2.2, 95% confidence interval [CI], 1.01-4.8, $p=0.047$). Patients who were discharged with greater than 10 medications were also more likely to demonstrate primary non-adherence (OR 2.3, 95% CI, 1.05-4.98, $p=0.036$). Patients with lower income were less likely to fill prescriptions in univariate analysis ($p=0.04$) but not multivariable analysis. Our study demonstrates that among patients hospitalized for acute cardiovascular events, primary medication non-adherence persisted despite discharge medication counseling. Targeted or multimodal approaches that address patient-specific barriers such as cost, social isolation, and polypharmacy, in addition to discharge counseling, may further facilitate adherence.

Introduction

Medication non-adherence after hospital discharge impacts morbidity and mortality in patients with cardiovascular disease.¹ Primary non-adherence, part of the spectrum of medication underuse, occurs when a patient receives a prescription but does not fill it.¹ Prior studies utilizing retrospective administrative data have found a prevalence of post-discharge primary non-adherence between 24-28%,^{1,2} similar to findings in a variety of outpatient populations.^{3,4}

One strategy for reduction in non-adherence is discharge medication counseling, which has been associated with improved post-discharge outcomes.¹ We evaluated the prevalence and predictors of refractory primary non-adherence in a cohort of patients hospitalized for acute cardiovascular conditions who received pharmacist counseling prior to discharge in order to guide future adherence interventions.

Methods

Setting & Participants

The present study represents a secondary analysis of data from the Pharmacist Intervention for Low Literacy in Cardiovascular Disease (PILL-CVD) study. PILL-CVD was a randomized, controlled trial that evaluated the effect of a tailored intervention consisting of pharmacist-assisted medication reconciliation, discharge counseling, low-literacy adherence aids, and follow-up phone calls in adults hospitalized for acute coronary syndromes or acute decompensated heart failure. Patients likely to be discharged home taking primary responsibility for their medication management were eligible. Full study methods and results, including inclusion and exclusion criteria, can be found elsewhere.⁵ The Institutional Review Boards of each site approved the study.

For the present analysis, patients were included if they had any new discharge prescriptions to fill and received the study intervention, including a post-discharge follow-up phone call with questions about filling discharge prescriptions.

Baseline Measures

Baseline data were obtained from medical records and patient interviews, including demographic information as well as survey data for cognitive impairment (Mini-Cog) and health literacy (s-TOFHLA).^{6,7}

Data were also collected related to medication use, including the number of scheduled and as-needed medications listed at discharge, self-reported pre-admission adherence, medication understanding, and medication management practices (e.g., use of a pillbox, refill reminders). Self-reported medication adherence was measured with the 4-item Morisky scale.⁸ Medication understanding was assessed with a tool previously developed by Marvanova et al.⁹

Outcome Measures

The primary outcome was the percentage of patients who reported not filling at least one discharge prescription on a telephone call that was conducted 1-4 days post-discharge. Patients were asked a dichotomous question about whether or not they filled all of their discharge prescriptions. Further characterization of the class or number of medications not filled was not performed. Patients were asked to provide a reason for not filling the prescriptions.

Analysis

We evaluated the prevalence and possible predictors of primary non-adherence including age, gender, race, marital status, education and income levels, insurance type, health literacy, cognition, presence of a primary care physician, number of listed discharge medications, pre-hospital medication adherence, medication understanding, and medication management practices using Pearson chi-square, Fisher's exact, or Wilcoxon rank sum tests as appropriate. Multiple logistic regression with backward elimination was performed to identify independent predictors, selected with p -values less than 0.1. We also evaluated reasons that patients cited for not filling prescriptions. Two-sided p -values <0.05 were considered statistically significant. All analyses were conducted using Stata, version 13.1 (StataCorp LP, College Station, TX).

Results

Of 851 patients in the PILL-CVD study, the present sample includes 341 patients who received the intervention, completed the post-discharge follow-up call, and had new discharge prescriptions to be filled. This represents 85% of patients who received the intervention.

The mean age of participants was 61.3 years, and 59.5% were male (Table 1). The majority was white (75.1%), and 88% had at least a high school education. Married or cohabitating patients represented 54.3% of the group. Just over half of the patients (54%) had an income of \$35K or greater. The primary source of insurance for 82.5% of patients was either Medicare or private insurance, and 7.4% of patients were self-pay. Most patients (80%) had adequate health literacy. The median Mini-Cog Score was 4 out of 5 (IQR=3-5), and 11% of

patients had scores indicating cognitive impairment. Just less than one-fourth of the patients (24.1%) had a Morisky Score of 8, indicating high self-reported adherence, and the median score of patients' understanding of medications (range of 0-3) was 2.5 (IQR=2.2-2.8), reflecting relatively high understanding. The median number of prescriptions on patients' discharge medications lists was 10 (IQR=8-13).

The prevalence of refractory primary non-adherence was 9.4%. In univariate analysis, single marital status, lower income, and having more than 10 total discharge medications were significantly associated with not filling medications ($p=0.018$, 0.04 , 0.016 ; Table 1). In multivariable analysis, single marital status and having more than 10 total discharge medications maintained significance when controlling for other patient characteristics. Patients who were single had higher odds of failing to fill discharge prescriptions compared to married or cohabitating individuals (odds ratio [OR] 2.2, 95% confidence interval [CI], 1.01-4.8, $p=0.047$). Patients with more than 10 discharge medications also had higher odds of failing to fill compared with patients who had fewer total medications (OR 2.3, 95% CI, 1.05-4.98, $p=0.036$).

Filling discharge prescriptions was not associated with health literacy, cognition, pre-hospital adherence, patients' medication understanding, or any of the surveyed medication management practices (Table 2). Patients' reasons for not filling included lack of time to go to the pharmacy, medications not being delivered or dispensed, or inability to afford prescriptions. Prescription cost was cited by 23.5% of patients who did not fill their prescriptions and provided a reason.

Discussion

Almost 1 in 10 patients hospitalized with cardiovascular disease demonstrated primary non-adherence refractory to an intervention including pharmacist discharge medication counseling. Being unmarried and having greater than 10 medications at discharge were significantly associated with higher primary non-adherence when controlling for other patient factors.

Patients with a co-habitant partner were significantly less likely to exhibit primary non-adherence, which may reflect higher levels of social support, including encouragement for disease self-management and/or support with tasks such as picking up medications from the pharmacy. Previous research has demonstrated that social support mediates outpatient medication adherence for heart failure patients.¹⁰

Similar to Jackevicius et al, we found that patients with more medications at discharge were less likely to fill their prescriptions.¹ These findings may reflect the challenges that patients face in adhering to complex treatment plans, which are associated with increased coordination and cost. Conversely, some prior studies have found that patients with fewer prescriptions were less likely to fill.^{11,12} These patients were often younger, thus potentially less conditioned to fill prescriptions, and unlike our cohort, these populations had consistent prescription coverage. Interventions for polypharmacy, which have been shown to improve

outcomes and decrease costs, especially in the geriatric population, may be of benefit for primary non-adherence as well.¹³

Additionally, patients with lower household incomes had higher rates of primary non-adherence, at least in univariate analysis. Medication cost and transportation limitations, which are more pronounced in lower income patients, likely play influential roles in this group. These findings build on prior literature that has found lower prescription cost to be associated with better medication adherence in a variety of settings.^{3,4,14}

Because the prevalence of primary non-adherence in this cohort is less than half of historical rates, we suspect the intervention did reduce unintentional non-adherence. However, regimen cost and complexity, transportation challenges, and ingrained medication beliefs likely remained barriers. It may be that a post-discharge phone call is able address unintended primary non-adherence in many cases. “Meds to beds” programs, where a supply of medications is provided to patients prior to discharge, could assist patients with limited transportation. Prior studies have also found reduced primary non-adherence when e-prescriptions are utilized.³

Establishing outpatient follow-up at discharge provides additional opportunities to address unanticipated adherence barriers. Since the efficacy of any adherence intervention depends on individual patient barriers, we recommend combining medication counseling with a targeted approach for patient-specific needs.

We note several limitations to our study. First, because we studied primary non-adherence that persisted despite an intervention, this cohort likely underestimates the prevalence of primary non-adherence and alters the associated patient characteristics found in routine practice (although counseling is becoming more common). Second, patient report is subject to biases that underestimate non-adherence, although this approach has been validated previously.¹⁵ Third, our outcome measure was unable to capture the spectrum of non-adherence that could provide a more nuanced look at predictors of post-discharge non-adherence. Fourth, we did not have patient copayment data to better characterize whether out of pocket costs or pharmacologic classes drove non-adherence. Finally, sample size may have limited the detection of other important factors, and the university setting may limit generalizability to cardiovascular patients in other practice environments. Future research should focus on intervention strategies that assess patients’ individual adherence barriers for a targeted or multimodal approach to improve adherence.

In conclusion, we found a prevalence of primary non-adherence of almost 1 in 10 patients who received pharmacist counseling. Non-adherence was associated with being single and those discharged with longer medication lists. Our results support existing literature that primary non-adherence is a significant problem in the post-discharge setting and substantiate the need for ongoing efforts to study and implement interventions for adherence after hospital discharge.

Acknowledgments

Funders: This material is based on work supported by the Office of Academic Affiliations, Department of Veterans Affairs, Veterans Affairs National Quality Scholars Program and with use of facilities at Veterans Affairs Tennessee Valley Healthcare System, Nashville (Dr. Wooldridge). The funding agency supported the work indirectly through provision of salary support and training for the primary author, but had no specific role in the design and conduct of the study, collection, management, analysis, and interpretation of the data, preparation, review, or approval of the manuscript, or decision to submit the manuscript for publication. This work was also supported by R01 HL089755 (Dr. Kripalani) and in part by K23 HL077597 (Dr. Kripalani), K08 HL072806 (Dr. Schnipper), and the Center for Clinical Quality and Implementation Research at Vanderbilt University Medical Center.

References

1. Jackevicius CA, Li P, Tu JV. Prevalence, predictors, and outcomes of primary nonadherence after acute myocardial infarction. *Circulation*. 2008; 117(8):1028–1036. [PubMed: 18299512]
2. Fallis BA, Dhalla IA, Klemensberg J, Bell CM. Primary medication non-adherence after discharge from a general internal medicine service. *PloS one*. 2013; 8(5):e61735. [PubMed: 23658698]
3. Fischer MA, Choudhry NK, Brill G, et al. Trouble getting started: predictors of primary medication nonadherence. *The American journal of medicine*. 2011; 124(11):1081.e1089–1022. [PubMed: 22017787]
4. Tamblyn R, Egale T, Huang A, Winslade N, Doran P. The incidence and determinants of primary nonadherence with prescribed medication in primary care: a cohort study. *Annals of internal medicine*. 2014; 160(7):441–450. [PubMed: 24687067]
5. Kripalani S, Roumie CL, Dalal AK, et al. Effect of a pharmacist intervention on clinically important medication errors after hospital discharge: a randomized trial. *Annals of internal medicine*. 2012; 157(1):1–10. [PubMed: 22751755]
6. Nurss, J.; Parker, R.; Williams, M.; Baker, D. Short test of Functional health literacy in adults. Peppercorn Books and Press; Snow Camp, NC: 1998.
7. Borson S, Scanlan JM, Watanabe J, Tu SP, Lessig M. Simplifying detection of cognitive impairment: comparison of the Mini-Cog and Mini-Mental State Examination in a multiethnic sample. *Journal of the American Geriatrics Society*. 2005; 53(5):871–874. [PubMed: 15877567]
8. Morisky DE, Green LW, Levine DM. Concurrent and predictive validity of a self-reported measure of medication adherence. *Medical care*. 1986; 24(1):67–74. [PubMed: 3945130]
9. Marvanova M, Roumie CL, Eden SK, Cawthon C, Schnipper JL, Kripalani S. Health literacy and medication understanding among hospitalized adults. *Journal of hospital medicine : an official publication of the Society of Hospital Medicine*. 2011; 6(9):488–493.
10. Wu JR, Frazier SK, Rayens MK, Lennie TA, Chung ML, Moser DK. Medication adherence, social support, and event-free survival in patients with heart failure. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 2013; 32(6):637–646.
11. Lagu T, Weiner MG, Eachus S, Tang SS, Schwartz JS, Turner BJ. Effect of patient comorbidities on filling of antihypertensive prescriptions. *The American journal of managed care*. 2009; 15(1): 24–30. [PubMed: 19146361]
12. Cheetham TC, Niu F, Green K, et al. Primary nonadherence to statin medications in a managed care organization. *Journal of managed care pharmacy : JMCP*. 2013; 19(5):367–373. [PubMed: 23697474]
13. Kojima G, Bell C, Tamura B, et al. Reducing cost by reducing polypharmacy: the polypharmacy outcomes project. *Journal of the American Medical Directors Association*. 2012; 13(9):818 e811–815. [PubMed: 22959733]
14. Shrank WH, Choudhry NK, Fischer MA, et al. The epidemiology of prescriptions abandoned at the pharmacy. *Annals of internal medicine*. 2010; 153(10):633–640. [PubMed: 21079218]
15. Haynes RB, Taylor DW, Sackett DL, Gibson ES, Bernholz CD, Mukherjee J. Can simple clinical measurements detect patient noncompliance? *Hypertension*. 1980; 2(6):757–764. [PubMed: 7007235]

Table 1Patient Characteristics[†]

Variable	Overall 341 (100.0%)	Filled Prescription 309 (90.6%)	Did Not Fill 32 (9.4%)	p-value
Age, years (N%)	69	63 (91.3)	6 (8.7)	0.745 [†]
18-49	128	114 (89.1)	14 (10.9)	
50-64	144	132 (91.7)	12 (8.3)	
65+				
Gender (N%)	203	189 (93.1)	14 (6.9)	0.056 [†]
Male	138	120 (87.0)	18 (13.0)	
Female				
Race (N%)	256	234 (91.4)	22 (8.6)	0.712 [†]
White	60	54 (90.0)	6 (10.0)	
African American	22	19 (86.4)	3 (13.6)	
Other				
Education (N%)	40	32 (80.0)	8 (20.0)	0.054 [†]
Less than high school	99	91 (91.9)	8 (8.1)	
High school	93	83 (89.2)	10 (10.8)	
13-15 years	109	103 (94.5)	6 (5.5)	
16 years				
Marital Status (N%)	156	135 (86.5)	21 (13.5)	0.018 ^{†*}
Separated/divorced/widowed/never married	185	174 (94.1)	11 (5.9)	
Married/cohabitating				
Income (N%)	58	48 (82.8)	10 (17.2)	0.040 ^{†*}
< 10K - <20K	86	76 (88.4)	10 (11.6)	
20K - 35K	40	36 (90.0)	4 (10.0)	
35K - < 50K	46	43 (93.5)	3 (6.5)	
50K - < 75K	83	81 (97.6)	2 (2.4)	
75K+				
Primary source of payment (N%)	34	28 (82.4)	6 (17.6)	0.272 [†]
Medicaid	145	131 (90.3)	14 (9.7)	
Medicare	132	123 (93.2)	9 (6.8)	
Private	25	22 (88.0)	3 (12.0)	
Self-pay				
Primary care physician (N%)	28	26 (92.9)	2 (7.1)	1.000 [‡]
None/Do not know	313	283 (90.4)	30 (9.6)	
Yes				
Site (N%)	172	151 (87.8)	21 (12.2)	0.071 [†]
Nashville, TN	169	158 (93.5)	11 (6.5)	
Boston, MA				

Note: Missing values are present in the following categories: race (n=3), income (n=28), insurance (n=5)

* p<0.05,

[†] Pearson Chi-Square test,

[‡] Fisher's exact test (2-sided)

Table 2

Patient Medication-Related Characteristics

Variable	Overall 341 (100.0%)	Filled Prescription 309 (90.6%)	Did Not Fill 32 (9.4%)	p- value
s-TOFHLA Score (range 0-36) (N%)	40	34 (85.0)	6 (15.0)	0.443 [†]
Inadequate (0-16)	27	25 (92.6)	2 (7.4)	
Marginal (17-22)	268	244 (91.0)	24 (9.0)	
Adequate (23-36)				
MiniCog Score (range 0-5) (N%)	304	276 (90.8)	28 (9.2)	0.764 [‡]
Not impaired (3-5)	37	33 (89.2)	4 (10.8)	
Impaired (0-2)				
Morisky Score (range 4-8) (N%)	249	224 (90.0)	25 (10.0)	0.517 [†]
Low/Moderate Self-Reported Adherence (4-7)	79	73 (92.4)	6 (7.6)	
High Self-Reported Adherence (8)				
Number of Discharge Medications [§] (range 1-26) (N%)	186	175 (94.1)	11 (5.9)	0.016 [†]
0-10 medications	155	134 (86.5)	21 (13.5)	
11+ medications				
Patient Responses to Medication Behavior Questions				
Patient associates medication taking time with daily events	253	229 (90.5)	24 (9.5)	0.913 [†]
Patient uses a pillbox to organize medicine	180	162 (90.0)	18 (10.0)	0.680 [†]
Friends or family help remind patient when it is time to take medicine	89	79 (88.8)	10 (11.2)	0.486 [†]
Patient writes down instructions for when to take medicine	60	55 (91.7)	5 (8.3)	0.758 [†]
Patient uses an alarm or a reminder that beeps when it is time to take medicine	8	6 (75.0)	2 (25.0)	0.167 [†]
Patient marks refill date on calendar	38	35 (92.1)	3 (7.9)	1.000 [‡]
Pharmacy gives or sends patient a reminder when it is time to refill medicine	94	84 (89.4)	10 (10.6)	0.624 [†]
Friends or family help patient to refill medicine	60	53 (88.3)	7 (11.7)	0.504 [†]

Note: Missing values are present in the following categories: s-TOFHLA (n=6), Morisky (n=13)

[§]Number on discharge medication list

[†]Pearson Chi-Square test,

[‡]Fisher's exact test (2-sided)