Published in final edited form as:

Supramol Chem. 2015 ; 27(10): 724–730. doi:10.1080/10610278.2015.1078898.

Pb, Sr and Ba calix[6]arene hexacarboxylic acid octahedral complexation: a dramatic effect of dealkylation

Birendra Babu Adhikari^a, Cuong-Alexander To^a, Tetsuo Iwasawa^b, and Michael P. Schramm^{a,*}

^aDepartment of Chemistry and Biochemistry, California State University Long Beach, Long Beach, CA, USA

^bDepartment of Materials Chemistry, Faculty of Science and Technology, Ryukoku University, Otsu, Shiga 520-2194, Japan

Abstract

Calix[6]arene hexacarboxylic acid binds instantly and with low symmetry to Pb, Sr and Ba. Later a highly symmetric up-down alternating conformation emerges. The solution structures are identical to their p-tert-butylcalix[6]arene hexacarboxylic acid counterparts. With either receptor an octahedral cage is formed around the metal. The transformation from low to high symmetry however proceeds at significantly faster rates for the de-t-butylated host.

Introduction

t-Butyl calix[6]arene hexaesters and hexaamides have enhanced selectivity towards the extraction and transport of a variety of ions including Cs(I) and Sr(II).^{1–5} Pb(II) is effectively transported by the hexaester.^{6, 7} t-Butyl calix[6]arene hexathioamides⁸ have been shown to be efficient towards Cu(II)⁹, Pb(II)⁷ and Ag(I).¹⁰ Variation in selectivity changed when several of these compounds lacked p-tert-butyl groups – and in many cases binding affinity was drastically enhanced.^{11, 12} Calix[n]arene phosphine oxides¹³ have also demonstrated enhanced binding affinity for Th(IV).¹⁴ Indeed work with calix[6]arenes towards nuclear waste separation remains an ongoing area of active research.¹⁵

Very recently we reported on the complexation of Pb(II), Sr(II) and Ba(II) with p-tert-butylcalix[6]arene hexacarboxylic acid **1** (figure 1).^{16, 17} In these reports we found that calixarene hexacarboxylic acid binds rapidly with all three metals and form low symmetry “lozenge” shaped complexes.¹⁸ These exhibit complex NMR signals with multiple types of methylene bridges for both the ether as well as the aryl linkages. Upon standing something remarkable occurred – all three metal-host complexes rearrange such that the individual phenyl rings alternate up-down so that the carboxylates form an octahedral cage with C_{3v} symmetry around the metal center. NMR and X-ray measurements for all three complexes

Department of Chemistry and Biochemistry, California State University Long Beach, 1250 Bellflower Blvd., Long Beach, 90840, CA, USA. Fax: (+)1-562-985-8557; michael.schramm@csulb.edu.

The content is solely the responsibility of the authors and does not necessarily represent the views of the Japan Society for the Promotion of Science, the National Cancer Institute, the National Institutes of Health, or NSF.

confirm the solution and solid-state structure of the host. The structures we reported have some semblance to valinomycin•K¹⁹ or Cram's sphereands.²⁰ With these metals – differences in time to rearrange were apparent. We demonstrated that several analogs of calixarene hexacarboxylic acid, such as those where acid was replaced with ether, di-ether, or ester resulted in initial low symmetry binding, but only those hosts with acid functional groups resulted in an octahedral cage for the metal. This behavior was attenuated when amine bases were added, presumably due to ion capping.²¹ The use of other bases to date has not had a marked impact on the behavior of these host-guest complexes, although our studies have not been exhaustive. A variety of other metals were tested and Pb, Sr, and Ba are a short list of metals that this host is selective for accommodation in an octahedral environment.

The remaining variable of interest to us was the effect of the t-butyl group.²² Is it possible that the t-butyl group has an influence on either binding or the rate of conversion from a low to high symmetry complex for these three metals?

Results and Discussion

Calix[6]arene hexacarboxylic acid **2** is a known compound.²¹ We prepared **2** through removal of *tert*-butyl group of *p*-*tert*-butylcalix[6]arene by adopting a procedure reported for *p*-*tert*-butylcalix[4]arene.²³ Alkylation and saponification as for **1**¹⁶ gave **2** (figure 2).

When **2** was mixed with 1 equivalent of Pb(ClO₄)₂•3H₂O in CDCl₃/CD₃CN immediate formation of one major species was obvious (figure 3), but some very small peaks that can be attributed to low-symmetry “lozenge” shaped species co-exist. After 30 minutes however the sample appears to be at equilibrium with a high symmetry geometry that is identical in pattern to the analogous *p*-*tert*-butyl host (figure 4).¹⁶ The complex remains soluble in a mixture chloroform/acetonitrile and is highly likely to be neutral. The exact location of all protons isn't known, but we surmise that 2 negative charges and 4 protons are distributed amongst the 6 carboxylates and are interchanging - indeed the symmetry of the solution state structure supports this assumption as well as exact mass measurements (See E.S.I.). The two sets of diastereotopic methylenes are split drastically. For the bridging methylene this isn't surprising, in a cone conformation calix[n]arenes place one H “down” and the other “out.” In our alternating up-down conformation this relationship remains. The second methylene group, a phenolic ether also undergoes drastic splitting; for **1**•Pb¹⁶ this result was explained with NOESY and X-ray data, one hydrogen points towards an adjacent aromatic proton, the other away. The same is now true for **2**•Pb(II) (See ESI for COSY, NOESY, HSQC spectra). In the instance of **2**, however the transformation is complete in 30 minutes, previously with the more hindered host **1**, the transformation takes approximately 15 hours.

Given the drastic reduction in time for the formation of the octahedral complex of **2**•Pb (1 hour) vs. **1**•Pb (15 hours), we anticipated that the rearrangement of **2**•Sr would also proceed more quickly than its *t*-butyl analog **1**•Sr. **1**•Sr took approximately 20 days to come to equilibrium. Sr(ClO₄)₂•3H₂O was again slow to entice host **2** into octahedral geometry, upon mixing several low concentration species are present (Figure 5) and several major broad peaks emerge. From 48 hours to 72 hours (latter not shown) the signals remain poorly

resolved and broad, but at 115 hours baseline resolution is achieved and we ascribe this to be an approximate time to arrive at equilibrium. A small peak at 7.0 ppm remains that is part of the free ligand **2**.

Finally, we conducted the same experiment with $\text{Ba}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$. Many sharp signals exist for **2**•Ba at the time of mixing, consistent with a low symmetry lozenge (figure 6). At 48 hours after mixing the sample is at equilibrium. This last result is most surprising because the high symmetry **1**•Ba sample took 38 days of time to come to equilibrium, both Pb and Sr were faster with host **1**. These results are summarized in Table 1 and we see that the t-butyl group of **1** has a marked effect on the time it takes complex **1** vs. **2** to arrive at equilibrium. In all cases equilibrium is arrived at an order of magnitude sooner. Surprisingly though, the behavior is not consistent – Ba and Sr switch their ranking as a function of host, always slower than Pb. It remains unclear why Ba and Sr don't have the same relative order of time to reach equilibrium.

We added 10% vol/vol of D_2O to test the stability of the complex and found that they rapidly degraded. First observations of NMR show a series of broadened peaks that promptly disappear (See ESI). No evidence of NMR signals could be found when **2** was directly mixed with Pb, Sr, or Ba in pure water.

Given the drastic change in the time it takes host **2** to rearrange, we conducted an initial screening program to uncover other metals that might be accommodated in this octahedral hole. As was the case with **1**, **2** fails to show evidence of complex formation with $\text{Ca}(\text{II})$, $\text{Y}(\text{III})$, $\text{Bi}(\text{III})$, or $\text{Gd}(\text{III})$ (See ESI for details). While new complexes weren't found it is clear that **2** remains a selective host.

Given the time differences to arrive at equilibrium, we conducted a competitive binding experiment using a 1:1:1 ratio of **1**, **2**, and $\text{Pb}(\text{II})$ (Figure 7). At the initial time of mixing, the **2**•Pb complex is easily identified as the predominant complex. Traces of **1**•Pb and the low symmetry complex of **1**•Pb can be identified. At the 16 hour mark, the two high symmetry complexes are present and the t-butyl **1**•Pb host is the major component. Surprisingly after 115 hours evidence of a roughly 1:1 mixture of **1**•Pb:**2**•Pb was apparent. Complex **2**•Pb more quickly converts from low to high symmetry, the dynamics of the system remain under investigation. Perhaps solubility differences are at work after more than 16 hours of resting while the system reverts to 1:1.

Next we performed a competitive binding experiment between host **2**, Pb and Ba. Mixing the three components in a 1:1:1 molar ratio, we see rapid formation of the high-symmetry Pb complex after 4 hours – the Pb complex is readily identified. After 115 hours, the Ba complex surpasses the concentration of Pb (Figure 8).

Host **1** and **2** now both serve as calix[6]arene hosts that display a remarkable conformational change when binding with Pb, Sr and Ba. In all combinations – initial low symmetry species are identified by NMR and given enough time high symmetry octahedral complexes are formed. These complexes display a unique alternating up-down arrangement of calix[6]arene phenyl groups so as to surround the metal center with 6 carboxylate groups.

We have clearly demonstrated the effect of removing the t-butyl groups of **1**; by using **2** the time to reorganization decreases by an order of magnitude (from weeks to days, or 15 hours to 30 minutes). The relative rates to equilibrium however change with differential behavior for Sr and Ba with **1** and **2**. Additionally, initial competitive binding experiments reveal some new details about stability and dynamics. Other combinations are currently being explored to thoroughly evaluate the competitive binding affinity and kinetics of Pb, Sr and Ba for both hosts **1** and **2**. At the end of these studies we hope to develop materials suitable for deployment towards the separation of Pb or Sr from complex waste streams.

Experimental

All spectra described above are provided in the E.S.I. along with complete characterization of all new compounds, some experimental procedures are cited above.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Japan Society of the Promotion of Science for an Invitational Fellowship (Long-term, L-15528, M.P.S.). The project described was supported by Grant Number SC2CA167636 from the National Cancer Institute. NMR instrumentation was provided for by the National Science Foundation (MRI CHE-1337559).

References

- [General perspectives] Lehn J-M. Supramolecular Chemistry: Concepts and Perspectives. VCH Weinheim, Germany 1995; Steed JW, Atwood JL. Supramolecular Chemistry: An Introduction. Wiley Chichester, U.K. 2000; Gokel GW. Molecular Recognition, Receptors for Cationic Guests. Comprehensive Supramolecular Chemistry. Lehn J-M, Atwood JL, Davies JED, MacNicol DD, Vogtle F. Pergamon Oxford, U.K. 1996; 1
- [General for cation recognition] Leray I, Valeur B. Eur J Org Chem. 2009; 3525–3535. Leray I, Valeur B. Coord Chem Rev. 2000; 205:3–40. de Silva AP, Gunaratne HQN, Gunlaugsson T, Huxley AJM, McCoy CP, Rademacher JT, Rice TE. Chem Rev. 1997; 97:1515–1566. [PubMed: 11851458] Calixarenes 2001. Asfari Z, Böhmer V, Harrowfield J, Vicens J. Kluwer Academic Publishers Dordrecht, The Netherlands 2001; Nendza M, Herbst T, Kussatz C, Gies A. Chemosphere. 1997; 35:1875–1885. [PubMed: 9353909]
- [Calixarenes for molecular recognition] Gutsche CD. Monographs in Supramolecular Chemistry, vol. 6, Calixarenes Revisited. The Royal Society of Chemistry Cambridge, U.K. 1998; Kim JS, Quang DT. Chem Rev. 2007; 107:3780–3799. [PubMed: 17711335] Ikeda A, Shinkai S. Chem Rev. 1997; 97:1713–1734. [PubMed: 11851464] Böhmer V. Angew Chem, Int Ed. 1995; 34:713–745. Valeur B, Leray I. Inorg Chim Acta. 2007; 360:765–774. Nolan EM, Lippard SJ. Chem Rev. 2008; 108:3443–3480. [PubMed: 18652512]
- [Calixarene-based sensing for Cs(I)] Kim SK, Vargas-Zuñiga GI, Hay BP, Young NJ, Delmau LH, Masselin C, Lee CH, Kim JS, Moyer BA, Lynch VM, Sessler JL. J Am Chem Soc. 2012; 134:1782–1792. [PubMed: 22191574] Casnati A, Pochini A, Ungaro R, Ugozzoli F, Arnaud F, Fanni S, Schwing MJ, Egberink RJM, de Jong F, Reinhoudt DN. J Am Chem Soc. 1995; 117:2767–2777. Ji HF, Dabestani R, Brown GM, Sachleben RA. Chem Commun. 2000:833–834. Ji HF, Dabestani R, Brown GM. J Am Chem Soc. 2000; 122:9306–9307. Talanova GG, Roper ED, Buie NM, Gorbunova MG, Bartsch RA, Talanov VS. Chem Commun. 2005:5673–5675. Roper ED, Talanov VS, Gorbunova MG, Bartsch RA, Talanova GG. Anal Chem. 2007; 79:1983–1989. [PubMed: 17266283] Arnaud-Neu F, Fanni S, Guerra L, McGregor W, Ziat K, Schwing-Weill MJ, Barrett G, McKervey MA, Marrs D, Seward EM. J Chem Soc, Perkin Trans. 1995; 2:113.

5. [Calixarene-based sensing for Sr(II)] Kim TH, Kim H, Lee JH, Kim JS. Bull Korean Chem Soc. 2008; 29:620–622. Casnati A, Baldini L, Pelizzi N, Rissanen K, Ugozzoli F, Ungaro R. J Chem Soc, Dalton Trans. 2000:3411–3415.
6. Minhas FT, Solangi IB, Memon S, Bhanger MI. Sep Sci Technol. 2014; 45:1448.
7. [Calixarene-based sensing for Pb(II)] Talanova GG, Hwang HS, Talanov VS, Bartsch RA. Chem Commun. 1998:419–420. Bok JH, Kim HJ, Lee JW, Kim SK, Choi JK, Vicens J, Kim JS. Tetrahedron Lett. 2006; 47:1237–1240. Metivier R, Leray I, Valeur B. Chem Commun. 2003:996–997. Buie NM, Talanov VS, Butcher RJ, Talanova GG. Inorg Chem. 2008; 47:3549–3558. [PubMed: 18345614] Liu JM, Bu JH, Zheng QY, Chen CF, Huang ZT. Tetrahedron Lett. 2006; 47:1905. Kim SK, Kim SH, Kim HJ, Lee SH, Lee SW, Ko J, Bartsch RA, Kim JS. Inorg Chem. 2005; 44:7866–7875. [PubMed: 16241136] Lee SH, Kim SK, Bok JH, Lee SH, Yoon J, Lee K, Kim JS. Tetrahedron Lett. 2005; 46:8163–8167. Choi JK, Lee A, Kim S, Ham S, No K, Kim JS. Org Lett. 2006; 8:1601–1604. [PubMed: 16597120]
8. [Calixarene-hexathioamide] Malinowska E, Brzozka Z, Kasiura K, Egberink RJM, Reinhoudt DN. Anal Chim Acta. 1994; 298:253–258. Arnaud-Neu F, Barrett G, Corry D, Cremin S, Ferguson G, Gallagher FJ, Harris JS, Anthony McKervey M, Schwing-Weill M-J. J Chem Soc, Perkin Trans. 1997; 2:575. Genorio B, He T, Meden A, Polanc S, Jamnik J, Tour JM. Langmuir. 2008; 24:11523–11532. [PubMed: 18816014] Wroblewski W, Brzozka Z. Anal Chim Acta. 1996; 326:163–168. Torma F, Gruen A, Bitter I, Toth K. Electroanalysis. 2009; 21:1961–1969. Bochenska M, Kulesza J, Chojnacki J, Arnaud-Neu F, Hubscher-Bruder V. J Inclusion Phenom Macrocyclic Chem. 2010; 68:75–83.
9. [Calixarene-based sensing for Cu(II)] Qazi MA, Ocak U, Ocak M, Memon S. Anal Chim Acta. 2013; 761:157–168. [PubMed: 23312327] Kim HJ, Kim SH, Kim JH, Anh LN, Lee JH, Lee CH, Kim JS. Tetrahedron Lett. 2009; 50:2782–2786. Ho IT, Chu JH, Chung WS. Eur J Org Chem. 2011:1472–1481. Kumar M, Babu JN, Bhalla V. J Inclusion Phenom Macrocyclic Chem. 2010; 66:139–145.
10. [Calixarene-based sensing for Ag(I)] Ki JS, Shon OJ, Rim JA, Kim SK, Yoon J. J Org Chem. 2002; 67:2348–2351. [PubMed: 11925252] Elcin S, Deligoz H. Tetrahedron. 2013; 69:6832–6838. Yafian MR, Burgard M, El-Bachiri A, Matt D, Wieser C, Dieleman CB. J Inclusion Phenom Mol Recognit Chem. 1997; 29:137–151.
11. Fanni S, Arnaud-Neu F, McKervey MA, Marie-José S-W, Ziat K. Tetrahedron Lett. 1996; 37:7975.
12. Casnati A, Barbosa S, Rouquette Hln, Schwing-Weill M-J, Arnaud-Neu FO, Dozol J-FO, Ungaro R. J Am Chem Soc. 2001; 123:12182–12190. [PubMed: 11734017]
13. [Calix_n_arene phosphine oxide] Miroshinichenko S, Ternova D, Billard I, Ouadi A, Klimchuk O, Kalchenko V. Phosphorus, Sulfur silicon Relat Elem. 2011; 186:903–905. Kostin GA, Us TV, Korda TM, Torgov VG, Kuratieva NV, Miroshinichenko SI, Kalchenko VI. J Inclusion Phenom Macrocyclic Chem. 2010; 68:131–137. Arnaud-Neu F, Browne JK, Byrne D, Marrs DJ, McKervey MA, O'Hagan P, Schwing-Weil MJ, Walker A. Chem Eur J. 1999; 5:175–186.
14. [Calixarene-based sensing for Th(IV)] Zhang YJ, Bhadbhade M, Gao JB, Karatchevtseva I, Price JR, Lumpkin GR. Inorg Chem Commun. 2013; 37:219–221. Lu YC, Bi YF, Bai Y, Liao WP. J Chem Technol Biotechnol. 2013; 88:1836–1840. Ramirez FD, Varbanov S, Padilla J, Buzil JCG. J Phys Chem B. 2008; 112:10976–10988. [PubMed: 18686994] Demirel N, Merdivan M, Pirincioglu, Hamamci C. Anal Chim Acta. 2003; 485:213–219. Malone JF, Marrs DJ, McKervey MA, O'Hagan P, Thompson N, Walker A, Arnaud-Neu F, Mauprivez O, Schwing-Weill MJ, Dozol JF, Rouquette H, Simon N. J Chem Soc, Chem Commun. 1995:2151–2153. Arnaud-Neu F, Cremin S, Harris S, MaKervey MA, Schwing-Weil MH, Schwinte P, Walker A. J Chem Soc, Dalton Trans. 1997:329–334. Arnaud-Neu F, Browne JK, Byrne D, Marrs DJ, McKervey MA, O'Hagan P, Schwing-Weill MJ, Walker A. Chem--Eur J. 1999; 5:2387.
15. [Nuclear waste separation] Casnati A, Pochini A, Ungaro R, Ugozzoli F, Arnaud F, Fanni S, Schwing MJ, Egberink RJM, De Jong F, Reinhoudt DN. J Am Chem Soc. 1995; 117:2767. Arnaud-Neu F, Asfari Z, Souley B, Vicens J. New J Chem. 1996; 20:453. Lamare V, Dozol JF, Fuangswasdi S, Arnaud-Neu F, Thuery P, Nierlich M, Asfari Z, Vicens J. J Chem Soc Perkin Trans. 1999; 2:271–284. Asfari Z, Bressot C, Vicens J, Hill C, Dozol JF, Rouquette H, Eymard S, Lamare V, Tournois B. Anal Chem. 1995; 67:3133–3139. Kim JS, Pang JH, Yu IY, Lee WK, Suh IH, Kim JK, Cho MH, Kim ET, Ra DY. J Chem Soc Perkin Trans. 1999; 2:837–846.

16. Adhikari BB, Ohto K, Schramm MP. Chem Commun. 2014; 50:1903.
17. Adhikari BB, Zhao X, Derakhshan S, Schramm MP. Dalton Trans. 2014; 43:15560. [PubMed: 25198172]
18. Otsuka H, Araki K, Shinkai S. Tetrahedron. 1995; 51:8757.

19. Neupert-Laves K, Dobler M. Helv Chim Acta. 1975; 58:432. [PubMed: 1158723]
20. Trueblood KN, Knobler CB, Maverick E, Helgeson RC, Brown SB, Cram DJ. J Am Chem Soc. 1981; 103:5594.
21. Le Gac S, Picron J-F, Reinaud O, Jabin I. Org Biomol Chem. 2011; 9:2387. [PubMed: 21327258]
22. We wish to thank astute reviewers and editorial staff at Dalton Transactions for this advice.
23. Kim SI, Shin TJ, Ree M, Hwang GT, Kim BH, Han H, Seo J. Journal of Polymer Science Part A: Polymer Chemistry. 1999; 37:2013.

Figure 1. *p*-tert-Butyl calix[6]arene hexacarboxylic acid **1**; its low symmetry “lozenge” solution structure with Pb, Sr and Ba; and a truncated solution (and solid state) structure after adopting a high symmetry C_{3V} conformation (based on NMR and 3 X-ray structures).

Figure 2.
Preparation of calix[6]arene hexacarboxylic acid **2**.

Figure 3.

^1H NMR (400 MHz) of 2.5 mM **2** in presence of 1 eq $\text{Pb}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ in $\text{CDCl}_3/\text{CD}_3\text{CN}$ (1:1). From bottom to top: ligand **2** (a), at the time of mixing with Pb (b), after 30 min (c), after 1 hr (d), and 4 hrs (e), respectively. The NMR signals are with reference to residual CH_3CN signal at 1.94 ppm. Red Arrows show faint traces of low symmetry “lozenge” conformation.

Figure 4. Conformational reorganization of **2** when mixed with Pb(II) with rendered model (MMFF) based on NMR data comparison to **1•Pb**, some protons omitted for clarity.

Figure 5.

^1H NMR (400 MHz) of 2.5 mM **2** in presence of 1 eq $\text{Sr}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ in $\text{CDCl}_3/\text{CD}_3\text{CN}$ (1:1). From bottom to top: at the time of mixing with Sr (a), after 8 hours (b), after 30 hrs (c), after 48 hrs (d), and after 115 hrs (e), respectively. The NMR signals are with reference to residual CH_3CN signal at 1.94 ppm. Red Arrows show evidence of low symmetry “lozenge” conformation.

Figure 6.

^1H NMR (400 MHz) of 2.5 mM **2** in presence of 1 eq $\text{Ba}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ in $\text{CDCl}_3/\text{CD}_3\text{CN}$ (1:1). From bottom to top: at the time of mixing (a), and after 8 hrs (b), 20 hrs (c), 30 hrs (d), and 48 hrs (e), respectively. The NMR signals are with reference to residual CH_3CN signal at 1.94 ppm. Red Arrows show some of the many peaks of low symmetry “lozenge” conformation.

Figure 7. ¹H NMR (400 MHz, in CDCl₃/CD₃CN) Expanded stack plot from bottom to top: **1**•Pb (a), **2**•Pb (b), an initially mixed sample of 5 mM of **1**+**2**+Pb (1:1:1) (c), the mixed sample after 16 hours (d), and 115 hours (e).

Figure 8. ^1H NMR (400 MHz, in $\text{CDCl}_3/\text{CD}_3\text{CN}$) Expanded stack plot from bottom to top: a) $2\cdot\text{Pb}$, b) $2\cdot\text{Ba}$, c) a sample of $2+\text{Pb}+\text{Ba}$ (1:1:1) after 4 hours, and d) after 115 hours.

Table 1

Times to arrive at high symmetry octahedral coordination for hosts **1** and **2** with Pb, Sr, and Ba. All species at 2.5 mM in CDCl₃/CD₃CN (1:1).

	Host 1	Host 2
Pb(II)	15 hours	30 minutes
Sr(II)	20 days	115 hours
Ba(II)	38 days	48 hours