Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/mjafi

Case Report

A rare case of epidermoid cyst of tongue

Lt Col Vivek Saxena^{a,*}, Maj Gen I.P. Majithia (Retd)^b^a Classified Specialist (Oral & Maxillofacial Surgery), Command Military Dental Centre (Central Command), Lucknow, India^b Ex-Commandant & Command Dental Adviser (Command Military Dental Centre (Central Command), Lucknow, India

ARTICLE INFO

Article history:

Received 10 May 2013

Accepted 27 January 2014

Available online 26 April 2014

Keywords:

Epidermoid

Tongue

Cysts

of life. However, when they are developed on the ventral surface of the tongue, their clinical appearance will be much later in childhood or at early adult life. Anatomic classification divides the epidermoid cysts of the floor of the mouth into three groups according to their relation to the muscles of the floor of the mouth: sublingual or median genioglossal cysts, located above the geniohyoid muscles; median geniohyoid cysts, located in the submental region between the geniohyoid and the mylohyoid muscles; and lateral cysts, located in the submaxillary region.⁵

We present the clinical, histological findings and management of epidermoid cyst on ventral surface of the tongue in a 12-year-old boy.

Introduction

Dermoid and epidermoid cysts are uncommon developmental cystic malformations termed as dysontogenetic cyst. Most clinicians and researchers believe that lingual dermoid and epidermoid cysts are a result of entrapped ectodermal tissue of the first and second branchial arches, which fuse during the third and fourth weeks in utero. A second theory suggests that these cysts may be a variant of the thyroglossal duct cyst with ectodermal elements predominating.¹

Dermoid and epidermoid cysts occur in the head and neck region with an incidence of 6.9–7%.^{2,3} About 11.5% of dermoid cysts of the head and neck appear in the floor of the mouth, the second most common location after the floor of the mouth being the cervical region and rarely in the tongue. They represent less than 0.01% of all oral cavity cysts.⁴ Usually lingual dermoid cysts are discovered at birth or at the first year

Case report

A 12-year-old male child patient reported to the dental centre with complains of nodular mass on ventral surface of tongue since 1 year and 6 months (Fig. 1). Patient also complained of difficulty in breathing, eating and swallowing. The nodule was painless and had progressively grown to its present size. Intra oral examination of the child revealed an ovoid solitary mass measuring about 5 cm × 4 cm on the ventral surface of the anterior two thirds of the tongue (Fig. 1). Edges of the mass could not be distinctly made out. Mucosa over the swelling was intact & normal but appeared stretched. The swelling was soft to cystic in consistency. There was no evidence of cervical lymphadenopathy. Transillumination was negative. Aspiration of the cystic mass yielded a viscous pale yellow fluid. Hence a provisional diagnosis of dermoid or epidermoid cyst was made. Routine haematological tests were carried out and found to be within normal limits. Computed tomography

* Tel.: +91 09457582795.

E-mail address: vivek484@gmail.com (V. Saxena).<http://dx.doi.org/10.1016/j.mjafi.2014.01.005>

0377-1237/© 2014, Armed Forces Medical Services (AFMS). All rights reserved.

Fig. 1 – Pre-op view of the lesion and CT sagittal view.

showed a well-circumscribed lesion involving the ventral aspect of the tongue with homogenous hypo-dense areas suggestive of a cystic lesion (Fig. 1). Surgical enucleation of the mass was done under general anaesthesia with nasotracheal intubation. A midline vertical incision was placed on the ventral surface of the tongue and the mass which appeared to be a thick walled tense cyst was gently separated from the tongue musculature and was excised completely (Fig. 2). Histopathological examination revealed the presence of pseudostratified ciliated squamous epithelium, with mononuclear inflammatory infiltrate and lymphoid follicles in the cyst wall, without any skin adnexa (Fig. 3). Thus the final diagnosis of epidermoid cyst was made. Post-operative period was uneventful and no recurrence was noted at 1-year follow-up period (Fig. 4).

Discussion

Epidermoid cysts occur primarily on facial skin or neck, and are most common in the midcheek and preauricular area. Intraoral epidermoid cysts are considered rare and most commonly occur in the floor of the mouth. New and Erich (1937) reported

24 (1.6%) epidermoid cysts occurring in the floor of the mouth out of 1495 cases of dermoid cysts seen at the Mayo Clinic.⁶

Clinically epidermoid cyst may be asymptomatic, often discovered by a painless swelling growing slowly and surrounding by a normal mucosa. The cyst can enlarge to a point that the patient present difficulty of articulation, mastication and deglutition and airway compromise. The differential diagnosis of a painless swelling in the midline of the tongue include cystic hygroma, lymphatic cyst, neurofibroma, haemangioma, lingual thyroid, enteric duplication cyst and dermoid cyst.^{7,8}

Most cases are suspected from the history of acne and of past cysts with infections, but less apparent epidermoid cysts will present in a manner similar to a lymph node enlargement. In various locations, they may be confused with entities more specific to that location. In the preauricular area, an epidermoid cyst may resemble a parotid tumour, in the mid-lateral neck, a branchial cyst, in the midline of the neck, a thyroglossal tract cyst, and in the submental triangle, a dermoid cyst.³

The imaging modalities include ultrasonography, computed tomography and magnetic resonance, which help in diagnosis. Ultrasonography is the most used technique because it is a low-cost, fast, reliable imaging method, without

Fig. 2 – Exposure & delineation of the lesion.

Fig. 3 – Micrograph of histopathological section.

X-ray exposure. Computed tomography presents better quality when compared to ultrasonography, because the image is more precise, showing the exact relation of the lesion to the muscles of the floor of the mouth. Magnetic resonance imaging (MRI) is an accurate, noninvasive modality for the diagnosis and follow-up evaluation for lingual dermoid cyst. It has the advantage of predicting the possible extension of the cyst through the floor of the mouth.⁹

Meyer described three histological variants of dermoid cyst – namely the true dermoid cyst, the epidermoid cyst and the teratoid variant. True dermoid cysts are cavities lined with epithelium showing keratinization and with identifiable skin appendages such as pilous follicles, and sudoriparous and

sebaceous glands on the cyst wall.¹⁰ The epidermoid cyst closely resembles a dermoid cyst, except that the former exhibits no dermal appendages. They are lined by orthokeratinized squamous epithelium and exhibit a lumen that is generally filled with desquamated keratin. The cyst wall consists of a narrow zone of compressed fibrous connective tissue that is generally free of inflammation. The teratoid cyst is most rare of the developmental cyst of the oral cavity and composed of elements derived from ectoderm, endoderm and mesoderm.¹¹

Treatment is complete removal of the cyst via an intraoral or extraoral approach. An intraoral approach is recommended by most authors for lingual cysts and sublingual cysts of small or moderate dimensions, whereas an extraoral approach is preferred for large sublingual cysts or geniohyoid cysts. In our case, an intra-oral approach with a mucosal incision through the ventral surface of the tongue was placed with a caution to avoid submandibular salivary gland duct openings. Prognosis is very good, with a very low incidence of recurrence.¹¹

Conclusion

In conclusion, occurrence of epidermoid cyst on ventral surface of tongue is an extremely rare phenomenon. Clinically they are painless, slowly growing mass causing difficulty in eating, swallowing and breathing. Imaging modalities help to delineate the extent of the lesion. Histopathological examination confirms the diagnosis of a cystic lesion lined by squamous epithelia without skin appendages. Complete surgical excision via an intraoral approach is the treatment of choice and recurrence is rare.

Conflicts of interest

The author has none to declare.

REFERENCES

1. Louis PJ, Hudson C, Reddi S. Lesion of floor of the mouth. *J Oral Maxillofac Surg.* 2002;60:804–807.
2. Koeller KK, Alamo L, Adair CF, Smirniotopoulos JG. Congenital cystic masses of the neck: radiological–pathologic correlation. *Radiographics.* 1999;19:121–146.
3. Dimov Zh, Dimov K, Kr'stev N, et al. Dermoid, epidermoid and teratoma cysts of the tongue and oral cavity. *Khirurgiia (Sofia).* 2000;56:30–32.
4. De Ponte FS, Brunelli A, Marchetti E, Bottini DJ. Sublingual epidermoid cyst. *J Craniofac Surg.* 2002;13:308–310.
5. Longo F, Maremonti P, Mangone GM, De Maria G, Califano L. Midline (dermoid) cysts of the floor of the mouth: report of 16 cases and review of surgical techniques. *Plast Reconstr Surg.* 2003;112:1560–1565.
6. New GB, Erich JB. Dermoid cysts of head and neck. *Surg Gynecol Obstet.* 1937;65:48–55.
7. Turetschek K, Hospodka H, Steiner E. Case report: epidermoid cyst of the floor of the mouth: diagnostic imaging by sonography, computed tomography and magnetic resonance imaging. *Br J Radiol.* 1995;68:205–207.

Fig. 4 – One year post-op view.

8. Gold BD, Sheinkopf DE, Levy B. Dermoid, epidermoid and teratomatous cysts of the tongue and the floor of the mouth. *J Oral Surg.* 1974;30:107–111.
9. Gleizal A, Abouchebel N, Lebreton F, Beziat JL. Dermoid cyst of the tongue: an association of dermoid cyst with bronchogenic epithelium. *J Craniomaxillofac Surg.* 2006;34:113–116.
10. Meyer I. Dermoid cyst of the mouth. *Oral Surg Med Oral Pathol.* 1955;8:1149–1164.
11. Calderon S. Concomitant sublingual and submental epidermoid cyst of the floor of the mouth. *J Oral Maxillofac Surg.* 1993;51:790–792.