Published in final edited form as:

Mol Oral Microbiol. 2016 February ; 31(1): 106–114. doi:10.1111/omi.12136.

***Aggregatibacter actinomycetemcomitans* leukotoxin induces cytosol acidification in LFA-1 expressing immune cells**

Nataliya Balashova¹, Anuradha Dhingra², Kathleen Boesze-Battaglia², and Edward T. Lally^{1,*}

¹Department of Pathology, University of Pennsylvania School of Dental Medicine, Philadelphia, Pennsylvania 19104, USA

²Department of Biochemistry, University of Pennsylvania School of Dental Medicine, Philadelphia, Pennsylvania 19104, USA

Summary

Studies have suggested that *Aggregatibacter actinomycetemcomitans* leukotoxin (LtxA) kills human LFA-1 (CD11a/CD18)-bearing immune cells through a lysosomal-mediated mechanism. Lysosomes are membrane-bound cellular organelles that contain an array of acid hydrolases that are capable of breaking down biomolecules. The lysosomal membrane bilayer confines the pH-sensitive enzymes within an optimal acidic (pH 4.8) environment thereby protecting the slightly basic cytosol (pH 6.8–7.5). In the current study, we have probed the effect of LtxA-induced cytolysis on lysosomal integrity in two different K562 erythroleukemia cell lines. K562-puro/LFA-1 cells were stably transfected with CD11a and CD18 cDNA to express LFA-1 on the cell surface while K562-puro, which does not express LFA-1, served as a control. Following treatment with 100 ng/ml LtxA cells were analyzed by live cell imaging in conjunction with time-lapse confocal microscopy and by flow cytometry. Using a pH sensitive indicator (pHrodo[®], Life Technologies) we demonstrated that the toxin causes a decrease in the intracellular pH in K562-puro/LFA-1 cells noticeable within the first 15 min of treatment. This process correlated with the disappearance of lysosomes in the cytosol as determined by both acridine orange and LysoTracker[®] Red DND-99 (Life Technologies) staining. These changes were not observed in K562-puro cells or when heat inactivated toxin was added to K562-puro/LFA-1. Our results suggest that LtxA induces lysosomal damage, cytosol acidification, which is followed by cell death in K562-puro/LFA-1 cells.

Keywords

bacterial pathogenesis; RTX-toxin; cytosolic pH; cytosol acidification; lysosome

Introduction

Aggregatibacter actinomycetemcomitans is a pioneer colonizer of the upper aerodigestive tract of both humans and non-human primates. In both man (Lamell *et al.*, 2000) and captive

* Address correspondence to: Edward T. Lally, Department of Pathology, University of Pennsylvania, School of Dental Medicine, 240 S. 40th St., 317 Levy Bldg., Philadelphia, PA 19104. Tel.: 215-898-5913; Fax: 215-898-2050; lally@dental.upenn.edu.

M. fascicularis (Beighton *et al.*, 1989) colonization of the tongue and buccal mucosa begins at 3 to 6 months of age. *A. actinomycetemcomitans*, a HACEK organism, is part of the group of causative organisms associated with Gram-negative endocarditis that are not due to illicit drug use (Das *et al.*, 1997; Meyer, 1989). However, the most-studied infection of *A. actinomycetemcomitans* is localized juvenile periodontitis (Zambon, 1985), which has been subsequently renamed localized aggressive periodontitis (LAP, (Armitage, 1999)). LAP is destructive inflammatory process of the periodontal ligament that affects the first molars and central incisors and results in rapid destruction of alveolar bone and subsequent tooth loss.

The organism has several virulence determinants including a 116 kDa leukotoxin (LtxA) that is a member of the RTX (Repeats in ToXin) family of cytolytic proteins (Welch, 1991). The interaction of LtxA with human immune cells is both complex and multifaceted with interactions that involve membrane hydrocarbons as well as cell-surface proteins. LtxA selectively kills human β 2-integrin lymphocyte function antigen-1 (LFA-1) bearing white blood cells (Lally *et al.*, 1997). While the binding of LtxA to LFA-1 on the surface of leukocytes is well established, our knowledge of the mechanisms by which LtxA causes intracellular changes leading to host cell death is only beginning to be understood. The ability of the toxin to destroy THP-1 lysosomes has recently been demonstrated (DiFranco *et al.*, 2012).

This work is aimed to further our understanding of the intracellular events leading to LtxA-induced cytolysis. Here we report LtxA-induced cytosolic acidification in K562 cells expressing LFA-1, which occurred concurrently with lysosomal rupture in these cells.

Materials and Methods

LtxA purification

A. actinomycetemcomitans strain JP2 (Tsai *et al.*, 1984) was grown in AAGM medium (Fine *et al.*, 1999). The bacteria were grown on solid AAGM for 48 h at 37 °C in the presence of 10% CO₂. Colonies were inoculated into AAGM broth and were incubated for 24 h. LtxA was purified by ammonium sulfate precipitation from the bacterial culture supernatants as described previously (Diaz *et al.*, 2006). Purified protein was resolved on a 4–20% SDS-PAGE and visualized by staining with Coomassie Brilliant Blue G-250 protein stain (Bio-Rad Laboratories). The protein concentration was measured with a nanodrop (Thermo Fisher Scientific).

Cell culture

Human erythroleukemia K562 cells obtained from ATCC were grown in RPMI 1640 medium containing 10% FBS. K562-derived cell line expressing CD11a and CD18 (K562-pEFpuro/LFA-1) was provided by Dr. Timothy A. Springer (Lu and Springer, 1997). To generate control K562-puro cell line, pEF1-puro/hpb1 (Addgene) was digested by *Bam*HI and *Pme*I, and the 6 kb pEF1puro fragment was eluted from gel. Two μ g of linearized pEF1-puro fragment was applied to electroporate 0.5×10^6 K562 cells at 155 V and 960 μ F in 0.4-cm path-length cuvettes (Bio-Rad Laboratories) using GenePulser X cell instrument (Bio-

Rad Laboratories). The stable cell lines transfected with pEF1puro or pEFpuro/LFA-1 were maintained in the complete medium containing 3 µg/ml puromycin.

Cytotoxicity assay

For toxicity tests, 2–500 ng/ml purified LtxA was added to 1×10^6 erythroleukemia cells in RPMI with 10% FBS and incubated for 3 or 24 h. The cell membrane permeability was determined with Trypan blue assay using Vi-cell Cell Viability Analyzer (Beckman Coulter). All reactions were run in technical duplicate; the assay was performed three independent times. The controls included LtxA inactivated at 65 °C for 30 min and untreated cells.

Confocal microscopy

For pHrodo staining 1×10^6 erythroleukemia cells were washed twice with Live Cell Imaging Solution (LCIS) (Life Technologies). The cells were incubated with 100 ng/ml LtxA for different periods of time and washed. The cells were then treated with a mix containing 5 mM of pHrodo™ Red AM, 10 µL of PowerLoad™ (Life Technologies) in 1 mL of LCIS for 37°C for 30 minutes. After the treatment the cells were washed twice with LCIS and imaged immediately. The pH calibration was performed using Intracellular pH Calibration Buffer Kit (Life technologies) according to the manufacturer's instructions. Briefly, the cell loading solution was prepared by diluting the valinomycin and nigericin stock solution in the desired cellular pH calibration buffer to a concentration of 10 µM each. The cell loading solution was then added to pHrodo™ Red AM labeled cells, incubated at 37°C for 5 minutes, and the cells were imaged. To analyze the effect of LtxA on cytosolic pH, K562 cells were labeled with pHrodo Red, treated with 100 ng/ml LtxA, and subjected to confocal imaging at different time points (0, 15, and 30 min).

For lysosomal staining, the cells were treated with 1 µM acridine orange (Sigma) for 15 min or 100 nM LysoTracker® Red DND-99 (Life Technologies) for 2 h at 37 °C. Then the cells were washed with LCIS and the cell nuclei were stained 1 µM Hoechst 33342 for 15 min (Image-iT™ LIVE Plasma Membrane and Nuclear Labeling Kit, Life technologies). The cells were washed with LCIS and treated with 100 ng/ml LtxA for different time intervals at 37 °C. Treatment with 1 mM Leu-Leu methyl ester (Leu-Leu-OMe) (Sigma) for 30 min at 37 °C was used to induce lysosomal rupture. The cells were then washed with LCIS and placed to attach for 20 min in ibiTreat 60 µ-dishes (Ibidi) coated with poly-L-lysine (Sigma), then they were washed again and covered with LCIS. The cells were examined using a Nikon A1R laser scanning confocal microscope with a 60X water objective (NA 1.2). The images were processed using Nikon's Elements software 4.1. Approximately 50 cells per image were analyzed in each experiment to identify the mean and maximum fluorescent intensity per cell by sorting non-saturated areas in 3 combined Z planes collected for each image. For live time lapse confocal microscopy images were collected every 2 min for 1 h.

Flow cytometry

To test LFA-1 expression, erythroleukemia cells (0.5×10^6 cells/run) were incubated with anti-human CD11a Alexa Fluor 594 clone HI111 (Biolegend) or anti-human CD11a FITC clone TS1/18 (Biolegend) (1:1000) for 1h at 25 °C and washed with PBS. FITC mouse IgG1

and Alexa Fluor 647 mouse IgG1 (Biolegend) served as isotype controls. To analyze cytosolic pH, K562 cells were labeled with pHrodo Green and treated with LtxA as for confocal imaging. At different periods of time (0, 15, and 30 min) the cells were analyzed by flow cytometry (509/533 nm). The fluorescence measurements were made with a BD LSR II flow cytometer (BD Biosciences). Ten thousand events were recorded per sample, and mean fluorescence intensities (MFI) were determined using WinList software.

Statistical Analysis

The data were analyzed using an unpaired Student's *t* test, with $p < 0.05$ considered to be statistically significant. The analysis was performed using SigmaPlot version 11.0 software.

Results

K562 cells as a model for study of LtxA-mediated cell death

K562 erythroleukemia cells can spontaneously develop characteristics similar to early stage erythrocytes, granulocytes, and monocytes (Lozzio and Lozzio, 1979; Lozzio and Lozzio, 1975). K562 cell lines transfected with different $\beta 2$ -integrins cDNA were used previously to study the function of the integrin proteins (Lu and Springer, 1997; Lub *et al.*, 1997; Sigal *et al.*, 2000) and for the identification of the receptors for RTX leukotoxins (Fett *et al.*, 2008; Munksgaard *et al.*, 2014; Reinholdt *et al.*, 2013). In our study of LtxA-mediated cytolysis, we utilized K562 cell line stably transfected with CD11a/CD18 cDNA (K562-puro/LFA-1) (Lu and Springer, 1997). We generated a K562 cell line stably transfected with pEF1-puro DNA to be used as a negative control. The expression of LFA-1 subunits (CD11a and CD18) on the surface of K562-puro/LFA-1 and their absence on the surface of K562-puro were confirmed by flow cytometry analysis (Fig. 1A). We determined the sensitivity of these two cell lines to LtxA in dose-dependent manner after 24 h of treatment. The dose-dependent killing curves for toxin concentrations from 2 to 500 ng/ml applied to both K562-puro/LFA-1 and K562-puro are provided in Fig. 1B. We selected 100 ng/ml LtxA for our experiments since it was a sub-maximal dose at which the effect of LtxA on K562-puro/LFA-1 was pronounced however K562-puro control cells were not significantly affected. This concentration was compatible with imaging studies. At the concentration of 100 ng/ml LtxA caused about 10% K562-puro/LFA-1 cell death in 3 h and more than 30% in 24 h while K562-puro cells were not significantly affected (Fig. 1B). Heat-inactivated toxin did not cause any K562-puro/LFA-1 cell lysis (data not shown). Thus, as expected, LFA-1 expression on the surface of K562-puro/LFA-1 significantly increased its susceptibility to LtxA.

LtxA causes intracellular acidification in K562-puro/LFA-1

Cytosolic pH plays an important role in maintenance of normal cell function. This parameter is tightly regulated within cells and varies depending on cell type (pH 6.8–7.5) (Deutsch *et al.*, 1982). Thus, we sought to determine if LtxA could change cytosolic pH in host leukocytes. We utilized pHrodoTM Red/Green AM fluorogenic probes (pKa 6.5) to measure cytosolic pH in live K562 cell lines. These membrane-permeable dyes are weakly fluorescent at neutral pH, but increase in fluorescence with a decrease of pH as the surrounding environment becomes more acidic. Using the valinomycin/nigericin system,

which equilibrates extracellular and cytosolic pH, the intracellular fluorescence intensity in pHrodo labeled K562-puro/LFA-1 cells was analyzed using confocal microscopy in the pH range (5.5 to 7.5) as shown in Fig. 2A, left panel. A steady increase in the fluorescence intensity in K562-puro/LFA-1 cells as the pH is decreased from 7.5 to 5.5. Due to the low pH of the endocytic vesicles and lysosomes, these cellular compartments were intensively stained with pHrodo, however, we observed acidification not only in acidic organelles but throughout the intracellular milieu. The fluorescence intensity values in pHrodo Green labeled K562-puro/LFA-1 cells were quantitated by flow cytometry analysis (indicated as mean fluorescent intensity, MFI; Fig. 2A, right panel).

Hence this technique enabled the quantification of cytosolic pH in K562-puro/LFA-1 cells, we then used pHrodo dyes to monitor this parameter in K562 cell lines after treatment with 100 ng/ml LtxA. We observed cytosolic pH decrease approx. 15 min after LtxA treatment of the cells prestained with pHrodo Red by confocal microscopy (Fig. 2B, left panel), which reached plateau by 30 min of the treatment with LtxA. Some loss of fluorescence signal occasionally was observed during later points (after 30 min) of the experiment. This could be because pHrodo dye might leak out of the cells or spontaneously released from dead cells, as well as it might be bleached by exposure to light during the time lapse confocal experiment. To confirm the confocal result, pHrodo labeled K562-puro/LFA-1 cells were subsequently treated with 100 ng/ml LtxA and analyzed by both confocal microscopy (Fig. 2B, left panel) and flow cytometry. MFI clearly demonstrated that the acidification occurred after the exposure to LtxA (Fig. 2B, right panel). Treatment of K562-puro cells did not result in significant changes in cytosolic pH of these cells as was observed by both confocal microscopy (Fig. 2C, left panel) and flow cytometry (Fig. 2C, right panel), as well as no changes in cytosolic pH were observed when K562-puro/LFA-1 cells were treated with heat inactivated LtxA (data not shown).

Ltxa causes lysosomal rupture in K562-puro/LFA-1

We hypothesized that the fast cytosol acidification observed in K562-puro/LFA-1 cells upon LtxA treatment could be due to the toxin induced lysosomal membrane damage and the release of the highly acidic lysosomal content into the intracellular space. Therefore, we wanted to see if LtxA was able to cause lysosomal damage or rupture in K562-puro/LFA-1 cells. In order to test this hypothesis we used acridine orange, a cell-permeant metachromatic fluorescent dye that stains cellular acidic organelles in red. Additionally, acridine orange is nucleic acid binding dye that emits green fluorescence when bound to dsDNA and red fluorescence when bound to ssDNA or RNA. As a positive control we utilized a lysosomotropic agent Leu-Leu-OMe, which is known to induce apoptosis of certain immune cells by causing rupture of the lysosomes and DNA fragmentation (Uchimoto *et al.*, 1999). When acridine orange stained cells were used to monitor lysosomal membrane integrity, red granular fluorescence was seen in untreated cells (Fig. 3A, B). In contrast, 1 mM Leu-Leu-OMe treated K562 cells showed significant change in red fluorescence intensity and distribution throughout the cytoplasm, suggesting that Leu-Leu-OMe induced lysosomal rupture. the treatment of K562-puro/LFA-1 cells with 100 ng/ml LtxA also caused changes in red fluorescence similar to those observed after Leu-Leu-OMe exposure. These changes were tracked over time in live cell imaging experiments and the

decrease in lysosomal fluorescence was detected as early as 4 min after toxin addition (Fig. 3A), indicating that LtxA is able to cause lysosomal degradation in K562-puro/LFA-1 cells. By contrast, no significant changes in fluorescence intensity were observed when the heat inactivated toxin was applied to K562-puro/LFA-1 (Fig. 3B). Using another lysosomal dye, LysoTracker® Red DND-99, we next followed changes in lysosomal properties of K562-puro/LFA-1 cells over time using live imaging in conjunction with time-lapse confocal microscopy. We detected a decrease in red fluorescence intensity within first 5 min of treatment and continued treatments resulted in further lysosomal degradation as indicated by the decrease in red fluorescence intensity (Fig. 3C). The loss of the fluorescence intensity was not observed by both lysosomal staining methods when K562-puro cells were used (Fig. 3D) suggesting that the expression of LFA-1 is a critical component required for LtxA to initiate cytosolic changes in host cells.

Discussion

Cellular homeostasis is dependent upon proper regulation of cytosolic pH. There are numerous mechanisms that can cause change in cytosolic pH, including metabolic acid production, leakage of acid through plasma and organelle membranes, and membrane transport processes. Imbalance in cytosolic pH affects many cellular functions, including ionic homeostasis (Park and Brown, 1995), apoptosis (Park, 1995), cell cycle, cell volume regulation (Ritter *et al.*, 1992), muscular contraction (Hao *et al.*, 1999), vascular function (Schulz and Munzel, 2011), concentration of intracellular messengers such as Ca^{2+} and cAMP (Brokaw, 1987; Speake and Elliott, 1998), and cytoskeletal interactions (Edmonds *et al.*, 1995).

Techniques to measure intracellular pH include H^+ permeable microelectrodes, nuclear magnetic resonance (NMR) analysis of cellular metabolites whose resonance frequency is influenced by pH, and using of weak acid fluorescent dyes, such as pHrodo™ AM (Kotyk, 1989; Wray, 1988). Here we present a quantitative methodology to assess cytosolic acidification in K562 cells in response to LtxA treatment by both confocal microscopy and flow cytometry. A shift in the fluorescence signal intensity in pHrodo labeled K562-puro/LFA-1 cells indicates cytosol acidification in these cells. An evaluation of both lysosomal rupture and cytosolic acidification using pHrodo in a single confocal experiment will result in induction of a technical artifact (pHrodo photobleaching). However, cytosolic acidification measured by pHrodo staining and changes in lysosomal structure/intensity detected using LysoTracker® Red DND-99 were observed in a very close time interval. Cytosol acidification becomes visible at approximately 15 after an addition of LtxA. We routinely observe the decrease in lysosomal intensity as early as 4 min after addition of LtxA (see Fig. 3) after staining with both LysoTracker® and acridine orange. Given that the decrease in cytosolic pH coincides with reduced lysosomal fluorescence intensity rupture could be the major reason for cytosolic acidification in these cells. Combination of cell imaging techniques with flow cytometry allowed us to analyze the effect of LtxA on both a whole population of cells and individual cells at the same time.

Lysosomes are fundamental for cell growth, and thus inhibition of the lysosomal function often leads to cell death. Besides that, lysosomal membrane and the subsequent leakage of

the highly acidic content into the cytosol may lead to the cell death. The lysosomal cell death may have necrotic and apoptotic features depending on the nature of the stimuli and the extent of the leakage (Aits and Jaattela, 2013). Extensive lysosomal damage may lead to inevitable necrotic cell death, while less extensive detriment of lysosomes may induce several apoptotic pathways, which can be attenuated by inhibition of by lysosomal proteases (cathepsins) (Kagedal *et al.*, 2001a; Kagedal *et al.*, 2001b; Kirkegaard and Jaattela, 2009). A number of organic and non-organic agents are known to cause lysosomal damage. These are lysosomotropic detergents such as imidazole, ciprofloxacin, Leu-leu-OMe (de Duve *et al.*, 1974); reactive oxygen species (Kurz *et al.*, 2008a; 2008b); some viral proteins (Lozach *et al.*, 2011); bacterial, fungal, and snake toxins (DiFranco *et al.*, 2012; Feofanov *et al.*, 2005; Ivanova *et al.*, 2012; Newman *et al.*, 2009); and other regulators (Aits and Jaattela, 2013).

Recent data indicate that LtxA was detected in lysosomes and could trigger lysosomal mediated cell death in THP-1 monocytes (DiFranco *et al.*, 2012). Our study shows that effect of LtxA on lysosomes in K562-puro/LFA-1 is similar to one induced in THP-1 cells. Additionally, our results demonstrate that a decrease in cytosolic pH of K562-puro/LFA-1 correlates well to the loss of lysosomal membrane integrity in these cells and is dependent upon expression of LFA-1. The killing rate of K562-puro/LFA-1 engineered cells is relatively low, compare to THP-1 cells, in which cell death of most of the cells was detected within first 3 h. This feature of K562-puro/LFA-1 gives us an advantage in imaging of the toxin at trafficking inside of host cells. K562-puro cells provide an excellent negative control to be used in this study. Moving forward the role of different integrins in the toxin trafficking will be investigated using this recombinant system.

Acknowledgments

This work was supported by the United States National Institute of Health grants R01DE009517 (ETL) and R01DE022465 (KBB). The authors thank the Penn Dental Medicine Live Cell Imaging Core facility and Dr. T.A. Springer for providing K562-puro/LFA-1 cells.

References

- Aits S, Jaattela M. Lysosomal cell death at a glance. *J Cell Sci.* 2013; 126:1905–12. [PubMed: 23720375]
- Armitage GC. Development of a classification system for periodontal diseases and conditions. *Ann Periodontol.* 1999; 4:1–6. [PubMed: 10863370]
- Beighton D, Taichman NS, Simpson DL, Dirienzo JM, Johnson NW. Intra-oral colonization of macaque monkeys by *Actinobacillus actinomycetemcomitans*. *Oral Microbiol Immunol.* 1989; 4:35–40. [PubMed: 2628866]
- Brokaw CJ. Regulation of sperm flagellar motility by calcium and cAMP-dependent phosphorylation. *J Cell Biochem.* 1987; 35:175–84. [PubMed: 2826504]
- Das M, Badley AD, Cockerill FR, Steckelberg JM, Wilson WR. Infective endocarditis caused by HACEK microorganisms. *Annu Rev Med.* 1997; 48:25–33. [PubMed: 9046942]
- de Duve C, de Barse T, Poole B, Trouet A, Tulkens P, Van Hoof F. Commentary. Lysosomotropic agents. *Biochem Pharmacol.* 1974; 23:2495–531. [PubMed: 4606365]
- Deutsch C, Taylor JS, Wilson DF. Regulation of intracellular pH by human peripheral blood lymphocytes as measured by ¹⁹F NMR. *Proc Natl Acad Sci U S A.* 1982; 79:7944–8. [PubMed: 6961462]

- Diaz R, Ghofaily LA, Patel J, Balashova NV, Freitas AC, Labib I, Kachlany SC. Characterization of leukotoxin from a clinical strain of *Actinobacillus actinomycetemcomitans*. *Microbial Pathogenesis*. 2006; 40:48–55. [PubMed: 16414241]
- DiFranco KM, Gupta A, Galusha LE, Perez J, Nguyen TV, Fineza CD, Kachlany SC. Leukotoxin (Leukothera(R)) targets active leukocyte function antigen-1 (LFA-1) protein and triggers a lysosomal mediated cell death pathway. *J Biol Chem*. 2012; 287:17618–27. [PubMed: 22467872]
- Edmonds BT, Murray J, Condeelis J. pH regulation of the F-actin binding properties of Dictyostelium elongation factor 1 alpha. *J Biol Chem*. 1995; 270:15222–30. [PubMed: 7797506]
- Feofanov AV, Sharonov GV, Astapova MV, Rodionov DI, Utkin YN, Arseniev AS. Cancer cell injury by cytotoxins from cobra venom is mediated through lysosomal damage. *Biochem J*. 2005; 390:11–8. [PubMed: 15847607]
- Fett T, Zecchinon L, Vanden Bergh P, Desmecht D. Mannheimia haemolytica leukotoxin-induced cytolysis of caprine (*Capra hircus*) leukocytes is mediated by the CD18 subunit of beta2-integrins. *Microb Pathog*. 2008; 45:337–42. [PubMed: 18775487]
- Fine DH, Furgang D, Schreiner HC, Goncharoff P, Charlesworth J, Ghazwan G, Fitzgerald-Bocarsly P, Figurski DH. Phenotypic variation in *Actinobacillus actinomycetemcomitans* during laboratory growth: implications for virulence. *Microbiology*. 1999; 145 (Pt 6):1335–47. [PubMed: 10411260]
- Hao W, Luo Z, Zheng L, Prasad K, Lafer EM. AP180 and AP-2 interact directly in a complex that cooperatively assembles clathrin. *J Biol Chem*. 1999; 274:22785–94. [PubMed: 10428863]
- Ivanova L, Egge-Jacobsen WM, Solhaug A, Thoen E, Faeste CK. Lysosomes as a possible target of enniatin B-induced toxicity in Caco-2 cells. *Chem Res Toxicol*. 2012; 25:1662–74. [PubMed: 22731695]
- Kagedal K, Johansson U, Ollinger K. The lysosomal protease cathepsin D mediates apoptosis induced by oxidative stress. *FASEB J*. 2001a; 15:1592–4. [PubMed: 11427496]
- Kagedal K, Zhao M, Svensson I, Brunk UT. Sphingosine-induced apoptosis is dependent on lysosomal proteases. *Biochem J*. 2001b; 359:335–43. [PubMed: 11583579]
- Kirkegaard T, Jaattela M. Lysosomal involvement in cell death and cancer. *Biochim Biophys Acta*. 2009; 1793:746–54. [PubMed: 18948147]
- Kotyk, ASaSJ. Intracellular pH and Its Measurement. CRC Press; Boca Raton, FL, USA: 1989.
- Kurz T, Terman A, Gustafsson B, Brunk UT. Lysosomes and oxidative stress in aging and apoptosis. *Biochim Biophys Acta*. 2008a; 1780:1291–303. [PubMed: 18255041]
- Kurz T, Terman A, Gustafsson B, Brunk UT. Lysosomes in iron metabolism, ageing and apoptosis. *Histochem Cell Biol*. 2008b; 129:389–406. [PubMed: 18259769]
- Lally ET, Kieba IR, Sato A, Green CL, Rosenbloom J, Korostoff J, Wang JF, Shenker BJ, Ortlepp S, Robinson MK, Billings PC. RTX toxins recognize a beta2 integrin on the surface of human target cells. *J Biol Chem*. 1997; 272:30463–9. [PubMed: 9374538]
- Lamell CW, Griffen AL, McClellan DL, Leys EJ. Acquisition and colonization stability of *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in children. *J Clin Microbiol*. 2000; 38:1196–9. [PubMed: 10699021]
- Lozach PY, Huotari J, Helenius A. Late-penetrating viruses. *Curr Opin Virol*. 2011; 1:35–43. [PubMed: 22440565]
- Lozzio BB, Lozzio CB. Properties and usefulness of the original K-562 human myelogenous leukemia cell line. *Leuk Res*. 1979; 3:363–70. [PubMed: 95026]
- Lozzio CB, Lozzio BB. Human chronic myelogenous leukemia cell-line with positive Philadelphia chromosome. *Blood*. 1975; 45:321–34. [PubMed: 163658]
- Lu CF, Springer TA. The alpha subunit cytoplasmic domain regulates the assembly and adhesiveness of integrin lymphocyte function-associated antigen-1. *J Immunol*. 1997; 159:268–78. [PubMed: 9200463]
- Lub M, van Vliet SJ, Oomen SP, Pieters RA, Robinson M, Figdor CG, van Kooyk Y. Cytoplasmic tails of beta 1, beta 2, and beta 7 integrins differentially regulate LFA-1 function in K562 cells. *Mol Biol Cell*. 1997; 8:719–28. [PubMed: 9247650]
- Meyer DJ. Favorable prognosis of patients with prosthetic valve endocarditis caused by gram-negative bacilli of the HACEK group. *Am J Med*. 1989; 85:104–107. [PubMed: 3291611]

- Munksgaard PS, Skals M, Reinholdt J, Poulsen K, Jensen MR, Yang C, Leipziger J, Vorup-Jensen T, Praetorius HA. Sialic acid residues are essential for cell lysis mediated by leukotoxin from *Aggregatibacter actinomycetemcomitans*. *Infect Immun*. 2014; 82:2219–28. [PubMed: 24643533]
- Newman ZL, Leppla SH, Moayeri M. CA-074Me protection against anthrax lethal toxin. *Infect Immun*. 2009; 77:4327–36. [PubMed: 19635822]
- Park HJ. Effects of intracellular pH on apoptosis in HL-60 human leukemia cells. *Yonsei Med J*. 1995; 36:473–9. [PubMed: 8599248]
- Park K, Brown PD. Intracellular pH modulates the activity of chloride channels in isolated lacrimal gland acinar cells. *Am J Physiol*. 1995; 268:C647–50. [PubMed: 7900771]
- Reinholdt J, Poulsen K, Brinkmann CR, Hoffmann SV, Stapulionis R, Enghild JJ, Jensen UB, Boesen T, Vorup-Jensen T. Monodisperse and LPS-free *Aggregatibacter actinomycetemcomitans* leukotoxin: interactions with human beta2 integrins and erythrocytes. *Biochim Biophys Acta*. 2013; 1834:546–58. [PubMed: 23234758]
- Ritter M, Woll E, Haussinger D, Lang F. Effects of bradykinin on cell volume and intracellular pH in NIH 3T3 fibroblasts expressing the ras oncogene. *FEBS Lett*. 1992; 307:367–70. [PubMed: 1322838]
- Schulz E, Munzel T. Intracellular pH: a fundamental modulator of vascular function. *Circulation*. 2011; 124:1806–7. [PubMed: 22025635]
- Sigal A, Bleijs DA, Grabovsky V, van Vliet SJ, Dwir O, Figdor CG, van Kooyk Y, Alon R. The LFA-1 integrin supports rolling adhesions on ICAM-1 under physiological shear flow in a permissive cellular environment. *J Immunol*. 2000; 165:442–52. [PubMed: 10861083]
- Speake T, Elliott AC. Modulation of calcium signals by intracellular pH in isolated rat pancreatic acinar cells. *J Physiol*. 1998; 506 (Pt 2):415–30. [PubMed: 9490869]
- Tsai CC, Shenker BJ, DiRienzo JM, Malamud D, Taichman NS. Extraction and isolation of a leukotoxin from *Actinobacillus actinomycetemcomitans* with polymyxin B. *Infect Immun*. 1984; 43:700–5. [PubMed: 6319288]
- Uchimoto T, Nohara H, Kamehara R, Iwamura M, Watanabe N, Kobayashi Y. Mechanism of apoptosis induced by a lysosomotropic agent, L-Leucyl-L-Leucine methyl ester. *Apoptosis*. 1999; 4:357–62. [PubMed: 14634338]
- Welch RA. Pore-forming cytolysins of gram-negative bacteria. *Mol Microbiol*. 1991; 5:521–8. [PubMed: 2046545]
- Wray S. Smooth muscle intracellular pH: measurement, regulation, and function. *Am J Physiol*. 1988; 254:C213–25. [PubMed: 3279796]
- Zambon JJ. *Actinobacillus actinomycetemcomitans* in human periodontal disease. *J Clin Periodontol*. 1985; 12:1–20. [PubMed: 3882766]

Figure 1.

A. Flow cytometry analysis of erythroleukemia cell lines K562-puro (green) and K562-puro/LFA-1 (red) with anti-human CD11a (HI111) and anti-human CD18 (TS1/18) antibody. The experiment was performed three independent times. Representative images are shown.

B. Dose-dependent determination of LtxA cytotoxicity for K562 cell lines. For toxicity tests, 2–500 ng/ml LtxA was added to 1×10^6 K562 cells in RPMI with 10% FBS. Toxicity of LtxA on K562 cell lines was measured using Trypan blue assay after 24 h. Untreated cells and cells treated with the toxin that was heat inactivated (HI LtxA) at 65 °C for 30 min served as controls. Average results of three independent experiments are displayed. Error bars indicate $\pm$ SEM.

Figure 2.

A. Left panel: Confocal images of K562 cells incubated with pHrodo™ Red AM and clamped with intracellular pH calibration buffer. K562 cells were incubated with 5 mM pHrodo™ Red AM for 30 minutes at room temperature. Calibration Buffer Kit (Life technologies) was used to clamp the intracellular pH with extracellular buffer at pH 5.5, 6.5, and 7.5. Right panel: Flow cytometry analysis of cytosolic pH calibration. K562-puro/LFA-1 cells were stained pHrodo™ Green AM. Calibration Buffer Kit was used to clamp the intracellular pH with extracellular buffer at pH 7.5 (blue), pH 6.5 (green), and pH 5.5 (red). Inset shows MFI value for each experimental condition. The experiment was performed three independent times. Representative images are shown. Error bars indicate $\pm$ SEM.

B. Left panel: Confocal images of K562-puro/LFA-1 cells stained with pHrodo™ Red AM (red). The images demonstrate cytosolic acidification in K562-puro/LFA-1 cells after 0, 15, and 30 min treatment with 100 ng/ml LtxA. Right panel: Flow cytometry analysis of K562-puro/LFA-1 cells stained with pHrodo Green AM after treatment with 100 ng/ml LtxA for 0, 15, and 30 min. Inset shows MFI value for each experimental condition. The experiment was performed three independent times. Representative images are shown. Average lysosomal fluorescence intensities per cell calculated in 50 cells at different time points are shown on the right.

C. The experiment as described in Fig. 2B performed on K562-puro cells. Scale bar = 10 μ m.

Figure 3.

A. The confocal images of live K562-puro/LFA-1 cells treated with acridine orange which stains cellular lysosomes red. The cells stained with acridine orange were untreated, treated with 100 ng/ml LtxA, 1 mM Leu-Leu-OMe, or 100 ng/ml HI-LtxA for 30 min and imaged. Representative images are shown. The lysosomal staining intensity was observed in K562-puro/LFA-1 cells during 60 min of treatment with 100 ng/ml LtxA using the time lapse confocal microscopy. Average lysosomal fluorescence intensities per cell calculated in 50 cells at different time points are shown on the right.

B. The confocal images of live K562-puro cells stained with acridine orange and treated as described above.

C. Lysosomes in K562-puro/LFA-1 cells were stained with LysoTracker® Red DND-99 (red) and nuclei were stained with Hoescht 33342 (blue). The lysosomal integrity was observed in the cells during 60 min after treatment with 100 ng/ml LtxA using the time lapse confocal microscopy. Representative images collected at 0, 17, 32, and 60 min of the treatment with LtxA are shown.

D. The experiment as described in Fig. 2B was performed on K562-puro cells. Representative images at 0 and 30 min of treatment with LtxA are shown. Average red fluorescence intensity per cell calculated in 50 cells are shown on the right. Error bars indicate $\pm$ SEM. * represents significant difference ($p < 0.05$) compare to untreated cells (t=0) lysosomal intensity. Scale bar = 10 μ m.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript