Published in final edited form as:

Int J STD AIDS. 2014 August ; 25(9): 669–675. doi:10.1177/0956462413516300.

The Etiology of Vaginal Symptoms in Rural Haiti

Claire C. Bristow, MSc¹, Tania Desgrottes, MD², Lauren Cutler, BA², David Cutler, MD¹,
Karthika Devarajan, MD², Oksana Ocheretina, PhD^{3,4}, Jean William Pape, MD^{3,4}, and Jeffrey
D. Klausner, MD MPH¹

¹University of California Los Angeles, Los Angeles CA, USA

²Dr. Henri Gerard Desgranges Foundation, Petit Goave, Haiti

³Weill Cornell Medical College, New York, NY, USA

⁴Les Centres GHESKIO, Port-au-Prince, Haiti

Abstract

Background—Vaginal symptoms are a common chief complaint amongst women visiting outpatient clinics in rural Haiti.

Methods—A systematic sample of 206 consecutive females over age 18 with gynecological symptoms underwent gynecologic examination and laboratory testing for chlamydia, gonorrhea, syphilis, HIV infection, trichomoniasis, candidiasis, and bacterial vaginosis.

Results—Among 206 women, 174 (84%) presented with vaginal discharge, 165 (80%) with vaginal itching, 123 (60%) with vaginal pain or dysuria, and 18 (9%) with non-traumatic vaginal sores or boils. Laboratory results were positive for *Chlamydia trachomatis* in 5.4% (11/203), syphilis in 3.5% (7/202), HIV in 1.0% (2/200), and *Neisseria gonorrhoeae* in 1.0% (2/203). Among those that had microscopy, hyphae suggestive of candidiasis were visualized in 2.2% (1/45) and no cases of trichomoniasis were diagnosed 0% (0/45). Bacterial vaginosis was diagnosed in 28.3% (13/46). The prevalence of chlamydia was 4.9 (95% CI: 1.3-17.7) times greater among those 25 years of age and under (10.8%) than those older (2.3%).

Conclusions—Chlamydia and bacterial vaginosis were the most common sexually transmitted infection and vaginal condition, respectively, in this study of rural Haitian adult women. The higher risk of chlamydia in younger women suggests education and screening programs in young women should be considered.

Keywords

Vaginitis; Haiti; Sexually Transmitted Infections; vaginal symptoms; International STIs

Author for correspondence: Claire C. Bristow, MSc, **Address:** UCLA Center for World Health, David Geffen School of Medicine, 10833 Le Conte Ave., CHS 13-154, Los Angeles, CA 90095, **Fax Number:** 310-825-3157, **Phone:** (415) 250-2681, ccbristow@gmail.com.

Disclosures include: No conflicts of interest exist in this study.

BACKGROUND

Manifestations of vaginal and sexually transmitted infections (STIs) are common in both rural and urban Haiti. Estimates of the prevalence of at least one STI among women attending antenatal services was shown to be 40% in data collected from 1996 in a rural region of Haiti and 47% in urban areas^{1,2}. A more recent analysis of programmatic data found that 11.3% (118 of 1042) of all female patients presenting for health care had vaginal symptoms, reportedly related to candidiasis, bacterial vaginosis, or STIs³.

Presentation of vaginal symptoms can have various etiologies that may include candidiasis, bacterial vaginosis, and STIs. The accurate diagnosis of vaginal infections and conditions can be complex, requiring multiple, sophisticated laboratory tests, without which the prevalence of these infections is unknown. Diagnostic tests may be constrained by limitations in laboratory and logistic support, particularly in resource-limited settings with a high STI burden. Failure to diagnose and cure persons with STIs is associated with patient morbidity and mortality and facilitates the continued transmission of infections to partners and from mother to fetus during pregnancy or at birth^{4,5}.

STIs and vaginal infections are related to many adverse health outcomes. Infections of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* are related to infertility, pelvic inflammatory disease, chronic pelvic pain, and ectopic pregnancy⁶. Syphilis infection and herpes simplex virus type 2 may facilitate human immunodeficiency virus (HIV) acquisition and transmission; genital ulcers interfere with the natural mucosal and epithelial barriers and cause inflammation providing a portal of entry for HIV⁷. In HIV co-infected patients, syphilis can increase transmission of HIV by increasing viral shedding and seminal viral load⁸. Furthermore, syphilis infection during pregnancy can lead to adverse birth outcomes, congenital syphilis, and greatly increases the risk of mother-to-child HIV transmission^{9,10}. Bacterial vaginosis increases the risk for the acquisition of infections including herpes simplex virus type 2, *Trichomonas vaginalis*, *Neisseria gonorrhoeae*, and *Chlamydia trachomatis*^{11,12,13}. In addition, bacterial vaginosis has also been associated with an increased risk of human immunodeficiency virus (HIV) acquisition and transmission^{14,15}.

Many STIs can be asymptomatic and can have long-term adverse effects if left untreated, but are not diagnosable without screening tests¹⁶. The current World Health Organization guidelines recommend syndromic management;¹⁷ yet, syndromic management is unlikely to enhance STI control¹⁸. Early identification and subsequent treatment of STIs is paramount to achieving an effective reduction in the disease burden¹⁹. In order to inform clinical management protocols and to identify pathogens for infection control interventions, the current study aimed to describe the distribution of etiologies of vaginal symptoms in Haitian women at a primary care center in rural Haiti.

MATERIALS AND METHODS

Study Design

This was a cross-sectional study designed to determine the etiology of vaginal symptoms among rural women in Haiti.

Setting

This study was completed at the Dr. Henri Gerard Desgranges Foundation clinic in Petit Goâve, a rural, under-served coastal town located 70 kilometers west of Haiti's capital, Port-au-Prince (www.hgdfoundation.org). The Dr. Henri Gerald Desgranges Foundation has been working in the Petit Goâve Community since 1994. It provides care through its outpatient clinic and immunization and prevention campaigns via its mobile health program.

Sample selection and study design

The systematic sample consisted of 206 consecutive females over age 18 complaining of vaginal itching, discharge, pain, or lesions or associated urinary symptoms presenting to the Henri Gerald Desgranges foundation clinic in March 2013. This was a period of increased clinical services and efforts were made in the community to encourage women with symptoms to seek care. All participants provided verbal informed consent.

All women underwent gynecologic examination including speculum and pelvic examination as well as specimen collection for laboratory analyses, by trained study clinicians. Data were collected on the women's pregnancy history, methods of contraception they had ever used including condoms, and current vaginal symptoms.

Laboratory Methods

Point-of-care testing was conducted in this study to decrease the time to treatment for some infections. Testing included Determine TP assay for syphilis (Abbott Laboratories, Tokyo, Japan) and the OraQuick ADVANCE® Rapid HIV-1/2 Antibody Test for HIV (OraSure Technologies, Bethlehem, Pennsylvania). Microscopic analyses for trichomonas, hyphae, and clue cells using saline and KOH mounts were used to diagnose trichomoniasis, candidiasis, and bacterial vaginosis, respectively. Bacterial vaginosis was defined using the Amsel criteria: at least 3 of elevated vaginal pH, increased vaginal discharge volume, positive wet mount for clue cells or positive whiff test²¹. The vaginal pH was determined using Hydriion pH test paper (Micro Essential Laboratory Inc, Brooklyn, New York, USA). Microscopy was only available for the first 1.5 days of the study period before the microscope malfunctioned and became unusable. Laboratory testing for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* was performed with the GeneXpert® CT/NG assay (Cepheid Inc., Sunnyvale, CA). Testing was encouraged for all women; however some participants refused HIV and syphilis testing.

As part of routine clinical gynecological examinations, vaginal swab specimens of the posterior and lateral vaginal walls were collected by the examining study clinician. Vaginal swabs were placed into specimen transport tubes, labeled with the unique study identifier of the participants, and stored for five days at ambient temperature. The specimens were then transported to l'Institut des Maladies Infectieuses et de Santé de la Reproduction clinical laboratory in Port-au-Prince, Haiti, where testing was performed using the GeneXpert® CT/NG assay. Test results were reported within 2 weeks to the clinic manager who contacted study participants with positive results who had not been treated empirically. Contacted study participants were treated and partners managed according to US CDC guidelines²².

Ethics

This study was reviewed and approved by the Institutional Review Board of the University of California Los Angeles.

Statistical Analysis

Descriptive statistics were used to summarize the prevalence of infections and risk factors among the study population. Basic comparative analyses using Cochran-Mantel-Haenszel tests between groups were used to identify statistically different levels of effect. Significance was defined at a p-value of less than or equal to 0.05. Data were entered from a paper form into a database using Epi Info 3.5.4 (CDC, Atlanta, GA, USA) and analyses were conducted using SAS software v9.3 (SAS Institute Inc., Cary, NC, USA).

RESULTS

Characteristics of participants

Between March 18th and March 22nd 2013, 206 women were enrolled in the study. The mean age in years of the women was 31.9 (SD 10.6) and 123 (61%) reported one or more pregnancies. Most participants (n=189, 92%) reported no condom use for contraception.

Patient symptoms and exam findings

The 206 women had at least one current vaginal symptom, including vaginal discharge (n=174, 84.5%), vaginal itching (n=165, 80.1%), vaginal pain or dysuria (n=123, 59.7%), and non-traumatic vaginal sores or boils (n=18, 8.7%). Clinical gynecological examinations revealed vaginal discharge in 64 women (32.0%) and cervical exudate in 5 women (2.0%).

Laboratory results

Among the 203 women with test results, 11 (5.4%) were positive for *Chlamydia trachomatis* and 2 (1.0%) were positive for *Neisseria gonorrhoeae*; both cases of gonorrhea infection occurred in women 25 years of age and under [see Figure].

Syphilis tests were positive in 3.5% (7/202) of the women. Of the 206 women, 19 (9.2%) were found to have at least one curable STI, chlamydia, gonorrhea or syphilis. Among the 200 tested 2 (1%) were found to be HIV positive.

Of those that had microscopy, hyphae suggestive of candidiasis were visualized in 2.2% (1/45) and bacterial vaginosis was diagnosed in 28.3% (13/46). Trichomoniasis was not diagnosed in any of the women tested (0/45).

Bivariate analysis

Age category had an observed effect on chlamydial and syphilis infections [see Table]. While chlamydia occurred primarily in those under 25 years of age, syphilis infection tended to occur in those over 25 years of age. Of the 11 chlamydia cases, 72.7% (n=8) were in women 25 years of age and younger, resulting in a risk of *Chlamydia trachomatis* in the younger women 4.9 (95% confidence interval: 1.3-17.7) times that of women over 25 years of age. Of the 7 cases of syphilis, 85.7% (n=6) occurred in women over 25 years of age. No

(0%) STIs were diagnosed in the 17 women that reported ever using condoms for contraception.

DISCUSSION

In this cross-sectional study of women in rural Haiti with vaginal symptoms, approximately 1 in 11 women had a curable STI and more than 3 times that had bacterial vaginosis or candidiasis. Chlamydial infection was the most common STI and bacterial vaginosis was the most common vaginal condition.

The higher risk of *Chlamydia trachomatis* in younger women is consistent with the epidemiology of the infection demonstrated in other studies²³. Conversely, those who tested positive for syphilis tended to be those over 25 years of age, in part, perhaps, because treponemal tests, such as the one used in this study, don't differentiate between past and current infections.

Syphilis infections are concerning in this setting because of the propensity of syphilis to adversely affect pregnancy. In syphilis-infected pregnant women, adverse birth outcomes are common and have been shown to be 4.5 times higher in those with untreated syphilis than those without syphilis²⁴. If syphilis remains untreated during pregnancy, it can lead to fetal loss or stillbirth or, in a live-born infant, neonatal death, prematurity, low birth weight, and congenital syphilis^{24,25}. It has been demonstrated that congenital syphilis rates in rural Haiti were among the highest in the world at 550 per 100,000 live births in 1998, highlighting the need for syphilis control strategies targeted to pregnant women and their partners¹. Congenital syphilis can be prevented by screening early in pregnancy, treating seropositive pregnant women, and preventing re-infection^{24,25}.

The high numbers of previously undiagnosed STIs observed in this study can continue to be transmitted in the community to sex partners unless treated. Failure to identify and cure persons with STIs impacts patient health, fertility, vulnerability to other infections such as HIV, and facilitates continued transmission⁵. In this study we found 2% of participants tested positive for HIV, which is consistent with Haitian national findings from a demographic health survey conducted in 2005–2006 where 2.3% of women aged 15 to 49 were HIV-positive (2.0% among males)²⁶.

The age-associated infections suggest education and age-targeted screening programs should be considered. While screening programs are warranted in the developed world to prevent adverse health outcomes, in developing countries there may be competing priorities, so further research is needed to determine the utility and cost effectiveness of such interventions for the most severely resource-limited settings.

The study had several limitations. The women included in the study were those that chose to seek healthcare for their vaginal symptoms, therefore our findings may not be representative of the overall local population. In addition, this study was done during a time of increased service delivery and women with symptoms were encouraged to seek services, therefore the findings may not be representative of the frequency of cases or infections during other clinic operating periods. However, there is strength in the systematic sampling method used to

enroll participants. The moderate sample size made it difficult to conduct conclusive between group comparisons. Additionally, the equipment malfunction made microscopy possible for only the first 1.5 days of the study, resulting in smaller sample sizes for those conditions diagnosed microscopically. Fortunately, many of the diagnostic tests did not require microscopy and we were able to test most patients for syphilis, gonorrhea, chlamydia, and HIV infection. The strengths of this study lay in the use of state-of-the art diagnostic tests for most infections using a combination of point-of-care tests and nucleic acid amplification test methods.

Although the World Health Organization recommends syndromic management for STI control in resource limited settings, this is not always effective as a public health strategy because of the high rate of asymptomatic STIs and the varying etiology of vaginal symptoms^{17,18}. Targeted screening programs, for improved clinical management, can be used to help mitigate STI epidemics and to reduce the burden of other vaginal infections. For optimal, comprehensive sexual and reproductive health diagnostics, a combination of tests are needed. Laboratory-based testing is not feasible in some resource-limited settings. Fortunately, tests that do not require extensive laboratory infrastructure including complex specimen collection and transport already exist for HIV infection, syphilis, bacterial vaginosis, trichomonas, and now most recently for chlamydia and gonorrhea as well. For *Chlamydia trachomatis* and *Neisseria gonorrhoeae*, the advent of on-site diagnostics with nucleic acid amplification tests has brought the technology of the laboratory to the clinic²⁷. For HIV and syphilis, rapid screening tests can be used to get a result within 20 minutes with a finger-prick blood test^{28,29}. Microscopy can be conducted in a clinical setting to identify infections of bacterial vaginosis, candidiasis, and trichomoniasis^{4,21,29}. In addition, test developers have created point-of-care or near-care screening tests that can detect multiple STIs with the same specimen using a single device, including the test used for detection of *chlamydia trachomatis* and *Neisseria gonorrhoeae* in this study, the GeneXpert® CT/NG. Combination tests simplify what is otherwise a complicated, costly, and labor intensive diagnostic process. These integrated, combination near-care or point-of-care tests are a major step towards the comprehensive prevention and control of STI transmission.

Conclusion

Public health strategies to prevent and control STIs should include early identification and treatment of infected individuals and their partners. The rates of infection suggest the need for better diagnostic testing including point-of-care diagnostics and improved understanding of chlamydia transmission and the impact of bacterial vaginosis is needed. Further examination is needed to better understand the efficacy of screening for STIs in limited-resource settings as a means of STI control. Additionally, studies addressing patient preferences for screening and prevention behaviors are needed. Cost effectiveness studies are necessary for the comparison of prevention efforts to estimate and compare the health and economic outcomes of alternative and additional methods. Information on preferences, efficacy, cost, and feasibility of prevention initiatives would help identify best practices for prevention, screening, and treatment to reduce the continued burden of sexual and reproductive health-related diseases in low-resource settings.

ACKNOWLEDGEMENTS

We would like to thank l'Institut des Maladies Infectieuses et de Santé de la Reproduction clinical laboratory in Port-au-Prince, Haiti for running GeneXpert CT/NG testing on the vaginal specimens, the Dr. Henri Gerard Desgranges Foundation donors, the UCLA Program in Global Health, and Dr. Daniel Fitzgerald of Weill Cornell Medical College. We would like to also thank the manufacturers of the diagnostic tests used in this study: Cepheid Inc., Abbott Laboratories, and OraSure Technologies.

This study was supported by the donation of test supplies by Cepheid, Inc. (Sunnyvale, CA, USA) and Orasure Technologies (Bethlehem, PA).

References

1. Fitzgerald DW, Behets F, Lucet C, Roberfroid D. The prevalence, burden, and control of syphilis in Haiti's rural Artibonite region. *Int J Infect Dis.* 1998; 2:127–131. [PubMed: 9531657]
2. Behets, Frieda M. -T.; Desormeaux, Julio; Joseph, Dania; Adrien, Mario; Coicou, Gessie; Dallabetta, Gina; Hamilton, Holli A.; Moeng, Stoffel; Davis, Homer; Cohen, Myron S.; Boulos, Reginald. Control of Sexually Transmitted Diseases in Haiti: Results and Implications of a Baseline Study among Pregnant Women Living in Cité Soleil Shantytowns. *The Journal of Infectious Diseases.* 1995; 172(3):764–771. [PubMed: 7658070]
3. Prentiss K, Prentiss T, Zervos T, King S, Zervos M. Short Term Medical Response in Haiti: Epidemiology and Model for Delivery of Care. *Epidemiol.* 2011 S2:002. doi: 10.4172/2161-1165.S2-002.
4. Shafii, T.; Radolf, JD.; Sanchez, PJ.; Schulz, KF.; Murphy, FK. Congenital Syphilis. Holmes, KK.; Sparling, PF.; Stamm, WE.; Piot, P.; Wasserheit, JN.; Corey, L.; Cohen, MS.; Watts, DH. Sexually Transmitted Diseases. McGraw-Hill Professional; Dec 14. 2007 Chapter 82
5. Klausner, JD. The sound of silence: missing the opportunity to save lives at birth. *WHO Bulletin;* Mar 1. 2013 Available at: <http://www.who.int/bulletin/volumes/91/3/13-118604>
6. Stamm, WE. Chlamydia trachomatis infection of the adult. Hook, EW.; Handsfield, HH. Gonococcal infections in the adult. Holmes, KK.; Sparling, PF.; Stamm, WE.; Piot, P.; Wasserheit, JN.; Corey, L.; Cohen, MS.; Watts, DH. Sexually Transmitted Diseases. McGraw-Hill Professional; Dec 14. 2007 Chapter 32 and 35
7. Boily M-C, Baggaley RF, Wang L, Masse B, White RG, Hayes RJ, et al. Heterosexual risk of HIV-1 infection per sexual act: systematic review and meta-analysis of observational studies. *Lancet Infectious Disease.* 2009; 9:118–29.
8. Quinn TC, Wawer MJ, Sewankambo N, et al. Rakai Project Study Group. Viral load and heterosexual transmission of human immunodeficiency virus type 1. *N Engl J Med.* 2000; 342(13): 921–929. [PubMed: 10738050]
9. Mwapasa V, Rogerson SJ, Kwiek JJ, Wilson PE, Milner D, Molyneux ME, et al. Maternal syphilis infection is associated with increased risk of mother-to-child transmission of HIV in Malawi. *AIDS.* 2006; 20:1869–77. [PubMed: 16954728]
10. Zetola NM, Klausner JD. Syphilis and HIV infection: an update. *Clin Infect Dis.* 2007; 44(9): 1222–8. [PubMed: 17407043]
11. Cherpes TL, Meyn LA, Krohn MA, Lurie JG, Hillier SL. Association between acquisition of herpes simplex virus type 2 in women and bacterial vaginosis. *Clin Infect Di.* 2003; 37:319–25.
12. Brotman RM, Klebanoff MA, Nansel TR, et al. Bacterial vaginosis assessed by Gram stain and diminished colonization resistance to incident gonococcal, chlamydial, and trichomonal genital infection. *J Infect Dis.* 2010; 202:1907–15. [PubMed: 21067371]
13. Wiesenfeld HC, Hillier SL, Krohn MA, Landers DV, Sweet RL. Bacterial vaginosis is a strong predictor of *Neisseria gonorrhoeae* and *Chlamydia trachomatis* infection. *Clin Infect Dis.* 2003; 36:663. [PubMed: 12594649]
14. Atashili J, Poole C, Ndumbe PM, Adimora AA, Smith JS. Bacterial vaginosis and HIV acquisition: a meta-analysis of published studies. *AIDS.* 2008; 22:1493. [PubMed: 18614873]

15. Cohen CR, Lingappa JR, Baeten JM, et al. Bacterial vaginosis associated with increased risk of female-to-male HIV-1 transmission: a prospective cohort analysis among African couples. *PLoS Med.* 2012; 9:e1001251. [PubMed: 22745608]
16. Smith Fawzi MC, Lambert W, Singler JM, Koenig SP, Léandre F, Nevil P, Bertrand D, Claude MS, Bertrand J, Salazar JJ, Louissaint M, Joanis L, Farmer PE. Prevalence and risk factors of STDs in rural Haiti: implications for policy and programming in resource-poor settings. *International Journal of STD & AIDS.* 2003; 14:848–853. [PubMed: 14678595]
17. WHO. Guidelines for the Management of Sexually Transmitted Infections. 2004 ISBN 92 4 154626 3. Available at: <http://www.who.int/hiv/pub/sti/pub6/en/>.
18. Clark JL, Lescano AG, Konda KA, Leon SR, Jones FR, Klausner JD, Coates TJ, Caceres CF. Syndromic management and STI control in urban Peru. *PLoS One.* 2009; 4(9):e7201. [PubMed: 19779620]
19. Klausner, JD. AIDS and Sexually Transmitted Disease Prevention and Control. In: Mayer, KH.; Pfizer, HF., editors. *The AIDS Pandemic: Impact on Science and Society.* Elsevier Academic Press; San Diego, CA: 2005.
20. Belotte, L.; Viel, E.; Michaud, G.; Mercier, P.; Verly, A.; Benavente, J.; Tassy, E.; Marsan, S.; Laraque, M. Ministry of Health and Population (MSPP): Guidelines and Norms for Family Planning, Maternal and Reproductive Health. National Strategy of Reproductive Health; Dec. 2010
21. Rathod SD, Krupp K, Klausner JD, Arun A, Reingold AL, Madhivanan P. Bacterial Vaginosis and Risk for *Trichomonas Vaginalis* Infection: A Longitudinal Analysis. *Sex Transm Dis.* 2011; 38(9): 882–886. [PubMed: 21844747]
22. U.S. Centers for Disease Control and Prevention. Sexually Transmitted Diseases Treatment Guidelines. *MMWR.* 2010; 13(RR-12):1–110.
23. Cuffini C, Bottiglieri M, Kiguen X, Alonso CE, Deimundo RV, Isa MB, Cannistraci R, Gonzalez S, Farinati A. Molecular Epidemiology of Genital *Chlamydia Trachomatis* Infection in Asymptomatic Adolescent-Young People. *Journal of Microbiology Research.* 2012; 2(4):114–117.
24. Gomez GB, Kamb ML, Newman LM, Mark J, Broutet N, Hawkes SJ. Untreated maternal syphilis and adverse outcomes of pregnancy: a systematic literature review and meta-analysis. *Bull World Health Organ.* 2013; 91:217–2. [PubMed: 23476094]
25. Santis MD, De Luca C, Mappa I, Spagnuolo T, Licameli A, Straface G, Scabia G. Syphilis Infection during Pregnancy: Fetal Risks and Clinical Management. *Infect Dis Obstet Gynecol.* 2012;430585. [PubMed: 22829747]
26. Cayemittes, M.; Placide, MF.; Mariko, S., et al. Enquête mortalité, morbidité et utilisation des services EMMUS–e Haïti 2005–2006. Institut Haïtien de l'Enfance, ORC Macro; Pétienville and Calverton: 2006.
27. Gaydos CA, Van Der Pol B, Jett-Goheen M, Barnes M, Quinn N, Clark C, Daniel GE, Dixon PB, Hook EW III, CT/NG Study Group. Performance of the Cepheid CT/NG Xpert Rapid PCR Test for the detection of *Chlamydia trachomatis* and *Neisseria gonorrhoeae*. *J. Clin. Microbiol.* Jun; 2013 51(6):1666–1672. doi: 10.1128/JCM.03461-12. [PubMed: 23467600]
28. Benzaken AS, Sabido M, Galban E, Pedroza V, Araujo AJ, Peeling RW, Mabey D. Field performance of a rapid point-of-care diagnostic test for antenatal syphilis screening in the Amazon region, Brazil. *International Journal of STD & AIDS.* 2011; 22(1):15–18. [PubMed: 21364061]
29. Tucker JD, Bien CH, Peeling RW. Point-of-care testing for sexually transmitted infections: recent advances and implications for disease control. *Curr Opin Infect Dis.* 2013; 26(1):73–9. [PubMed: 23242343]

Short summary

A study of women presenting to a rural clinic with vaginal symptoms in Haiti found several different causes of symptoms including candidiasis, bacterial vaginosis, chlamydia, and gonorrhea.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure.
Distribution of vaginal infections and conditions in the study population. Petit Goâve, Haiti, 2013 (N=206)

Select characteristics of study population and frequencies of cervical and vaginal infections and conditions, Petit Gove, Haiti, 2013.

Table

Characteristic ^{†‡}	Chlamydia (n=206), n (%)	Gonorrhea (n=203), n (%)	Syphilis (n=202), n (%)	At least 1 bacterial STI [*] (n=199), n (%)	HIV (n=200), n (%)	Bacterial Vaginosis (n=46), n (%)	Candidiasis (n=45), n (%)
Age 25 years and under	74(35.9)	8(10.8)	1(1.4)	10(13.5)	1(1.4)	5(6.8)	0(0)
Age over 25 years	132(64.1)	3(2.3)	6(4.5)	9(6.8)	1(0.8)	8(6.1)	1(0.8)
Never been pregnant	80(39.2)	4(5.0)	0(0)	5(6.3)	0(0)	5(6.3)	0(0)
At least 1 pregnancy	124(60.8)	7(5.6)	7(5.6)	14(11.3)	2(1.6)	8(6.5)	1(0.8)
No reported condom use	189(91.8)	11(5.8)	7(3.7)	19(10.1)	2(1.1)	11(5.8)	1(0.5)
Reported ever using condoms	17(8.3)	0(0)	0(0)	0(0)	0(0)	2(11.8)	0(0)
Normal vaginal discharge volume	142(68.9)	7(4.9)	4(2.8)	12(8.5)	1(0.7)	5(3.5)	1(0.7)
Increased vaginal discharge volume	64(31.1)	4(6.3)	3(4.7)	7(10.9)	1(1.6)	8(12.5)	0(0)
No cervical exudate	201(97.6)	10(5.0)	6(3.0)	17(8.5)	2(1.0)	12(6.0)	1(0.5)
Cervical exudate	5(2.4)	1(20.0)	1(20.0)	2(40.0)	0(0)	1(20.0)	0(0)
Reported sores or boils	18(8.7)	1(5.6)	1(5.6)	2(11.1)	1(5.6)	2(11.1)	0(0)
No reported sores or boils	188(91.3)	10(5.3)	6(3.2)	17(9.1)	1(0.5)	11(5.9)	1(0.5)
TOTAL, n		11(5.4)	7(3.5)	19(9.6)	2(1.0)	13(28.2)	1(2.2)

Denominators may not sum to 206 because of inadequate sample or test refusal

^{*} At least one of chlamydia, gonorrhea, and/or syphilis of those that were tested for all three[†] Includes only those that gave a response, so there may be some missing values[‡] Percentages may not sum to 100 because of rounding