Published in final edited form as:

Child Maltreat. 2016 February ; 21(1): 16–25. doi:10.1177/1077559515620853.

Characteristics of Child Maltreatment and Adolescent Marijuana Use: A Prospective Study

Howard Dubowitz,

Department of Pediatrics, University of Maryland School of Medicine, Baltimore

Richard Thompson,

Richard H. Calica Center for Innovation in Children and Family Services, Juvenile Protective Association

Amelia M. Arria,

Department of Behavioral and Community Health, University of Maryland School of Public Health

Diana English,

School of Social Work, University of Washington

Richard Metzger, and

Department of Pediatrics, University of Maryland School of Medicine, Baltimore

Jonathan B. Kotch

Department of Maternal and Child Health, Gillings School of Global Public Health, University of North Carolina at Chapel Hill

Abstract

There has been increasing acceptance of marijuana use in the US in recent years, and rates among adolescents have risen. At the same time, marijuana use during adolescence has been linked to an array of health and social problems. Maltreated children are at risk for marijuana use, but the relationships among characteristics of maltreatment and marijuana use are unclear. In this paper we examine how the type and the extent of maltreatment are related to the level of adolescent marijuana use. Data analyses were conducted on a subsample of maltreated adolescents ($n = 702$) from the Longitudinal Studies of Child Abuse and Neglect (LONGSCAN) project. Approximately half the sample had used marijuana, and maltreatment was associated with its use. Multivariate regression models showed that being male, extensive maltreatment, and peer marijuana use were associated with Heavy Use of marijuana. These findings suggest the importance of comprehensively assessing children's maltreatment experiences and their peers' drug use to help prevent or address possible marijuana use in these high-risk adolescents.

Keywords

Child maltreatment; marijuana; adolescence

Although there is a recent trend toward legalization of marijuana use among adults that might imply little concern for marijuana use as a public health issue, marijuana has been linked to several adverse health outcomes (Hall, 2014; Volkow, Baler, Compton & Weiss, 2014), including dependence (Swift, Coffey, Carlin, Degenhardt & Patton, 2008), cognitive dysfunction (Meier, et al., 2012), and impaired driving (Hartman & Huestis, 2013). Marijuana use has also been associated with mental health problems such as panic disorder (Zvolensky, Coughle, Johnson, Bonn-Miller & Bernstein, 2010), anxiety (Degenhardt et al., 2012; Kedzior & Laeber, 2014), and psychosis (Fergusson, Horwood & Ridder, 2005; Moore, 2007).

Although marijuana use as compared to non-use confers risk for various adverse outcomes in young adulthood (Ellickson, Martino & Collins, 2004), it is especially problematic when use begins early in adolescence or is frequent (Brook, Lee, Brown, Finch & Brook, 2011; Flory, Lynam, Lilich, Leukefeld & Clayton, 2004; Meier, et al., 2012; Solowij, Stephens, et al., 2002; Windle & Wiesner, 2004). Adolescent marijuana users are at risk for academic difficulties (Lynskey & Hall, 2000), with users less likely than non-users to complete high school or undertake postsecondary education, even after controlling for possible confounding variables (Homel, Thompson & Leadbeater, 2014; Horwood et al., 2010; Palamar et al., 2014). Brain imaging studies have found structural abnormalities in gray matter density, volume, and shape of the nucleus accumbens and amygdala in 18–25 year old marijuana users (Gilman et al., 2014). This finding is not unexpected given that the developing brain appears to be highly vulnerable to the effects of substance use and that the brain continues to develop into the mid-20s (Giedd, 2004). Other studies have reported a variety of neurocognitive problems among marijuana users (Hanson et al., 2010; Schweinsburg, Brown & Tapert, 2008; Solowij et al., 2011). In the longer term, chronic heavier patterns of marijuana use predict risk for a number of psychosocial outcomes including diminished relationship quality (Brook et al, 2008), greater need for economic assistance and occupational difficulties (Fergusson & Boden, 2008; Green & Ensminger, 2006; Brook, Lee, Finch, Seltzer & Brook, 2013), and lower satisfaction with life (Schulenberg et al., 2005).

Despite the evidence regarding the effects of marijuana, recent national surveys have shown decreases in perceived risk concerning marijuana (Johnston, O'Malley, Bachman, Schulenberg & Miech, 2014a), owing in part to the relaxation of state laws regarding the legality of marijuana use (Schuermeyer et al., 2014). In tandem with decreasing perceived risk, recent data show increased marijuana use by adolescents. By age 18, nearly half of American adolescents report having tried marijuana (Johnston, O'Malley, Bachman, Schulenberg & Miech, 2014b). In 2013, 46% of high school seniors had smoked marijuana at some point, making it the most commonly used illicit drug among youth (Johnston et al., 2014a).

Among the research studies on risk and protective factors associated with marijuana use, several have focused on the impact of child maltreatment, including physical and sexual abuse as well as neglect. Adults who retrospectively report having been maltreated as children are at elevated risk for drug involvement compared to individuals who have not reported maltreatment (Green et al. 2010; Wu, Schairer, Dellor & Grella, 2010; McLaughlin

et al., 2010, Kilpatrick et al., 2000). Moreover, prospective studies have found child maltreatment to be associated with drug use during adolescence (Oshri, Rogosuch, Burnette & Cicchetti, 2011; Horan & Widom, 2015) and young adulthood (Widom, Marmorsetin & White, 2006; Lo & Cheng, 2007; Lansford, Dodge, Pettit & Bates, 2010; Thornberry et al., 2014). Wilson & Widom (2009) observed significant relationships between child abuse and neglect and subsequent illicit drug use in young adult women, but not men. Maltreatment has been identified as a significant predictor of cannabis dependence (Handley, Rogosch, Guild & Cicchetti, 2015; Green et al., 2010), even after adjustment for genetic vulnerability (Duncan et al., 2008).

Child maltreatment is not, however, always associated with drug use, raising the question of what explains the variability in outcomes. First, the timing of maltreatment has been suggested to predict psychosocial and health outcomes. One study found that maltreatment during adolescence, as opposed to early childhood, was associated with substance use and other maladaptive behaviors (Thornberry, Ireland & Smith, 2001). Oshri et al. (2012) reported that the severity of maltreatment in early childhood was related to adolescent marijuana abuse and dependence, via poor adaptation in childhood and externalizing problems.

Dodge and colleagues (2009) described a dynamic cascade model of the development of early-onset drug use that supports the aggravating effect of early maltreatment. In contrast, other research has suggested that contemporaneous or recent maltreatment might be more predictive of substance use in adolescence and young adulthood. It has been suggested that adolescence is a period of heightened vulnerability to stress (Jankord et al., 2011; McCormick, Mathews, Thomas & Waters, 2010), which might contribute to their risk taking behavior (Steinberg, 2010). Persistent maltreatment that occurs during both childhood and adolescence might compound the risk for subsequent drug use (Ireland, Smith & Thornberry, 2002)

Second, the type of maltreatment might also be a factor in the variation in drug use outcomes (Moran, Vuchinich & Hall, 2004; Bailey & McCloskey, 2005). Neglect is the most common type of maltreatment reported to Child Protective Services (CPS). Of the 679,000 substantiated reports in 2013, 82% of the children were neglected, 18% were physically abused, 9% were sexually abused and 9% were psychologically maltreated (U.S. Department of Health and Human Services, 2013). Just as different types of maltreatment have been associated with different child outcomes (Lau et al., 2005), physical abuse appears to be especially associated with drug use (Huang et al., 2011; Lo & Cheng, 2007; Wilson & Widom, 2009;), but others found significant relationships with both physical and sexual abuse, but not neglect (Appleyard, Berlin, Rosanbalm & Dodge, 2011). In contrast, Cheng and Lo (2010) found that sexual abuse, but not physical abuse or neglect, predicted cocaine, crack and heroin use. In one of their studies, no relationship was found between childhood maltreatment and marijuana use after controlling for marijuana use prior to official CPS reports, parental closeness and parent monitoring (Cheng & Lo, 2011). The influence of peers on adolescents' drug use (Glaser, Shelton & van den Bree, 2010; Monahan, Rhew, Hawkins & Brown, 2014; Tucker, de la Haye, Kennedy, Green & Pollard,

2014) requires taking this into account when examining the influence of child maltreatment on later drug use.

Despite strong support for the overall link between maltreatment and drug use, methodological limitations of previous research leave important gaps in our understanding of those characteristics of maltreatment that influence drug use. Research to date offers a mixed, uncertain picture of how characteristics of maltreatment influence adolescent drug use. Our goal in this multisite prospective study of children at risk of or reported for maltreatment was to examine whether the type, extent and timing of their maltreatment experiences were associated with the level of their marijuana use during adolescence. To overcome some of the limitations of prior research, we utilized data from a large, multi-wave and multi-site prospective study of high-risk children.

The current study design affords the opportunity to fill important gaps in prior research on maltreatment and marijuana use. As indicated earlier, little research has focused on the relationship between maltreatment and marijuana use, and it has typically relied on retrospective recall in adults, failed to examine characteristics of maltreatment (e.g., Khoury, Tang, Bradley, Cubells & Ressler, 2010), or examined outcomes in adulthood (Fergusson et al., 2008; Huang et al., 2011). In particular, prior research has rarely examined all types of maltreatment as predictors; in particular, the role of neglect has been neglected (e.g., Khoury et al., 2010). Several studies have examined the relationship between maltreatment and “substance use” as a more general class, without focusing on marijuana (Fergusson et al., 2008; Moran et al., 2004).

Method

Sample

The analytic sample was derived from the Longitudinal Studies of Child Abuse and Neglect (LONGSCAN), a consortium of five prospective studies of the antecedents and consequences of child maltreatment (Runyan et al., 1998). Although each subsample consisted of families at high risk for maltreatment as indicated by their potential or current involvement with CPS, they were deliberately varied to represent those at risk for maltreatment or already identified as maltreated. Participants were included in this sample if there were data from at least one interview indicating marijuana use or data at 18 that they had never used marijuana. Of the original LONGSCAN sample of 1354 children, 1041 (77%) met the inclusion criteria. Excluded participants were more likely to have been maltreated (73.5%) than those included (67.4%), $X^2(1) = 4.10, p = .04$. Excluded participants were more likely to be White (30.4%) and less likely to be Hispanic (16.0%) or Black (44.9%) compared to those included, 24.9%, 12.6% and 55.8% respectively ($X^2(3) = 11.61, p = .009$). LONGSCAN included children who had been maltreated as well as others who were not. Analyses were finally conducted on the maltreated subsample ($n = 702$).

Procedure

The consortium began common data collection when the children were 4 to 6 years old and gathered data from the caregivers and children every two years until 18 years of age.

(Runyan et al., 1998). CPS data were obtained by each site from the local or state agency at regular intervals. Each site employed procedures approved by its local Institutional Review Board. Families were paid a nominal amount for their participation to compensate for their time and travel.

Measures

Maltreatment—A participant was defined as maltreated if the review of CPS data reports reviewed by the LONGSCAN project (Runyan et al., 1998) indicated one or more reports from birth to 18 years of age. English, Bangdiwala & Runyan (2005) describe in detail the identification of maltreatment from CPS records. In sum, trained observers at each site reviewed CPS records every 2 years and collected information about allegations of abuse and neglect. These allegations were categorized according to a modified version of the Maltreatment Classification Scheme (Barnett, Manly & Cichetti, 1993). The decision to use allegations instead of substantiated reports was based on earlier research on LONGSCAN participants. Hussey et al. (2005) found that children with substantiated and unsubstantiated CPS reports had similar behavioral and developmental outcomes. Other researchers have also suggested that there is not a useful distinction between substantiated and unsubstantiated cases (Kohl, Jonson-Reid, & Drake 2009).

Characteristics of Maltreatment—Maltreatment experiences were further characterized for those in the maltreated subsample ($n = 702$) (Table 2). The Type of maltreatment was classified as Physical Abuse, Sexual Abuse, Emotional Maltreatment, or Neglect (English, Graham, Litrownik, Everson & Bangdiwala, 2005). Each type of maltreatment was coded dichotomously, and children could be classified as having experienced more than one type of maltreatment.

The Chronicity of maltreatment was defined by the number of developmental periods during which at least one report was made: 0 – 5, 6 – 11, and 12 – 18. The Frequency of maltreatment was defined as either low (1 – 5 reports) or high (6 or more) between the ages of 0 and 18 years. Chronicity and Frequency as well as Timing of CPS reports were highly correlated, so we created a summary variable – Extent of child maltreatment. Individuals with Limited Maltreatment (44%, $n=309$) were defined as those who had CPS reports in only one developmental period, and had low Frequency of reports (mean = 2.0). Those who experienced Extensive Maltreatment (56%, $n= 393$) were maltreated in more than one developmental period (median = 2) or had high Frequency of CPS reports (mean = 7.2).

Peer Marijuana Use—At ages 12, 14, and 16, participants were asked whether any of their friends used marijuana ($n = 911$). If at any interview participants indicated Yes, they were counted as having had Peer Use ($n = 437$, 48%). If during at least two interviews participants indicated that their friends did not use marijuana, (and never described their friends as using), they were counted as No Peer Use ($n = 474$, 52%).

Level of Adolescent Marijuana Use—Participants were first classified as having used marijuana if use was reported at any of the age 12, 14, 16, and 18 interviews. Non-use was defined by indicating at 18 that they had never used marijuana (Never Used Marijuana).

Adolescents who never indicated marijuana use and who did not complete the age 18 interview were considered to have missing data and were excluded. Those who reported marijuana use were further categorized by level of use. Heavy use was defined as reports in at least one interview of 1) use on more than 20 days in the last year or 2) use more than 10 times in the last month or 3) meeting criteria for Substance Dependence or Abuse per the National Institute of Mental Health Diagnostic Interview Schedule for Children version IV at age 14 or 18.

Data Analyses

The goal of the present study was to understand the maltreatment characteristics that predicted the level of marijuana use during adolescence. A correlation matrix including the maltreatment variables as well as peer use (see Table 3) shows that the correlations were modest to moderate, ranging from -.09 to .48. Thus, multicollinearity was of minimal concern and allowed us to develop a model that included multiple dimensions of childhood maltreatment. Multinomial logistic regression analyses were conducted, with Site and Race/Ethnicity as covariates. Site included the 5 research centers with Northwest as the reference. Race/ethnicity was coded in four categories (Black, Hispanic, Other and White), with White as the reference group. Peer Use was used as a covariate in some models with No Peer Use as the reference. In a preliminary analysis of the covariates as predictors of Level of Marijuana Use, Participant Sex was found to be significantly related to Level of Use and therefore was included in each model. Next, a model was developed that included all variables except Peer Use (Model 1) followed by a model that included Peer Use (Model 2).

Results

Child maltreatment was associated with marijuana use. Specifically, a greater percentage of maltreated youth reported any marijuana use (53%) compared to those who had not been maltreated (36%, $p = <.001$). Maltreated youth were more likely to report both Some Use (31% vs. 22%) as well as Heavy Use (22% vs. 14%). Table 1 summarizes characteristics of the maltreated subsample ($n = 702$). Females were less likely than males to engage in Heavy Use (16% vs. 28%, $p = .001$). Marijuana use did not differ significantly according to race/ethnicity or study site.

The bivariate relationships between maltreatment characteristics and marijuana use are shown in Table 2. Regarding Maltreatment Type, Physical Abuse was significantly associated with Heavy Use of marijuana, and Sexual Abuse primarily with Some Use. More Extensive child maltreatment (CM) was significantly associated with Heavy Use. Emotional maltreatment and neglect did not predict marijuana use. The Tolerance ranged from .63 to .99 and the VIF ranged from 1.01 to 1.44, indicating that multicollinearity among the maltreatment variables was of little concern.

All of the characteristics of maltreatment and covariates were jointly examined (Table 4). In Model 1, Sexual Abuse predicted Some Use of marijuana, and Extensive CM was associated with Heavy Use. Females were more likely than males to be reported for sexual abuse (38% vs. 17%), $X^2(1) = 40.35$, $p < .001$, and they are more likely to have Some Use than Heavy Use. When Peer Use was added in the regression (Model 2), it is strongly predictive of

marijuana use, and being male and having experienced Extensive CM also are associated with Heavy Use of marijuana.

Discussion

This study adds important information to our understanding of the characteristics and predictors of marijuana use in vulnerable populations of children who are at-risk or reported for maltreatment. Although not generalizable to the broader US population of adolescents, these data are valuable because “at-risk and/or maltreated adolescents have an increased likelihood of using illicit drugs” (Braciszewski & Stout, 2012). As such, studying marijuana use within such high-risk populations has the potential to better elucidate the relative contributions of maltreatment and other risk factors such as peer use. As well, reported maltreatment is associated with a host of risk behaviors, and the pattern of predictors for each risk behavior is somewhat different (Courtney & Dworsky, 2006).

This study is consistent with earlier research and extends it by providing information about the effects of different characteristics of reported maltreatment and their relative influence in the context of peer use. Maltreatment was associated with marijuana use in bivariate models, but did not retain significance in models that adjusted for peer marijuana use. In terms of type of maltreatment, sexual abuse predicted marijuana use. This finding is consistent with earlier research (Fergusson Boden & Horwood, 2008; Moran et al., 2004) which found that physical abuse and sexual abuse (and these combined) predicted substance abuse or dependence, although not specifically marijuana use. Physical abuse has also been found to predict earlier onset of substance use in young adolescents (Lansford et al., 2010). Those who experienced Extensive maltreatment were more likely to report Heavy Use of marijuana. This finding is consistent with research suggesting the importance of both early (e.g., Dodge et al., 2009) and recent (i.e., adolescent) exposure to maltreatment (e.g., Thornberry Ireland & Smith, 2001). It appears likely that persistent exposure to maltreatment carries an especially high level of risk for adolescent substance use (Thornberry et al., 2014). Indeed, Extensive CM was the one characteristic that significantly predicted adolescent marijuana use, after taking into account demographic variables and peer use.

Future research should focus on better understanding the mechanisms by which physical and sexual abuse influence youth adolescent risk taking behaviors in general and marijuana use in particular. It is likely that adolescent (and earlier child) emotional functioning is a key contributor, as it has been shown to influence other effects of child maltreatment on risk behaviors (e.g., Oberlander et al., 2011; Thompson & Neilson, 2014). However, more detail is needed in terms of the specific pathways from particular characteristics of maltreatment, to particular deficits in emotional or psychological functioning, to risk for marijuana use. As well, it is likely that for many adolescents, marijuana use is only one of a cluster of risk behaviors that are part of a larger syndrome of risk-taking (Monshouwer et al., 2012). From this perspective, peer use may have a more complex relationship with marijuana use than elucidated here (Monahan et al., 2014). Peer use appears to mitigate or possibly mediate the effects of reported maltreatment in these analyses. However, in terms of understanding this relationship, it is likely that maltreatment is one link in a chain that comprise a risk

“caravan” for use of illicit drugs (Layne, Warren, Watson & Shalev, 2007). The social development model described by Catalano and colleagues (1996) incorporating the effects of empirical predictors, or risk factors, in the development of antisocial behavior, may help illustrate the roles peers played in this study. The model also incorporates protective factors for example, peers not using marijuana may moderate the effects of child maltreatment. This potential protective role was not evident in this study.

The results of the current study are largely consistent with several previous investigations showing an increased risk of problematic marijuana use among maltreated children as compared to their non-maltreated counterparts (Duncan, 2008; Handley, 2015; Lo & Cheng, 2007; Oshri, 2011; Widom, 2006; Wright, 2013). This was true regardless of whether youth were sampled from the general population or identified from administrative records, and whether marijuana problems were operationalized utilizing a standardized instrument such as the DISC or a frequency measure such as in the current study. Findings regarding the dimensions of maltreatment that are related to marijuana use are mixed, with some studies observing a significant relationship with sexual abuse (Duncan, 2008) and others with physical abuse (Lo and Cheng, 2007). The current finding that maltreatment that occurred in multiple developmental periods is more strongly predictive of marijuana use than limited exposure to maltreatment supports prior research findings indicating that the severity or chronicity of maltreatment might have greater explanatory value rather than the type of maltreatment (Oshri, 2011; Lo and Cheng, 2007). The present study also highlights the utility of the ecological-transactional framework as a model for understanding the complex relationship between childhood maltreatment and substance use (Oshri, 2011), where the interaction between individual-level factors with characteristics of the broader environment can increase the risk of expression of various adverse outcomes of maltreatment. The current study revealed the importance of peer drug use on risk for marijuana use frequency and found that the prospective association between the extent of maltreatment and marijuana use was robust even after including peer drug use. This finding extends what was observed by Huang (2011), where both maltreatment and peer drug use were significantly associated with drug-related problems. Future research is needed to attain an even more complete picture of the various mechanisms by which child maltreatment exerts its adverse effects. For example, Handley (2015) observed that neighborhood disadvantage conferred risk for marijuana dependence symptoms more strongly for maltreated children than non-maltreated children.

In interpreting these findings, there are several limitations that should be kept in mind. First, it is important to acknowledge that the assessment of maltreatment based on CPS reports has clear limitations including bias as to who gets reported and investigated. As well, the Extent of maltreatment is important to consider, although the chronicity component used relatively broad developmental periods (0–5, 6–11, 12–17). Despite the usefulness of these periods in prior LONGSCAN research (e.g., Flaherty et al., 2009; Thompson et al., 2012), future work should examine this in a more nuanced way (English, Graham, Litrownik, Everson & Bangdiwala, 2005). Similar refinements could be made in the measurement of other variables. For example, missing data on peer marijuana use was addressed by coding those with no peer use at at least two time points as “no peer use,” which was an imperfect, but necessary, compromise. Given the uncertain chronology of events, it was not possible to

ascertain whether predictors directly triggered the use of marijuana, or whether there were other intervening factors.

With these caveats, this study's findings, especially given earlier research, strongly suggest that children exposed to physical and sexual abuse are at increased risk of marijuana use. Child-serving professionals should begin to integrate the findings from this and other studies into discussions about effective interventions to decrease the likelihood of vulnerable populations engaging in illicit drug use – in this case, marijuana. Prior research suggests the importance of primary prevention and early intervention. Adolescents who use substances have been found to increase their use during adolescence (Brown et al., 2008; Maggs & Schulenberg, 2005). This may be part of a risk-taking or emotional coping approach common among adolescents with maltreatment experiences (Thompson et al., 2011). Alternatively, recent research with veterans has highlighted the possibility that use of marijuana may be a form of self-medication of some symptoms of post-traumatic stress disorder (Bonn-Miller, Vujanovic & Drescher, 2011), and our findings are consistent with this possibility as well. In any case, earlier studies have found that maltreatment and other adverse childhood experiences (ACEs) increase the likelihood of future substance use well into adulthood (Aarons et al., 2008; Fergusson et al., 2008; Dube et al., 2003; Huang et al., 2011). Therefore, it is important to disrupt these potential trajectories at the earliest possible stage.

Since many adolescents who have experienced maltreatment and use substances are involved in the child welfare system, these findings have both policy and practice implications. The findings emphasize the importance of both comprehensive and accurate assessments including the Extent of a child's maltreatment as well as possible peer marijuana use. Such assessments are necessary to guide effective prevention and/or intervention for adolescents already using marijuana or considering its use. Maltreated males are at particularly high risk for Heavy Use of marijuana, and likely require added support, interventions and monitoring. There are also opportunities for screening and early intervention to detect and address marijuana use and related problems for maltreated adolescents in the child welfare system, particularly when there are such opportunities for children in Independent Living programs. Pediatric primary health care visits offer another opportunity where more frequent visits may be scheduled to support adolescents and to monitor their health risk behaviors.

Acknowledgments

This research was funded by grants from the Office of Child Abuse and Neglect, Administration on Children and Families, US DHHS (Grants Nos. 90CA1401, 90CA156901, 90CA1681, and 90CA1749), and the National Institute on Drug Abuse (Grant no. 5R01DA031189-04).

References

- Aarons GA, Monn AR, Hazen AL, Connelly CD, Leslie LK, Landsverk JA, Hough RL, Brown SA. Substance involvement among youths in child welfare: the role of common and unique risk factors. *American Journal of Orthopsychiatry*. 2008; 78(3):340–9. [PubMed: 19123753]

- Appleyard K, Berlin LJ, Rosanbalm KD, Dodge KA. Preventing early child maltreatment: Implications from a longitudinal study of maternal abuse history, substance use problems, and offspring victimization. *Prevention Science*. 2011; 12(2):139–149. [PubMed: 21240556]
- Bailey JA, McCloskey LA. Pathways to adolescent substance use among sexually abused girls. *Journal of Abnormal Child Psychology*. 2005; 33:39–53. [PubMed: 15759590]
- Barnett, D.; Manly, JT.; Cicchetti, D. Defining child maltreatment: The interface between policy and research. In: Cicchetti, D.; Toth, SL., editors. *Child abuse, child development, and social policy*. Norwood, NJ: Ablex; 1993. p. 7-74.
- Bonn-Miller MO, Vujanovic AA, Drescher KD. Cannabis use among military veterans after residential treatment for posttraumatic stress disorder. *Psychology of Addictive Behaviors*. 2011; 25:485–491. [PubMed: 21261407]
- Brook JS, Lee JY, Brown EN, Finch SJ, Brook DW. Developmental trajectories of marijuana use from adolescence to adulthood: Personality and social role outcomes. *Psychological Reports*. 2011; 108(2):339–357. [PubMed: 21675549]
- Brook JS, Lee JY, Finch SJ, Seltzer N, Brook DW. Adult work commitment, financial stability, and social environment as related to trajectories of marijuana use beginning in adolescence. *Substance Abuse*. 2013; 34(3):298–305. [PubMed: 23844962]
- Brook JS, Pahl K, Cohen P. Associations between marijuana use during emerging adulthood and aspects of the significant other relationship in young adulthood. *Journal of Child and Family Studies*. 2008; 17(1):1–12. [PubMed: 23667303]
- Braciszewski JM, Stout RL. Substance use among current and former foster youth: A systematic review. *Children and Youth Services Review*. 2012; 34(12):2337–2344. [PubMed: 23504534]
- Brown SA, McGue M, Maggs J, Schulenberg J, Hingson R, Swartzwelder S, Martin C, Chung T, Tapert SF, Sher K, Winters KC, Lowman C, Murphy S. A developmental perspective on alcohol and youths 16 to 20 years of age. *Pediatrics*. 2008; 121(Suppl 4):S290–310. [PubMed: 18381495]
- Catalano RF, Kosterman R, Hawkins JD, Newcomb MD, Abbott RD. Modeling the Etiology of Adolescent Substance Use: A Test of the Social Development Model. *J Drug Issues*. 1996; 26(2):429–455. [PubMed: 17848978]
- Cheng TC, Lo CC. Drug use among maltreated adolescents receiving child welfare services. *Children and Youth Services Review*. 2010; 32(12):1735–1739.
- Cheng TC, Lo CC. A longitudinal analysis of some risk and protective factors in marijuana use by adolescents receiving child welfare services. *Children and Youth Services Review*. 2011; 33(9):1667–1672.
- Courtney ME, Dworsky A. Early outcomes for young adults transitioning from out-of-home care in the USA. *Child and Family Social Work*. 2006; 11:209–219.
- Degenhardt L, Coffey C, Romaniuk H, Swift W, Carlin JB, Hall WD, Patton GC. The persistence of the association between adolescent cannabis use and common mental disorders into young adulthood. *Addiction*. 2012; 108(1):124–33. [PubMed: 22775447]
- Dodge KA, Malone PS, Lansford JE, Miller S, Pettit GS, Bates JE. A dynamic cascade model of the development of substance-use onset. *Monographs of the Society for Research in Child Development*. 2009; 74:1–134.
- Dube SR, Felitti VJ, Dong M, Giles WH, Anda RF. The impact of adverse childhood experiences on health problems: Evidence from four birth cohorts dating back to 1900. *Preventive Medicine*. 2003; 37:268–277. [PubMed: 12914833]
- Duncan AE, Sartor CE, Scherrer JF, Grant JD, Heath AC, Nelson EC, Bucholz KK. The association between cannabis abuse and dependence and childhood physical and sexual abuse: Evidence from an offspring of twins design. *Addiction*. 2008; 103(6):990–997. [PubMed: 18482422]
- Ellickson PL, Martino SC, Collins RL. Marijuana use from adolescence to young adulthood: Multiple developmental trajectories and their associated outcomes. *Health Psychology*. 2004; 23(3):299–307. [PubMed: 15099171]
- English DJ, Bangdiwala SI, Runyan DK. The dimensions of maltreatment: introduction. *Child Abuse & Neglect*. 2005; 29(5):441–460. [PubMed: 15970319]

- English DJ, Graham JC, Litrownik AJ, Everson M, Bangdiwala SI. Defining maltreatment chronicity: Are there differences in child outcomes? *Child Abuse and Neglect*. 2005; 29:575–595. [PubMed: 15970326]
- Fergusson DM, Boden JM. Cannabis use and later life outcomes. *Addiction*. 2008; 103:969–976. [PubMed: 18482420]
- Fergusson DM, Boden JM, Horwood LJ. Exposure to childhood sexual and physical abuse and adjustment in early adulthood. *Child Abuse and Neglect*. 2008; 32:607–619. [PubMed: 18565580]
- Fergusson DM, Horwood J, Ridder EM. Tests of causal linkages between cannabis use and psychotic symptoms. *Addiction*. 2005; 100:354–366. [PubMed: 15733249]
- Flaherty EG, Thompson R, Litrownik AJ, Zolotor AJ, Dubowitz H, Runyan DK, English DJ, Everson MD. Adverse Childhood Exposures and Reported Child Health at Age 12. *Academic Pediatrics*. 2009; 9(3):150–156. [PubMed: 19450774]
- Flory K, Lynam D, Milich R, Leukefeld C, Clayton R. Early adolescent through young adult alcohol and marijuana use trajectories: Early predictors, young adult outcomes, and predictive utility. *Development and Psychopathology*. 2004; 16(1):193–213. [PubMed: 15115071]
- Giedd JN. Structural magnetic resonance imaging of the adolescent brain. *Annals of the New York Academy of Sciences*. 2004; 1021:77–85. [PubMed: 15251877]
- Gilman JM, Kuster JK, Sang L, Myung Joo L, Byoung Woo K, Makris N, Breiter HC. Cannabis use is quantitatively associated with nucleus accumbens and amygdala abnormalities in young adult recreational users. *Journal of Neuroscience*. 2014; 34:5529–5538. [PubMed: 24741043]
- Glaser B, Shelton KH, van den Bree MB. The moderating role of close friends in the relationship between conduct problems and adolescent substance use. *Journal of Adolescent Health*. 2010; 47:35–42. [PubMed: 20547290]
- Green KM, Ensminger ME. Adult social behavioral effects of heavy adolescent marijuana use among African Americans. *Developmental Psychology*. 2006; 42:1168–1178. [PubMed: 17087550]
- Green JG, McLaughlin KA, Berglund PA, Gruber MJ, Sampson NA, Zaslavsky AM, Kessler RC. Childhood adversities and adult psychiatric disorders in the National Comorbidity Survey Replication I: Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*. 2010; 67(2):113–123. [PubMed: 20124111]
- Hall W. What has research over the past two decades revealed about the adverse health effects of recreational cannabis use? *Addiction*. 2014; 110:19–35. [PubMed: 25287883]
- Handley ED, Rogosch FA, Guild DK, Cicchetti D. Neighborhood Disadvantage and Adolescent Substance Use Disorder: The Moderating Role of Maltreatment. *Child Maltreatment*. 2015 May 5. pii: 1077559515584159 [Epub ahead of print].
- Hanson KL, Winward JL, Schweinsburg AD, Medina KL, Brown SA, Tapert SF. Longitudinal study of cognition among adolescent marijuana users over three weeks of abstinence. *Addictive Behaviors*. 2010; 35(11):970–976. [PubMed: 20621421]
- Hartman RL, Huestis MA. Cannabis effects on driving skills. *Clinical Chemistry*. 2013; 59(3):478–492. [PubMed: 23220273]
- Hommel J, Thompson K, Leadbeater B. Trajectories of marijuana use in youth ages 15–25: Implications for postsecondary education experiences. *Journal of Studies on Alcohol and Drugs*. 2014; 75(4): 674–683. [PubMed: 24988266]
- Horan JM, Widom CS. Does age of onset of risk behaviors mediate the relationship between child abuse and neglect and outcomes in middle adulthood? *Journal of Youth and Adolescence*. 2015; 44(3):670–82. [PubMed: 25104419]
- Horwood LJ, Fergusson DM, Hayatbakhsh MR, Najman JM, Coffey C, Patton GC, Silins E, Hutchinson DM. Cannabis use and educational achievement: Findings from three Australasian cohort studies. *Drug and Alcohol Dependence*. 2010; 110:247–253. [PubMed: 20456872]
- Huang S, Trapido E, Fleming L, Arheart K, Crandall L, French M, Malcolm S, Prado G. The long-term effects of childhood maltreatment experiences on subsequent illicit drug use and drug-related problems in young adulthood. *Addictive Behaviors*. 2011; 36:95–102. [PubMed: 20947260]
- Hussey JM, Marshall JM, English DJ, Knight ED, Lau AS, Dubowitz H, Kotch JB. Defining maltreatment according to substantiation: distinction without a difference? *Child Abuse & Neglect*. 2005; 29(5):479–492. [PubMed: 15970321]

- Ireland TO, Smith CA, Thornberry TP. Developmental issues in the impact of child maltreatment on later delinquency and drug use. *Criminology*. 2002; 40(2):359–400.
- Jankord R, Solomon MB, Albertz J, Flak JN, Zhang R, Herman JP. Stress vulnerability during adolescent development in rats. *Endocrinology*. 2011; 152:629–638. [PubMed: 21106877]
- Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE.; Miech, RA. Monitoring the Future: National survey results on drug use, 1975–2013: Volume I: Secondary school students. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2014a. Available at: http://www.monitoringthefuture.org/pubs/monographs/mtf-vol1_2013.pdf
- Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE.; Miech, RA. Demographic subgroup trends among adolescents in the use of various licit and illicit drugs: 1975–2013. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2014b. Monitoring the Future Occasional Paper No. 81 Available at: <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ81.pdf>
- Jonson-Reid M, Kohl PL, Drake B. Child and adult outcomes of chronic child maltreatment. *Pediatrics*. 2012; 129(5):839–845. [PubMed: 22529281]
- Kedzior KK, Laeber LT. A positive association between anxiety disorders and cannabis use or cannabis use disorders in the general population- a meta-analysis of 31 studies. *BMC Psychiatry*. 2014; 14:136. [PubMed: 24884989]
- Khoury L, Tang YL, Bradley B, Cubells JF, Ressler KJ. Substance use, childhood traumatic experience, and posttraumatic stress disorder in an urban civilian population. *Depression and Anxiety*. 2010; 27:1077–1086. [PubMed: 21049532]
- Kilpatrick DG, Acierno R, Saunders B, Resnick HS, Best CL, Schnurr PP. Risk factors for adolescent substance abuse and dependence: Data from a national sample. *Journal of Consulting and Clinical Psychology*. 2000; 68:19–30. [PubMed: 10710837]
- Kohl PL, Jonson-Reid M, Drake B. Time to leave substantiation behind: findings from a national probability study. *Child Maltreatment*. 2009; 14(1):17–26. [PubMed: 18971346]
- Lansford JE, Dodge KA, Pettit GS, Bates JE. Does physical abuse in early childhood predict substance use in adolescence and early adulthood? *Child Maltreatment*. 2010; 15:190–194. [PubMed: 20019026]
- Lau AS, Leeb RT, English D, Graham JC, Briggs EC, Brody KE, Marshall JM. What's in a name? A comparison of methods for classifying predominant type of maltreatment. *Child Abuse & Neglect*. 2005; 29:533–551. [PubMed: 15970324]
- Layne, CM.; Warren, J.; Watson, P.; Shalev, A. Risk, vulnerability, resistance and resilience: Towards an integrative conceptualization of posttraumatic adaption. In: Friedman, MJ.; Keanne, TM.; Resick, PA., editors. *Handbook of PTSD: Science and Practice*. New York: Guilford; 2007. p. 497-520.
- Lo CC, Cheng TC. The impact of childhood maltreatment on young adults' substance abuse. *American Journal of Drug and Alcohol Abuse*. 2007; 33:139–146. [PubMed: 17366254]
- Lynskey M, Hall W. The effects of adolescent cannabis use on educational attainment: A review. *Addiction*. 2000; 95(11):1621–1630. [PubMed: 11219366]
- Maggs JL, Schulenberg JE. Initiation and course of alcohol consumption among adolescents and young adults. *Recent Developments in Alcoholism*. 2005; 17:29–47. [PubMed: 15789858]
- McCormick CM, Mathews IZ, Thomas C, Waters P. Investigations of HPA function and the enduring consequences of stressors in adolescence in animal models. *Brain and Cognition*. 2010; 72:73–85. [PubMed: 19616355]
- McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, Kessler RC. Childhood adversities and adult psychiatric disorders in the National Comorbidity Survey Replication II: Associations with persistence of DSM-IV disorders. *Archives of General Psychiatry*. 2010; 67:124–132. [PubMed: 20124112]
- Meier MH, Caspi A, Ambler A, Harrington H, Houts R, Keefe RSE, Moffitt TE. Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proceedings of the National Academy of Sciences*. 2012; 109(40):e2657–2664.

- Monahan KC, Rhew IC, Hawkins JD, Brown EC. Adolescent pathways to co-occurring problem behavior: The effects of peer delinquency and peer substance use. *Journal of Research on Adolescence*. 2014; 24:630–645. [PubMed: 25506186]
- Monshouwer K, Harakeh Z, Lugtig P, Huizink A, Creemers HE, Reijneveld SA, De Winter AF, Van Oort F, Ormel J, Vollebergh WAM. Predicting transitions in low and high levels of risk Behavior from early to middle Adolescence: The TRAILS Study. *Journal of Abnormal Child Psychology*. 2012; 40:923–931. [PubMed: 22427248]
- Moore TH, Zammit S, Lingford-Hughes A, Barnes TR, Jones PB, Burke M, Lewis G. Cannabis use and risk of psychotic or affective mental health outcomes: A systematic review. *Lancet*. 2007; 370:319–328. [PubMed: 17662880]
- Moran PB, Vuchinich S, Hall NK. Associations between types of maltreatment and substance use during adolescence. *Child Abuse & Neglect*. 2004; 28:565–574. [PubMed: 15159070]
- Oberlander SE, Wang Y, Thompson R, Lewis T, Proctor LJ, Isbell P, English DE, Dubowitz H, Litrownik AJ, Black MM. Childhood maltreatment, emotional distress, and early adolescent sexual intercourse: Multi-informant perspectives on parental monitoring. *Journal of Family Psychology*. 2011; 25:885–894. [PubMed: 21928888]
- Oshri A, Rogosch FA, Burnette ML, Cicchetti D. Developmental pathways to adolescent cannabis abuse and dependence: Child maltreatment, emerging personality, and internalizing versus externalizing psychopathology. *Psychology of Addictive Behaviors*. 2011; 25:634–644. [PubMed: 21534646]
- Palamar JJ, Fenstermaker M, Kamboukos D, Ompad DC, Cleland CM, Weitzman M. Adverse psychosocial outcomes associated with drug use among US high school seniors: A comparison of alcohol and marijuana. *American Journal of Drug and Alcohol Abuse*. 2014; 40(6):438–446. [PubMed: 25169838]
- Runyan DK, Curtis PA, Hunter WM, Black MM, Kotch JB, Bangdiwala S, Dubowitz H, English D, Everson MD, Landsverk J. LONGSCAN: A consortium for longitudinal studies of maltreatment and the life course of children. *Aggression and Violent Behavior*. 1998; 3:275–285.
- Schuermeier J, Salomonsen-Sautel S, Price RK, Balan S, Thurstone C, Min SJ, Sakai JT. Temporal trends in marijuana attitudes, availability and use in Colorado compared to non-medical marijuana states: 2003–11. *Drug and Alcohol Dependence*. 2014; 140:145–155. [PubMed: 24837585]
- Schulenberg J, Merline A, Johnston LD, O'Malley PM, Bachman JG, Laetz VB. Trajectories of marijuana use during the transition to adulthood: The big picture based on national panel data. *Journal of Drug Issues*. 2005; 35(2):255–279. [PubMed: 16534532]
- Schweinsburg AD, Brown SA, Tapert SF. The influence of marijuana use on neurocognitive functioning in adolescents. *Current Drug Abuse Reviews*. 2008; 1(1):99–111. [PubMed: 19630709]
- Solowij N, Stephens RS, Roffman RA, Babor T, Kadden R, Miller M, Vendetti J. Cognitive functioning of long-term heavy cannabis users seeking treatment. *Journal of the American Medical Association*. 2002; 287(9):1123–1131. [PubMed: 11879109]
- Solowij N, Jones K, Rozman M, Davis S, Ciarrochi J, Heaven PL, Yücel M. Verbal learning and memory in adolescent cannabis users, alcohol users and non-users. *Psychopharmacology*. 2011; 216(1):131–144. [PubMed: 21328041]
- Steinberg L. A dual systems model of adolescent risk-taking. *Developmental Psychobiology*. 2010; 52:216–224. [PubMed: 20213754]
- Swift W, Coffey C, Carlin JB, Degenhardt L, Patton GC. Adolescent cannabis users at 24 years: Trajectories to regular weekly use and dependence in young adulthood. *Addiction*. 2008; 103(8):1361–1370. [PubMed: 18855826]
- Thompson R, Tabone JK, Litrownik AJ, Briggs EC, Hussey JM, English DJ, Dubowitz H. Early adolescent risk behavior outcomes of childhood externalizing behavioral trajectories. *Journal of Early Adolescence*. 2011; 31:234–257.
- Thompson R, Litrownik AJ, Isbell P, Everson MD, English DJ, Dubowitz H, Proctor LJ, Flaherty EG. Adverse experiences and suicidal ideation in adolescence: Exploring the link using the LONGSCAN samples. *Psychology of Violence*. 2012; 2:211–225.

- Thompson R, Neilson EC. Early parenting: The roles of maltreatment, trauma symptoms, and future expectations. *Journal of Adolescence*. 2014; 37:1099–1108. [PubMed: 25150891]
- Thornberry TP, Henry KL, Ireland TO, Smith CA. The causal impact of childhood-limited maltreatment and adolescent maltreatment on early adult adjustment. *Journal of Adolescent Health*. 2010; 46:359–365. [PubMed: 20307825]
- Thornberry TP, Ireland TO, Smith CA. The importance of timing: the varying impact of childhood and adolescent maltreatment on multiple problem outcomes. *Development and Psychopathology*. 2001; 13:957–979. [PubMed: 11771916]
- Thornberry TP, Matsuda M, Greenman SJ, Augustyn MB, Henry KL, Smith CA, Ireland TO. Adolescent risk factors for child maltreatment. *Child Abuse & Neglect*. 2014; 38(4):706–722. [PubMed: 24075569]
- Tucker JS, de la Haye K, Kennedy DP, Green HD, Pollard MS. Peer influence on marijuana use in different types of friendships. *Journal of Adolescent Health*. 2014; 54:67–73. [PubMed: 24054813]
- U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau. *Child Maltreatment 2012*. 2013. Available from: <http://www.acf.hhs.gov/programs/cb/research-data-technology/statistics-research/child-maltreatment>
- Volkow ND, Baler RD, Compton WM, Weiss SRB. Adverse health effects of marijuana use. *New England Journal of Medicine*. 2014; 370(23):2219–2227. [PubMed: 24897085]
- Widom CS, Marmorstein NR, White HR. Childhood victimization and illicit drug use in middle adulthood. *Psychology of Addictive Behaviors*. 2006; 20:394–403. [PubMed: 17176174]
- Wilson HW, Widom CS. A prospective examination of the path from child abuse and neglect to illicit drug use in middle adulthood: The potential mediating role of four risk factors. *Journal of Youth and Adolescence*. 2009; 38:340–354. [PubMed: 19636749]
- Windle M, Wiesner M. Trajectories of marijuana use from adolescence to young adulthood: Predictors and outcomes. *Development and Psychopathology*. 2004; 16(4):1007–1027. [PubMed: 15704825]
- Wright EM, Fagan AA, Pinchevsky GM. The effects of exposure to violence and victimization across life domains on adolescent substance use. *Child Abuse & Neglect*. 2013; 37(11):899–909. [PubMed: 23743232]
- Wu NS, Schairer LC, Dellor E, Grella C. Childhood trauma and health outcomes in adults with comorbid substance abuse and mental health disorders. *Addictive Behaviors*. 2010; 35:68–71. [PubMed: 19775820]
- Zvolensky MJ, Cogle JR, Johnson KA, Bonn-Miller MO, Bernstein A. Marijuana use and panic psychopathology among a representative sample of adults. *Experimental and Clinical Psychopharmacology*. 2010; 18(2):129–134. [PubMed: 20384424]

Table 1

Sample Characteristics by Level of Marijuana Use among Maltreated Youth (n = 702).

Characteristic	n (%)	Level of Marijuana Use			<i>p</i> ^a
		Never (%) (n=329)	Some (%) (n=219)	Heavy (%) (n=154)	
Site					.40
East	91 (13)	54	25	21	
Midwest	75 (11)	51	27	23	
South	92 (13)	50	33	17	
Southwest	251 (36)	41	33	26	
Northwest	193 (28)	48	33	19	
Sex of Participant					.001
Male	332 (47)	43	29	28	
Female	370 (53)	50	34	16	
Race/Ethnicity					.052
Black	334 (48)	51	30	22	
White	209 (30)	47	31	19	
Other ^b	105 (15)	41	28	31	
Hispanic	18 (8)	33	43	24	

Note.^a $\chi^2 (3,k)$ of Marijuana Use (Never/Some/Heavy) $\times$ Characteristic (*k*).^bMixed, Native American, Asian/Pacific Islander.

Table 2
Bivariate Relationships between Characteristics of Maltreatment and Level of Marijuana Use (n = 702).

Characteristic	n (%)	Level of Marijuana Use			<i>p</i> ^a
		Never (%)	Some (%)	Heavy (%)	
Physical Abuse					.01
No	313 (45)	53	28	18	
Yes	389 (55)	42	33	25	
Sexual Abuse					<.01
No	506 (72)	49	28	23	
Yes	196 (28)	40	41	19	
Neglect					.56
No	81 (12)	42	32	26	
Yes	621 (88)	48	31	21	
Emotional Maltreatment					.14
No	316 (45)	51	29	20	
Yes	386 (55)	44	33	24	
Extent of CM ^b					<.001
Limited	309 (44)	55	28	18	
Extensive	393 (56)	40	34	25	

Note.

^a χ^2 (3, *k*) of Marijuana Use (No /Some/Heavy) x Characteristic (*k*).

^b Extent defines Limited as CPS reports in one developmental period, and fewer than 5 reports. Extensive represents CPS reports in more than one developmental period, and more than 5 reports.

Table 3

Correlation matrix of the Maltreatment and Peer Use Predictor Variables

Variable	Physical Abuse	Sexual Abuse	Neglect	Emotional Abuse	Extent of CM	Peer Use
Physical Abuse	1					
Sexual Abuse	.22**	1				
Neglect	-.09*	-.03	1			
Emotional Maltreatment	.36**	.16**	.03	1		
Extent of CM	.48**	.30**	.18**	.40**	1	
Peer Use	.04	.07	-.07	.04	.12**	1

*
 $p < .05$ **
 $p < .01$

Table 4

The Relationships among Maltreatment Characteristics and Level of Adolescent Marijuana Use, Controlling for Site, Participant Sex, and Race/Ethnicity (n = 702).

Variable	Model 1			Model 2 ^b		
	Likelihood Ratio X ² (df)	Some Use ^d AOR (CI)	Heavy Use AOR (CI)	Likelihood Ratio X ² (df)	Some Use AOR (CI)	Heavy Use AOR (CI)
Physical Abuse	0.20(2)			1.10 (2)		
Yes		1.08 (0.71–1.65)	1.10 (0.68–1.77)		1.25 (0.80–1.95)	1.23 (0.72–2.08)
No		reference	reference		reference	reference
Sexual Abuse	7.32 (2) *			6.82 (2) *		
Yes		1.62 (1.07–2.45) *	0.88 (0.54–1.46)		1.52 (0.98–2.36)	0.80 (0.47–1.37)
No		reference	reference		reference	reference
Neglect	2.78 (2)			0.48 (2)		
Yes		0.78 (0.44–1.37)	0.59 (0.31–1.10)		0.89 (0.49–1.63)	0.78 (0.39–1.58)
No		reference	reference		reference	reference
Emotional Maltreatment	0.13 (2)			0.02 (2)		
Yes		1.01 (0.68–1.50)	1.08 (0.69–1.69)		1.01 (0.66–1.54)	1.03 (0.63–1.69)
No		reference	reference		reference	reference
Extent of CM	9.52 (2) **			7.52 (2) *		
Extensive		1.40 (0.90–2.17)	2.16 (1.31–3.58) **		1.31 (0.83–2.08)	2.28 (1.31–3.95) **
Limited		reference	reference		reference	reference
Sex of Participant	12.37 (2) **			6.50 (2) *		
Female		0.89 (0.62–1.29)	0.49 (0.32–0.74) ***		0.88 (0.60–1.28)	0.56 (0.36–0.88) *
Male		reference	reference		reference	reference
Site	2.54 (2)			2.15 (2)		
Race/Ethnicity	2.99 (2)			5.10 (2)		
Peer Use				92.77 (2) ***		
Yes					3.57 (2.46–5.18) ***	6.80 (4.30–10.76) ***
No					reference	reference

Note. AOR = Adjusted Odds Ratio. CI = 95% confidence interval.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

No Use is the reference category.
N= 617 because of reduced number with Peer Use data.
 p
*
 $p < .05$
**
 $p < .01$

 $p < .001$