Published in final edited form as:

Tetrahedron Lett. 2016 January 20; 57(3): 423–425. doi:10.1016/j.tetlet.2015.12.038.

Lanthanide and asymmetric catalyzed syntheses of sterically hindered 4-isoxazolyl-1,4-dihydropyridines and 4-isoxazolyl-quinolones

Scott A. Steiger^{a,d}, Chun Li^b, Charles F. Campana^c, and Nicholas R. Natale^{a,*}

^aMedicinal Chemistry Graduate Program, University of Montana, Missoula MT 59812, USA

^bDepartment of Chemistry Ithaca College, Ithaca NY, USA

^cBruker AXS Inc., 5465 East Cheryl Parkway, Madison, WI 53711, USA

Multi-component organic cyclizations,¹ exemplified by the Hantzsch pyridine synthesis have been known for over a century.² The concept of scaffolds and scaffold hopping in medicinal chemistry is much more recent, however, the number of potentially useful applications of scaffolds is arithmetically increased by their judicious combination, and especially valuable are those combinations which are attainable by efficient stereoselective methodology.

Interest in the usefulness of Hantzsch esters was exponentially enhanced by the discovery that specific 4-aryldihydropyridines (DHPs) possessed robust biological activity as calcium channel antagonists, and thus are useful as antihypertensive medicines.³ In many cases, there is a pronounced enantioselectivity of action,⁴ yet as often observed the agents in general medical practice have been obtained by chromatographic or classical resolution. The corresponding 4-aryl quinolones have been found to have useful activity as inhibitors of TGF β Signaling,^{5,6} and as agents which reduce cellular tau levels which represents an important target in Alzheimer's disease,^{7,8} and are the current subject of intensive pre-clinical development. Early advances in stoichiometric adjuvant asymmetric synthesis of chiral DHPs were accomplished first by Meyers and Natale,⁹ and later by Enders,¹⁰ and Dondoni.¹¹ The critical breakthrough to catalytic multi-component asymmetric DHP synthesis was achieved initially by Gong,¹² and extended to quinolones by Gestwicki.⁷ We herein report, to the best of our knowledge, the first facile preparation of the most sterically hindered 4-isoxazolyl-1,4-dihydropyridines to date, the first examples of 4-isoxazolyl-quinolones, as well as their highly stereoselective organocatalytic asymmetric synthesis.

*Corresponding author. Tel 406-243-4132; fax 406-243-5228. nicholas.natale@umontana.edu (N.R. Natale).

^dPresent address: Office of Commercialization, Washington State University P.O. Box 641802, Pullman, WA 99164-1802, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary Material

Full experimental details and characterization for all new compounds. Crystal structure determination, atomic coordinates and hydrogen-bond geometry for **2a**, **2b**, and **RS-3a**, including unit cells. SYBYL transition state modeling of the organocatalytic asymmetric synthesis.

The Isoxazolyl-dihydropyridines are characterized by their complex conformational dynamics,¹³ which we have demonstrated can lead to divergent structure activity relationships (SAR) for different biomolecular targets.^{14,15} In previous studies we observed that increasing the size of substituents on the isoxazole lead to vanishingly low yields, even under forcing conditions.¹⁶ As an example, naphthyl groups appended to isoxazoles, even separated by an ethylene spacer, produced typical single digit yields in the Hantzsch synthesis, even under elevated pressure. Previous reports on the use of lanthanide catalysts are noteworthy for their relatively mild reaction conditions, and we sought to test the scope and limitations of this promising methodology.

The Ytterbium catalyzed¹⁶ method produced moderate to good yields of IDHPs unavailable using standard methods. The single crystal x-ray diffractometry (sc-xrd) of **2a** and **2b** illustrates the steric crowding of the C-3 isoxazole substituent. Characteristic of IDHPs, in the unit cell, intermolecular hydrogen bonding is observed between the isoxazole ring N and the DHP N-H. In the anthryl example, **2c**, significant anisotropy is observed in the ¹H NMR for *both* the C-2 and C-6 methyl groups of the DHP moiety, consistent with an analogous conformation in solution.

Single crystal x-ray diffractometry and solid state conformation

The Isoxazolyl-quinolone was crystallized and its structure determined by x-ray diffractometry. The heterocyclic axis at the quinolone C-4 is oriented *O-endo*, that is, with the isoxazolyl oxygen towards the dihydropyridine moiety, and the 3'-aryl *exo* to the quinolone. The Quinolone 3-ester adopts a *synperiplanar* conformation, the quinolone locks the C-5 ketone in the *antiperiplanar* conformer. In the unit cell, there is an intermolecular hydrogen bond between the quinolone N-1 hydrogen, and the quinolone ring carbonyl oxygen.

The asymmetric synthesis was performed according to Gestwicki,⁶ from the isoxazole aldehydes using the organocatalysis of BINOL phosphates as pioneered by Gong. Chemical yields were moderate to good, and optical purities as established by HPLC-CSP were excellent (>90% e.e.). Absolute configurations were assigned by analogy to the work of Schade,⁷ who established the absolute configuration of the (+) enantiomer as (*R*)- by single crystal x-ray diffractometry. The quinolone absolute configuration is probably driven by the relative reactivity rates of ketoester versus cyclic 1,3-diketone, and therefore arises from initial Knoevenagel condensation by the ketoester, followed by Michael addition of the diketone. In contrast, the IDHP in Entry **2j**, the Knoevenagel product was preformed with the triazole, followed by condensation with the ketoester. Therefore, that the absolute configuration of the entries in Table 2 are all (–), (*S*)- is fortuitous. We would not expect the corresponding multi-component synthesis of IDHPs analogous to **2j** to proceed with significant stereoselectivity, unless the initial Knoevenagel adduct was similarly preformed.

We are intrigued by the potential of the hindered 4-isoxazole dihydropyridines and 4-isoxazolylquinolones for theranostic applications,¹⁸ and look forward to detailed study of their conformational dynamics and bioactivity. We will report on our progress in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

The authors are grateful to the Center for Structural and Functional Neuroscience for NIH NCRR P20RR015583 and NINDS R15 NS038444 for support of this work.

References

1. Seayad, J.; List, B. Catalytic Asymmetric Multicomponent Reactions. In: Zhu, Y.; Bienayme, H., editors. Chapter 9 in Multicomponent Reactions. Wiley-VCH: Weinheim; 2005. p. 277-299.
2. (a) Hantzsch A. Condensationprodukte aus Aldehydammoniak und Ketonartigen Verbindungen. Chemische Berichte. 1881; 14(2):1637-1638. (b) Hantzsch A. Justus Liebigs Ann. Chemie. 1882; 215:1-82.
3. Lemke, TL.; Williams, DA.; Roche, VF.; Zito, SW. Foye's Principles of Medicinal Chemistry. Seventh. Vol. 24. Lippincott: Williams and Wilkins; 2013.
4. Triggle DJ, Langs DA, Janis RA. Med. Res. Rev. 1989; 9:123-180. [PubMed: 2654521]
5. Schade D, Lanier M, Willems E, Okolotowicz K, Bushway P, Wahlquist C, Gilley C, Mercola M, Cashman JR. J. Med. Chem. 2012; 55:9946-9957. [PubMed: 23130626]
6. Langle D, Marquardt V, Heider E, Vigante B, Duburs G, Luntana I, Flotgen D, Golz C, Strohmann C, Koch O, Schade D. Design, synthesis and 3D-QSAR studies of novel 1,4-dihydropyridines as TGF β /Smad inhibitors. European J. Med. Chem. 2015; 95:249-266. [PubMed: 25817775]
7. Evans CG, Gestwicki JE. Org. Lett. 2009; 11:2957-2959. [PubMed: 19527003]
8. Evans CG, Jinwal UK, Makley LN, Dickey CA, Gestwicki JE. Chem Commun. 2011; 47(1):529-531.
9. Meyers AI, Natale NR, Wettlaufer DG, Rafii S, Clardy JC. Tetrahedron Lett. 1981; 5:123-126.
10. Enders D, Muller S, Demir AS. Tetrahedron Lett. 1988; 29:6437-6440.
11. Dondoni A, Massi A. Mol. Divers. 2003; 6:261-270. [PubMed: 15068090]
12. Jiang J, Yu J, Sun X, Rao Q, Gong L. Angew. Chem. Int. Ed. 2008; 47:2458-2462.
13. Natale NR, Steiger SA. Future Med. Chem. 2014; 6:923-943. [PubMed: 24962283]
14. Zamponi G, Stotz SC, Staples RJ, Rogers TA, Nelson JK, Hulubei V, Blumenfeld A, Natale NR. J. Med. Chem. 2003; 46:87-96. [PubMed: 12502362]
15. Hulubei V, Meikrantz SB, Quincy David A, Houle Tina, McKenna John I, Rogers Mark E, Steiger Scott, Natale NR. Bioorg. Med. Chem. 2012; 20:6613-6620. [PubMed: 23063517]
16. Natale NR, Rogers ME, Staples R, Triggle DJ, Rutledge A. Lipophilic 4-isoxazolyl-1,4-dihydropyridines: synthesis and structure activity relationship. J. Med. Chem. 1999; 42(16):3087-3093. [PubMed: 10447952]
17. Wang M, Sheng J, Zhang L, Han J-W, Fan Z-Y, Tian H, Qian C-T. Facile Yb(OTf)₃-promoted one-pot synthesis of polyhydroquinoline derivatives through Hantzsch reaction. Tetrahedron. 2005; 6:1539-1543.
18. Natale, NR. Learning from the Hantzsch Synthesis. In: Natale, NR., editor. Chemical Innovation. American Chemical Society; 2000 Nov. p. 22-28. [<http://pubs.acs.org/subscribe/archive/ci/30/i11/html/11natale.html>]

Figure 1. 50% probability ellipsoids ORTEPs of (A) Naphthyl-2a. R = 5.1%. (B) Methoxy-Naphthyl-2b. R = 9%. (C) IQ (±)-3a, R = 5.2%.

Scheme 1.

Lanthanide catalyzed synthesis of sterically hindered 3-aryl-Isoxazolyl-Dihydropyridines **2**, and asymmetric organocatalysis of dihydroquinolones **3**

Table 1Lanthanide catalyzed synthesis of sterically hindered IDHPs **2a–i**.

Entry	Ar	% yield	HRMS calc'd	Found
2a	1'-Naphthyl	64.04	501.2026	501.2019
2b	1'-(2'-methoxy-Naphthyl)	50.21	491.2182	491.2205
2c	9'-Anthryl	41.98	511.2233	511.2228
2d	3,4-Bis-benzyloxy-phenyl	67.95	469.1975	469.1990
2e	3-Phenoxy- phenyl	60.98	501.2026	501.2019
2f	4-Methoxy- phenyl	60.96	441.2026	441.2057
2g	2,4-Dimethoxy- phenyl	63.98	623.2757	623.2709
2h	3,4-Dimethoxy- phenyl	53.19	623.2757	623.2807
2i	4-Biphenyl	46.23	487.2233	487.2256

Table 2

Asymmetric organocatalytic synthesis of IDHP **2j** and 4-isoxazolyl-Quinolones, **3**. All examples gave optical purities of >90% e.e by HPLC-CSP (Supporting data).

Entry	Ar	R	Catalyst	Yield	[α] _D
RS- 3a	-Ph	-CH ₂ CH ₃	Yb(OTf) ₃	30.25	
(-)- 3a			(R)-TRIP	25.92	-11.2
RS- 3b	- <i>o</i> -Br-Ph	-CH ₂ CH ₃	Yb(OTf) ₃	71.60	
(-)- 3b			(R)-TRIP	64.35	-13.5
RS- 3c	- <i>m</i> -Br-Ph	-CH ₂ CH ₃	Yb(OTf) ₃	62.07	
(-)- 3c			(R)-TRIP	60.40	-2.36
RS- 3d	- <i>p</i> -Br-Ph	-CH ₂ CH ₃	Yb(OTf) ₃	67.19	
(-)- 3d			(R)-TRIP	65.34	-0.032
RS- 2j	- <i>p</i> -Br-Ph	-4-CH ₂ (N-phenethyl)triazole	Yb(OTf) ₃	31.56	
(-)- 2j			(R)-TRIP	21.32	-2.68