Published in final edited form as:

Angew Chem Int Ed Engl. 2015 November 23; 54(48): 14531–14534. doi:10.1002/anie.201507928.

An Epitope-Specific Respiratory Syncytial Virus Vaccine based on an Antibody Scaffold

Xiaozhou Luo^[+],

Department of Chemistry, The Scripps Research Institute 10550 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Dr. Tao Liu^[+],

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Dr. Ying Wang,

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Dr. Haiqun Jia,

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Yuhan Zhang,

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Dawna Caballero,

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Dr. Juanjuan Du,

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Dr. Rongsheng E. Wang,

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Dr. Danling Wang,

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Peter G. Schultz, and

Department of Chemistry, The Scripps Research Institute 10550 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Correspondence to: Peter G. Schultz, schultz@scripps.edu; Feng Wang, fwang@calibr.org.

[+]⁺These authors contributed equally to this work.

Supporting information for this article is given via a link at the end of the document.

Dr. Feng Wang

California Institute for Biomedical Research (Calibr) 11119 N. Torrey Pines Road, La Jolla, CA 92037 (USA)

Peter G. Schultz: schultz@scripps.edu; Feng Wang: fwang@calibr.org

Abstract

Respiratory syncytial virus (RSV) is a leading cause of lower respiratory tract infections in children. We have generated an epitope-specific RSV vaccine by grafting a neutralizing epitope (F-epitope) in its native conformation into an immunoglobulin scaffold. The resulting antibody fusion exhibited strong binding affinity to Motavizumab, an RSV neutralizing antibody, and effectively induced potent neutralizing antibodies in mice. This work illustrates the potential of the immunoglobulin molecule as a scaffold to present conformationally constrained B-cell epitopes.

Keywords

antibodies; RSV vaccine; conformation; polypeptide; protein engineering

Respiratory syncytial virus (RSV) is a negative-sense, single-stranded RNA virus that belongs to the *Paramyxoviridae* family. RSV is the cause of more than 30 million lower respiratory tract infections and more than 60 thousand deaths worldwide.^[1] It is also the major cause of hospitalization in children under 5 years of age. Young children at risk are currently treated with multiple injections of the antibody Palivizumab; a neutralizing antibody directed against an epitope of the RSV F protein. A vaccine based approach would be far more effective in reducing RSV infections, especially in developing countries where treatment with anti-RSV antibodies is difficult. However, in early clinical trials, a formalin-inactivated whole virus vaccine caused enhanced disease severity upon natural RSV infection. DNA-based, live-attenuated or whole-inactivated virus-based and protein subunit vaccines, have been explored during the past 50 years without success,^[2] partially because of the poor induction of neutralizing antibodies.^[3] Immunization with linear peptides corresponding to the neutralizing epitope failed to elicit neutralizing antibodies in a mouse study, which underlines the critical role of the native conformation of the epitope.^[4] Recently it was shown that protective antibodies can be elicited with an epitope-focused vaccination strategy, wherein the active structural epitope was grafted onto a computationally designed protein scaffold which stabilized its wild-type helix-turn-helix conformation.^[5] This work highlights the importance of presenting epitopes in their native conformation in a vaccine candidate. Moreover, it encourages the development of additional protein scaffolds to present foreign epitopes to the immune system in defined conformations.

We previously identified a family of natural bovine antibodies (Figure 1A) with an ultralong heavy chain complementary determining region 3 (CDR3) consisting of an extended β -sheet structure terminated in a disulfide bonded “knob” domain.^[6] On the basis of this structure we have designed both β -sheet and helical CDR architectures that present fused growth factors, cytokines and bioactive peptides in their native conformations. The resulting antibody fusion proteins retain their biological activity but have increased stability and serum half-lives.^[7] Here, we demonstrate that a similar approach can be used to present

an immunogenic RSV epitope in its active helical conformation. The resulting fusion protein elicits protective neutralizing antibodies in mice.

Of the eleven proteins encoded in the RSV genome^[8], antibodies directed against the fusion (F) and attachment (G) glycoproteins confer neutralization and protection to RSV in animal models.^[9] Because the RSV F glycoprotein is conserved among RSV A and B strains, antibodies targeting the F protein have the potential to provide cross protection.^[10] The crystal structure of the neutralizing antibody Motavizumab (Mota, a more potent second generation Palivizumab) complexed to the F protein reveals a 22-amino acid epitope in a helix-turn-helix conformation (Figure 1B).^[11] A stable helix-turn-helix mimic of the epitope (SELLSKINDMPITNDQKKLMSN) has been shown to elicit neutralizing antibodies against RSV virus in rhesus macaques.^[5] Similarly, we reasoned that one can graft this F-epitope into an appropriately designed CDR of an antibody and maintain its native conformation. The resulting antibody-epitope fusion is expected to have a long circulating half-life, present antigen in a bivalent fashion, and minimize induction of off-target antibodies.

To test whether one can substitute the Mota-binding F-epitope into an antibody CDR and retain its active conformation, we fused this peptide to various CDRs of Herceptin, an anti-Her2 antibody with low immunogenicity used clinically for the treatment of breast cancer.^[12] Previously we have fused various conformationally constrained peptides, such as CXCR4 receptor antagonist peptide, glucagon-like peptide receptor agonist peptide, and protease inhibitor peptides to various CDRs of Herceptin in their biological active conformations.^[7a, 13] The engineered F-epitope (EpiS) consisted of the 22-amino acid Mota-binding peptide core, flanked by short helices consisting of 8 N-terminal and 10 C-terminal residues to further stabilize the fused helical peptide (Figure 1C). In order to identify the best site for presenting the epitope, we screened different CDR and constant loops in the antibody scaffold. The EpiS epitope was inserted between R98 and D108 of CDRH3 (hH3-EpiS), P53 and G56 of CDRH2 (hH2-EpiS), Q27 and V29 of CDRL1 (hL1-EpiS), Y92 and P95 of CDRL3 (hL3-EpiS) and between G200 and S202 of light chain constant loop3 (hLoop3-EpiS) (Figure 1C). EpiS is predicted to have T-cell epitopes using the IEDB analysis resource consensus tool (Supporting Information, Figure S1), therefore, we reasoned the antibody EpiS fusion is likely to elicit a T-cell dependent antibody response. All fusion proteins were transiently expressed in FreeStyle 293 cells as secreted proteins and purified by protein G chromatography. hH3-EpiS, hL3-EpiS and hLoop3-EpiS were produced in good yields of 8.4 mg/L, 6.7 mg/L and 8.5 mg/L, respectively; These purified antibody fusions were characterized by SDS-PAGE and MS (Figure S2).

To confirm that the epitope retained its correct conformation when grafted into the antibody scaffold, we determined the binding affinities of all purified fusion proteins to Mota. A qualitative enzyme-linked immunosorbent assay (ELISA) was performed and the results revealed that all fusion proteins bound tightly with Mota while Herceptin itself showed no binding (Figure 2A). The apparent binding constant of hH3-EpiS with Mota was determined by a quantitative ELISA to be 154 ± 18 pM (Figure 2B), which is consistent with the binding affinity of F-epitope to Mota (reported values of 30 pM to 800 pM).^[5] A K_d value of 240

pM obtained from biolayer interferometry (BLI) further validated the high affinity of hH3-EpiS to Mota (Figure S3).

Next, we examined this immunoglobulin-based vaccine in a mouse model by monitoring antibody titers in response to immunization with the immunoglobulin-epitope fusion. Because human antibodies are immunogenic in mice,^[14] a mouse immunoglobulin scaffold was used to present the F protein epitope. OKT3 is a fully mouse antibody with good stability, which makes it a suitable surrogate for immunization studies in mice. hLoop3-EpiS was translated into OKT3 based on sequence alignment by inserting the EpiS epitope between T199 and T201 in Loop3 to yield mLoop3-EpiS. The engineered antibody was transiently expressed in FreeStyle 293 cells with a yield of 3.0 mg/L. The purified antibody fusion was characterized by SDS-PAGE and MS (Figure S2). The apparent binding constant of mLoop3-EpiS with Mota was assessed by a quantitative ELISA to be 80 ± 13 pM (Figure 2B), which was further validated by BLI ($k_{\text{on}} = 1.3 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, k_{off} is below the detection limit) (Figure S3).

Two groups of BALB/c mice ($n=10$) were immunized three times with either 10 μg mLoop3-EpiS or 10 μg of wild-type OKT3 as a negative control on a biweekly schedule in weeks 0, 2 and 4. Serum samples from weeks 3, 5 and 7 were collected, and their antibody titers against the mLoop3-EpiS, RSV F protein and OKT3 were measured by ELISA at 1:100 dilution. Antibodies against the fusion protein mLoop3-EpiS were observed from week 3 onward and continuously increased in weeks 5 and 7, while there was no detectable antibody response towards the OKT3 control up to week 7 (Figure 3A and S4). The antibodies elicited by mLoop3-EpiS cross-react with RSV F glycoprotein by ELISA (Figure 3A). Finally, we demonstrated the neutralizing activity of our vaccine in an RSV virus protection assay. A widely used RSV neutralization assay was adopted to determine whether antibodies generated by this epitope-focused fusion protein can neutralize the active virus.^[15] 10 μL RSV at 10^6 cfu/mL pretreated with 10 μL of serum dilutions (1:10, 1:100, 1:1000) from the vaccinated or control groups were allowed to infect a monolayer Hep-2 cells for four hours before the medium was removed and the cells were overlaid with DMEM/F12 in 0.3% agarose. After 6 days, the monolayers were fixed and stained. Plaques were counted and the neutralizing rates were calculated accordingly. Significant differences were observed in RSV infectivity with 1:10 and 1:100 serum dilutions of the vaccinated versus control group. (Figure 3B).^[16] This result demonstrates that the antibodies raised to the antibody-epitope fusion protein bind the intact viruses and neutralize their infectivity *in vitro*. The presence of neutralizing antibodies is a key step in evaluation of a vaccine candidate and has been proven to be crucial for the protection of the respiratory tract against RSV infection.^[17] The titers of neutralizing antibodies present in the vaccinated mice strongly suggest that the antibody-epitope vaccine will protect in animal models of RSV infection. To this end we will next investigate the magnitude and durability of these responses as well as the *in vivo* protection in non-human primates.

In conclusion, we have demonstrated that the RSV F-epitope peptide can be fused to the immunoglobulin domain which supports its native-like helix-turn-helix conformation. The fusion protein is capable of eliciting a strong antibody response with cross-reactivity towards the wild-type RSV F antigen. Moreover, serum from immunized mice contain RSV

neutralizing antibodies which block RSV infection *in vitro*. This work suggests that the immunoglobulin scaffold can serve as a universal and non-immunogenic scaffold for the development of next generation epitope-focused vaccine, including fusion of two or more epitopes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institutes of Health (NIH) Grant 5R01 GM062159-13 (P.G.S.). This manuscript is number 29194 of The Scripps Research Institute.

References

1. a) Nair H, Nokes DJ, Gessner BD, Dherani M, Madhi SA, Singleton RJ, O'Brien KL, Roca A, Wright PF, Bruce N, Chandran A, Theodoratou E, Sutanto A, Sedyaningsih ER, Ngama M, Munywoki PK, Kartasasmita C, Simoes EAF, Rudan I, Weber MW, Campbell H. *Lancet*. 2010; 375:1545. [PubMed: 20399493] b) Glezen WP, Taber LH, Frank AL, Kasel JA. *Am J Dis Child*. 1986; 140:543. [PubMed: 3706232] c) Simoes EAF. *Lancet*. 1999; 354:847. [PubMed: 10485741]
2. Gomez RS, Guisile-Marsollier I, Bohmwald K, Bueno SM, Kalergis AM. *Immunol Lett*. 2014; 162:237. [PubMed: 25268876]
3. a) Murphy BR, Prince GA, Walsh EE, Kim HW, Parrott RH, Hemming VG, Rodriguez WJ, Chanock RM. *J Clin Microbiol*. 1986; 24:197. [PubMed: 3755730] b) Murphy BR, Walsh EE. *J Clin Microbiol*. 1988; 26:1595. [PubMed: 2459154]
4. Lopez JA, Andreu D, Carreno C, Whyte P, Taylor G, Melero JA. *J Gen Virol*. 1993; 74:2567. [PubMed: 7506298]
5. Correia BE, Bates JT, Loomis RJ, Baneyx G, Carrico C, Jardine JG, Rupert P, Correnti C, Kalyuzhnyi O, Vittal V, Connell MJ, Stevens E, Schroeter A, Chen M, MacPherson S, Serra AM, Adachi Y, Holmes MA, Li YX, Klevit RE, Graham BS, Wyatt RT, Baker D, Strong RK, Crowe JE, Johnson PR, Schief WR. *Nature*. 2014; 507:201. [PubMed: 24499818]
6. Wang F, Ekiert DC, Ahmad I, Yu WL, Zhang Y, Bazirgan O, Torkamani A, Raudsepp T, Mwangi W, Criscitiello MF, Wilson IA, Schultz PG, Smider VV. *Cell*. 2013; 153:1379. [PubMed: 23746848]
7. a) Liu T, Fu GS, Luo XZ, Liu Y, Wang Y, Wang RSE, Schultz PG, Wang F. *J Am Chem Soc*. 2015; 137:4042. [PubMed: 25775396] b) Wang RSE, Liu T, Wang Y, Cao Y, Du JT, Luo XZ, Deshmukh V, Kim CH, Lawson BR, Tremblay MS, Young TS, Kazane SA, Wang F, Schultz PG. *J Am Chem Soc*. 2015; 137:3229. [PubMed: 25699419] c) Liu T, Zhang Y, Liu Y, Wang Y, Jia HQ, Kang MC, Luo XZ, Caballero D, Gonzalez J, Sherwood L, Nunez V, Wang DL, Woods A, Schultz PG, Wang F. *Proc Natl Acad Sci U S A*. 2015; 112:1356. [PubMed: 25605877]
8. Collins PL, Dickens LE, Bucklerwhite A, Olmsted RA, Spriggs MK, Camargo E, Coelingh KVV. *Proc Natl Acad Sci U S A*. 1986; 83:4594. [PubMed: 3460060]
9. a) Taylor G, Stott EJ, Bew M, Fernie BF, Cote PJ. *Lancet*. 1983; 2:976. [PubMed: 6138541] b) Walsh EE, Schlesinger JJ, Brandriss MW. *Infect Immun*. 1984; 43:756. [PubMed: 6363300]
10. Agenbach E, Tiemessen CT, Venter M. *Virus Genes*. 2005; 30:267. [PubMed: 15744582]
11. McLellan JS, Chen M, Kim A, Yang YP, Graham BS, Kwong PD. *Nat Struct Mol Biol*. 2010; 17:248. [PubMed: 20098425]
12. Ponce R, Abad L, Amaravadi L, Gelzleichter T, Gore E, Green J, Gupta S, Herzyk D, Hurst C, Ivens IA, Kawabata T, Maier C, Mounho B, Rup B, Shankar G, Smith H, Thomas P, Wierda D. *Regul Toxicol Pharmacol*. 2009; 54:164. [PubMed: 19345250]

13. a) Liu T, Liu Y, Wang Y, Hull M, Schultz PG, Wang F. J Am Chem Soc. 2014; 136:10557. [PubMed: 25041362] b) Zhang Y, Zou HF, Wang Y, Caballero D, Gonzalez J, Chao E, Welzel G, Shen WJ, Wang DL, Schultz PG, Wang F. Angew Chem Int Ed. 2015; 54:2126.
14. Eidem JK, Rasmussen IB, Lunde E, Gregers TF, Rees AR, Bogen B, Sandlie I. J Immunol Methods. 2000; 245:119. [PubMed: 11042289]
15. McKimm-Breschkin JL. J Virol Methods. 2004; 120:113. [PubMed: 15234816]
16. Johnson TR, Rangel D, Graham BS, Brough DE, Gall JG. Mol Ther. 2014; 22:196. [PubMed: 23752342]
17. Prince GA, Horswood RL, Chanock RM. J Virol. 1985; 55:517. [PubMed: 4020957]

Figure 1.

Vaccine design. (A) Crystal structure of bovine antibody BLV1H12 (PDB code 4K3D) shows an ultralong CDR3 with a disulfide cross-linked “knob” domain on top of a solvent-exposed β -strand “stalk”. (B) Crystal structure of Mota in complex with its peptide epitope (dark, enlarged) from RSV F protein (PDB code 3IXT). (C) Map of the key elements of EpiS and its fusion to Herceptin. Numbers indicate the fusion sites.

Figure 2.

Binding affinities of the antibody-epitope fusion proteins with Mota. (A) ELISA analysis for hH3-EpiS, hL3-EpiS, hLoop3-EpiS and mLoop3-EpiS fusion proteins binding with Mota. (B) ELISA determination of the apparent K_d for hH3-EpiS and mLoop3-EpiS with Mota, wild-type Herceptin as a control. A plate coated with Mota Fab at 1 μ g per well was blocked with 2% BSA in PBS and incubated with (A) different fusion proteins, or wild-type Herceptin and OKT3 as controls, at 0.5 μ g/mL or (B) increasing concentration of hH3-EpiS, mLoop3-EpiS or Herceptin, followed by the addition of HRP-conjugated anti-human IgG (Fc specific) or anti-mouse IgG (Fc specific). After incubation and wash, 100 μ L QuantaBlu fluorogenic ELISA substrate was added to each well and the signals were monitored using a plate reader (EX 325 nm/EM 420 nm). Assays were performed in triplicate and error bars represent the standard deviation.

Figure 3.

(A) Mouse vaccination study. Immune responses 7 weeks after the first immunization (3 weeks after the third immunization) with mLoop3-EpiS showed significant differences in titers against the fusion protein mLoop3-EpiS and the RSV F protein, but no difference for OKT3-WT. A plate coated with Mota Fab at 1 μ g per well was blocked with 2% BSA in PBS and incubated with 1:100 diluted serum samples at room temperature for 1 hour, followed by the addition of HRP-conjugated anti-mouse IgG (Fc specific). After incubation and wash, 100 μ L QuantaBlu fluorogenic ELISA substrate was added to each well and the signals were monitored using a plat reader (EX 325 nm/EM 420 nm). X axis represents the coating agents. * P <0.05, ** P <0.01 vs. control group; n=10 indicates the number of mice per group. (B) Serum from mice immunized with mLoop3-EpiS significantly block the viral infection of Hep-2 cells. Serum dilutions (1:10, 1:100, 1:1000) were mixed 1:1 with RSV virus and the mixture was inoculated on Hep-2 cell monolayer. After 4 hour incubation, the inoculum was removed and 3 ml overlay of DMEM/F12 with 0.3% ICN immunodiffusion grade agarose was added. The plates were fixed after 7 day incubation and the plaques were counted after visualized by the addition of 0.05% neutral red. *** P <0.001 vs. control group; n=10 indicates the number of mice per group.