Published in final edited form as:

Angew Chem Int Ed Engl. 2015 November 23; 54(48): 14447–14451. doi:10.1002/anie.201508331.

Gold(I)-Catalyzed Enantioselective Desymmetrization of 1,3-Diols *via* Intramolecular Hydroalkoxylation of Allenes^{**}

Dr Weiwei Zi and Prof. Dr. F. Dean Toste

Department of Chemistry, University of California, Berkeley, Berkeley, CA 94720 (USA)

F. Dean Toste: fdtoste@berkeley.edu

Abstract

A gold(I)-catalyzed enantioselective desymmetrization of 1,3-diols was achieved via intramolecular hydroalkoxylation of allenes. The catalyst system 3-F-dppe(AuCl)₂/(R)-C₈-TRIPAg proved to be specifically efficient to promote the desymmetrizing cyclization of 2-aryl-1,3-diols, which have proven challenging substrates in previous reports. Multisubstituted tetrahydrofurans were prepared in good yield with good enantioselectivity and diastereoselectivity by this method.

Graphical Abstract

A chiral anion mediated enantioselective gold(I)-catalyzed desymmetrization of 1,3-diols by intramolecular allene hydroalkoxylation was developed. Subtle tuning of the both the chiral phosphate (X^{*}) and the achiral phosphine ligand (L) components of the catalyst system allowed for the preparation of oxygen heterocycles containing two stereocenters in high enantio- and diastereoselectivity

Keywords

Gold; desymmetrization; 1,3-diols; enantioselective; hydroalkoxylation; allenes

^{**}We gratefully acknowledge NIHGMS (RO1 GM073932) for financial support. We acknowledge College of Chemistry CheXray (NIH Shared Instrumentation Grant S10-RR027172) and Antonio DiPasquale for X-ray crystallographic data. W. Z. thanks Mark Levin for helpful discussions during the preparation of this manuscript.

Correspondence to: F. Dean Toste, fdtoste@berkeley.edu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201508331>.

Asymmetric desymmetrization of prochiral 1,3-diols provides an indirect but powerful way to form stereogenic centers, especially for the synthesis of chiral all-carbon quaternary centers.^[1] During the past decade, substantial effort has been made towards the development of methods for the enantioselective intermolecular desymmetrization of such substrates,^[2,3] On the other hand, intramolecular asymmetric desymmetrization reactions of 1,3-diols to generate two or more stereogenic centers have only recently been realized via organocatalysis.^[4] Sun described a chiral phosphoric acid-catalyzed intramolecular transacetalization of 1,3-diols to form tetrahydrofuran skeletons with high efficiency and stereoselectivity.^[4a] Yeung reported enantioselective desymmetrizing bromoetherification reactions of olefinic 1,3-diols catalyzed by a quinidine derived amino-thiocarbamate^[4b] or a C2 symmetric sulfide.^[4c]

In contrast, transition-metal catalyzed reactions for the intramolecular asymmetric desymmetrization of 1,3-diols have been far less explored.^[5] Palladium-catalyzed asymmetric C-O bond formation, including allylic alkylation reaction^[5a] and Ullmann-type coupling reaction^[5b], were designed for this purpose; however, only modest enantioselectivities were obtained. A copper-catalyzed consecutive desymmetrization and kinetic resolution sequence was designed to obtain highly enantioenriched products.^[5c]

Gold-catalyzed asymmetric hydrofunctionalization of allenes has drawn extensive interest during the last decade.^[6] These reactions serve as atom-economical and highly stereoselective methods to assemble commonly encountered heterocycle motifs found in natural products and bioactive molecules. Despite these achievements, no successful method has been established to construct heterocycles incorporating more than one stereocenter via these transformations. In 2007, our group reported an enantioselective hydroalkoxylation of allenes enabled by a chiral counter anion strategy.^[7] Herein we disclose our recent efforts towards a gold-catalyzed desymmetrizing hydroalkoxylation of allenes that implements this same tactic, and provides access to multisubstituted tetrahydrofurans containing two stereogenic centers with good enantioselectivity and excellent diastereoselectivity.

Previous studies have demonstrated that the chiral induction induced by a chiral counteranion can be sharply modulated by achiral ligands.^[8] Thus, our exploration commenced with the examination of varying achiral phosphine ligands and their synergistic properties in combination with Ag-(S)-TRIP in the desymmetrization of 1,3-diols. The fact that 2-aryl-1,3-diol (R = Ar, Figure 1) derived substrates were absent in Sun and Yeung's reports,^[4] prompted us to employ 2-phenyl substituted 1,3-diol **1a** as the model substrate (Table 1). Various mono- and bis-phosphines ligands for gold were first investigated. Excellent diastereoselectivity and reactivity was obtained with monophosphine ligands (Ph₃P, t-Bu₃P); however, the enantioselectivity was very low (entries 1 and 2). Switching to the a bisphosphine ligand, diphenylphosphinomethane (dppm), produced an improvement in the enantioselectivity of the reaction, but with a slight loss of diastereoselectivity (d.r. = 15:1, 50% ee). Other bisphosphine ligands were subsequently examined (entries 4 and 5).^[9] Among them, diphenylphosphinoethane (dppe) proved optimal not only in enantioselectivity but also in diastereoselectivity (>25:1 d.r., 57% ee).

Further optimization of the ligand focused on the adjustment of its electronic properties. Neither strongly electron-donating groups (4-MeO) nor strongly electron-withdrawing groups (4-CF₃) were beneficial to the reaction (entries 3, 6 and 7). However, a mild electron withdrawing group (4-F) increased the ee to 66% (entry 8). Additional enhancement of the enantioselectivity was observed by employing 3-F-dppe as the ligand instead of 4-F-dppe (70% ee, entry 9). The structure of the counter anion was also examined, with (R)-C₈-TRIP providing the desired product **2a** in 92% yield with 87% ee as a single diastereoisomer (entry 12). Finally, performing the reaction at increased dilution and low temperature resulted in further improvements to the enantioselectivity (entries 13, 14).

With the optimal reaction conditions in hand, we next examined the substrate scope of this reaction (Table 2). The allene moiety (R₁) was investigated, with six-membered ring substituent on the allene affording the desired product **2b** in 95% yield with 90% ee (entry 2). By contrast, slightly poorer results were obtained for the five-membered ring containing substrate (entry 3). Substrates with an electron-withdrawing group (F, Cl, Br, CF₃), an electron-donating group (EtO, Me), or two substituents on the aryl ring all were well-tolerated (entries 4 to 14). Additionally, reactions for substrates bearing a naphthalene or indole proceeded smoothly under the standard reaction conditions (entries 15 and 16).

We next turned our attention to substrates varying at the nascent desymmetrized quaternary carbon. When a benzyl group was introduced, modified conditions (−35 °C, 60 hrs) were needed to maintain good enantioselectivity (entry 17). Substrate with a cinnamyl group was also tested, giving the hydroalkoxylation product **2r** in 87% yield, without any [2 + 2] adduct formed under this catalysts system (entry 18).^[10] Similarly, no hydroarylation of allene was observed during the preparation of **2s** by this method (entry 19).^[11] These results feathered the counterion controlled reactivity in homogeneous gold catalysis.^[12] Finally, the reaction was easily extended to substrates incorporating a protected amine functionality at the C₂ position. While prolonged reaction time was required, the desired tetrahydrofuran with a protected amine moiety was isolated in 74% yield with 87% enantioselectivity (entry 20). Attempts to prepare tetrahydropyran product **2u** by this desymmetrization method under the standard conditions (−10 °C for 24 hrs) resulted in low conversion (< 10%). Elevated reaction temperature could expedite the transformation, however, modest diastereoselectivity and enantioselectivity was obtained (d.r. = 3.8:1, 69% ee, entry 21).^[13]

The absolute stereochemistry of the hydroalkoxylation product **1a** was determined by transformation to its sulfonylated derivative **3**. X-ray crystallographic analysis of **3** confirmed its structure and disclosed the absolute configuration as 2S, 4S.^[14] Synthetic transformation of the desymmetrization products are illustrated in Scheme 1. The alkene moiety in **2a** was readily converted to the corresponding aldehyde, which was transformed to α,β-unsaturated ester using the Horner-Wadsworth-Emmons olefination reaction. Moreover, the hydroxyl in **2a** was extended to an ester by oxidation/olefination sequence. Additionally, spirocyclic ether **2ma** was preped through an Ullmann type C-O bond formation reaction of the adduct (**2m**) bearing a halogen group at ortho position on the phenyl ring.

To gain insight into the mechanism, some control experiments were conducted. First, when the gold catalyst was omitted, no reaction occurred, excluding a chiral silver-catalyzed hydroalkoxylation as an operative mechanism [Eq. (1)].^[15] Second, the ratio of Au/Ag was investigated. Although excess (R)-C₈-TRIPAg had no obvious effect to the enantioselectivity, we found that when the ratio of Au/Ag was 2:1, the enantiomeric excess dropped dramatically to 78% [Eq. (2)].^[16] These observations raise the possibility that both gold centers in the catalyst play an important role in enantioinduction. While it is tempting to invoke a mechanism involving dual activation of the allene leading to a *gem*-diaurated intermediate, these species are generally believed to be less reactive towards protodeauration and, therefore, lie off the catalytic cycle.^[17] Moreover, we observed a clear linear relationship^[18] ($r^2 = 0.996$) between the enantiomeric excess of the phosphate catalyst and that of the product (Figure 4). Therefore, it seems most plausible that the second gold phosphate center in the dinuclear catalyst is providing a structural or steric effect^[19] rather than dual chiral induction.

In summary, we have achieved the first gold(I)-catalyzed enantioselective desymmetrization of 1,3-diols by intramolecular hydroalkoxylation of allenes. With this method, multisubstituted oxygen heterocycles, bearing all-carbon quaternary stereogenic centers are easily accessed in good yield with high enantioselectivity and diastereoselectivity. Notably, in addition to selecting the appropriate chiral anion, tuning of the steric and electronic properties of the achiral phosphine supporting ligand proved crucial to identifying the ideal catalyst system. This reaction represents an important further example of power of the chiral anion strategy for transition metal catalysis.^[20]

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. For recent reviews on on desymmetrization of diols: Müller CE, Schreiner PR. *Angew Chem Int Ed*. 2011; 50:6012–6042. Enríquez-García Á, Kündig EP. *Chem Soc Rev*. 2012; 41:7803–7831. [PubMed: 23007376]
2. a) Trost BM, Mino T. *J Am Chem Soc*. 2003; 125:2410–2411. [PubMed: 12603126] b) T. Honjo, M. Nakao, S. Sano, M. Shiro, K. Yamaguchi, Y. Sei, Y. Nagao, *Org. Lett*. 2007, 9, 509–512; c) Jung B, Kang SH. *Proc Natl Acad Sci U S A*. 2007; 104:1471–1475. [PubMed: 17242359] d) B. Jung, M. S. Hong, S. H. Kang, *Angew. Chem., Int. Ed*. 2007, 46, 2616–2618; e) Lee JY, You YS, Kang SH. *J Am Chem Soc*. 2011; 133:1772–1774. [PubMed: 21265528] f) Z. You, A. H. Hoveyda, M. L. Snapper, *Angew. Chem. Int. Ed*. 2009, 47, 547–550; g) Meng SS, Liang Y, Cao KS, Zou L, Lin XB, Yang H, Houk KN, Zheng WH. *J Am Chem Soc*. 2014; 136:12249–12252. [PubMed: 25126836]
3. For application towards natural product syntheses, see: Celindro NC, Kim TW, Kang SH. *Chem Commun*. 2012; 48:6295–6297. Lee W, Youn JH, Kang SH. *Chem Commun*. 2013; 49:5231–5233. Kim HC, Kang SH. *Angew Chem, Int Ed*. 2009; 48:1827–1829.
4. a) Chen Z, Sun J. *Angew Chem, Int Ed*. 2013; 52:13593–13596. b) D. W. Tay, G. Y. C. Leung, Y.-Y. Yeung, *Angew. Chem., Int. Ed*. 2014, 53, 5161–5164; c) Ke Z, Tan CK, Chen F, Yeung YY. *J Am Chem Soc*. 2014; 136:5627–5630. [PubMed: 24697791]
5. a) Kubiak A, Porwaski S, Zawisza A. *Tetrahedron: Asymmetry*. 2013; 24:212–216. b) Yang W, Yan J, Long Y, Zhang S, Liu J, Zeng Y, Cai Q. *Org Lett*. 2013; 15:6022–6025. [PubMed: 24261632] c) Y. S. You, T. W. Kim, S. H. Kang, *Chem. Commun.*, 2013, 49, 9669–9671; During

the preparation of this manuscript, a copper-catalyzed intramolecular desymmetrization of 1,3-diols was reported, see: d) Yang W, Liu Y, Zhang S, Cai Q. *Angew Chem Int Ed*. 2015; 54:8805–8808.

6. For selected examples on asymmetric hydrofunctionalization of allenes, see: Aikawa K, Kojima M, Mikami K. *Adv Synth Catal*. 2010; 352:3131–3135. Butler KL, Tragni M, Widenhoefer RA. *Angew Chem, Int Ed*. 2012; 51:5175–5178. Zhang Z, Widenhoefer RA. *Angew Chem, Int Ed*. 2007; 46:283–285. Zhang Z, Bender CF, Widenhoefer RA. *J Am Chem Soc*. 2007; 129:14148–14149. [PubMed: 17967025] Liu C, Widenhoefer RA. *Org Lett*. 2007; 9:1935–1938. [PubMed: 17428061] LaLonde RL, Wang ZJ, Mba M, Lackner AD, Toste FD. *Angew Chem, Int Ed*. 2010; 49:598–601. LaLonde RL, Sherry BD, Kang EJ, Toste FD. *J Am Chem Soc*. 2007; 129:2452–2453. [PubMed: 17290999]
7. a) Hamilton GL, Kang EJ, Mba M, Toste FD. *Science*. 2007; 317:496–499. [PubMed: 17656720] b) R. L. LaLonde, Z. J. Wang, M. Mba, A. D. Lackner, F. D. Toste *Angew. Chem. Int. Ed*. 2010, 49, 598–601. For reviews on asymmetric gold catalysis: c) Pradal A, Toullec PY, Michelet V. *Synthesis*. 2011; 10:1501–1514. d) S. Sengupta, X. Shi, *ChemCatChem* 2010, 2, 609–619; e) Wang YM, Lackner AD, Toste FD. *Acc Chem Res*. 2014; 47:889–901. [PubMed: 24228794]
8. For recent reviews on chiral anion in asymmetric catalysis: Lacour J, Hebbe-Viton V. *Chem Soc Rev*. 2003; 32:373–382. [PubMed: 14671792] Mahlau M, List B. *Angew Chem, Int Ed*. 2012; 28:518–533. Phipps RJ, Hamilton GL, Toste FD. *Nature Chem*. 2013; 6:159–164. [PubMed: 24451593]
9. Sterically demanding chiral bisphosphine ligands ((R)- or (S)-DTMB-Segphos and (R)- or (S)-MeO-DTBM-Biphep) were also examined in combination with Ag-(S)-TRIP, resulting in lower enantioselectivity and poor diastereoselectivity.
10. Luzung MR, Mauleón P, Toste FD. *J Am Chem Soc*. 2007; 129:12402–12403. [PubMed: 17887681]
11. a) Liu C, Widenhoefer RA. *Org Lett*. 2007; 9:1935–1938. [PubMed: 17428061] b) Toups KL, Liu GT, Widenhoefer RA. *J Organomet Chem*. 2009; 694:571–575. [PubMed: 20305794]
12. For review on counterion effects in homogeneous gold catalysis: Jia M, Bandini M. *ACS Catal*. 2015; 5:1638–1652.
13. Attempts to optimize the reaction results for **2u** by adjustment of Achiral bisphosphine gold or Chiral silver phosphate all meet failure.
14. For supplementary crystallographic data see supporting information.
15. Wang Y, Zhang K, Hong R. *J Am Chem Soc*. 2012; 134:4096–4099. [PubMed: 22329839]
16. In stark contrast, Mikami and coworkers found that dicationic BIPHEPgold(I) phosphate catalysts gave lower enantioselectivity in intramolecular hydroalkoxylation reactions (see ref 6a).
17. a) Brown TJ, Weber D, Gagné MR, Widenhoefer RA. *J Am Chem Soc*. 2012; 134:9134–9137. [PubMed: 22621418] b) G. Seidel, C. W. Lehmann, A. Fürstner, A. *Angew. Chem. Int. Ed*. 2010, 49, 8466–8470; c) Weber D, Jones TD, Adduci LL, Gagné MR. *Angew Chem Int Ed*. 2012; 51:2452–2456.
18. For some reviews on the subject of NLEs, see: Girard C, Kagan HB. *Angew Chem, Int Ed*. 1998; 37:2922–2959. Inanaga J, Furuno H, Hayano T. *Chem Rev*. 2002; 102:2211–2226. [PubMed: 12059267]
19. a) Kim JH, Park SW, Park SR, Lee S, Kang EJ. *Chem Asian J*. 2011; 6:1982–1986. [PubMed: 21735554] b) Brooner REM, Widenhoefer RA. *Organometallics*. 2012; 31:768–771. c) Bandini M, Bottoni A, Chiarucci M, Cera G, Miscone GP. *J Am Chem Soc*. 2012; 134:20690–20700. [PubMed: 23193975]
20. For examples of chiral anion strategy in transition metal catalysis, see: Mukherjee S, List B. *J Am Chem Soc*. 2007; 129:11336–11337. [PubMed: 17715928] Liao S, List B. *Angew Chem, Int Ed*. 2010; 49:628–631. Rauniyar V, Wang ZJ, Burks HE, Toste FD. *J Am Chem Soc*. 2011; 133:8486–8489. [PubMed: 21561153] Ohmatsu K, Ito M, Kunieda T, Ooi T. *Nature Chem*. 2012; 4:473–477. [PubMed: 22614382] Nelson HM, Williams BD, Miró J, Toste FD. *J Am Chem Soc*. 2015; 137:3213–3216. [PubMed: 25723255]

A. Transacetalization (Sun, 2013) **B. Bromoalkoxylation (Yeung, 2014)** **C. This work**

Figure 1.
Strategies for enantioselective intramolecular desymmetrization of 1,3-diols

Figure 2.
Absolute stereochemistry of **3** determined by X-ray crystallography.

Figure 3.
Relationship between the optical purity of product **2d** and catalyst using 3-F-dppe(AuCl)₂/ H_8 -TripAg in toluene.

Scheme 1.

Synthetic transformation of the products. Reaction conditions: a) O₃, CH₂Cl₂, -78°C then Me₂S from 78°C to room temperature. b) (EtO)₂P(O)CH₂CO₂Me, NaH, THF. c) Dess-Martin periodinane, NaHCO₃, CH₂Cl₂. d) CuI, NaH, toluene, 120°C. See supporting information for details.

Scheme 2.
Control experiments.

Table 1

Optimization of the reaction conditions.^[a]

entry ^[a]	Ligand	Anion	d.r. ^[b]	yield (%) ^[c]	ee (%) ^[d]
1	Ph3P	(S)-TRIP	>25:1	90	-28
2	t-BuP3	(S)-TRIP	>25:1	>95	-4
3	dppm	(S)-TRIP	15:1	87	-50
4	dppe	(S)-TRIP	>25:1	93	-57
5	dppp	(S)-TRIP	18:1	>95	-41
6	4-MeO-dppe	(S)-TRIP	>25:1	>95	-39
7	4-CF3-dppe	(S)-TRIP	14:1	84	-28
8	4-F-dppe	(S)-TRIP	>25:1	>95	-66
9	3-F-dppe	(S)-TRIP	>25:1	>95	-70
10	3-F-dppe	(R)-H8-TRIP	18:1	>95	81
11 ^e	3-F-dppe	(R)-C8-TCYP	>25:1	>95	83
12	3-F-dppe	(R)-C8-TRIP	>25:1	94	87
13 ^f	3-F-dppe	(R)-C8-TRIP	>25:1	92	91
14 ^g	3-F-dppe	(R)-C8-TRIP	>25:1	>95	93

^[a] Reaction conditions: 2.5 mol % L(AuCl)₂ or 5.0 mol % LAuCl, 5 mol % AgX*, 0.05 mmol substrate in 0.5 mL toluene (C = 0.1 M), room temperature, 2 hrs.^[b] d.r. was determined by ¹H NMR analysis of the crude product.^[c] ¹H NMR yield based on internal standard.^[d] determined by chiral HPLC.^[e] full conversion after 12 h.^[f] C = 0.01 M.^[g] Reaction was run at -10°C with 20 mg 4Å molecular sieve as additive.

Table 2

Scope of substrates.^{[a],[b]}

^[a] Reaction conditions: 2.5 mol % 3-F-dppe(AuCl)₂, 5 mol % (R)-C₈-TRIPAg, 0.1 mmol substrate in 10 mL toluene (C = 0.01 M), 40 mg 4 Å molecular sieve, -10°C for 24 hrs. See supporting information for details.

^[b] d.r. was determined by ¹H NMR analysis of the crude product and ee was determined by chiral HPLC. Isolated yield.

^[c] - 35°C for 60 hrs.

^[d] - 10°C for 72 hrs.

^[e] 20°C for 24 hrs.