Published in final edited form as:

J Cell Biochem. 2016 February ; 117(2): 267–278. doi:10.1002/jcb.25283.

Hypoxia-Inducible Factor-1: A critical player in the survival strategy of stressed cells

Shuyang Chen¹ and Nianli Sang^{1,2,3}

¹Department of Biology and Graduate Program of Biological Sciences, College of Arts & Sciences, Drexel University, Philadelphia, PA, USA

²Department of Pathology and Laboratory Medicine, Drexel University College of Medicine, Philadelphia, PA, USA

³Sydney Kimmel Cancer Center, Philadelphia, PA, USA

Abstract

HIF-1 activation has been well known as an adaptive strategy to hypoxia. Recently it became clear that hypoxia was often accompanied by insufficient supply of glucose or amino acids as a common result of poor circulation that frequently occurs in solid tumors and ischemic lesions, creating a mixed nutrient insufficiency. In response to nutrient insufficiency, stressed cells elicit survival strategies including activation of AMPK and HIF-1 to cope with the stress. Particularly, in solid tumors, HIF-1 promotes cell survival and migration, stimulates angiogenesis, and induces resistance to radiation and chemotherapy. Interestingly, radiation and some chemotherapeutics are reported to trigger the activation of AMPK. Here we discuss the recent advances that may potentially link the stress responsive mechanisms including AMPK activation, ATF4 activation and the enhancement of Hsp70/Hsp90 function to HIF-1 activation. Potential implication and application of the stress-facilitated HIF-1 activation in solid tumors and ischemic disorders will be discussed. A better understanding of HIF-1 activation in cells exposed to stresses is expected to facilitate the design of therapeutic approaches that specifically modulate cell survival strategy.

Keywords

AMPK; HDAC4; HDAC5; HIF-1; Hsp70; Hsp90; Stress response; Hypoxia

Introduction

Hypoxia-inducible factor –1 (HIF-1) was first discovered in the studies of hypoxia-triggered elevation of human erythropoietin (EPO), a glycoprotein hormone made in renal cells and fetal hepatocytes which stimulates bone marrow hematopoietic cell proliferation to produce red blood cells [Semenza and Wang, 1992]. Early purification work using DNA affinity chromatography and subsequent biochemical characterization revealed that HIF-1 is a heterodimer consisting of HIF-1 α and HIF-1 β [Wang and Semenza, 1995]. HIF-1 α protein levels increase dramatically when oxygen concentration decreases [Wang et al., 1995]. As a

transcription factor, activated HIF-1 enters the nucleus, recruits coactivator p300/CBP, and reprograms gene expression to alter cellular processes, facilitating cell survival under transient hypoxic condition. Since the identification of HIF-1, more than 1,000 HIF-1 targets have been identified by a variety of approaches [Manalo et al., 2005; Schodel et al., 2011; Xia et al., 2009]. Metabolically, activation of HIF-1 switches the ATP producing mode from aerobic respiration to the oxygen-independent fermentation, a more rapid, but less efficient way of glucose utilization. Consequently, HIF-1 drives the overexpression of glucose transporter 1 (GLUT1), which increases glucose uptake to compensate the low efficiency. Increased lactate dehydrogenase (LDH) accelerates the conversion of pyruvate and NADH to lactate and NAD⁺, respectively, ensuring an undisrupted glycolytic process and maintaining an intracellular redox balance. HIF-1 also stimulates the expression of carbonic anhydrase 9 (CA-IX), which maintains intracellular pH by exporting protons to the extracellular matrix.

In this review, we will briefly summarize the activation of HIF-1 by hypoxia to set the platform for the subsequent discussion of the control of HIF-1 α translation and posttranslational processing as a survival strategy of stressed cells (Figure 1). Integrating knowledge in literature and recent findings, we propose a crucial role of HIF-1 as a facilitator of adaptation to a variety of cellular stresses such as proliferative biosynthesis, nutrient insufficiency and radiation therapy.

Oxygen Sensing Pathways Activate HIF-1 by Hypoxic Stabilization of Mature HIF-1 α

As a general final electron acceptor in oxidative phosphorylation and other redox reactions, molecular oxygen is essential for mammalian cell survival, energy homeostasis and other cellular processes. A constant supply of oxygen is indispensable for maintaining normal tissue functions, particularly cardiac and brain function. Hypoxia, a state in which oxygen supply is insufficient, has been involved in the pathogenesis of multiple diseases with top leading mortality rates, including heart diseases, chronic respiration disease, stroke, anemia and bleeding [Kochanek et al., 2014]. Hypoxia also contributes to tumor progression, metastasis and resistance to radiation and chemotherapy [Kaelin and Ratcliffe, 2008; Semenza, 2003; Semenza, 2012]. The identification of HIF-1 as a major regulator of adaptation to hypoxia gave us a better understanding of the molecular mechanisms underlying oxygen sensing pathways, oxygen homeostasis, energy homeostasis and disease pathology [Doedens et al., 2013; Keith and Simon, 2007; Semenza, 2014; Takubo et al., 2010; Zhang et al., 2011].

Structurally, HIF-1 is a basic helix-loop-helix (bHLH) transcription factor of the PER-ARNT-SIM (PAS) subfamily. HIF-1 α and HIF-1 β are structurally similar, each containing a bHLH domain for DNA binding, a PAS domain for hetero-dimerization and transactivation domain for transcriptional activity regulation [Semenza, 2003] (Figure 2). HIF-1 β , also known as aryl hydrocarbon nuclear translocator (ARNT), is a stable protein, serving as a common dimerization partner for multiple transcription factors, while the protein level and functionality of HIF-1 α determine the transactivation activity of HIF-1. Like HIF-1 β , HIF-1 α is also constitutively translated; unlike HIF-1 β , HIF-1 α protein is subject to

degradation through an oxygen-dependent mechanism (Figure 2). In the presence of adequate molecular oxygen, HIF-1 α protein will be recognized by three prolyl hydroxylases (PHDs), which hydroxylate HIF-1 α at two prolyl residues (P402 and P564) [Bruick and McKnight, 2001; Epstein et al., 2001]. The hydroxylated HIF-1 α is recognized by the von Hippel Lindau protein (VHL), a ubiquitin ligase, leading to HIF-1 α polyubiquitination. The ubiquitinated HIF-1 α will be finally degraded by the 26S proteasome [Huang et al., 1998; Salceda and Caro, 1997].

The PHDs that determine the stability of mature HIF-1 α belong to a dioxygenase family [Schofield and Ratcliffe, 2004]. These enzymes require molecular oxygen and α -ketoglutarate as co-substrates, as well as Fe²⁺ and ascorbate (vitamin C) as co-factors for enzymatic activity (Figure 2) [Hirsilä et al., 2003]. Mutation of succinate dehydrogenase (SDH) and fumarate hydratase (FH), two enzymes of the tricarboxylic acid cycle, was found to block PHDs activity and increase HIF-1 α protein levels by increasing the concentration of succinate and fumarate respectively, which act as PHD inhibitors [Isaacs et al., 2005; Pollard et al., 2005; Selak et al., 2005]. Moreover, reports showed that mutations in the cytosolic isocitrate dehydrogenase (IDH1,2) may result in a decrease of α -ketoglutarate, thus stabilizing HIF-1 α protein [Zhao et al., 2009]. More recent reports demonstrate that mutations in (IDH1 or IDH2) result in the formation of abnormal metabolite (R)-2-hydroxyglutarate, which stimulates PHD activity, thus decreasing HIF-1 α protein levels in gliomas [Koivunen et al., 2012]. However, clinical studies by directly imaging HIF-1 target gene expression in tumors indicate that the relationship between IDH mutations and HIF-1 activity *in vivo* remains unclear [Metellus et al., 2011]. In addition, HIF-1 α transcriptional activity also is regulated by an oxygen-dependent hydroxylation event. Factor inhibiting HIF-1 (FIH) [Mahon et al., 2001], another oxygen-dependent hydroxylase, modifies the asparagine residue (N803) in the carboxyl terminal activation domain and disrupts its interaction with p300/CBP [Lando et al., 2002; Sang et al., 2002]. In summary, the oxygen-dependent hydroxylation of HIF-1 α forms the biochemical basis for the conventional oxygen sensing pathway (Figure 1), representing the initiation of a physiological adaptation to hypoxia.

Proliferative Signaling Pathways Activate HIF-1 by Accelerated Translation

Proliferating cells usually demand more molecular oxygen to support active biosynthesis. Growth factors, which activate the major signaling pathways such as mitogen activated protein kinase (MAPK) and insulin-stimulated PI3K pathways for cell proliferation, also increase HIF-1 α protein synthesis to promote glucose utilization and ATP production [Kaelin and Ratcliffe, 2008; Semenza, 2003; Semenza, 2012]. In solid tumors, activation of oncogenes (Ras, PI3K/AKT) and loss of tumor suppressors (PTEN, LKB1 and TSC2/1) may activate HIF-1 by enhancing HIF-1 α protein synthesis (Figure 3). As in the translation of other proteins, mechanistic target of rapamycin (mTOR, also known as mammalian target of rapamycin), as part of mTOR complex 1 (mTORC1), is the key regulator that transmits growth signals to translational machinery, which is often dysregulated in cancers [Benjamin et al., 2011]. As an integrator of growth signaling pathways and nutrient status, mTOR stimulates protein translation by phosphorylating ribosomal protein S6 kinase (S6K), eukaryotic elongation factor 2 kinase (EEF2K) and eukaryotic translation initiation factor 4E

(EIF4E)-binding protein 1 (4EBP1), leading to increased protein translation [Wouters and Koritzinsky, 2008]. Particularly, insulin/IGF and EGF, the most important cellular signaling pathways that regulate cell proliferation, stimulate mTORC1 [Harris and Lawrence, 2003]. The mechanism by which these signal pathways activate mTORC1 has been highlighted by the discovery that mTOR is suppressed by the tuberous sclerosis complex (TSC) gene products TSC2/1 [Inoki et al., 2005], and TSC2 is phosphorylated and inhibited by AKT [Manning et al., 2002; Potter et al., 2002], which is activated by PI3K [Kumar et al., 2005]. Either stimulating any of the Ras, PI3K, AKT and mTOR signaling pathways or inhibiting TSC2 enhances HIF-1 α translation [Bernardi et al., 2006; Blancher et al., 2001; Brugarolas et al., 2004; Brugarolas et al., 2003; Chen et al., 2001; Hudson et al., 2002; Majumder et al., 2004]. On the contrary, phosphatase and tensin homolog (PTEN), a suppressor of PI3K [Carracedo and Pandolfi, 2008], was found to be a negative regulator of HIF-1 α translation [Zundel et al., 2000].

On the another hand, mTOR signaling is inhibited by activation of AMPK, which phosphorylates TSC2 [Kimura et al., 2003]. Upstream to the AMPK, the LKB1 tumor suppressor gene encodes a serine/threonine kinase [Hawley et al., 2003], and the LKB1-AMPK pathway inactivates mTOR signaling [Shaw et al., 2004]. Protein levels of HIF-1 α and its target GLUT1 are increased in LKB1-deficient and AMPK-deficient fibroblasts, and the epithelia of gastrointestinal hamartomas of Lkb1 $^{+/-}$ mice also show increased HIF-1 α and GLUT1 compared with the surrounding normal tissue [Shackelford et al., 2009]. It is important to note that as the energy sensor, AMPK is activated by low glucose, low glutamine and hypoxia [Davie et al., 2015; Jibb and Richards, 2008; Laderoute et al., 2006; Mungai et al., 2011]. Therefore, nutrients insufficiency represents another fundamental parameter determining rate of cell growth and proliferation as well as HIF-1 α translation [Harris and Lawrence, 2003].

Stresses Facilitate HIF-1 Activation by Enhancing Posttranslational Processing of HIF-1 α

At the cellular level, HIF-1 activation drives the utilization of glucose and rapid production of ATP, a process that is not only critical for proliferating cells, but also facilitates cells to survive stress conditions such as nutrient insufficiency and cell damage caused by a variety of physical, chemical and mechanical factors. How HIF-1 is activated under stress conditions has been revealed by the analysis of a variety of compounds that destabilize HIF-1 α and/or inhibit HIF-1 function (Table 1). While not fully understood, efforts to dissect the molecular mechanisms underlying the repressive effects of drugs on HIF-1 indicate the involvement of the molecular chaperone heat shock protein 90kD (Hsp90) [Isaacs et al., 2002] and heat shock protein 70kD (Hsp70) [Kong et al., 2006; Luo et al., 2010]. They form a quality control (QC) system of HIF-1 α and are responsible for the posttranslational processing and stabilization of nascent HIF-1 α (Figure 4). Hsp90 is one of the most abundant cytosolic proteins, counting 1–2% of total amount of proteins inside mammalian cells [Borkovich et al., 1989]. Hsp90 assists varieties of proteins (known as clients), including key regulatory proteins such as steroid hormone receptors, transcription factors, and kinases, to fold properly into their active conformation by consuming energy

from ATP hydrolysis, which buffers proteostasis against environmental stress [Whitesell and Lindquist, 2005]. Usually Hsp90 does not recognize client proteins in the nascent state; it interacts with other co-chaperones, which guide the client recognition and delivery of client proteins. Hsp70 is a well-known co-chaperone that works with Hsp90 to form the chaperone cycle [Pratt and Toft, 2003; Wegele et al., 2004]. The monomeric Hsp70 recognizes short hydrophobic motifs in client proteins, which are commonly exposed in nascent polypeptide chains. The newly synthesized client proteins then associate with Hsp70 to form an early complex, leading to the delivery of client proteins to Hsp90 to form the mature complex [Pratt and Toft, 2003; Taipale et al., 2010; Wegele et al., 2004]. The Hsp90 oligomeric double ring structure encloses the entire nascent client protein and ensures the correct folding [Pratt and Toft, 2003; Taipale et al., 2010; Wegele et al., 2004]. The dynamic interaction between HIF-1 α and Hsp90 was found to facilitate the stabilization of HIF-1 α [Katschinski et al., 2004], indicating HIF-1 α is one of the client proteins of the Hsp70/Hsp90 system. This was further supported by the evidence that Hsp90 inhibitors, such as 17-allylaminogeldanamycin (17-AAG), induce proteasomal degradation of HIF-1 α in the absence of oxygen and in a VHL-independent manner [Isaacs et al., 2002; Kong et al., 2006; Mabeesh et al., 2002; Zagzag et al., 2003].

Another group of compounds, the histone deacetylase (HDAC) inhibitors, including trichostatin A (TSA), SAHA (vorinostat), sodium butyrate and LAQ824, were also reported to induce proteasomal degradation of HIF-1 α in a VHL-independent manner similar to Hsp90 inhibitors [Fath et al., 2006; Qian et al., 2006] (Table 1). Those HDACs with suppressive effect on HIF-1 usually non-selectively inhibit all Class I and Class II HDACs, including HDAC1–9 [Chen and Sang, 2011; Yang and Seto, 2007]. Recently, a member of Class IIa HDACs, HDAC5, has been identified as the specific HDAC member that facilitates the stabilization and nuclear accumulation of HIF-1 α protein [Chen et al., 2015]. HDAC5 dynamically shuttles between the nucleus and the cytosol and has relatively weak enzymatic activity on the histones [Fischle et al., 2002; McKinsey et al., 2001]. New findings reveal that Hsp70 is a cytosolic substrate of HDAC5 [Chen et al., 2015], and deacetylation of the Hsp70 by HDAC5 promotes HIF-1 α interaction with Hsp90, thus facilitating the rapid nuclear accumulation of HIF-1 α [Chen et al., 2015]. Retrospectively, both HDAC4 and HDAC6 were proposed to destabilize HIF-1 α and repress HIF-1 function [Geng et al., 2011; Kong et al., 2006; Qian et al., 2006]. HDAC4 is highly homologous to HDAC5 [McKinsey et al., 2001]. We found that the knockdown of either HDAC4 or HDAC5 reduces HIF-1 α protein levels and suppresses HIF-1 function (Figure 5A, B). Moreover, expression of an HDAC4 mutant S265/266A, which is exclusively localized in the cytosol, is sufficient to stabilize HIF-1 α (Figure 5C). This indicates that cells expressing HDAC4 might facilitate HIF-1 α stabilization through a similar mechanism. However, hypoxia only increases the protein level of HDAC5 [Chen et al., 2015], not HDAC4 (Figure 5A). Their tissue-specific expression levels in different tissues also vary dramatically; HDAC4 is highly expressed in myeloid, whereas HDAC5 is highly expressed in heart, neuron, muscle, lung, and placenta, which represent the tissues most sensitive to physiological fluctuation of oxygen levels and metabolic stress [Chang et al., 2004]. Therefore, both HDAC4 and HDAC5 may be involved in the regulation of posttranslational processing of HIF-1 α by regulating the Hsp70/Hsp90 system, but it is likely that HDAC5

plays a more important role in facilitating HIF-1 α folding in response to hypoxic stress. HDAC6 inhibition was proposed to increase the acetylation level of Hsp90, thus impairing Hsp90 function [Fath et al., 2006; Kovacs et al., 2005]. However, romidepsin (FK228,depsipeptide), an HDAC inhibitor that has very weak repressive effect on HDAC6, effectively destabilizes HIF-1 α , making it possible that it also destabilizes HIF-1 α through inhibiting HDAC5 or HDAC4, leading to Hsp70 hyperacetylation [To et al., 2011; Wang et al., 2007].

AMPK, the master regulator of cellular energy homeostasis, is activated by nutrient insufficiency including low glucose, low glutamine and hypoxia [Davie et al., 2015; Jibb and Richards, 2008; Laderoute et al., 2006; Mungai et al., 2011]. AMPK is a highly conserved serine or threonine heterotrimeric kinase consisting of α , β , and γ subunits [Winder and Hardie, 1999]. Nutrient insufficiency causes an increase of intracellular AMP; the binding of AMP to AMPK- γ subunit exposes the catalytic domain on AMPK- α , thus activating AMPK [Adams et al., 2004]. The canonical function of AMPK activation is to maintain the intracellular ATP homeostasis by stimulating ATP producing catabolic pathways but inhibiting ATP consuming biosynthetic pathways. Interestingly, AMPK is also activated when cells are challenged with cell damage agents such as increased work load, heat, radiation and pathogenic microbials, indicating that AMPK activation may have a universal and critical role in cells' survival strategy upon stresses. As an ATP conserving strategy, AMPK activation inhibits general protein translation by suppressing mTOR signaling pathways, including HIF-1 α [Shackelford and Shaw, 2009]. However, HIF-1 is the major driver of glucose utilization, which provides rapid ATP generation for cells to survive and sustain certain key functions under stressful conditions [Chen et al., 2015; Davie et al., 2015]. It had been unclear until recently how HIF-1 function is sustained under the stress conditions. In fact, activated AMPK phosphorylates and promotes the nuclear export of HDAC5 and perhaps HDAC4 as well [McKinsey et al., 2001]. In the cytosol, HDAC4 and HDAC5 stabilize HIF-1 α by deacetylating Hsp70, promoting the transfer of HIF-1 α from Hsp70 to Hsp90 for the completion of the maturation process. Therefore, while the AMPK may reduce the translation of HIF-1 α , it compensates it by enhancing the efficiency of posttranslational folding of HIF-1 α , thus ensuring HIF-1 activation under stressful conditions [Chen et al., 2015].

It is important to point out that for normal cells under physiological conditions, a basal level of Hsp70/Hsp90 activity may be sufficient to process newly synthesized proteins. However, under conditions that new proteins are translated in large quantities (proliferative biosynthesis, overexpressing recombinant proteins), an enhanced capacity of Hsp70/Hsp90 may become crucial for the processing of newly synthesized critical proteins such as HIF-1 α . Hence, suppression of the enhancement of Hsp70/Hsp90 function may disrupt the formation of correctly folded, functional active HIF-1 α , leading to disposal of misfolded HIF-1 α by a quality control system. This disposal appears to involve a hydroxylation-independent and ubiquitination-independent proteasome degradation (UIP) [Kong et al., 2006; Lao et al., 2012; Liang et al., 2006], hence being mechanistically distinguishable from the oxygen-triggered degradation (Figure 1). Blocking this process with proteasome inhibitors or overexpressing HIF-1 α increases the total levels of HIF-1 α , but may not significantly activate HIF-1 function [Chen et al., 2015; Kaluz et al., 2006], demonstrating

the importance of the posttranslational processing mediated by Hsp70/Hsp90. Therefore, the stress-activated AMPK-HDAC5-Hsp70/Hsp90-HIF-1 pathway may function as a critical but universal survival strategy for cells to cope with a variety of stresses, including but not limited to hypoxia.

In addition to the enhanced Hsp70/Hsp90 function, selective translation of ATF4, a transcription factor often activated during endoplasmic reticulum stress and unfolded protein response, also occurs as a result of cellular stress and mTORC1 inhibition [Rutkowski and Kaufman, 2003]. It has been shown that the activation of ATF4 is critical for cell adaptation to nutrient depletion [Ye et al., 2010]. Well known target genes activated by ATF4 are DDIT3 (CHOP) and stanniocalcin 2 (STC2). However, it remains unclear if there is a link among the ATF4-pathway, AMPK activation and HIF-1 activation. Finally, it is reported that a stress-activated protein kinase, JNK1, is required for HIF-1 α stabilization in hypoxic tissues [Antoniou et al., 2009], providing another link between cell stress and HIF-1 activation.

Relevance of HIF-1 Activation in Tumors

In solid tumors, oncogenic pathways drive rapid cell proliferation, which demands for more nutrients, particularly molecular oxygen, glucose and amino acids. However, the abnormal vascular structure limits the nutrient delivery, resulting in nutrient insufficiency [Chen and Sang, 2011]. While hypoxia is the major cause of HIF-1 α stabilization, recently it became clear that the lack of glucose or amino acids might also contribute to HIF-1 activation. It has been well known that HIF-1 activation plays an important role for hypoxic cell survival and migration [Vaupel, 2009]. HIF-1 α overexpression and HIF-1 activation have been observed in varieties of solid tumors [Semenza, 2004; Semenza, 2006; Vaupel et al., 2007; Zhong et al., 2000]. Of the more than 1,000 HIF-1 target genes that have been identified [Manalo et al., 2005; Schodel et al., 2011; Xia et al., 2009], many are implicated in adaptive strategies including metabolic reprogramming, survival, angiogenesis, and migration; enhanced survival and migration directly contribute to tumor metastasis and resistance to radiation and chemotherapy [Harada et al., 2012; Hockel and Vaupel, 2001; Lee et al., 2009; Semenza, 2003; Semenza, 2004; Semenza, 2010]. Accordingly, it has been proposed that repressing HIF-1 function may improve the therapeutic outcome of tumors [Semenza, 2003; Semenza, 2009; Semenza, 2010].

Oncogenic activation of HIF-1 triggered by both hypoxia and hypoxia independent mechanisms plays a critical role in metabolic transformation and tumor progression. The aerobic glycolysis caused by oncogenic activation of HIF-1 is best known as the Warburg effect. In addition to metabolic adaptation as discussed in introduction, angiogenesis and lymphogenesis are promoted by HIF-1. Deletion of HIF-1 α dramatically inhibits solid tumor development *in vivo*, and reduces expression of pro-angiogenic factors, such as VEGF, which promotes the formation of capillaries, and fibroblast growth factor 1 (FGF-1), which primarily stimulates the formation of arterial vessels [Hirota and Semenza, 2006; Ryan et al., 1998; Ryan et al., 2000]. HIF-1 activation also promotes cancer invasion and metastasis, a major obstacle to successful cancer therapy [Fidler, 2003; Huber et al., 2005; Polyak and Weinberg, 2009; Singh and Settleman, 2010]. HIF-1 α overexpression is associated with

more aggressive and invasive behaviors of human cancers [Bos et al., 2001; Zhong et al., 1999].

HIF-1 activation results in radiation resistance in solid tumors [Hockel and Vaupel, 2001]. Radiation reacts with intracellular molecules and molecular oxygen to cause DNA damage by forming oxygen free radicals [Hall and Giaccia, 2006]. In mouse model, HIF-1 α null tumor xenografts show increased radiation sensitivity [Aebersold et al., 2001]. Clinical data also revealed that HIF-1 α overexpression increased risk of failure to achieve complete remission after radiation therapy [Aebersold et al., 2001]. One study reports that cancer cells require HIF-1 activation for metastasis after surviving radiation [Harada et al., 2012].

While radiation resistance may involve multiple pathways including the established ATR/ATM-DNA repair response [Choudhury et al., 2006], radiation also increases HIF-1 activity and promotes endothelial cell survival [Moeller et al., 2004], which may form another adaptive strategy for tumor cells to survive the radiation. How radiation activates HIF-1 remains unclear; it has been reported that radiation activates AMPK and the ATF4 pathway [Cao and Wan, 2009; Fels and Koumenis, 2006; Kim et al., 2014; Lee et al., 2008; Sanli et al., 2010; Sanli et al., 2012; Zannella et al., 2011; Zhang et al., 2010], and overexpression of STC2, an ATF4 target, positively correlates to radiation resistance [Lin et al., 2014; Smith et al., 2006]. It is interesting to note that an intriguing similarity exists between the radiation-triggered and the nutrient insufficiency-triggered cellular responses: both induce cell cycle inhibition, activation of AMPK, activation of ATF4 pathways [Davie et al., 2015; Meng et al., 2010; Qie et al., 2012], and importantly, activation of HIF-1. However, a mechanistic link among those responses remains unclear. Further investigation is needed to address if this similarity explains the radiation resistance of tumor cells exposed to nutrient insufficiency.

Considering the role of HIF-1 in cells' survival strategy, researchers are actively exploring HIF-1 as a target for cancer therapy either alone or as a sensitizer for chemoradiation [Kong et al., 2005; Tan et al., 2005; Welsh et al., 2004]. Current strategies include suppressing HIF-1 α translation, increasing HIF-1 α degradation and inhibiting HIF-1 transcription activity [Semenza, 2007]. For example, mTOR inhibitors have been tested to inhibit HIF-1 α translation [Majumder et al., 2004; Thomas et al., 2006]. Other compounds including HDACIs [Drummond et al., 2005; Johnstone and Licht, 2003; Marks et al., 2001], Hsp90 inhibitors [Miyata, 2005; Neckers and Neckers, 2005], proteasome inhibitors [Bazzaro et al., 2006; Ishii et al., 2007; Joazeiro et al., 2006; Sunwoo et al., 2001] and microtubule inhibitors [Escuin et al., 2005; Mabjeesh et al., 2003; Newcomb et al., 2006] were found to destabilize HIF-1 α or inhibit HIF-1 function (Table 1). Echinomycin, an antibiotic peptide, suppresses HIF-1 function by reducing HIF-1 α binding to DNA [Kong et al., 2005]; PX478 and 103D5R showed the ability to inhibit HIF-1 function through unknown mechanisms [Kong et al., 2005; Tan et al., 2005; Welsh et al., 2004]. While these approaches have demonstrated to be effective, severe adverse effects on normal tissues limit their clinical application. Particularly, several compounds directly targeting the Hsp70/Hsp90 system or HIF-1 have proven effective for cancers. However, since a basal level of Hsp70/Hsp90 activity may be required for normal cells under physiological conditions to process newly synthesized proteins, it would not be surprise to find that directly and completely disrupting

the Hsp70/Hsp90 has various side effects on normal tissues. Since tumor cells frequently experience nutrient insufficiency and therapy-elicited stresses; refolding or disposal of stress-triggered misfolded or damaged proteins may effectively overwhelm the basal level of Hsp70/Hsp90 function. Accordingly, it is expected that targeting the AMPK-HDAC5 or AMPK-HDAC4-enhanced Hsp70/Hsp90 activity may specifically disrupt the survival strategy of tumor cells exposed to nutrient insufficiency, radiation and chemotherapy.

Relevance of HIF-1 Activation in Ischemic Disorders

Nutrient insufficiency, and particularly hypoxia, is the initiator and major contributing factor of the pathology of ischemic disorders. As opposed to solid tumors, ischemic lesions will greatly benefit from a successful survival strategy to cope with the nutrient insufficiency. Animal studies showed that HIF-1 function plays an important role in ischemia-induced vascular remodeling [Bosch-Marce et al., 2007; Patel et al., 2005; Rey et al., 2009; Rey et al., 2011; Rivard et al., 1999]. HIF-1 α $-/-$ mice were found to be lethal, but HIF-1 α $+/-$ mice showed reduced blood flow recovery and increased tissue damage after femoral artery ligation compared with wildtype littermates [Bosch-Marce et al., 2007]. Intramuscularly expressing a constitutively active form of HIF-1 α [Kelly et al., 2003] not only protects mice from age-dependent impairment of ischemia-induced vascular remodeling [Bosch-Marce et al., 2007], but also improves the recovery of blood flow in a rabbit model [Patel et al., 2005]. In addition, HIF-1 activation also mediates the protection induced by ischemic preconditioning [Murry et al., 1986]. After being exposed to short periods of ischemia and reperfusion, the heart suffered less injury from a subsequent prolonged ischemic insult, shown by a smaller infarct size. Loss of one allele of HIF-1 α abolishes this protection [Cai et al., 2008], demonstrating the importance of HIF-1 activation in this protection. HIF-1 may also protect the heart from pressure overload heart failure [Sano et al., 2007]. Hypertension could form a compensatory left ventricular hypertrophy to maintain the ejection fraction at an early stage, but it will progress to an uncompensated state with a decreased ejection fraction and eventually lead to heart failure [Levy et al., 1990]. Tissue-specific knockout of HIF-1 α in cardiomyocytes rapidly induced cardiac hypertrophy [Sano et al., 2007]. Interestingly, HDAC5 knockout impairs heart adaptation to increased load, partly phenocopies heart-specific HIF-1 α knockout. Along with the fact that hypoxia upregulates HDAC5, these findings indicate that the AMPK-HDAC5 pathway may be also involved in HIF-1 α stabilization and HIF-1 activation in ischemic lesions of normal tissues.

Perspectives

In summary, nutrient insufficiency frequently occurs in solid tumors and ischemic lesions. Activation of the AMPK pathway is a common response to nutrient insufficiency, which triggers multiple responses to maintain the ATP homeostasis. One of the AMPK-triggered cascades of events is the phosphorylation and cytosolic shuttling of HDAC4 and HDAC5, which enhances Hsp70/Hsp90 function. This facilitates the posttranslational processing and nuclear localization of HIF-1 α , ensuring a rapid activation of HIF-1 as a survival strategy. The role of this AMPK-facilitated HIF-1 activation in tumor metastasis and resistance to radiation remains unclear. However, the identification of cytosolic HDAC4 and HDAC5 as key players in cells' survival strategy provides new drug target to improve the specificity of

cancer therapy; targeting the conditionally required cytosolic HDAC4 and HDAC5 is expected to have less adverse effects on normal tissues than non-selective HDACIs, AMPK inhibitors and Hsp90 inhibitors. The involvement of AMPK-HDAC5 or AMPK-HDAC4 pathway in HIF-1 activation under other conditions, such as ischemic disorders, immune response, wound healing and neuron-degenerative disorders, remains to be explored.

Acknowledgement

We thank our colleagues C. Yin and N.G. Dahake for helpful discussion and proofreading the manuscript prior to submission. This work is supported in part by grant R01-CA129494 (to NS) from NCI, National Institutes of Health (NIH) and a contract Grant from Ivygene (DE, USA). S.C. is a recipient of a Ph.D. dissertation fellowship from Drexel University.

References

- Adams J, Chen ZP, Van Denderen BJ, Morton C, Parker MW, Witters LA, Stapleton D, Kemp BE. Intracellular control of AMPK via the gamma1 subunit AMP allosteric regulatory site. *Protein Sci.* 2004; 13:155–165. [PubMed: 14691231]
- Aebbersold DM, Burri P, Beer KT, Laissue J, Djonov V, Greiner RH, Semenza GL. Expression of hypoxia-inducible factor-1alpha: a novel predictive and prognostic parameter in the radiotherapy of oropharyngeal cancer. *Cancer Res.* 2001; 61:2911–2916. [PubMed: 11306467]
- Antoniou X, Sclip A, Ploia C, Colombo A, Moroy G, Borsello T. JNK contributes to Hif-1 α regulation in hypoxic neurons. *Molecules.* 2009; 15:114–127. [PubMed: 20110876]
- Bazzaro M, Lee MK, Zoso A, Stirling WL, Santillan A, Shih Ie M, Roden RB. Ubiquitin-proteasome system stress sensitizes ovarian cancer to proteasome inhibitor-induced apoptosis. *Cancer Res.* 2006; 66:3754–3763. [PubMed: 16585202]
- Benjamin D, Colombi M, Moroni C, Hall MN. Rapamycin passes the torch: a new generation of mTOR inhibitors. *Nat. Rev. Drug Discov.* 2011; 10:868–880. [PubMed: 22037041]
- Berhoerster K, Haerter M, Ellinghaus P, Heisler I, Beck H, Greschat S, Thierauch K-H, Wilhelm S, von Ahsen O, Hess-Stumpp H. BAY 87-2243 targets hypoxia-induced activation and stabilization of HIF-1 {alpha}: A novel approach to overcome resistance mechanisms in cancer therapy. *Cancer Res.* 2011; 71:4509.
- Bernardi R, Guernah I, Jin D, Grisendi S, Alimonti A, Teruya-Feldstein J, Cordon-Cardo C, Simon MC, Raffi S, Pandolfi PP. PML inhibits HIF-1 α translation and neoangiogenesis through repression of mTOR. *Nature.* 2006; 442:779–785. [PubMed: 16915281]
- Blancher C, Moore JW, Robertson N, Harris AL. Effects of ras and von Hippel-Lindau (VHL) gene mutations on hypoxia-inducible factor (HIF)-1 α , HIF-2 α , and vascular endothelial growth factor expression and their regulation by the phosphatidylinositol 3-kinase/Akt signaling pathway. *Cancer Res.* 2001; 61:7349–7355. [PubMed: 11585776]
- Borkovich KA, Farrelly FW, Finkelstein DB, Taulien J, Lindquist S. hsp82 is an essential protein that is required in higher concentrations for growth of cells at higher temperatures. *Mol. Cell. Biol.* 1989; 9:3919–3930. [PubMed: 2674684]
- Bos R, Zhong H, Hanrahan CF, Mommers EC, Semenza GL, Pinedo HM, Abeloff MD, Simons JW, van Diest PJ, van der Wall E. Levels of hypoxia-inducible factor-1 alpha during breast carcinogenesis. *J. Natl. Cancer Inst.* 2001; 93:309–314. [PubMed: 11181778]
- Bosch-Marce M, Okuyama H, Wesley JB, Sarkar K, Kimura H, Liu YV, Zhang H, Strazza M, Rey S, Savino L, Zhou YF, McDonald KR, Na Y, Vandiver S, Rabi A, Shaked Y, Kerbel R, Lavalley T, Semenza GL. Effects of aging and hypoxia-inducible factor-1 activity on angiogenic cell mobilization and recovery of perfusion after limb ischemia. *Circ. Res.* 2007; 101:1310–1318. [PubMed: 17932327]
- Brugarolas J, Lei K, Hurley RL, Manning BD, Reiling JH, Hafen E, Witters LA, Ellisen LW, Kaelin WG. Regulation of mTOR function in response to hypoxia by REDD1 and the TSC1/TSC2 tumor suppressor complex. *Genes Dev.* 2004; 18:2893–2904. [PubMed: 15545625]

- Brugarolas JB, Vazquez F, Reddy A, Sellers WR, Kaelin WG. TSC2 regulates VEGF through mTOR-dependent and-independent pathways. *Cancer Cell*. 2003; 4:147–158. [PubMed: 12957289]
- Bruick RK, McKnight SL. A conserved family of prolyl-4-hydroxylases that modify HIF. *Science*. 2001; 294:1337–1340. [PubMed: 11598268]
- Cai Z, Zhong H, Bosch-Marce M, Fox-Talbot K, Wang L, Wei C, Trush MA, Semenza GL. Complete loss of ischaemic preconditioning-induced cardioprotection in mice with partial deficiency of HIF-1 alpha. *Cardiovasc. Res*. 2008; 77:463–470. [PubMed: 18006459]
- Cao C, Wan Y. Parameters of protection against ultraviolet radiation - induced skin cell damage. *J. Cell. Physiol*. 2009; 220:277–284. [PubMed: 19360745]
- Carracedo A, Pandolfi P. The PTEN-PI3K pathway: of feedbacks and cross-talks. *Oncogene*. 2008; 27:5527–5541. [PubMed: 18794886]
- Chang SR, McKinsey TA, Zhang CL, Richardson JA, Hill JA, Olson EN. Histone deacetylases 5 and 9 govern responsiveness of the heart to a subset of stress signals and play redundant roles in heart development. *Mol. Cell. Biol*. 2004; 24:8467–8476. [PubMed: 15367668]
- Chen C, Pore N, Behrooz A, Ismail-Beigi F, Maity A. Regulation of glut1 mRNA by hypoxia-inducible factor-1 Interaction between H-ras and hypoxia. *Journal of Biological Chemistry*. 2001; 276:9519–9525. [PubMed: 11120745]
- Chen S, Sang N. Histone deacetylase inhibitors: the epigenetic therapeutics that repress hypoxia-inducible factors. *J. Biomed. Biotechnol*. 2011; 2011:197946. [PubMed: 21151670]
- Chen S, Yin C, Lao T, Liang D, He D, Wang C, Sang N. AMPK-HDAC5 Pathway Facilitates Nuclear Accumulation of HIF-1alpha and Functional Activation of HIF-1 by Deacetylating Hsp70 in the Cytosol. *Cell Cycle*. 2015; 14
- Cho DC, Cohen MB, Panka DJ, Collins M, Ghebremichael M, Atkins MB, Signoretti S, Mier JW. The efficacy of the novel dual PI3-kinase/mTOR inhibitor NVP-BEZ235 compared with rapamycin in renal cell carcinoma. *Clin. Cancer Res*. 2010; 16:3628–3638. [PubMed: 20606035]
- Choudhury, A.; Cuddihy, A.; Bristow, RG. *Sem. Rad. Oncol*. Elsevier; 2006. Radiation and new molecular agents part I: targeting ATM-ATR checkpoints, DNA repair, and the proteasome editor^editors; p. 51-58.
- Davie E, Forte GM, Petersen J. Nitrogen regulates AMPK to control TORC1 signaling. *Curr. Biol*. 2015; 25:445–454. [PubMed: 25639242]
- Doedens AL, Phan AT, Stradner MH, Fujimoto JK, Nguyen JV, Yang E, Johnson RS, Goldrath AW. Hypoxia-inducible factors enhance the effector responses of CD8(+) T cells to persistent antigen. *Nat. Immunol*. 2013; 14:1173–1182. [PubMed: 24076634]
- Drummond DC, Noble CO, Kirpotin DB, Guo Z, Scott GK, Benz CC. Clinical development of histone deacetylase inhibitors as anticancer agents. *Annu. Rev. Pharmacol. Toxicol*. 2005; 45:495–528. [PubMed: 15822187]
- Epstein AC, Gleadle JM, McNeill LA, Hewitson KS, O'Rourke J, Mole DR, Mukherji M, Metzen E, Wilson MI, Dhanda A, Tian YM, Masson N, Hamilton DL, Jaakkola P, Barstead R, Hodgkin J, Maxwell PH, Pugh CW, Schofield CJ, Ratcliffe PJ. C. elegans EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. *Cell*. 2001; 107:43–54. [PubMed: 11595184]
- Escuin D, Kline ER, Giannakakou P. Both microtubule-stabilizing and microtubule-destabilizing drugs inhibit hypoxia-inducible factor-1alpha accumulation and activity by disrupting microtubule function. *Cancer Res*. 2005; 65:9021–9028. [PubMed: 16204076]
- Eskandani M, Abdolalizadeh J, Hamishehkar H, Nazemiyeh H, Barar J. Galbanic acid inhibits HIF-1α expression via EGFR/HIF-1α pathway in cancer cells. *Fitoterapia*. 2015; 101:1–11. [PubMed: 25510323]
- Fang J, Xia C, Cao Z, Zheng JZ, Reed E, Jiang B-H. Apigenin inhibits VEGF and HIF-1 expression via PI3K/AKT/p70S6K1 and HDM2/p53 pathways. *FASEB J*. 2005; 19:342–353. [PubMed: 15746177]
- Fath DM, Kong X, Liang D, Lin Z, Chou A, Jiang Y, Fang J, Caro J, Sang N. Histone deacetylase inhibitors repress the transactivation potential of hypoxia-inducible factors independently of direct acetylation of HIF-alpha. *J. Biol. Chem*. 2006; 281:13612–13619. [PubMed: 16543236]

- Fels DR, Koumenis C. The PERK/eIF2 α /ATF4 module of the UPR in hypoxia resistance and tumor growth. *Cancer Biol. Ther.* 2006; 5:723–728. [PubMed: 16861899]
- Fidler IJ. The pathogenesis of cancer metastasis: the 'seed and soil' hypothesis revisited. *Nat. Rev. Cancer.* 2003; 3:453–458. [PubMed: 12778135]
- Fischle W, Dequiedt F, Hendzel MJ, Guenther MG, Lazar MA, Voelter W, Verdin E. Enzymatic activity associated with class II HDACs is dependent on a multiprotein complex containing HDAC3 and SMRT/N-CoR. *Mol. Cell.* 2002; 9:45–57. [PubMed: 11804585]
- Geng H, Harvey CT, Pittsenbarger J, Liu Q, Beer TM, Xue C, Qian DZ. HDAC4 protein regulates HIF1 α protein lysine acetylation and cancer cell response to hypoxia. *J. Biol. Chem.* 2011; 286:38095–38102. [PubMed: 21917920]
- Hall, EJ.; Giaccia, AJ. *Radiobiology for the radiologist*. Vol. ix. Philadelphia: Lippincott Williams & Wilkins; 2006. p. 546
- Harada H, Inoue M, Itasaka S, Hirota K, Morinibu A, Shinomiya K, Zeng L, Ou G, Zhu Y, Yoshimura M, McKenna WG, Muschel RJ, Hiraoka M. Cancer cells that survive radiation therapy acquire HIF-1 activity and translocate towards tumour blood vessels. *Nat. Commun.* 2012; 3:783. [PubMed: 22510688]
- Harris, TE.; Lawrence, JC, Jr. TOR signaling. *Sci.* 2003. STKE 2003:re15
- Hawley SA, Boudeau J, Reid JL, Mustard KJ, Udd L, Mäkelä TP, Alessi DR, Hardie DG. Complexes between the LKB1 tumor suppressor, STRAD α/β and MO25 α/β are upstream kinases in the AMP-activated protein kinase cascade. *J. Biol.* 2003; 2:28. [PubMed: 14511394]
- Hirota K, Semenza GL. Regulation of angiogenesis by hypoxia-inducible factor 1. *Crit. Rev. Oncol Hematol.* 2006; 59:15–26. [PubMed: 16716598]
- Hirsilä M, Koivunen P, Günzler V, Kivirikko KI, Myllyharju J. Characterization of the human prolyl 4-hydroxylases that modify the hypoxia-inducible factor. *J. Biol. Chem.* 2003; 278:30772–30780. [PubMed: 12788921]
- Hockel M, Vaupel P. Tumor hypoxia: definitions and current clinical, biologic, and molecular aspects. *J. Natl. Cancer Inst.* 2001; 93:266–276. [PubMed: 11181773]
- Huang LE, Gu J, Schau M, Bunn HF. Regulation of hypoxia-inducible factor 1 α is mediated by an O₂-dependent degradation domain via the ubiquitin-proteasome pathway. *Proc. Natl. Acad. Sci. U S A.* 1998; 95:7987–7992. [PubMed: 9653127]
- Huber MA, Kraut N, Beug H. Molecular requirements for epithelial-mesenchymal transition during tumor progression. *Curr. Opin. Cell Biol.* 2005; 17:548–558. [PubMed: 16098727]
- Hudson CC, Liu M, Chiang GG, Otterness DM, Loomis DC, Kaper F, Giaccia AJ, Abraham RT. Regulation of hypoxia-inducible factor 1 α expression and function by the mammalian target of rapamycin. *Mol. Cell. Biol.* 2002; 22:7004–7014. [PubMed: 12242281]
- Inoki K, Corradetti MN, Guan KL. Dysregulation of the TSC-mTOR pathway in human disease. *Nat. Genet.* 2005; 37:19–24. [PubMed: 15624019]
- Isaacs JS, Jung YJ, Mimnaugh EG, Martinez A, Cuttitta F, Neckers LM. Hsp90 regulates a von Hippel Lindau-independent hypoxia-inducible factor-1 α -degradative pathway. *J. Biol. Chem.* 2002; 277:29936–29944. [PubMed: 12052835]
- Isaacs JS, Jung YJ, Mole DR, Lee S, Torres-Cabala C, Chung YL, Merino M, Trepel J, Zbar B, Toro J, Ratcliffe PJ, Linehan WM, Neckers L. HIF overexpression correlates with biallelic loss of fumarate hydratase in renal cancer: novel role of fumarate in regulation of HIF stability. *Cancer Cell.* 2005; 8:143–153. [PubMed: 16098467]
- Ishii Y, Waxman S, Germain D. Targeting the ubiquitin-proteasome pathway in cancer therapy. *Anticancer Agents Med. Chem.* 2007; 7:359–365. [PubMed: 17504161]
- Jibb LA, Richards JG. AMP-activated protein kinase activity during metabolic rate depression in the hypoxic goldfish, *Carassius auratus*. *J. Exp. Biol.* 2008; 211:3111–3122. [PubMed: 18805810]
- Joazeiro CA, Anderson KC, Hunter T. Proteasome inhibitor drugs on the rise. *Cancer Res.* 2006; 66:7840–7842. [PubMed: 16861477]
- Johnstone RW, Licht JD. Histone deacetylase inhibitors in cancer therapy: is transcription the primary target? *Cancer Cell.* 2003; 4:13–18. [PubMed: 12892709]
- Kaelin WG Jr, Ratcliffe PJ. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol. Cell.* 2008; 30:393–402. [PubMed: 18498744]

- Kaluz S, Kaluzova M, Stanbridge EJ. Proteasomal inhibition attenuates transcriptional activity of hypoxia-inducible factor 1 (HIF-1) via specific effect on the HIF-1 α C-terminal activation domain. *Mol. Cell. Biol.* 2006; 26:5895–5907. [PubMed: 16847340]
- Katschinski DM, Le L, Schindler SG, Thomas T, Voss AK, Wenger RH. Interaction of the PAS B domain with HSP90 accelerates hypoxia-inducible factor-1 α stabilization. *Cell. Physiol. Biochem.* 2004; 14:351–360. [PubMed: 15319539]
- Keith B, Simon MC. Hypoxia-inducible factors, stem cells, and cancer. *Cell.* 2007; 129:465–472. [PubMed: 17482542]
- Kelly BD, Hackett SF, Hirota K, Oshima Y, Cai Z, Berg-Dixon S, Rowan A, Yan Z, Campochiaro PA, Semenza GL. Cell type-specific regulation of angiogenic growth factor gene expression and induction of angiogenesis in nonischemic tissue by a constitutively active form of hypoxia-inducible factor 1. *Circ. Res.* 2003; 93:1074–1081. [PubMed: 14576200]
- Kim EJ, Lee Y-J, Kang S, Lim Y-B. Ionizing radiation activates PERK/eIF2 α /ATF4 signaling via ER stress-independent pathway in human vascular endothelial cells. *Int. J. Rad. Biol.* 2014; 90:306–312. [PubMed: 24456547]
- Kimura N, Tokunaga C, Dalal S, Richardson C, Yoshino K, Hara K, Kemp BE, Witters LA, Mimura O, Yonezawa K. A possible linkage between AMP-activated protein kinase (AMPK) and mammalian target of rapamycin (mTOR) signalling pathway. *Genes Cells.* 2003; 8:65–79. [PubMed: 12558800]
- Kochanek KD, Murphy SL, Xu J, Arias E. Mortality in the United States. 2014; 2013 NCHS Data Brief:1–8.
- Koh MY, Spivak-Kroizman T, Venturini S, Welsh S, Williams RR, Kirkpatrick DL, Powis G. Molecular mechanisms for the activity of PX-478, an antitumor inhibitor of the hypoxia-inducible factor-1 α . *Mol. Cancer Therapeutics.* 2008; 7:90–100.
- Koivunen P, Lee S, Duncan CG, Lopez G, Lu G, Ramkissoon S, Losman JA, Joensuu P, Bergmann U, Gross S, Travins J, Weiss S, Looper R, Ligon KL, Verhaak RG, Yan H, Kaelin WG Jr. Transformation by the (R)-enantiomer of 2-hydroxyglutarate linked to EGLN activation. *Nature.* 2012; 483:484–488. [PubMed: 22343896]
- Kong D, Park EJ, Stephen AG, Calvani M, Cardellina JH, Monks A, Fisher RJ, Shoemaker RH, Melillo G. Echinomycin, a small-molecule inhibitor of hypoxia-inducible factor-1 DNA-binding activity. *Cancer Res.* 2005; 65:9047–9055. [PubMed: 16204079]
- Kong X, Lin Z, Liang D, Fath D, Sang N, Caro J. Histone deacetylase inhibitors induce VHL and ubiquitin-independent proteasomal degradation of hypoxia-inducible factor 1 α . *Mol. Cell. Biol.* 2006; 26:2019–2028. [PubMed: 16507982]
- Kovacs JJ, Murphy PJ, Gaillard S, Zhao X, Wu JT, Nicchitta CV, Yoshida M, Toft DO, Pratt WB, Yao TP. HDAC6 regulates Hsp90 acetylation and chaperone-dependent activation of glucocorticoid receptor. *Mol. Cell.* 2005; 18:601–607. [PubMed: 15916966]
- Kumar V, Zhang M-X, Swank MW, Kunz J, Wu G-Y. Regulation of dendritic morphogenesis by Ras-PI3K–Akt–mTOR and Ras–MAPK signaling pathways. *J. Neurosci.* 2005; 25:11288–11299. [PubMed: 16339024]
- Kurebayashi J, Otsuki T, Kurosumi M, Soga S, Akinaga S, Sonoo H. A Radicicol Derivative, KF58333, Inhibits Expression of Hypoxia - inducible Factor - 1 α and Vascular Endothelial Growth Factor, Angiogenesis and Growth of Human Breast Cancer Xenografts. *Cancer Sci.* 2001; 92:1342–1351.
- Laderoute KR, Amin K, Calaoagan JM, Knapp M, Le T, Orduna J, Foretz M, Viollet B. 5'-AMP-activated protein kinase (AMPK) is induced by low-oxygen and glucose deprivation conditions found in solid-tumor microenvironments. *Mol. Cell. Biol.* 2006; 26:5336–5347. [PubMed: 16809770]
- Lando D, Peet DJ, Whelan DA, Gorman JJ, Whitelaw ML. Asparagine hydroxylation of the HIF transactivation domain a hypoxic switch. *Science.* 2002; 295:858–861. [PubMed: 11823643]
- Lang SA, Moser C, Gaumann A, Klein D, Glockzin G, Popp FC, Dahlke MH, Piso P, Schlitt HJ, Geissler EK. Targeting heat shock protein 90 in pancreatic cancer impairs insulin-like growth factor-I receptor signaling, disrupts an interleukin-6/signal-transducer and activator of

- transcription 3/hypoxia-inducible factor-1 α autocrine loop, and reduces orthotopic tumor growth. *Clin. Cancer Res.* 2007; 13:6459–6468. [PubMed: 17975158]
- Lao T, Chen S, Sang N. Two mutations impair the stability and function of ubiquitin - activating enzyme (E1). *J. Cell. Physiol.* 2012; 227:1561–1568. [PubMed: 21678405]
- Laughner E, Taghavi P, Chiles K, Mahon PC, Semenza GL. HER2 (neu) signaling increases the rate of hypoxia-inducible factor 1 α (HIF-1 α) synthesis: novel mechanism for HIF-1-mediated vascular endothelial growth factor expression. *Mol. Cell. Biol.* 2001; 21:3995–4004. [PubMed: 11359907]
- Lee H-J, Lee Y-J, Kang C-M, Bae S, Jeoung D, Jang J-J, Lee S-S, Cho C-K, Lee Y-S. Differential Gene Signatures in Rat Mammary Tumors Induced by DMBA and Those Induced by Fractionated γ Radiation. *Rad. Res.* 2008; 170:579–590.
- Lee K, Qian DZ, Rey S, Wei H, Liu JO, Semenza GL. Anthracycline chemotherapy inhibits HIF-1 transcriptional activity and tumor-induced mobilization of circulating angiogenic cells. *Proc. Natl. Acad. Sci. U S A.* 2009; 106:2353–2358. [PubMed: 19168635]
- Lee YM, Kim S-H, Kim H-S, Son MJ, Nakajima H, Kwon HJ, Kim K-W. Inhibition of hypoxia-induced angiogenesis by FK228, a specific histone deacetylase inhibitor, via suppression of HIF-1 α activity. *Biochem. Biophys. Res. Commun.* 2003; 300:241–246.
- Levy D, Garrison RJ, Savage DD, Kannel WB, Castelli WP. Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N. Engl. J. Med.* 1990; 322:1561–1566. [PubMed: 2139921]
- Liang D, Kong X, Sang N. Effects of histone deacetylase inhibitors on HIF-1. *Cell Cycle.* 2006; 5:2430–2435. [PubMed: 17102633]
- Lin S, Guo Q, Wen J, Li C, Lin J, Cui X, Sang N, Pan J. Survival analyses correlate stanniocalcin 2 overexpression to poor prognosis of nasopharyngeal carcinomas. *J. Exp. Clin. Cancer Res.* 2014; 33:1–11. [PubMed: 24383517]
- Liu YV, Baek JH, Zhang H, Diez R, Cole RN, Semenza GL. RACK1 competes with HSP90 for binding to HIF-1 α and is required for O 2-independent and HSP90 inhibitor-induced degradation of HIF-1 α . *Mol. Cell.* 2007; 25:207–217. [PubMed: 17244529]
- Luo W, Zhong J, Chang R, Hu H, Pandey A, Semenza GL. Hsp70 and CHIP selectively mediate ubiquitination and degradation of hypoxia-inducible factor (HIF)-1 α but not HIF-2 α . *J. Biol. Chem.* 2010; 285:3651–3663. [PubMed: 19940151]
- Mabjeesh NJ, Escuin D, LaVallee TM, Pribluda VS, Swartz GM, Johnson MS, Willard MT, Zhong H, Simons JW, Giannakakou P. 2ME2 inhibits tumor growth and angiogenesis by disrupting microtubules and dysregulating HIF. *Cancer Cell.* 2003; 3:363–375. [PubMed: 12726862]
- Mabjeesh NJ, Post DE, Willard MT, Kaur B, Van Meir EG, Simons JW, Zhong H. Geldanamycin induces degradation of hypoxia-inducible factor 1 α protein via the proteasome pathway in prostate cancer cells. *Cancer Res.* 2002; 62:2478–2482. [PubMed: 11980636]
- Mahon PC, Hirota K, Semenza GL. FIH-1: a novel protein that interacts with HIF-1 α and VHL to mediate repression of HIF-1 transcriptional activity. *Genes Dev.* 2001; 15:2675–2686. [PubMed: 11641274]
- Majumder PK, Febbo PG, Bikoff R, Berger R, Xue Q, McMahon LM, Manola J, Brugarolas J, McDonnell TJ, Golub TR. mTOR inhibition reverses Akt-dependent prostate intraepithelial neoplasia through regulation of apoptotic and HIF-1-dependent pathways. *Nature Med.* 2004; 10:594–601. [PubMed: 15156201]
- Manalo DJ, Rowan A, Lavoie T, Natarajan L, Kelly BD, Ye SQ, Garcia JG, Semenza GL. Transcriptional regulation of vascular endothelial cell responses to hypoxia by HIF-1. *Blood.* 2005; 105:659–669. [PubMed: 15374877]
- Manning BD, Tee AR, Logsdon MN, Blenis J, Cantley LC. Identification of the tuberous sclerosis complex-2 tumor suppressor gene product tuberlin as a target of the phosphoinositide 3-Kinase/Akt pathway. *Mol. Cell.* 2002; 10:151–162. [PubMed: 12150915]
- Marks PA, Richon VM, Breslow R, Rifkind RA. Histone deacetylase inhibitors as new cancer drugs. *Curr. Opin. Oncol.* 2001; 13:477–483. [PubMed: 11673688]
- McKinsey TA, Zhang CL, Olson EN. Identification of a signal-responsive nuclear export sequence in class II histone deacetylases. *Mol. Cell. Biol.* 2001; 21:6312–6321. [PubMed: 11509672]

- Meng M, Chen S, Lao T, Liang D, Sang N. Nitrogen anabolism underlies the importance of glutaminolysis in proliferating cells. *Cell Cycle*. 2010; 9:3921–3932. [PubMed: 20935507]
- Metellus P, Colin C, Taieb D, Guedj E, Nanni-Metellus I, de Paula AM, Colavolpe C, Fuentes S, Dufour H, Barrie M. IDH mutation status impact on in vivo hypoxia biomarkers expression: new insights from a clinical, nuclear imaging and immunohistochemical study in 33 glioma patients. *J. Neuro-oncol*. 2011; 105:591–600.
- Miyata Y. Hsp90 inhibitor geldanamycin and its derivatives as novel cancer chemotherapeutic agents. *Curr. Pharm. Des.* 2005; 11:1131–1138. [PubMed: 15853661]
- Moeller BJ, Cao Y, Li CY, Dewhirst MW. Radiation activates HIF-1 to regulate vascular radiosensitivity in tumors: role of reoxygenation, free radicals, and stress granules. *Cancer Cell*. 2004; 5:429–441. [PubMed: 15144951]
- Mungai PT, Waypa GB, Jairaman A, Prakriya M, Dokic D, Ball MK, Schumacker PT. Hypoxia triggers AMPK activation through reactive oxygen species-mediated activation of calcium release-activated calcium channels. *Mol. Cell. Biol.* 2011; 31:3531–3545. [PubMed: 21670147]
- Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation*. 1986; 74:1124–1136. [PubMed: 3769170]
- Neckers L, Neckers K. Heat-shock protein 90 inhibitors as novel cancer chemotherapeutics - an update. *Expert. Opin. Emerg. Drugs*. 2005; 10:137–149. [PubMed: 15757409]
- Newcomb EW, Lukyanov Y, Schnee T, Ali MA, Lan L, Zagzag D. Noscipine inhibits hypoxia-mediated HIF-1 α expression and angiogenesis in vitro: a novel function for an old drug. *Int. J. Oncol.* 2006; 28:1121–1130. [PubMed: 16596228]
- Patel TH, Kimura H, Weiss CR, Semenza GL, Hofmann LV. Constitutively active HIF-1 α improves perfusion and arterial remodeling in an endovascular model of limb ischemia. *Cardiovasc. Res.* 2005; 68:144–154. [PubMed: 15921668]
- Pollard PJ, Briere JJ, Alam NA, Barwell J, Barclay E, Wortham NC, Hunt T, Mitchell M, Olpin S, Moat SJ, Hargreaves IP, Heales SJ, Chung YL, Griffiths JR, Dalgleish A, McGrath JA, Gleeson MJ, Hodgson SV, Poulson R, Rustin P, Tomlinson IP. Accumulation of Krebs cycle intermediates and over-expression of HIF1 α in tumours which result from germline FH and SDH mutations. *Hum. Mol. Genet.* 2005; 14:2231–2239. [PubMed: 15987702]
- Polyak K, Weinberg RA. Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. *Nat. Rev. Cancer*. 2009; 9:265–273. [PubMed: 19262571]
- Pore N, Jiang Z, Gupta A, Cerniglia G, Kao GD, Maity A. EGFR tyrosine kinase inhibitors decrease VEGF expression by both hypoxia-inducible factor (HIF)-1-independent and HIF-1-dependent mechanisms. *Cancer Res.* 2006; 66:3197–3204. [PubMed: 16540671]
- Potter CJ, Pedraza LG, Xu T. Akt regulates growth by directly phosphorylating Tsc2. *Nat. Cell Biol.* 2002; 4:658–665. [PubMed: 12172554]
- Pratt WB, Toft DO. Regulation of signaling protein function and trafficking by the hsp90/hsp70-based chaperone machinery. *Exp. Biol. Med.* (Maywood). 2003; 228:111–133. [PubMed: 12563018]
- Qian DZ, Kachhap SK, Collis SJ, Verheul HM, Carducci MA, Atadja P, Pili R. Class II histone deacetylases are associated with VHL-independent regulation of hypoxia-inducible factor 1 α . *Cancer Res.* 2006; 66:8814–8821. [PubMed: 16951198]
- Qie S, Liang D, Yin C, Gu W, Meng M, Wang C, Sang N. Glutamine depletion and glucose depletion trigger growth inhibition via distinctive gene expression reprogramming. *Cell Cycle*. 2012; 11:3679–3690. [PubMed: 22935705]
- Rey S, Lee K, Wang CJ, Gupta K, Chen S, McMillan A, Bhise N, Levchenko A, Semenza GL. Synergistic effect of HIF-1 α gene therapy and HIF-1-activated bone marrow-derived angiogenic cells in a mouse model of limb ischemia. *Proc. Natl. Acad. Sci. U S A*. 2009; 106:20399–20404. [PubMed: 19948968]
- Rey S, Luo W, Shimoda LA, Semenza GL. Metabolic reprogramming by HIF-1 promotes the survival of bone marrow-derived angiogenic cells in ischemic tissue. *Blood*. 2011; 117:4988–4998. [PubMed: 21389314]
- Rivard A, Fabre JE, Silver M, Chen D, Murohara T, Kearney M, Magner M, Asahara T, Isner JM. Age-dependent impairment of angiogenesis. *Circulation*. 1999; 99:111–120. [PubMed: 9884387]

- Rutkowski DT, Kaufman RJ. All roads lead to ATF4. *Developmental cell*. 2003; 4:442–444. [PubMed: 12689582]
- Ryan HE, Lo J, Johnson RS. HIF-1 alpha is required for solid tumor formation and embryonic vascularization. *EMBO J*. 1998; 17:3005–3015. [PubMed: 9606183]
- Ryan HE, Poloni M, McNulty W, Elson D, Gassmann M, Arbeit JM, Johnson RS. Hypoxia-inducible factor-1alpha is a positive factor in solid tumor growth. *Cancer Res*. 2000; 60:4010–4015. [PubMed: 10945599]
- Salceda S, Caro J. Hypoxia-inducible factor 1alpha (HIF-1alpha) protein is rapidly degraded by the ubiquitin-proteasome system under normoxic conditions. Its stabilization by hypoxia depends on redox-induced changes. *J. Biol. Chem*. 1997; 272:22642–22647. [PubMed: 9278421]
- Sang N, Fang J, Srinivas V, Leshchinsky I, Caro J. Carboxyl-terminal transactivation activity of hypoxia-inducible factor 1 alpha is governed by a von Hippel-Lindau protein-independent, hydroxylation-regulated association with p300/CBP. *Mol. Cell. Biol*. 2002; 22:2984–2992. [PubMed: 11940656]
- Sanli T, Rashid A, Liu C, Harding S, Bristow RG, Cutz J-C, Singh G, Wright J, Tsakiridis T. Ionizing radiation activates AMP-activated kinase (AMPK): a target for radiosensitization of human cancer cells. *Int. J. Rad. Oncol*. 2010; 78:221–229.
- Sanli T, Storozhuk Y, Linher-Melville K, Bristow RG, Laderout K, Viollet B, Wright J, Singh G, Tsakiridis T. Ionizing radiation regulates the expression of AMP-activated protein kinase (AMPK) in epithelial cancer cells: modulation of cellular signals regulating cell cycle and survival. *Radiother. Oncol*. 2012; 102:459–465. [PubMed: 22225791]
- Sano M, Minamino T, Toko H, Miyauchi H, Orimo M, Qin Y, Akazawa H, Tateno K, Kayama Y, Harada M, Shimizu I, Asahara T, Hamada H, Tomita S, Molkentin JD, Zou Y, Komuro I. p53-induced inhibition of Hif-1 causes cardiac dysfunction during pressure overload. *Nature*. 2007; 446:444–448. [PubMed: 17334357]
- Schodel J, Oikonomopoulos S, Ragoussis J, Pugh CW, Ratcliffe PJ, Mole DR. High-resolution genome-wide mapping of HIF-binding sites by ChIP-seq. *Blood*. 2011; 117:e207–e217. [PubMed: 21447827]
- Schofield CJ, Ratcliffe PJ. Oxygen sensing by HIF hydroxylases. *Nat Rev Mol Cell Biol*. 2004; 5:343–354. [PubMed: 15122348]
- Selak MA, Armour SM, MacKenzie ED, Boulahbel H, Watson DG, Mansfield KD, Pan Y, Simon MC, Thompson CB, Gottlieb E. Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase. *Cancer Cell*. 2005; 7:77–85. [PubMed: 15652751]
- Semenza GL. Targeting HIF-1 for cancer therapy. *Nat. Rev. Cancer*. 2003; 3:721–732. [PubMed: 13130303]
- Semenza GL. Intratumoral hypoxia, radiation resistance, and HIF-1. *Cancer Cell*. 2004; 5:405–406. [PubMed: 15144945]
- Semenza GL. Development of novel therapeutic strategies that target HIF-1. *Expert. Opin. Ther. Targets*. 2006; 10:267–280. [PubMed: 16548775]
- Semenza GL. Evaluation of HIF-1 inhibitors as anticancer agents. *Drug discovery today*. 2007; 12:853–859. [PubMed: 17933687]
- Semenza GL. Regulation of cancer cell metabolism by hypoxia-inducible factor 1. *Semin Cancer Biol*. 2009; 19:12–16. [PubMed: 19114105]
- Semenza GL. Defining the role of hypoxia-inducible factor 1 in cancer biology and therapeutics. *Oncogene*. 2010; 29:625–634. [PubMed: 19946328]
- Semenza GL. Hypoxia-inducible factors in physiology and medicine. *Cell*. 2012; 148:399–408. [PubMed: 22304911]
- Semenza GL. Hypoxia-inducible factor 1 and cardiovascular disease. *Annu. Rev. Physiol*. 2014; 76:39–56. [PubMed: 23988176]
- Semenza GL, Wang GL. A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Mol. Cell. Biol*. 1992; 12:5447–5454. [PubMed: 1448077]
- Shackelford DB, Shaw RJ. The LKB1-AMPK pathway: metabolism and growth control in tumour suppression. *Nat. Rev. Cancer*. 2009; 9:563–575. [PubMed: 19629071]

- Shackelford DB, Vasquez DS, Corbeil J, Wu S, Leblanc M, Wu C-L, Vera DR, Shaw RJ. mTOR and HIF-1 α -mediated tumor metabolism in an LKB1 mouse model of Peutz-Jeghers syndrome. *Proc. Natl. Acad. Sci. USA*. 2009; 106:11137–11142. [PubMed: 19541609]
- Shaw RJ, Bardeesy N, Manning BD, Lopez L, Kosmatka M, DePinho RA, Cantley LC. The LKB1 tumor suppressor negatively regulates mTOR signaling. *Cancer Cell*. 2004; 6:91–99. [PubMed: 15261145]
- Singh A, Settleman J. EMT, cancer stem cells and drug resistance: an emerging axis of evil in the war on cancer. *Oncogene*. 2010; 29:4741–4751. [PubMed: 20531305]
- Smith DJ, Jaggi M, Zhang W, Galich A, Du C, Sterrett SP, Smith LM, Balaji K. Metallothioneins and resistance to cisplatin and radiation in prostate cancer. *Urology*. 2006; 67:1341–1347. [PubMed: 16765200]
- Staab A, Loeffler J, Said HM, Diehlmann D, Katzer A, Beyer M, Fleischer M, Schwab F, Baier K, Einsele H, Flentje M, Vordermark D. Effects of HIF-1 inhibition by chetomin on hypoxia-related transcription and radiosensitivity in HT 1080 human fibrosarcoma cells. *BMC Cancer*. 2007; 7:213. [PubMed: 17999771]
- Sunwoo JB, Chen Z, Dong G, Yeh N, Crawl Bancroft C, Sausville E, Adams J, Elliott P, Van Waes C. Novel proteasome inhibitor PS-341 inhibits activation of nuclear factor-kappa B, cell survival, tumor growth, and angiogenesis in squamous cell carcinoma. *Clin. Cancer Res*. 2001; 7:1419–1428. [PubMed: 11350913]
- Taipale M, Jarosz DF, Lindquist S. HSP90 at the hub of protein homeostasis: emerging mechanistic insights. *Nat. Rev. Mol. Cell. Biol*. 2010; 11:515–528. [PubMed: 20531426]
- Takubo K, Goda N, Yamada W, Iriuchishima H, Ikeda E, Kubota Y, Shima H, Johnson RS, Hirao A, Suematsu M, Suda T. Regulation of the HIF-1 α level is essential for hematopoietic stem cells. *Cell Stem Cell*. 2010; 7:391–402. [PubMed: 20804974]
- Tan C, de Noronha RG, Roecker AJ, Pyrzynska B, Khwaja F, Zhang Z, Zhang H, Teng Q, Nicholson AC, Giannakakou P, Zhou W, Olson JJ, Pereira MM, Nicolaou KC, Van Meir EG. Identification of a novel small-molecule inhibitor of the hypoxia-inducible factor 1 pathway. *Cancer Res*. 2005; 65:605–612. [PubMed: 15695405]
- Thomas GV, Tran C, Mellinghoff IK, Welsbie DS, Chan E, Fueger B, Czernin J, Sawyers CL. Hypoxia-inducible factor determines sensitivity to inhibitors of mTOR in kidney cancer. *Nat. Med*. 2006; 12:122–127. [PubMed: 16341243]
- To KK, Robey R, Zhan Z, Bangiolo L, Bates SE. Upregulation of ABCG2 by romidepsin via the aryl hydrocarbon receptor pathway. *Mol. Cancer Res*. 2011; 9:516–527. [PubMed: 21357443]
- Vaupel P. Prognostic potential of the pre-therapeutic tumor oxygenation status. *Adv. Exp. Med. Biol*. 2009; 645:241–246. [PubMed: 19227477]
- Vaupel P, Hockel M, Mayer A. Detection and characterization of tumor hypoxia using pO₂ histography. *Antioxid. Redox Signal*. 2007; 9:1221–1235. [PubMed: 17536958]
- Wan X, Shen N, Mendoza A, Khanna C, Helman LJ. CCI-779 inhibits rhabdomyosarcoma xenograft growth by an antiangiogenic mechanism linked to the targeting of mTOR/Hif-1 α /VEGF signaling. *Neoplasia*. 2006; 8:394–401. [PubMed: 16790088]
- Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc. Natl. Acad. Sci. U S A*. 1995; 92:5510–5514. [PubMed: 7539918]
- Wang GL, Semenza GL. Purification and characterization of hypoxia-inducible factor 1. *J. Biol. Chem*. 1995; 270:1230–1237. [PubMed: 7836384]
- Wang Y, Wang SY, Zhang XH, Zhao M, Hou CM, Xu YJ, Du ZY, Yu XD. FK228 inhibits Hsp90 chaperone function in K562 cells via hyperacetylation of Hsp70. *Biochem. Biophys. Res. Commun*. 2007; 356:998–1003. [PubMed: 17397803]
- Wegele H, Muller L, Buchner J. Hsp70 and Hsp90—a relay team for protein folding. *Rev. Physiol. Biochem. Pharmacol*. 2004; 151:1–44. [PubMed: 14740253]
- Welsh S, Williams R, Kirkpatrick L, Paine-Murrieta G, Powis G. Antitumor activity and pharmacodynamic properties of PX-478, an inhibitor of hypoxia-inducible factor-1 α . *Mol. Cancer Ther*. 2004; 3:233–244. [PubMed: 15026543]

- Whitesell L, Lindquist SL. HSP90 and the chaperoning of cancer. *Nat. Rev. Cancer*. 2005; 5:761–772. [PubMed: 16175177]
- Winder WW, Hardie DG. AMP-activated protein kinase, a metabolic master switch: possible roles in type 2 diabetes. *Am. J. Physiol*. 1999; 277:E1–E10. [PubMed: 10409121]
- Wouters BG, Koritzinsky M. Hypoxia signalling through mTOR and the unfolded protein response in cancer. *Nat. Rev. Cancer*. 2008; 8:851–864. [PubMed: 18846101]
- Xia X, Lemieux ME, Li W, Carroll JS, Brown M, Liu XS, Kung AL. Integrative analysis of HIF binding and transactivation reveals its role in maintaining histone methylation homeostasis. *Proc. Natl. Acad. Sci. U S A*. 2009; 106:4260–4265. [PubMed: 19255431]
- Yang XJ, Seto E. HATs and HDACs: from structure, function and regulation to novel strategies for therapy and prevention. *Oncogene*. 2007; 26:5310–5318. [PubMed: 17694074]
- Ye J, Kumanova M, Hart LS, Sloane K, Zhang H, De Panis DN, Bobrovnikova-Marjon E, Diehl JA, Ron D, Koumenis C. The GCN2-ATF4 pathway is critical for tumour cell survival and proliferation in response to nutrient deprivation. *EMBO J*. 2010; 29:2082–2096. [PubMed: 20473272]
- Zagzag D, Nomura M, Friedlander DR, Blanco C, Gagner JP, Nomura N, Newcomb EW. Geldanamycin inhibits migration of glioma cells in vitro: A potential role for hypoxia-inducible factor (HIF-1 α) in glioma cell invasion. *J. Cell. Physiol*. 2003; 196:394–402. [PubMed: 12811834]
- Zannella VE, Cojocari D, Hilgendorf S, Vellanki RN, Chung S, Wouters BG, Koritzinsky M. AMPK regulates metabolism and survival in response to ionizing radiation. *Radiother. Oncol*. 2011; 99:293–299. [PubMed: 21715037]
- Zhang B, Wang Y, Pang X, Su Y, Ai G, Wang T. ER stress induced by ionising radiation in IEC-6 cells. *Int. J. Rad. Biol*. 2010; 86:429–435. [PubMed: 20470193]
- Zhang Z, Yan J, Chang Y, ShiDu Yan S, Shi H. Hypoxia inducible factor-1 as a target for neurodegenerative diseases. *Curr. Med. Chem*. 2011; 18:4335–4343. [PubMed: 21861815]
- Zhao S, Lin Y, Xu W, Jiang W, Zha Z, Wang P, Yu W, Li Z, Gong L, Peng Y, Ding J, Lei Q, Guan KL, Xiong Y. Glioma-derived mutations in IDH1 dominantly inhibit IDH1 catalytic activity and induce HIF-1 α . *Science*. 2009; 324:261–265. [PubMed: 19359588]
- Zhong H, Chiles K, Feldser D, Laughner E, Hanrahan C, Georgescu MM, Simons JW, Semenza GL. Modulation of hypoxia-inducible factor 1 α expression by the epidermal growth factor/phosphatidylinositol 3-kinase/PTEN/AKT/FRAP pathway in human prostate cancer cells: implications for tumor angiogenesis and therapeutics. *Cancer Res*. 2000; 60:1541–1545. [PubMed: 10749120]
- Zhong H, De Marzo AM, Laughner E, Lim M, Hilton DA, Zagzag D, Buechler P, Isaacs WB, Semenza GL, Simons JW. Overexpression of hypoxia-inducible factor 1 α in common human cancers and their metastases. *Cancer Res*. 1999; 59:5830–5835. [PubMed: 10582706]
- Zundel W, Schindler C, Haas-Kogan D, Koong A, Kaper F, Chen E, Gottschalk AR, Ryan HE, Johnson RS, Jefferson AB. Loss of PTEN facilitates HIF-1-mediated gene expression. *Genes Dev*. 2000; 14:391–396. [PubMed: 10691731]

Figure 1. Summary of the synthesis, posttranslational processing and hypoxia-controlled degradation of HIF-1α

After translation, nascent HIF-1α interacts with the molecule chaperone Hsp70/Hsp90 system for processing and maturation. Proteins fail the posttranslational folding processes will be degraded by a quality control system using a ubiquitination-independent proteasome degradation pathway (UIP). The protein levels of correctly folded functional HIF-1α will be regulated by the oxygen-dependent hydroxylation-ubiquitination mechanism.

Figure 2. Schematic structure of HIF-1α and the conventional oxygen-dependent inhibition of HIF-1

Figure 3. Signaling pathways that controls the rate of HIF-1 α translation

Activation of oncogenes or loss of tumor suppressors promotes HIF-1 α translation, which is suppressed by mTOR inhibition.

Figure 4. Proposed model of AMPK-HDAC5 enhanced, Hsp70/Hsp90 executed posttranslational processing of HIF-1α

AMPK serves as a sensor of multiple types of stresses, which promotes the cytosolic localization of HDAC4 and HDAC5, depending on which is expressed in a specific type of cells. HDAC4 and HDAC5 catalyzed deacetylation of Hsp70 enhances the efficiency of posttranslational processing of HIF-1α, minimizing the pre-mature degradation.

Figure 5. Cytosolic HDAC4 and HDAC5 contribute to hypoxic stabilization of HIF-1α and functional activation of HIF-1

Knockdown of either HDAC4 or HDAC5 attenuates hypoxia triggered HIF-1α accumulation. Hep3B cells were transfected with indicated siRNA. After 42 h, cells were exposed to hypoxia (1% O₂) for 6 h. **(A)** Western blotting showing the protein levels of HIF-1α, HDAC4 and HDAC5. The protein levels of α-tubulin were determined as a loading control. Note that hypoxia enhanced HDAC5 but not HDAC4. **(B)** Total RNA was collected, and quantitative real time PCR was performed following reverse transcription to determine the mRNA levels of CA-IX as an indicator of HIF-1 function. **(C)** Cytosolic HDAC4 mutant (S265/266A) is sufficient to stabilize HIF-1α. Hep3B cells were cotransfected with 2 μg of HA-HIF-1α together with 2 μg of either control vector, Flag-HDAC4(WT), or cytosolic mutant HDAC4-S265/266A. HIF-1α and Flag-tagged HDAC4 were detected by Western blotting.

Table 1

Summary of Compounds That Repress HIF-1

Mechanism	Drug Classification	Dugs	Potential Implication in the QC Model
Repressing HIF-1α translation	EGFR inhibitor	Erlotinib [Pore et al., 2006] Galbanic acid [Eskandani et al., 2015] Gefitinib [Pore et al., 2006]	Translation driven by oncogenic and proliferative signals
	Her2/ErbB2 inhibitor	Herceptin [Laughner et al., 2001]	
	PI3K inhibitor	Apigenin [Fang et al., 2005] NVP-BEZ235 [Cho et al., 2010]	
	mTOR inhibitor	CCI-779 [Wan et al., 2006] Everolimus [Majumder et al., 2004] Rapamycin [Laughner et al., 2001]	
	?	PX-478[Koh et al., 2008; Welsh et al., 2004]	
Reduce HIF-1α stability	Hsp90 inhibitors	17-AAG [Isaacs et al., 2002; Kong et al., 2006; Liu et al., 2007] 17-DMAG [Lang et al., 2007] Geldanamycin [Mabjeesh et al., 2002; Zagzag et al., 2003] KF58333 [Kurebayashi et al., 2001]	Hsp70/Hsp90 function is needed for posttranslational stabilization of HIF-1 α
	HDAC inhibitors	FK228 [Lee et al., 2003] LAQ824 [Qian et al., 2006] LMK235 [Chen et al., 2015] SAHA [Kong et al., 2006] TSA [Kong et al., 2006]	
	Microtubule-disrupting drugs	2ME2 [Escuin et al., 2005; Mabjeesh et al., 2003] Colchicine [Escuin et al., 2005] Discodermolide [Escuin et al., 2005] Epothilone B [Escuin et al., 2005] Taxotere [Escuin et al., 2005] Vincristine [Escuin et al., 2005]	
	Electron transfer chain disrupting	BAY 87-2243 [Berhoerster et al., 2011]	
Decrease HIF-1 transactivation	HDAC inhibitors	SAHA [Fath et al., 2006] TSA [Fath et al., 2006]	Cytosolic deacetylases facilitate Hsp70/Hsp90 function, promoting formation of mature HIF-1 α
	Proteasome inhibitors	Velcade [Kaluz et al., 2006]	
	p300/CBP inhibitors	Chetomin [Staab et al., 2007]	