Published in final edited form as:

J Cell Biochem. 2016 January ; 117(1): 9–19. doi:10.1002/jcb.25280.

Chromosomes at Work: Organization of Chromosome Territories in the Interphase Nucleus

Andrew Fritz¹, A. Rasim Barutcu², Lori Martin-Buley¹, André J. vanWijnen³, Sayyed K. Zaidi¹, Anthony N. Imbalzano², Jane B. Lian¹, Janet L. Stein¹, and Gary S. Stein^{1,*}

¹University of Vermont Cancer Center, Department of Biochemistry, University of Vermont College of Medicine, 89 Beaumont Avenue, Burlington, VT 05405, USA

²Department of Cell and Developmental Biology, University of Massachusetts Medical School, 55 Lake Avenue North, Worcester, MA 01655, USA

³Departments of Orthopedic Surgery and Biochemistry & Molecular Biology, Mayo Clinic, Rochester, MN, USA

Abstract

The organization of interphase chromosomes in chromosome territories (CTs) was first proposed more than one hundred years ago. The introduction of increasingly sophisticated microscopic and molecular techniques, now provide complementary strategies for studying CTs in greater depth than ever before. Here we provide an overview of these strategies and how they are being used to elucidate CT interactions and the role of these dynamically regulated, nuclear-structure building blocks in directly supporting nuclear function in a physiologically responsive manner.

Introduction

Genomic organization is coordinated by nuclear processes that include DNA replication, chromatin remodeling, RNA transcription and splicing. In the disease state, the frequent dysfunction and compromised organization of regulatory machinery suggests a strong role of genomic organization in pathology. Although the nucleus is not subdivided by membrane-bound compartments, the regulatory machinery for transcription, replication, and repair are architecturally organized in subnuclear domains with specific functions. This compartmentalization supports a direct relationship between nuclear structure and function that is dynamically regulated in a physiologically responsive manner. Prominent domains within the nucleus, such as the nucleolus and heterochromatin, were identified long before their functions were elucidated. Other architectural components of the nucleus, including the nuclear matrix, nuclear lamina, Cajal bodies, speckles, and the Barr body were identified once advances in microscopy became available and molecular approaches were introduced [Hemmerich et al., 2011; Mehta et al., 2007; Zhu and Brangwynne, 2015].

During mitosis, the genome is organized into easily recognized X-shaped chromosomes, however, this configuration does not persist during the G1, S, and G2 phases of the cell

*To whom correspondence should be addressed: gary.stein@uvm.edu.

cycle. During interphase, the genome, like the other nuclear domains, is organized in discrete bodies. Chromosomes relax into nuclear domains that are referred to as chromosome territories (CTs; Figure 1A), [Cremer and Cremer, 2001; Cremer et al., 2006; Cremer et al., 1981; Lichter et al., 1988; Manuelidis, 1985; Schardin et al., 1985; Stack et al., 1977; Zorn et al., 1979; Zorn et al., 1976]. This realization that genomic structure and function are dynamic and highly integrated has been driven by technologies that include fluorescence in situ hybridization (FISH, [Solovei and Cremer, 2010]) and, more recently, chromatin conformation capture (3C) techniques [Barutcu et al., 2015; de Wit and de Laat, 2012]. Here, we provide an overview of microscopic and molecular techniques that have advanced understanding of nuclear organization and a synopsis of the role of CTs in genomic organization and expression.

Origins and Reestablishment of the Chromosome Territory Paradigm

The existence of CTs was first suggested by Theodor Boveri in 1885 and was supported by Carl Rabl in 1909 [Cremer and Cremer, 2010]. These early observations and predictions of genome compartmentalization in the interphase nucleus were superseded when electron microscopy became available and evidence of chromosome intermingling during interphase was directly observed. This led to the “spaghetti” model of the interphase nucleus that persisted for decades—in which chromatin fibers from different chromosomes are interwoven. It wasn’t until the late 1970s that the concept of chromosome territories began to re-emerge due to compelling evidence from several lines of inquiry [Cremer and Cremer, 2010]. One such line utilized a laser microbeam to inflict DNA damage in a small region within the nucleus and then determine whether the damaged DNA was distributed throughout the genome or was localized to specific chromosomes. Investigators found that the damage was confined to just a few chromosomes, and that the reverse experiment, in which laser-UV-microirradiation was used to cause damage to a small portion of the metaphase plate produced daughter-cell nuclei with mirror-image DNA-damage visualized in interphase; a finding that strongly supports the CT model [Zorn et al., 1979; Zorn et al., 1976]. The development of chromosome paints that consist of fluorescent-dye labeled chromosome-specific probes enabled direct visualization of individual chromosomes in interphase nuclei [Lichter et al., 1988; Manuelidis, 1985; Schardin et al., 1985]. These findings were further corroborated by experiments in which fluorescent nucleotide analogs were used to uniformly pulse-label replicating DNA. After several rounds of replication/independent assortment of chromosomes, individual CTs could be visualized in live cells [Visser and Aten, 1999; Zink and Cremer, 1998]. Within these CT, subdomains were proposed to occur via 6–12 chromatin loops of approximately 50–200 kb arranged in a rosette pattern composed of approximately 1 Mbp of DNA [Berezney, 2002; Jackson and Pombo, 1998; Ma et al., 1998]. The data obtained using chromatin capture techniques has extended this concept to include more precise characterization of the intrachromosomal interactions defined by so-called topologically associating domains (TADs) that represent discrete domains of relatively high DNA-contact frequencies [Barutcu et al., 2015; Dekker et al., 2013]. Chromosome organization in CTs during interphase is now widely accepted in the scientific community.

FISH and Chromatin Conformation Capture—Complementary Approaches for Chromosome Territory Delineation

Currently, two main techniques are used to study chromosome territory organization: fluorescence in situ hybridization (FISH) and chromatin conformation capture. FISH is a microscopic method to visualize specific DNA or RNA sequences within the nucleus by hybridizing complementary fluorescent-dye labeled DNA or RNA probes. Chromatin conformation capture comprises several related molecular biology techniques that involve crosslinking nuclei to “capture” intra- and inter-chromosomal interactions, fragmenting DNA with restriction enzymes, ligating the interacting fragments, purifying the DNA, and finally identifying the captured sequences by deep-sequencing, microarray analysis, or PCR.

FISH has been used extensively to identify chromosomal aberrations (e.g., translocations or deletions), and characterize CTs, and other nuclear bodies and microenvironments. These analyses are enabled by an increasingly sophisticated array of 3-D computational image-analysis tools, such as NEMO [Iannucci et al., 2010], eFISHent [Fritz et al., 2014b], and TANGO [Ollion et al., 2013]. Chromosome conformation capture strategies include a compendium of techniques that include 3C, 4C, 5C, Hi-C, and ChIA-PET [Barutcu et al., 2015]. These methods provide a global snapshot of chromosome interactions. Used together, FISH and chromosome conformation capture techniques can complement the limitations inherent to each strategy on its own; they can be used to validate each other for CT analyses. Table 1 summarizes the strengths and limitations of FISH microscopy and chromosome conformation capture strategies for analyzing CTs.

A spectrum of increasingly complex chromatin conformations impacting gene expression

The configuration of chromatin is coincident with the coordinated expression of genes under different physiological conditions, such as cell cycle, differentiation state, and disease status. This has been shown to be critical for the transcription of individual genes that rely on enhancer-promoter interactions that may cover over 100 kb of linear genomic distance (Figure 1B) [Kadauke and Blobel, 2009; Krivega and Dean, 2012]. Examples of this include the β -globin locus [Carter et al., 2002; Palstra et al., 2003; Tolhuis et al., 2002], IFNG [Eivazova and Aune, 2004], MHC class II [Majumder and Boss, 2010], and CFTR [Gheldof et al., 2010] genes. Upon transcription, these genes and their respective enhancers are in close proximity. This conformation, termed active chromatin hubs (ACHs), is mediated through the binding of transcription factors, and is only present when these genes are expressed. In hematopoietic stem cells (HSCs), the transcription factor Runx1 is required for the interaction between the CD34 gene promoter and its downstream regulatory element [Levantini et al., 2011]. Transcription factor-mediated chromatin looping was demonstrated in erythroid cells. In order to adopt an ACH formation, the Beta-globin gene and its locus of control require specific transcription factors, including EKLF1 and GATA1 [Drissen et al., 2004]. Additionally, the global chromatin organizer and transcription factor, SATB1 was also shown to form an ACH in the Rag1 and Rag2 genes in thymocyte development [Hao et al., 2015].

Physical proximity is not only essential for the expression of individual genes; multiple genes that are coexpressed may congregate into ACHs when active (Figure 1C) [Osborne et al., 2004a; Osborne et al., 2007; Spilianakis et al., 2005]. For example, the active alpha-globin genes loop into an ACH with several nearby housekeeping genes within a 130kb domain [Zhou et al., 2006]. These enhancer-promoter and local DNA clusters might represent what are now recognized as topologically associating domains (TADs), defined as preferential interactions that occur over approximately 100 kb to 1 Mbp genomic sequence lengths [Barutcu et al., 2015; Dekker et al., 2013; Lieberman-Aiden et al., 2009]. Although chromosomal contacts within TADs are more prevalent, interactions occur between genes that are located up to ~40 mbp apart in different TADs. In murine erythroid cells, the highly transcribed hemoglobin beta gene, Hbb-b1, and other active genes distally located in sequence, colocalize at shared transcription factories [Osborne et al., 2004b]. Clusters of active genes in open chromatin (A-type compartments) or inactive genes in closed chromatin (B-type compartments) were later established using 3C assays [Cremer et al., 2015; Dekker, 2014].

This clustered chromatin organization is created through boundaries between neighboring genes or gene clusters with different expression programs and/or nearby heterochromatic microenvironments [Giles et al., 2010]. Specifically, certain DNA sequences and protein factors may act as insulators, providing a barrier to block the inappropriate spreading of heterochromatin or contact with enhancers that belong to other genes. One such site, discovered via DNase-sensitivity analysis, is known as 52HS4; it marks the separation between the beta-globin genes and an upstream heterochromatic region [Chung et al., 1997; Chung et al., 1993]. A well-known factor involved in this insulator function is known as CCCTC-binding factor (CTCF). One function of CTCF is to prevent the inappropriate enhancer activation of genes. For example, CTCF is critical for monoallelic expression in imprinting. CTCF binds to the imprinted control region (ICR) of the maternal allele, thus preventing distal enhancers from activating Igf2. In contrast the paternal-allele ICR DNA is methylated, thereby occluding CTCF and allowing the enhancer to activate Igf2 transcription [Bell and Felsenfeld, 2000; Hark et al., 2000; Kanduri et al., 2000]. Genome-wide, CTCF and cohesin complexes have been shown to demarcate TAD boundaries [Phillips-Cremins and Corces, 2013; Seitan et al., 2013; Sofueva et al., 2013; Zuin et al., 2014] and are involved in the evolutionary conservation and alteration of TADs [Vietri Rudan et al., 2015].

Chromatin alterations can be visualized microscopically. For example, actively expressed genes project out of the CT in chromatin loops (Figure 1D), but remain inside their respective CTs when expression is not in progress [Chambeyron and Bickmore, 2004; Volpi et al., 2000; Williams et al., 2002]. This has been demonstrated for the major histocompatibility complex on CT6 [Volpi et al., 2000], HOX genes on CT11 [Chambeyron and Bickmore, 2004], and the epidermal differentiation complex on CT1 [Mahy et al., 2002a; Williams et al., 2002]. On a broader level it was suggested that actively expressed genes are more often found on the periphery of CT [Zirbel et al., 1993]. However, other investigations have demonstrated that both active and inactive genes are found at the CT boundary [Clemson et al., 2006; Kurz et al., 1996] and that genes are evenly distributed throughout the CT regardless of their expression level [Kupper et al., 2007; Mahy et al.,

2002b]. Moreover, re-localization of the HOX genes to the CT periphery is not required upon their activation in different cell types [Morey et al., 2007; Morey et al., 2009]. More recently, with the adoption of chromatin capture techniques, it was shown that more active domains are more likely to be found on the periphery of CTs [Nagano et al., 2013], offering credence to an early hypothesis that lost favor before the use of molecular strategies.

The presence of active domains at CT boundaries suggests the possibility that specific coregulated genes may interact at CT interfaces. Interchromosomal interactions have been detected between Myc and Igh [Osborne et al., 2007] and human erythroid-specific genes [Brown et al., 2008]. Interestingly, Spilianakis and colleagues also found that, not only do the regulatory regions of the Locus Control Region (LCR) and TH2 interact on chromosome 11, but they also form an interchromosomal cluster with the IFN- γ gene on chromosome 10 [Spilianakis et al., 2005].

Colocalization has also been shown to preserve gene inactivity. For example downregulation of the lamin b receptor mediates clustering of olfactory genes at heterochromatic foci [Clowney et al., 2012]. The localization of heterochromatin at specific nuclear microenvironments such as the nuclear periphery, perinucleolar regions [Nemeth et al., 2010; van Koningsbruggen et al., 2010], and within polycomb bodies in drosophila [Tolhuis et al., 2011] results in the clustering of inactive genes [Guelen et al., 2008].

Chromosome Territory neighborhoods, a network of interacting chromosomes

In addition to specific interchromosomal interactions between genes, heterologous CT-neighborhood arrangements have been established [Bolzer et al., 2005; Nagele et al., 1995]. This suggests that the colocalization of genes between chromosomes may be established not only by genes projecting out of their CT on chromatin loops, but also through repositioning of entire CTs. Preferential interchromosomal associations measured between pairwise heterologous CTs suggest an overall nonrandom organization of CTs with respect to each other [Bolzer et al., 2005; Brianna Caddle et al., 2007; Khalil et al., 2007; Kuroda et al., 2004; Marella et al., 2009a; Marella et al., 2009c; Mayer et al., 2005; Zeitz et al., 2009]. This nonrandom organization of chromosomes in neighborhoods is relevant for the establishment of cellular identity. As cells differentiate, entire CTs are repositioned [Kuroda et al., 2004; Marella et al., 2009c] and individual genes within CTs exhibit altered interchromosomal interactions; the pluripotency-related gene Nanog is an example [Apostolou et al., 2013]. In, and unique CT interaction profiles are found in different cell and tissue lineages [Marella et al., 2009a; Mayer et al., 2005; Parada et al., 2004; Tanabe et al., 2002b; Zeitz et al., 2009]. These profiles, in turn, suggest that genomic organization could explain the high frequency of particular translocations prevalent in different cancer types (discussed below).

Although seminal, microscopic studies of interchromosomal organization have limitations. First, the intranuclear locations of CTs are probable rather than absolute, and therefore efforts to map CTs are challenging. One such limitation is in the determination of the CT positions. The majority of studies that utilize FISH to determine CT organization have

primarily focused on their gravity centers, thus interactions at the interfaces of CTs are not captured. This is significant because studies that have focused on CT-border interactions reveal high variability in the distance between CT-centers of interacting chromosomes [Fritz et al., 2014b; Pliss et al., 2014]. Furthermore, due to technical limitations, most studies have analyzed only three chromosomes per nucleus. Recent investigations, however, using a variation on FISH known as reFISH to sequentially label up to 9 CT, have determined specific CT neighborhoods for each individual homolog within the nucleus [Fritz et al., 2014b; Pliss et al., 2014]. Regardless, technical limitations in microscopic studies restrict their ability to provide a full understanding of the complex nature of nuclear organization. Hi-C, on the contrary, can provide a more comprehensive view of intra- and interchromosomal interactions. However, the data represent a snapshot in time and the average of millions of interactions within a population. In a heterogeneous population, Hi-C cannot differentiate stable CT interactions in a subpopulation of cells from a series of dynamic interactions that occur in most cells at different times. The probable, yet variable, nature of interchromosomal CT interactions is likely due to the fact that the genome as a whole is far more complex than any particular gene family or network of gene regulation.

Radial nuclear positioning, centering the genome

Genome organization is a fundamental epigenetic mediator of gene expression [Berezney, 2002; Berezney et al., 2005; Bickmore, 2013; Cremer and Cremer, 2010; Kumaran et al., 2008; Meaburn and Misteli, 2007; Misteli, 2004; Misteli, 2007; Stein G, 2003; Stein G, 2008]. It is generally accepted that the three dimensional positions of CTs are nonrandom relative to the nuclear periphery. Initial studies indicated that CT positioning correlates with gene activity, with higher gene activity levels associated with more interior CT positioning [Boyle et al., 2001; Kreth et al., 2004]. The inactive-X CT is a clear example; it is located closer to the nuclear periphery than its active counterpart [Dyer et al., 1989]. Later studies suggested that gene density influences the nonrandom radial organization of somatic CTs [Boyle et al., 2001]. Because gene-dense chromosomes generally exhibit higher levels of transcription than gene-poor chromosomes [Versteeg et al., 2003], gene activity may be the underlying basis for the radial positioning of the genome. Interestingly, this paradigm holds true not just for entire CTs, but also for individual genes and gene complexes within chromosomes; expressed alleles are generally found further from the nuclear periphery than alleles that are not expressed [Dietzel et al., 2004; Fedorova and Zink, 2009; Finlan et al., 2008; Fraser and Bickmore, 2007; Pickersgill et al., 2006; Reddy et al., 2008; Takizawa et al., 2008; Therizols et al., 2014]. This could result from the preferential localization of tightly packed chromatin at the nuclear periphery [Belmont et al., 1993; Busch, 1966]. While tethering to the nuclear lamina results in repression of transcription [Amendola and van Steensel, 2014], some genes do not display this phenomenon when recruited to the nuclear periphery [Kumaran and Spector, 2008]. It is important to note that transcription can and does occur near the nuclear periphery and conversely, inactive genes are sometimes located in the nuclear interior [Cremer et al., 2015; Misteli, 2013].

Another possible principle governing nuclear organization is born out by studies suggesting that chromosome size is the major determinant of the nonrandom radial positioning of CTs. These investigations have found that longer chromosomes are located closer to the nuclear

periphery [Sun et al., 2000]. Efforts to reconcile the findings from these studies point to cell cycle and nuclear shape as possible effectors. For example, CT are repositioned in cycling fibroblasts compared to cells in G0 [Mehta et al., 2010]. Other studies have attributed differences in nuclear shape between cell types as a key determinant of whether chromosome size or gene densities drive radial CT positioning [Bolzer et al., 2005]. CT size–radial position relationships are more prevalent in ellipsoidal fibroblast nuclei, whereas studies in more spherical lymphocyte nuclei find a higher correlation between gene density and radial arrangement which is conserved across evolution [Tanabe et al., 2002a; Tanabe et al., 2005].

Interactions between chromosome territories and other nuclear compartments

A more mechanistic perspective examines nonrandom organization of CTs established through functional interactions with other nuclear structures. For example, the nuclear lamina that lines the nuclear envelope have been proposed to have a major role in organizing the genome [Filesi et al., 2005; Goldman et al., 2005]. Lamina-associated domains (LADs) within chromosomes tether CTs to the nuclear periphery (Figure 1D–E) and are found with greater frequency in CTs that are peripherally positioned [Bridger et al., 2007; Guelen et al., 2008]. Consistent with observations about gene activity and CT radial positioning, LADs are typically found in repressed heterochromatin microenvironments. The condensation of DNA into tight heterochromatin substantially hinders access to DNA by nuclear machinery for transcription, replication, and DNA/chromosome modification. Some studies have suggested that recruitment to a peripheral heterochromatic microenvironment may have functional implications for replication [Li et al., 2001], transcription [Finlan et al., 2008; Kumaran et al., 2008; Reddy et al., 2008] and genome stability [Oza and Peterson, 2010]. These ideas are supported by the observation that disruption of the nuclear lamina in patients with progeria results in perturbation of telomere length [Gonzalez-Suarez et al., 2009], radial CT positioning [Mehta et al., 2011], and the condensation and compartmentalization of peripheral heterochromatin [Goldman et al., 2004; Nikolova et al., 2004; Taimen et al., 2009].

While the nuclear lamina is thought to exert its largest influence on CTs near the nuclear periphery, other nuclear bodies and microenvironments mediate genomic organization further inside the nucleus. For example, the interchromosomal interactions between the human erythroid-specific genes colocalize at nuclear speckles [Brown et al., 2008]. However, one of the most extensively studied prominent domains impacting nuclear organization is the nucleolus. They are formed by the congregation of the acrocentric chromosomes 13, 14, 15, 21 and 22, specifically their nucleolar organizer regions (NORs), which are comprised of tandem arrays of ribosomal RNA (rRNA) genes [Carmo-Fonseca et al., 2000; Prieto and McStay, 2005]. Although some NORs remain associated with components of the RNA polymerase I transcriptional machinery through mitosis (so-called competent NORs), others are not “bookmarked” and rRNA transcription factors assemble and disassemble with each cell cycle [Schlesinger et al., 2009; Smirnov et al., 2006]. During interphase, competent NORs tend to be located closer to nucleoli than their non-competent

counterparts [Kalmarova et al., 2007], however both types of NOR-bearing CTs are frequently associated with nucleoli [Kalmarova et al., 2007; Sullivan et al., 2001]. Detailed biochemical and DNA-sequence analysis of nucleoli indicates that not only the acrocentric NOR-bearing chromosomes interact with nucleoli, but all chromosomes have nucleolar-associating domains (NADs). These NADs share sequence similarity with LADs and localize to the perinucleolar heterochromatin (Figure 1D–E) [Nemeth et al., 2010].

Despite their substantial impact on the nuclear landscape, the number of nucleoli in any given nucleus varies from one to five. As a result, NOR-bearing CTs display heterogeneous nucleolar-interaction profiles within a population of cells. In nuclei with multiple nucleoli, a dominant nucleolus has more contributing NOR-bearing CTs than the others. Furthermore, even nuclei with the same number of nucleoli may exhibit different patterns of association with NOR-bearing CT. For example, a different complement of NOR-bearing CT are found to associate with each nucleolus in cells that contain only two nucleoli. This association of CTs with nucleoli at the nuclear interior, may help to explain the correlation between short sequence length and interior radial positioning [Bolzer et al., 2005; Heride et al., 2010; Raska et al., 2006]. The fact that shorter NOR-bearing CT are preferentially located further from the nuclear periphery than their longer NOR-bearing counterparts [Pliss et al., 2014] suggests that CT size outweighs the influence of nucleolar association in determining radial position.

Chromosome territory organization in the cell cycle

As cell exits mitosis, chromosomes de-compact and gene expression resumes—ostensibly recapitulating the program of the parent cells to maintain their cellular lineage. This is accomplished in part through a process known as bookmarking, in which key transcription factors are retained on chromosomes through mitosis [Delcuve et al., 2008]. Following mitotic de-compactation, bookmarking enables the rapid resumption of cell-specific gene expression or continued genes repression in subsequent generations [Kadauke and Blobel, 2013]. An interesting example of this is the residency of transcription factors at rDNA. In mesenchymal stem cells (MSCs), which are not lineage committed, Myc bookmarking of rDNA promotes rRNA expression in progeny cells. This bookmarking could enable immediate post-mitosis resumption of gene expression that is needed to fuel the rapid cell cycle progression characteristic of MSCs. However, in cells derived from MSCs, that have a longer cell cycle, various transcription factors act to reduce rRNA expression; these include Runx1 in osteoblasts, MyoD in myoblasts, and C/EBP in adipocytes [Zaidi et al., 2010].

Following mitosis, in early G1, in vivo–labeled chromosome domains are mobile relative to the nucleus and each other [Csink and Henikoff, 1998; Lucas and Cervantes, 2002; Walter et al., 2003] whereas during the rest of the cell cycle, the large scale arrangement of chromatin in the cell nucleus exhibits a degree of stability [Abney et al., 1997; Bornfleth et al., 1999; Chubb et al., 2002; Gerlich et al., 2003; Lucas and Cervantes, 2002; Shelby et al., 1996; Zink and Cremer, 1998]. This observation suggests that the breakdown of the nuclear envelope and its reassembly at the end of mitosis could provide an opportunity for remodeling of nuclear organization in subsequent generations. Outside of mitosis, overall CT morphologies were demonstrated to be maintained for up to four hours in the cell cycle

[Edelmann et al., 2001; Muller et al., 2010], however alterations were detected in internal CT organization [Muller et al., 2010]. CT morphology was found to be altered, however, in studies that examined CTs in G1 compared to S-phase nuclei [Sehgal et al., 2014]. Along with this altered CT morphology, cell-type specific alterations in the interchromosomal organization of specific CT were demonstrated [Fritz et al., 2014a].

On a smaller scale, chromatin remodeling has been shown to control progression through the cell cycle by regulating transcription of essential cell-cycle genes [Cao et al., 1997; Vignali et al., 2000] and DNA replication. For example, during S phase, loosening of chromatin structure enables the replication machinery to access the DNA [Raynaud et al., 2014]. At a global level, as cells progress from early S to mitosis, chromatin becomes increasingly condensed and less sensitive to DNase I [Pfeffer et al., 1991]. Using Hi-C on synchronized HeLa cells, it was demonstrated that the intrachromosomal TADs within specific CTs are largely, but not entirely, maintained across the cell cycle [Naumova et al., 2013]. This work also determined that TADs are broken down in mitosis and are re-established in early G1 [Naumova et al., 2013]. However, the degree to which the organization within specific TADs is altered during interphase from early G1 to late G2 remains unclear.

Chromosome territory organization in cancer

Changes in nuclear structure have long been used as a major diagnostic tool to detect cancer [Zaidi et al., 2007; Zink et al., 2004]. Cancer nuclei are generally larger and more irregularly shaped than nuclei in healthy cells [Zink et al., 2004]; nuclear compartmentalization is also altered. For example, some cancers are characterized by an increase in the numbers and sizes of nucleoli, possibly as a result of an increased requirement for protein synthesis [Derenzini et al., 2000; Maggi and Weber, 2005]. Partial loss of heterochromatin compartmentalization is another characteristic change [Zink et al., 2004]. Although it is known that nuclear architecture is altered in cancer, less is known about the concomitant changes in CT organization. Studies of nuclear organization in cancer are complicated by the fact that comparisons are often made between cells from different origins and the karyotypes of each cancer cell type are highly variable—karyotype heterogeneity within a population of cells further complicates analyses. Even in cells with similar karyotypes, cell-to-cell variability in CT organization has been demonstrated using microscopy and single-cell Hi-C [Nagano et al., 2013]. Finally, even single-cell Hi-C, cannot currently distinguish the data from individual somatic chromosomes; data represent an average of both CT homologs [Nagano et al., 2013]. When determining the organization of CT this is of particular importance. For example, in chromosomes that are involved in translocations, altered contact frequencies could be found within the translocated CT, the non-translocated CT, or distributed between both homologs of the CT.

Interchromosomal interactions have been suggested to play a role in cancer progression. Cell-type differences in CT organization may present an explanation for the high frequency of specific translocations that occur in the progression of several different cancers. Increased translocation frequency has been determined between specific CTs that are in closer proximity [Brianna Caddle et al., 2007; Folle, 2008; Parada et al., 2002; Roix et al., 2003; Soutoglou et al., 2007] and have higher degrees of intermingling [Branco and Pombo, 2006].

Within cancer nuclei, it was demonstrated that translocated CT regions retain the preferential interaction partners they had before the translocation [Parada et al., 2002]. A wide-scale alteration in interactions between non-translocated CTs was determined in MCF10A cells versus their metastatic counterpart MCF10CA1a cells [Fritz et al., 2014b]. Interestingly, the inactive X CT was found to have fewer interchromosomal interactions in MCF10A, but not in MCF10CA1a cells. This increase in interchromosomal interactions was coincident with increased CT volume and global gene expression within the X chromosome [Fritz et al., 2014b]. Changes in CT interaction profiles extend to the gene level in cancer; it was shown that the specific genes that participate in interchromosomal interactions are different in breast cancer versus normal breast cells [Zeitz et al., 2013].

Conclusion

Although progress has been made in understanding how CTs influence and/or respond to nuclear organization, further work to fully elucidate the rules that govern the functional relationships between genomic organization, and genetic and epigenetic regulation in the context of nuclear microenvironments is needed. Due to the complexity of genome-wide regulatory networks, high-throughput biochemical techniques will be vital for illuminating these functional relationships. The dynamic nature of structural and functional relationships across the cell cycle (e.g., from mitotic bookmarking of genes with their respective regulatory proteins to domain assembly during interphase) as well as their perturbation in disease also require further study. Architectural signatures that reveal specificity of localization and/or colocalization for regulatory domains that may be altered could represent targets for the detection and treatment of disease states. Microscopy remains an important approach for high-throughput genome-wide studies; used in combination with emerging molecular chromatin conformation capture techniques, and ever more sophisticated and powerful analysis methods, it is likely that ongoing investigation will yield ever more precise and detailed information on the role of CT organization in biology.

Acknowledgments

Contract Grant Sponsor: National Institutes of Health; Contract Grant Numbers: P01 CA082834

References

- Abney JR, Cutler B, Fillbach ML, Axelrod D, Scalettar BA. Chromatin dynamics in interphase nuclei and its implications for nuclear structure. *J Cell Biol.* 1997; 137:1459–68. [PubMed: 9199163]
- Amendola M, van Steensel B. Mechanisms and dynamics of nuclear lamina-genome interactions. *Curr Opin Cell Biol.* 2014; 28:61–8. [PubMed: 24694724]
- Apostolou E, Ferrari F, Walsh RM, Bar-Nur O, Stadtfeld M, Cheloufi S, Stuart HT, Polo JM, Ohsumi TK, Borowsky ML, Kharchenko PV, Park PJ, Hochedlinger K. Genome-wide chromatin interactions of the Nanog locus in pluripotency, differentiation, and reprogramming. *Cell Stem Cell.* 2013; 12:699–712. [PubMed: 23665121]
- Barutcu AR, Fritz AJ, Zaidi SK, vanWijnen AJ, Lian JB, Stein JL, Nickerson JA, Imbalzano AN, Stein GS. C-ing the Genome: A Compendium of Chromosome Conformation Capture Methods to Study Higher-Order Chromatin Organization. *J Cell Physiol.* 2015
- Bell AC, Felsenfeld G. Methylation of a CTCF-dependent boundary controls imprinted expression of the Igf2 gene. *Nature.* 2000; 405:482–5. [PubMed: 10839546]

- Belmont AS, Zhai Y, Thilenius A. Lamin B distribution and association with peripheral chromatin revealed by optical sectioning and electron microscopy tomography. *J Cell Biol.* 1993; 123:1671–85. [PubMed: 8276889]
- Berezney R. Regulating the mammalian genome: the role of nuclear architecture. *Advan Enzyme Regul.* 2002; 42:39–52. [PubMed: 12123705]
- Berezney R, Malyavantham KS, Pliss A, Bhattacharya S, Acharya R. Spatio-temporal dynamics of genomic organization and function in the mammalian cell nucleus. *Adv Enzyme Regul.* 2005; 45:17–26. [PubMed: 16139341]
- Bickmore WA. The spatial organization of the human genome. *Annu Rev Genomics Hum Genet.* 2013; 14:67–84. [PubMed: 23875797]
- Bolzer A, Kreth G, Solovei I, Koehler D, Saracoglu K, Fauth C, Muller S, Eils R, Cremer C, Speicher MR, Cremer T. Three-dimensional maps of all chromosomes in human male fibroblast nuclei and prometaphase rosettes. *PLoS Biol.* 2005; 3:e157. [PubMed: 15839726]
- Bornfleth H, Edelmann P, Zink D, Cremer T, Cremer C. Quantitative motion analysis of subchromosomal foci in living cells using four-dimensional microscopy. *Biophys J.* 1999; 77:2871–86. [PubMed: 10545385]
- Boyle S, Gilchrist S, Bridger JM, Mahy NL, Ellis JA, Bickmore WA. The spatial organization of human chromosomes within the nuclei of normal and emerin-mutant cells. *Hum Mol Genet.* 2001; 10:211–9. [PubMed: 11159939]
- Branco MR, Pombo A. Intermingling of chromosome territories in interphase suggests role in translocations and transcription-dependent associations. *PLoS Biol.* 2006; 4:e138. [PubMed: 16623600]
- Brianna Caddle L, Grant JL, Szatkiewicz J, van Hase J, Shirley BJ, Bewersdorf J, Cremer C, Arneodo A, Khalil A, Mills KD. Chromosome neighborhood composition determines translocation outcomes after exposure to high-dose radiation in primary cells. *Chromosome Res.* 2007; 15:1061–73. [PubMed: 18060570]
- Bridger JM, Foeger N, Kill IR, Herrmann H. The nuclear lamina. Both a structural framework and a platform for genome organization. *FEBS J.* 2007; 274:1354–61. [PubMed: 17489093]
- Brown JM, Green J, das Neves RP, Wallace HA, Smith AJ, Hughes J, Gray N, Taylor S, Wood WG, Higgs DR, Iborra FJ, Buckle VJ. Association between active genes occurs at nuclear speckles and is modulated by chromatin environment. *J Cell Biol.* 2008; 182:1083–97. [PubMed: 18809724]
- Busch H. The cell nucleus. *Nature.* 1966; 211:1347–48.
- Cao Y, Cairns BR, Kornberg RD, Laurent BC. Sfh1p, a component of a novel chromatin-remodeling complex, is required for cell cycle progression. *Mol Cell Biol.* 1997; 17:3323–34. [PubMed: 9154831]
- Carmo-Fonseca M, Mendes-Soares L, Campos I. To be or not to be in the nucleolus. *Nat Cell Biol.* 2000; 2:E107–12. [PubMed: 10854340]
- Carter D, Chakalova L, Osborne CS, Dai YF, Fraser P. Long-range chromatin regulatory interactions in vivo. *Nat Genet.* 2002; 32:623–6. [PubMed: 12426570]
- Chambeyron S, Bickmore WA. Chromatin decondensation and nuclear reorganization of the HoxB locus upon induction of transcription. *Genes Dev.* 2004; 18:1119–30. [PubMed: 15155579]
- Chen B, Gilbert LA, Cimini BA, Schnitzbauer J, Zhang W, Li GW, Park J, Blackburn EH, Weissman JS, Qi LS, Huang B. Dynamic imaging of genomic loci in living human cells by an optimized CRISPR/Cas system. *Cell.* 2013; 155:1479–91. [PubMed: 24360272]
- Chubb JR, Boyle S, Perry P, Bickmore WA. Chromatin motion is constrained by association with nuclear compartments in human cells. *Curr Biol.* 2002; 12:439–45. [PubMed: 11909528]
- Chung JH, Bell AC, Felsenfeld G. Characterization of the chicken beta-globin insulator. *Proc Natl Acad Sci U S A.* 1997; 94:575–80. [PubMed: 9012826]
- Chung JH, Whiteley M, Felsenfeld G. A 5' element of the chicken beta-globin domain serves as an insulator in human erythroid cells and protects against position effect in *Drosophila*. *Cell.* 1993; 74:505–14. [PubMed: 8348617]
- Clemson CM, Hall LL, Byron M, McNeil J, Lawrence JB. The X chromosome is organized into a gene-rich outer rim and an internal core containing silenced nongenic sequences. *Proc Natl Acad Sci U S A.* 2006; 103:7688–93. [PubMed: 16682630]

- Clowney EJ, LeGros MA, Mosley CP, Clowney FG, Markenskoff-Papadimitriou EC, Myllys M, Barnea G, Larabell CA, Lomvardas S. Nuclear aggregation of olfactory receptor genes governs their monogenic expression. *Cell*. 2012; 151:724–37. [PubMed: 23141535]
- Cremer T, Cremer C. Chromosome territories, nuclear architecture and gene regulation in mammalian cells. *Nat Rev Genet*. 2001; 2:292–301. [PubMed: 11283701]
- Cremer T, Cremer M. Chromosome territories. *Cold Spring Harb Perspect Biol*. 2010; 2:1–22.
- Cremer T, Cremer M, Dietzel S, Muller S, Solovei I, Fakan S. Chromosome territories--a functional nuclear landscape. *Curr Opin Cell Biol*. 2006; 18:307–16. [PubMed: 16687245]
- Cremer T, Cremer M, Hubner B, Strickfaden H, Smeets D, Popken J, Sterr M, Markaki Y, Rippe K, Cremer C. The 4D nucleome: Evidence for a dynamic nuclear landscape based on co-aligned active and inactive nuclear compartments. *FEBS Lett*. 2015
- Cremer T, Peterson SP, Cremer C, Berns MW. Laser microirradiation of Chinese hamster cells at wavelength 365 nm: effects of psoralen and caffeine. *Radiat Res*. 1981; 85:529–43. [PubMed: 7010411]
- Csink AK, Henikoff S. Large-scale chromosomal movements during interphase progression in *Drosophila*. *J Cell Biol*. 1998; 143:13–22. [PubMed: 9763417]
- Dekker J. Two ways to fold the genome during the cell cycle: insights obtained with chromosome conformation capture. *Epigenetics Chromatin*. 2014; 7:25. [PubMed: 25435919]
- Dekker J, Marti-Renom MA, Mirny LA. Exploring the three-dimensional organization of genomes: interpreting chromatin interaction data. *Nat Rev Genet*. 2013; 14:390–403. [PubMed: 23657480]
- Delcuve GP, He S, Davie JR. Mitotic partitioning of transcription factors. *J Cell Biochem*. 2008; 105:1–8. [PubMed: 18459122]
- Derenzini M, Trere D, Pession A, Govoni M, Sirri V, Chieco P. Nucleolar size indicates the rapidity of cell proliferation in cancer tissues. *J Pathol*. 2000; 191:181–6. [PubMed: 10861579]
- Dietzel S, Zolghadr K, Hepperger C, Belmont AS. Differential large-scale chromatin compaction and intranuclear positioning of transcribed versus non-transcribed transgene arrays containing beta-globin regulatory sequences. *J Cell Sci*. 2004; 117:4603–14. [PubMed: 15331668]
- Drissen R, Palstra RJ, Gillemans N, Splinter E, Grosveld F, Philipsen S, de Laat W. The active spatial organization of the beta-globin locus requires the transcription factor EKLF. *Genes Dev*. 2004; 18:2485–90. [PubMed: 15489291]
- Dyer KA, Canfield TK, Gartler SM. Molecular cytological differentiation of active from inactive X domains in interphase: implications for X chromosome inactivation. *Cytogenet Cell Genet*. 1989; 50:116–20. [PubMed: 2776476]
- Edelmann P, Bornfleth H, Zink D, Cremer T, Cremer C. Morphology and dynamics of chromosome territories in living cells. *Biochim Biophys Acta*. 2001; 1551:M29–39. [PubMed: 11553419]
- Eivazova ER, Aune TM. Dynamic alterations in the conformation of the *Ifng* gene region during T helper cell differentiation. *Proc Natl Acad Sci U S A*. 2004; 101:251–6. [PubMed: 14691261]
- Fedorova E, Zink D. Nuclear genome organization: common themes and individual patterns. *Curr Opin Genet Dev*. 2009; 19:166–71. [PubMed: 19321336]
- Filesi I, Gullotta F, Lattanzi G, D'Apice MR, Capanni C, Nardone AM, Columbaro M, Scarano G, Mattioli E, Sabatelli P, Maraldi NM, Biocca S, Novelli G. Alterations of nuclear envelope and chromatin organization in mandibuloacral dysplasia, a rare form of laminopathy. *Physiol Genomics*. 2005; 23:150–8. [PubMed: 16046620]
- Finlan LE, Sproul D, Thomson I, Boyle S, Kerr E, Perry P, Ylstra B, Chubb JR, Bickmore WA. Recruitment to the nuclear periphery can alter expression of genes in human cells. *PLoS Genet*. 2008; 4:e1000039. [PubMed: 18369458]
- Folle GA. Nuclear architecture, chromosome domains and genetic damage. *Mutat Res*. 2008; 658:172–83. [PubMed: 17921046]
- Fraser P, Bickmore W. Nuclear organization of the genome and the potential for gene regulation. *Nature*. 2007; 447:413–7. [PubMed: 17522674]
- Fritz AJ, Stojkovic B, Ding H, Xu J, Bhattacharya S, Berezney R. Cell type specific alterations in interchromosomal networks across the cell cycle. *PLoS Comput Biol*. 2014a; 10:e1003857. [PubMed: 25275626]

- Fritz AJ, Stojkovic B, Ding H, Xu J, Bhattacharya S, Gaile D, Berezney R. Wide-scale alterations in interchromosomal organization in breast cancer cells: defining a network of interacting chromosomes. *Hum Mol Genet.* 2014b
- Gavrilov AA, Gushchanskaya ES, Strelkova O, Zhironkina O, Kireev II, Iarovaia OV, Razin SV. Disclosure of a structural milieu for the proximity ligation reveals the elusive nature of an active chromatin hub. *Nucleic Acids Res.* 2013; 41:3563–75. [PubMed: 23396278]
- Gerlich D, Beaudouin J, Kalbfuss B, Daigle N, Eils R, Ellenberg J. Global chromosome positions are transmitted through mitosis in mammalian cells. *Cell.* 2003; 112:751–64. [PubMed: 12654243]
- Gheldof N, Smith EM, Tabuchi TM, Koch CM, Dunham I, Stamatoyannopoulos JA, Dekker J. Cell-type-specific long-range looping interactions identify distant regulatory elements of the CFTR gene. *Nucleic Acids Res.* 2010; 38:4325–36. [PubMed: 20360044]
- Giles KE, Gowher H, Ghirlando R, Jin C, Felsenfeld G. Chromatin boundaries, insulators, and long-range interactions in the nucleus. *Cold Spring Harb Symp Quant Biol.* 2010; 75:79–85. [PubMed: 21047907]
- Goldman RD, Goldman AE, Shumaker DK. Nuclear lamins: building blocks of nuclear structure and function. *Novartis Found Symp.* 2005; 264:3–16. discussion 16–21, 227–30. [PubMed: 15773744]
- Goldman RD, Shumaker DK, Erdos MR, Eriksson M, Goldman AE, Gordon LB, Gruenbaum Y, Khuon S, Mendez M, Varga R, Collins FS. Accumulation of mutant lamin A causes progressive changes in nuclear architecture in Hutchinson-Gilford progeria syndrome. *Proc Natl Acad Sci U S A.* 2004; 101:8963–8. [PubMed: 15184648]
- Gonzalez-Suarez I, Redwood AB, Perkins SM, Vermolen B, Lichtensztejn D, Grotsky DA, Morgado-Palacin L, Gapud EJ, Sleckman BP, Sullivan T, Sage J, Stewart CL, Mai S, Gonzalo S. Novel roles for A-type lamins in telomere biology and the DNA damage response pathway. *EMBO J.* 2009; 28:2414–27. [PubMed: 19629036]
- Guelen L, Pagie L, Brasset E, Meuleman W, Faza MB, Talhout W, Eussen BH, de Klein A, Wessels L, de Laat W, van Steensel B. Domain organization of human chromosomes revealed by mapping of nuclear lamina interactions. *Nature.* 2008; 453:948–51. [PubMed: 18463634]
- Hao B, Naik AK, Watanabe A, Tanaka H, Chen L, Richards HW, Kondo M, Taniuchi I, Kohwi Y, Kohwi-Shigematsu T, Krangel MS. An anti-silencer- and SATB1-dependent chromatin hub regulates Rag1 and Rag2 gene expression during thymocyte development. *J Exp Med.* 2015; 212:809–24. [PubMed: 25847946]
- Hark AT, Schoenherr CJ, Katz DJ, Ingram RS, Levorse JM, Tilghman SM. CTCF mediates methylation-sensitive enhancer-blocking activity at the H19/Igf2 locus. *Nature.* 2000; 405:486–9. [PubMed: 10839547]
- Hemmerich P, Schmiedeberg L, Diekmann S. Dynamic as well as stable protein interactions contribute to genome function and maintenance. *Chromosome Res.* 2011; 19:131–51. [PubMed: 21046224]
- Heride C, Ricoul M, Kieu K, von Hase J, Guillemot V, Cremer C, Dubrana K, Sabatier L. Distance between homologous chromosomes results from chromosome positioning constraints. *J Cell Sci.* 2010; 123:4063–75. [PubMed: 21084563]
- Iannuccelli E, Mompert F, Gellin J, Lahbib-Mansais Y, Yerle M, Boudier T. NEMO: a tool for analyzing gene and chromosome territory distributions from 3D-FISH experiments. *Bioinformatics.* 2010; 26:696–7. [PubMed: 20080510]
- Jackson DA, Pombo A. Replicon clusters are stable units of chromosome structure: evidence that nuclear organization contributes to the efficient activation and propagation of S phase in human cells. *J Cell Biol.* 1998; 140:1285–95. [PubMed: 9508763]
- Kadauke S, Blobel GA. Chromatin loops in gene regulation. *Biochim Biophys Acta.* 2009; 1789:17–25. [PubMed: 18675948]
- Kadauke S, Blobel GA. Mitotic bookmarking by transcription factors. *Epigenetics Chromatin.* 2013; 6:6. [PubMed: 23547918]
- Kalmarova M, Smirnov E, Masata M, Koberna K, Ligasova A, Popov A, Raska I. Positioning of NORs and NOR-bearing chromosomes in relation to nucleoli. *J Struct Biol.* 2007; 160:49–56. [PubMed: 17698369]

- Kanduri C, Pant V, Loukinov D, Pugacheva E, Qi CF, Wolffe A, Ohlsson R, Lobanenkov VV. Functional association of CTCF with the insulator upstream of the H19 gene is parent of origin-specific and methylation-sensitive. *Curr Biol*. 2000; 10:853–6. [PubMed: 10899010]
- Khalil A, Grant JL, Caddle LB, Atzema E, Mills KD, Arneodo A. Chromosome territories have a highly nonspherical morphology and nonrandom positioning. *Chromosome Res*. 2007; 15:899–916. [PubMed: 17926137]
- Kreth G, Finsterle J, von Hase J, Cremer M, Cremer C. Radial arrangement of chromosome territories in human cell nuclei: a computer model approach based on gene density indicates a probabilistic global positioning code. *Biophys J*. 2004; 86:2803–12. [PubMed: 15111398]
- Krivega I, Dean A. Enhancer and promoter interactions-long distance calls. *Curr Opin Genet Dev*. 2012; 22:79–85. [PubMed: 22169023]
- Kumaran RI, Spector DL. A genetic locus targeted to the nuclear periphery in living cells maintains its transcriptional competence. *J Cell Biol*. 2008; 180:51–65. [PubMed: 18195101]
- Kumaran RI, Thakar R, Spector DL. Chromatin dynamics and gene positioning. *Cell*. 2008; 132:929–34. [PubMed: 18358806]
- Kupper K, Kolbl A, Biener D, Dittrich S, von Hase J, Thormeyer T, Fiegler H, Carter NP, Speicher MR, Cremer T, Cremer M. Radial chromatin positioning is shaped by local gene density, not by gene expression. *Chromosoma*. 2007; 116:285–306. [PubMed: 17333233]
- Kuroda M, Tanabe H, Yoshida K, Oikawa K, Saito A, Kiyuna T, Mizusawa H, Mukai K. Alteration of chromosome positioning during adipocyte differentiation. *J Cell Sci*. 2004; 117:5897–903. [PubMed: 15537832]
- Kurz A, Lampel S, Nickolenko JE, Bradl J, Benner A, Zirbel RM, Cremer T, Lichter P. Active and inactive genes localize preferentially in the periphery of chromosome territories. *J Cell Biol*. 1996; 135:1195–205. [PubMed: 8947544]
- Lakadamyali M, Cosma MP. Advanced microscopy methods for visualizing chromatin structure. *FEBS Lett*. 2015
- Levantini E, Lee S, Radomska HS, Hetherington CJ, Alberich-Jorda M, Amabile G, Zhang P, Gonzalez DA, Zhang J, Basseres DS, Wilson NK, Koschmieder S, Huang G, Zhang DE, Ebralidze AK, Bonifer C, Okuno Y, Gottgens B, Tenen DG. RUNX1 regulates the CD34 gene in haematopoietic stem cells by mediating interactions with a distal regulatory element. *EMBO J*. 2011; 30:4059–70. [PubMed: 21873977]
- Li F, Chen J, Izumi M, Butler MC, Keezer SM, Gilbert DM. The replication timing program of the Chinese hamster beta-globin locus is established coincident with its repositioning near peripheral heterochromatin in early G1 phase. *J Cell Biol*. 2001; 154:283–92. [PubMed: 11470818]
- Lichter P, Cremer T, Borden J, Manuelidis L, Ward DC. Delineation of individual human chromosomes in metaphase and interphase cells by in situ suppression hybridization using recombinant DNA libraries. *Hum Genet*. 1988; 80:224–34. [PubMed: 3192212]
- Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO, Sandstrom R, Bernstein B, Bender MA, Groudine M, Gnirke A, Stamatoyanopoulos J, Mirny LA, Lander ES, Dekker J. Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science*. 2009; 326:289–93. [PubMed: 19815776]
- Long BR, Robinson DC, Zhong H. Subdiffractive microscopy: techniques, applications, and challenges. *Wiley Interdiscip Rev Syst Biol Med*. 2014; 6:151–68. [PubMed: 24443323]
- Lucas JN, Cervantes E. Significant large-scale chromosome territory movement occurs as a result of mitosis, but not during interphase. *Int J Radiat Biol*. 2002; 78:449–55. [PubMed: 12065049]
- Ma H, Samarabandu J, Devdhar RS, Acharya R, Cheng PC, Meng C, Berezney R. Spatial and temporal dynamics of DNA replication sites in mammalian cells. *J Cell Biol*. 1998; 143:1415–25. [PubMed: 9852140]
- Maggi LB Jr, Weber JD. Nucleolar adaptation in human cancer. *Cancer Invest*. 2005; 23:599–608. [PubMed: 16305988]
- Mahy NL, Perry PE, Bickmore WA. Gene density and transcription influence the localization of chromatin outside of chromosome territories detectable by FISH. *J Cell Biol*. 2002a; 159:753–63. [PubMed: 12473685]

- Mahy NL, Perry PE, Gilchrist S, Baldock RA, Bickmore WA. Spatial organization of active and inactive genes and noncoding DNA within chromosome territories. *J Cell Biol.* 2002b; 157:579–89. [PubMed: 11994314]
- Majumder P, Boss JM. CTCF controls expression and chromatin architecture of the human major histocompatibility complex class II locus. *Mol Cell Biol.* 2010; 30:4211–23. [PubMed: 20584980]
- Manuelidis L. Individual interphase chromosome domains revealed by in situ hybridization. *Hum Genet.* 1985; 71:288–93. [PubMed: 3908288]
- Marella NV, Bhattacharya S, Mukherjee L, Xu J, Berezney R. Cell type specific chromosome territory organization in the interphase nucleus of normal and cancer cells. *J Cell Physiol.* 2009a; 221:130–8. [PubMed: 19496171]
- Marella NV, Malyavantham KS, Wang J, Matsui S, Liang P, Berezney R. Cytogenetic and cDNA microarray expression analysis of MCF10 human breast cancer progression cell lines. *Cancer Res.* 2009b; 69:5946–53. [PubMed: 19584277]
- Marella NV, Seifert B, Nagarajan P, Sinha S, Berezney R. Chromosomal rearrangements during human epidermal keratinocyte differentiation. *J Cell Physiol.* 2009c; 221:139–46. [PubMed: 19626667]
- Markaki Y, Smeets D, Fiedler S, Schmid VJ, Schermelleh L, Cremer T, Cremer M. The potential of 3D-FISH and super-resolution structured illumination microscopy for studies of 3D nuclear architecture: 3D structured illumination microscopy of defined chromosomal structures visualized by 3D (immuno)-FISH opens new perspectives for studies of nuclear architecture. *Bioessays.* 2012; 34:412–26. [PubMed: 22508100]
- Mayer R, Brero A, von Hase J, Schroeder T, Cremer T, Dietzel S. Common themes and cell type specific variations of higher order chromatin arrangements in the mouse. *BMC Cell Biol.* 2005; 6:44. [PubMed: 16336643]
- Meaburn KJ, Misteli T. Cell biology: chromosome territories. *Nature.* 2007; 445:379–781. [PubMed: 17251970]
- Mehta IS, Amira M, Harvey AJ, Bridger JM. Rapid chromosome territory relocation by nuclear motor activity in response to serum removal in primary human fibroblasts. *Genome Biol.* 2010; 11:R5. [PubMed: 20070886]
- Mehta IS, Eskiw CH, Arican HD, Kill IR, Bridger JM. Farnesyltransferase inhibitor treatment restores chromosome territory positions and active chromosome dynamics in Hutchinson-Gilford progeria syndrome cells. *Genome Biol.* 2011; 12:R74. [PubMed: 21838864]
- Mehta IS, Figgitt M, Clements CS, Kill IR, Bridger JM. Alterations to nuclear architecture and genome behavior in senescent cells. *Ann N Y Acad Sci.* 2007; 1100:250–63. [PubMed: 17460187]
- Misteli T. Spatial positioning; a new dimension in genome function. *Cell.* 2004; 119:153–6. [PubMed: 15479633]
- Misteli T. Beyond the sequence: cellular organization of genome function. *Cell.* 2007; 128:787–800. [PubMed: 17320514]
- Misteli T. The cell biology of genomes: bringing the double helix to life. *Cell.* 2013; 152:1209–12. [PubMed: 23498929]
- Morey C, Da Silva NR, Perry P, Bickmore WA. Nuclear reorganisation and chromatin decondensation are conserved, but distinct, mechanisms linked to Hox gene activation. *Development.* 2007; 134:909–19. [PubMed: 17251268]
- Morey C, Kress C, Bickmore WA. Lack of bystander activation shows that localization exterior to chromosome territories is not sufficient to up-regulate gene expression. *Genome Res.* 2009; 19:1184–94. [PubMed: 19389823]
- Muller I, Boyle S, Singer RH, Bickmore WA, Chubb JR. Stable morphology, but dynamic internal reorganisation, of interphase human chromosomes in living cells. *PLoS One.* 2010; 5:e11560. [PubMed: 20644634]
- Nagano T, Lubling Y, Stevens TJ, Schoenfelder S, Yaffe E, Dean W, Laue ED, Tanay A, Fraser P. Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. *Nature.* 2013; 502:59–64. [PubMed: 24067610]
- Nagele R, Freeman T, McMorrow L, Lee HY. Precise spatial positioning of chromosomes during prometaphase: evidence for chromosomal order. *Science.* 1995; 270:1831–5. [PubMed: 8525379]

- Naumova N, Imakaev M, Fudenberg G, Zhan Y, Lajoie BR, Mirny LA, Dekker J. Organization of the mitotic chromosome. *Science*. 2013; 342:948–53. [PubMed: 24200812]
- Nemeth A, Conesa A, Santoyo-Lopez J, Medina I, Montaner D, Peterfia B, Solovei I, Cremer T, Dopazo J, Langst G. Initial genomics of the human nucleolus. *PLoS Genet*. 2010; 6:e1000889. [PubMed: 20361057]
- Nikolova V, Leimena C, McMahon AC, Tan JC, Chandar S, Jogia D, Kesteven SH, Michalick J, Otway R, Verheyen F, Rainer S, Stewart CL, Martin D, Feneley MP, Fatkin D. Defects in nuclear structure and function promote dilated cardiomyopathy in lamin A/C-deficient mice. *J Clin Invest*. 2004; 113:357–69. [PubMed: 14755333]
- Ollion J, Cochenne J, Loll F, Escude C, Boudier T. TANGO: a generic tool for high-throughput 3D image analysis for studying nuclear organization. *Bioinformatics*. 2013; 29:1840–1. [PubMed: 23681123]
- Osborne C, Wilson P, Tripathy D. Oncogenes and tumor suppressor genes in breast cancer: potential diagnostic and therapeutic applications. *Oncologist*. 2004a; 9:361–77. [PubMed: 15266090]
- Osborne CS, Chakalova L, Brown KE, Carter D, Horton A, Debrand E, Goyenechea B, Mitchell JA, Lopes S, Reik W, Fraser P. Active genes dynamically colocalize to shared sites of ongoing transcription. *Nat Genet*. 2004b; 36:1065–71. [PubMed: 15361872]
- Osborne CS, Chakalova L, Mitchell JA, Horton A, Wood AL, Bolland DJ, Corcoran AE, Fraser P. Myc dynamically and preferentially relocates to a transcription factory occupied by Igh. *PLoS Biol*. 2007; 5:e192. [PubMed: 17622196]
- Oza P, Peterson CL. Opening the DNA repair toolbox: localization of DNA double strand breaks to the nuclear periphery. *Cell Cycle*. 2010; 9:43–9. [PubMed: 20016273]
- Palstra RJ, Tolhuis B, Splinter E, Nijmeijer R, Grosveld F, de Laat W. The beta-globin nuclear compartment in development and erythroid differentiation. *Nat Genet*. 2003; 35:190–4. [PubMed: 14517543]
- Parada LA, McQueen PG, Misteli T. Tissue-specific spatial organization of genomes. *Genome Biol*. 2004; 5:R44. [PubMed: 15239829]
- Parada LA, McQueen PG, Munson PJ, Misteli T. Conservation of relative chromosome positioning in normal and cancer cells. *Curr Biol*. 2002; 12:1692–7. [PubMed: 12361574]
- Pfeffer U, Di Vinci A, Geido E, Vidali G, Giaretti W. Cell cycle dependent alterations of chromatin structure in situ as revealed by the accessibility of the nuclear protein AF-2 to monoclonal antibodies. *J Cell Physiol*. 1991; 149:567–74. [PubMed: 1744180]
- Phillips-Cremins JE, Corces VG. Chromatin insulators: linking genome organization to cellular function. *Mol Cell*. 2013; 50:461–74. [PubMed: 23706817]
- Pickersgill H, Kalverda B, de Wit E, Talhout W, Fornerod M, van Steensel B. Characterization of the *Drosophila melanogaster* genome at the nuclear lamina. *Nat Genet*. 2006; 38:1005–14. [PubMed: 16878134]
- Pliss A, Fritz AJ, Stojkovic B, Ding H, Mukherjee L, Bhattacharya S, Xu J, Berezney R. Non-random Patterns in the Distribution of NOR-bearing Chromosome Territories in Human Fibroblasts: A Network Model of Interactions. *J Cell Physiol*. 2014
- Prieto JL, McStay B. Nucleolar biogenesis: the first small steps. *Biochem Soc Trans*. 2005; 33:1441–3. [PubMed: 16246141]
- Raska I, Shaw PJ, Cmarko D. Structure and function of the nucleolus in the spotlight. *Curr Opin Cell Biol*. 2006; 18:325–34. [PubMed: 16687244]
- Raynaud C, Mallory AC, Latrasse D, Jegu T, Bruggeman Q, Delarue M, Bergounioux C, Benhamed M. Chromatin meets the cell cycle. *J Exp Bot*. 2014
- Reddy KL, Zullo JM, Bertolino E, Singh H. Transcriptional repression mediated by repositioning of genes to the nuclear lamina. *Nature*. 2008; 452:243–7. [PubMed: 18272965]
- Roix JJ, McQueen PG, Munson PJ, Parada LA, Misteli T. Spatial proximity of translocation-prone gene loci in human lymphomas. *Nat Genet*. 2003; 34:287–91. [PubMed: 12808455]
- Schardin M, Cremer T, Hager HD, Lang M. Specific staining of human chromosomes in Chinese hamster x man hybrid cell lines demonstrates interphase chromosome territories. *Hum Genet*. 1985; 71:281–7. [PubMed: 2416668]

- Schlesinger S, Selig S, Bergman Y, Cedar H. Allelic inactivation of rDNA loci. *Genes Dev.* 2009; 23:2437–47. [PubMed: 19833769]
- Sehgal N, Fritz AJ, Morris K, Torres I, Chen Z, Xu J, Berezney R. Gene density and chromosome territory shape. *Chromosoma.* 2014; 123:499–513. [PubMed: 25106753]
- Seitan VC, Faure AJ, Zhan Y, McCord RP, Lajoie BR, Ing-Simmons E, Lenhard B, Giorgetti L, Heard E, Fisher AG, Flicek P, Dekker J, Merkenschlager M. Cohesin-based chromatin interactions enable regulated gene expression within preexisting architectural compartments. *Genome Res.* 2013; 23:2066–77. [PubMed: 24002784]
- Shelby RD, Hahn KM, Sullivan KF. Dynamic elastic behavior of alpha-satellite DNA domains visualized in situ in living human cells. *J Cell Biol.* 1996; 135:545–57. [PubMed: 8909532]
- Smirnov E, Kalmarova M, Koberna K, Zemanova Z, Malinsky J, Masata M, Cvackova Z, Michalova K, Raska I. NORs and their transcription competence during the cell cycle. *Folia Biol (Praha).* 2006; 52:59–70. [PubMed: 17089916]
- Sofueva S, Yaffe E, Chan WC, Georgopoulou D, Vietri Rudan M, Mira-Bontenbal H, Pollard SM, Schroth GP, Tanay A, Hadjur S. Cohesin-mediated interactions organize chromosomal domain architecture. *EMBO J.* 2013; 32:3119–29. [PubMed: 24185899]
- Solovei I, Cremer M. 3D-FISH on cultured cells combined with immunostaining. *Methods Mol Biol.* 2010; 659:117–26. [PubMed: 20809307]
- Soutoglou E, Dorn JF, Sengupta K, Jasin M, Nussenzweig A, Ried T, Danuser G, Misteli T. Positional stability of single double-strand breaks in mammalian cells. *Nat Cell Biol.* 2007; 9:675–82. [PubMed: 17486118]
- Spilianakis CG, Lalioti MD, Town T, Lee GR, Flavell RA. Interchromosomal associations between alternatively expressed loci. *Nature.* 2005; 435:637–45. [PubMed: 15880101]
- Stack SM, Brown DB, Dewey WC. Visualization of interphase chromosomes. *J Cell Sci.* 1977; 26:281–99. [PubMed: 562895]
- Stein GZS, Braastad C, Montecino M, van Wijnen A, Choi J, Stein J, Lian J, Javed A. Functional architecture of the nucleus: organizing the regulatory machinery for gene expression, replication and repair. *TRENDS in Cell Biology.* 2003; 13:584–592. [PubMed: 14573352]
- Stein GZS, Stein J, Lian J, van Wijnen A, Montecino M, Young D, Javed A, Pratap J, Choi J, Ali S, Pande S, Hassan M. Genetic and Epigenetic Regulation in Nuclear Microenvironments for Biological Control in Cancer. *Journal of Cellular Biochemistry.* 2008; 104:2016–2026. [PubMed: 18615590]
- Sullivan GJ, Bridger JM, Cuthbert AP, Newbold RF, Bickmore WA, McStay B. Human acrocentric chromosomes with transcriptionally silent nucleolar organizer regions associate with nucleoli. *EMBO J.* 2001; 20:2867–74. [PubMed: 11387219]
- Sun HB, Shen J, Yokota H. Size-dependent positioning of human chromosomes in interphase nuclei. *Biophys J.* 2000; 79:184–90. [PubMed: 10866946]
- Taimen P, Pfliegerhaer K, Shimi T, Moller D, Ben-Harush K, Erdos MR, Adam SA, Herrmann H, Medalia O, Collins FS, Goldman AE, Goldman RD. A progeria mutation reveals functions for lamin A in nuclear assembly, architecture, and chromosome organization. *Proc Natl Acad Sci U S A.* 2009; 106:20788–93. [PubMed: 19926845]
- Takizawa T, Gudla PR, Guo L, Lockett S, Misteli T. Allele-specific nuclear positioning of the monoallelically expressed astrocyte marker GFAP. *Genes Dev.* 2008; 22:489–98. [PubMed: 18281462]
- Tanabe H, Habermann FA, Solovei I, Cremer M, Cremer T. Non-random radial arrangements of interphase chromosome territories: evolutionary considerations and functional implications. *Mutat Res.* 2002a; 504:37–45. [PubMed: 12106644]
- Tanabe H, Kupper K, Ishida T, Neusser M, Mizusawa H. Inter- and intra-specific gene-density-correlated radial chromosome territory arrangements are conserved in Old World monkeys. *Cytogenet Genome Res.* 2005; 108:255–61. [PubMed: 15545738]
- Tanabe H, Muller S, Neusser M, von Hase J, Calcagno E, Cremer M, Solovei I, Cremer C, Cremer T. Evolutionary conservation of chromosome territory arrangements in cell nuclei from higher primates. *Proc Natl Acad Sci U S A.* 2002b; 99:4424–9. [PubMed: 11930003]

- Therizols P, Illingworth RS, Courilleau C, Boyle S, Wood AJ, Bickmore WA. Chromatin decondensation is sufficient to alter nuclear organization in embryonic stem cells. *Science*. 2014; 346:1238–42. [PubMed: 25477464]
- Tolhuis B, Blom M, Kerkhoven RM, Pagie L, Teunissen H, Nieuwland M, Simonis M, de Laat W, van Lohuizen M, van Steensel B. Interactions among Polycomb domains are guided by chromosome architecture. *PLoS Genet*. 2011; 7:e1001343. [PubMed: 21455484]
- Tolhuis B, Palstra RJ, Splinter E, Grosveld F, de Laat W. Looping and interaction between hypersensitive sites in the active beta-globin locus. *Mol Cell*. 2002; 10:1453–65. [PubMed: 12504019]
- van Koningsbruggen S, Gierlinski M, Schofield P, Martin D, Barton GJ, Ariyurek Y, den Dunnen JT, Lamond AI. High-resolution whole-genome sequencing reveals that specific chromatin domains from most human chromosomes associate with nucleoli. *Mol Biol Cell*. 2010; 21:3735–48. [PubMed: 20826608]
- Versteeg R, van Schaik BD, van Batenburg MF, Roos M, Monajemi R, Caron H, Bussemaker HJ, van Kampen AH. The human transcriptome map reveals extremes in gene density, intron length, GC content, and repeat pattern for domains of highly and weakly expressed genes. *Genome Res*. 2003; 13:1998–2004. [PubMed: 12915492]
- Vietri Rudan M, Barrington C, Henderson S, Ernst C, Odom DT, Tanay A, Hadjir S. Comparative Hi-C reveals that CTCF underlies evolution of chromosomal domain architecture. *Cell Rep*. 2015; 10:1297–309. [PubMed: 25732821]
- Vignali M, Hassan AH, Neely KE, Workman JL. ATP-dependent chromatin-remodeling complexes. *Mol Cell Biol*. 2000; 20:1899–910. [PubMed: 10688638]
- Visser AE, Aten JA. Chromosomes as well as chromosomal subdomains constitute distinct units in interphase nuclei. *J Cell Sci*. 1999; 112(Pt 19):3353–60. [PubMed: 10504340]
- Volpi EV, Chevret E, Jones T, Vatcheva R, Williamson J, Beck S, Campbell RD, Goldsworthy M, Powis SH, Ragoussis J, Trowsdale J, Sheer D. Large-scale chromatin organization of the major histocompatibility complex and other regions of human chromosome 6 and its response to interferon in interphase nuclei. *J Cell Sci*. 2000; 113(Pt 9):1565–76. [PubMed: 10751148]
- Walter J, Schermelleh L, Cremer M, Tashiro S, Cremer T. Chromosome order in HeLa cells changes during mitosis and early G1, but is stably maintained during subsequent interphase stages. *J Cell Biol*. 2003; 160:685–97. [PubMed: 12604593]
- Williams RR, Broad S, Sheer D, Ragoussis J. Subchromosomal positioning of the epidermal differentiation complex (EDC) in keratinocyte and lymphoblast interphase nuclei. *Exp Cell Res*. 2002; 272:163–75. [PubMed: 11777341]
- Williamson I, Berlivet S, Eskeland R, Boyle S, Illingworth RS, Paquette D, Dostie J, Bickmore WA. Spatial genome organization: contrasting views from chromosome conformation capture and fluorescence in situ hybridization. *Genes Dev*. 2014; 28:2778–91. [PubMed: 25512564]
- Zaidi SK, Young DW, Javed A, Pratap J, Montecino M, van Wijnen A, Lian JB, Stein JL, Stein GS. Nuclear microenvironments in biological control and cancer. *Nat Rev Cancer*. 2007; 7:454–63. [PubMed: 17522714]
- Zaidi SK, Young DW, Montecino MA, Lian JB, van Wijnen AJ, Stein JL, Stein GS. Mitotic bookmarking of genes: a novel dimension to epigenetic control. *Nat Rev Genet*. 2010; 11:583–9. [PubMed: 20628351]
- Zeitzi MJ, Ay F, Heidmann JD, Lerner PL, Noble WS, Steelman BN, Hoffman AR. Genomic interaction profiles in breast cancer reveal altered chromatin architecture. *PLoS One*. 2013; 8:e73974. [PubMed: 24019942]
- Zeitzi MJ, Mukherjee L, Bhattacharya S, Xu J, Berezney R. A probabilistic model for the arrangement of a subset of human chromosome territories in WI38 human fibroblasts. *J Cell Physiol*. 2009; 221:120–9. [PubMed: 19507193]
- Zhou GL, Xin L, Song W, Di LJ, Liu G, Wu XS, Liu DP, Liang CC. Active chromatin hub of the mouse alpha-globin locus forms in a transcription factory of clustered housekeeping genes. *Mol Cell Biol*. 2006; 26:5096–105. [PubMed: 16782894]
- Zhu L, Brangwynne CP. Nuclear bodies: the emerging biophysics of nucleoplasmic phases. *Curr Opin Cell Biol*. 2015; 34:23–30. [PubMed: 25942753]

- Zink D, Cremer T. Cell nucleus: chromosome dynamics in nuclei of living cells. *Curr Biol.* 1998; 8:R321–4. [PubMed: 9560332]
- Zink D, Fischer AH, Nickerson JA. Nuclear structure in cancer cells. *Nat Rev Cancer.* 2004; 4:677–87. [PubMed: 15343274]
- Zirbel RM, Mathieu UR, Kurz A, Cremer T, Lichter P. Evidence for a nuclear compartment of transcription and splicing located at chromosome domain boundaries. *Chromosome Res.* 1993; 1:93–106. [PubMed: 8143096]
- Zorn C, Cremer C, Cremer T, Zimmer J. Unscheduled DNA synthesis after partial UV irradiation of the cell nucleus. Distribution in interphase and metaphase. *Exp Cell Res.* 1979; 124:111–9. [PubMed: 499376]
- Zorn C, Cremer T, Cremer C, Zimmer J. Laser UV microirradiation of interphase nuclei and post-treatment with caffeine. A new approach to establish the arrangement of interphase chromosomes. *Hum Genet.* 1976; 35:83–9. [PubMed: 1002167]
- Zuin J, Dixon JR, van der Reijden MI, Ye Z, Kolovos P, Brouwer RW, van de Corput MP, van de Werken HJ, Knoch TA, van IWF, Grosveld FG, Ren B, Wendt KS. Cohesin and CTCF differentially affect chromatin architecture and gene expression in human cells. *Proc Natl Acad Sci U S A.* 2014; 111:996–1001. [PubMed: 24335803]

Figure 1. A spectrum of increasingly complex chromatin conformations impacting gene expression

Whereas mitotic chromosomes are condensed into broadly recognized X-shaped entities, during interphase they relax and occupy distinct domains within the nucleus termed chromosome territories (CT). Seven CTs labeled with fluorescence in situ hybridization are shown (A). Within chromosomes, the configuration is coordinated with gene expression. Looping of regulatory elements within individual genes is essential for their expression mediated through transcription factor binding and CTCF (B). Moreover, actively expressed genes cluster in what are known as active chromatin hubs (C). Genes have also been shown to extend out of their main CT bodies in order to form interchromosomal interactions with coregulated genes (D). These interactions are mediated in part through interactions with other nuclear structures, such as the nucleoli and the nuclear lamina via lamina associated domains (LADs) or nucleolar associated domains (NADs, D-E).

Table 1

FISH and Chromatin Conformation Capture Techniques for Analyzing Chromosome Territories.

	FISH	Chromatin Conformation Capture
Throughput	<p>Slow, low throughput: microscopy and image analysis are time consuming.</p> <ul style="list-style-type: none"> FISH is ideal for analysis of individual CTs. 	<p>Hi-C is a relatively complex procedure, however, results represent the average of millions of cells and contacts across the entire genome.</p>
Process Considerations	<p>Target sequence information required. With the most advanced equipment, it is theoretically possible to use 5 fluorescently labeled probes simultaneously, however, most studies query only two or three probes at a time to avoid technical difficulties.</p>	<p>Hi-C and ChIA-PET do not require target sequence information; both techniques provide genome-wide chromosome-contact frequency data. Other chromatin conformation capture techniques (3C, 4C, 5C) do require target sequence information.</p>
	<p>Conventional FISH is in fixed nuclei (data are from individual cells at the time they were prepared for microscopy).</p> <ul style="list-style-type: none"> Stage specific markers can be used to identify cells at different points in the cell cycle. Live-cell FISH has been achieved using catalytically inactive CRISPR/cas9 labeled with EGFP and tiled sgRNAs [Chen et al., 2013]. 	<p>Crosslinking fixes molecular interactions within nuclei (Data represent the average frequency of chromosome interactions in a cell population at the time nuclei were crosslinked)</p> <ul style="list-style-type: none"> Cell populations may be heterogeneous. Values in Hi-C are relative rather than absolute. Particular cell types and/or cell-cycle stage cells can be enriched via sorting prior to analysis. Single-cell Hi-C averages interactions from both somatic CT homologs [Nagano et al., 2013].
	<p>Denaturation may alter nuclear structure.</p> <ul style="list-style-type: none"> It is important to verify that nuclear structure remains intact after processing. This can be done by visualizing colocalization of EdU before and after the FISH procedure [Fritz et al., 2014b]. 	<p>Chromatin decondensation in digestion and SDS steps could result in indirect ligation between fragments located hundreds of nanometers apart in nuclei [Gavrilov et al., 2013; Williamson et al., 2014]. Interactions in different regions of the genome may not be captured uniformly since different proteins have different binding affinities [Williamson et al., 2014].</p> <ul style="list-style-type: none"> May not be a concern in relatively low-resolution Hi-C studies.
	<p>Adequacy of hybridization efficiency is a concern.</p> <ul style="list-style-type: none"> Evaluate by labeling chromosomes in spontaneous metaphase cells using CT paints. 	<p>Repetitive sequences limit accurate mapping to chromosomes—data on interactions in these areas is typically not available.</p>
Position information	<p>Ideal for determining the position of individual CTs in the nucleus and/or relative to nuclear compartments.</p> <ul style="list-style-type: none"> Specificity of antibody to detect nuclear compartment should be determined. 	<p>Does not provide information on position relative to the nuclear periphery or other nuclear structures.</p> <ul style="list-style-type: none"> ChIA-PET can detect contacts mediated by protein components, however, genes may functionally and cooperatively associate with a third nuclear structure, without interacting at a molecular level—these contacts would not be detectable (e.g., associations localized to opposite sides of a nucleolus).
Data analysis	<p>Analysis programs measure volumes of nuclei, CT or other cellular domains; overlap/colocalization volumes; distances between CT, genes and/or domains; and the degree of irregularity in shape.</p>	<p>Requires sophisticated analysis of next generation sequencing, but provides genome-wide specific DNA interactions.</p> <ul style="list-style-type: none"> Appropriate normalization is crucial.
Resolution	<p>Conventional microscopy resolution is constrained by the diffraction limit of light.</p> <ul style="list-style-type: none"> Super resolution microscopy [Lakadamyali and Cosma, 2015; 	<p>Resolution is dependent on choice of restriction enzyme (4 or 6 bp cutter).</p>

	FISH	Chromatin Conformation Capture
	Markaki et al., 2012] can increase resolution up to 20 fold over conventional microscopy [Long et al., 2014]	<ul style="list-style-type: none">• In Hi-C, the effective resolution for cis interactions will exceed that of trans interactions with an equal number of sequencing reads. [Belton et al., 2012].• Capture Hi-C can increase the effective resolution [Barutcu et al., 2015].

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript