CASE REPORT

A rare presentation of bilateral spontaneous superficial femoral arteriovenous fistulas

Inyong Hwang,¹ Kavita Bomb,² Raza Askari,³ Rami N Khouzam⁴¹Medicine, Division of Cardiovascular Diseases, University of Tennessee Health Science Center, Memphis, Tennessee, USA²University of Tennessee Health Science Center, Memphis, Tennessee, USA³Medicine, Department of Cardiology, University of Tennessee Health Science Center, Memphis, Tennessee, USA⁴Medicine, Division of Cardiovascular Diseases, University of Tennessee Health Science Center, Memphis, Tennessee, USA

Correspondence to

Dr Rami N Khouzam, khouzamrami@yahoo.com

Accepted 18 December 2015

SUMMARY

Spontaneous arteriovenous fistulas are a rare clinical entity with very few cases reported in the literature. Prompt diagnosis and treatment is crucial and can prevent further complications in such patients. We report a case of a patient who presented with progressive bilateral claudication, discolouration of feet, hypaesthesia and non-healing ulcers. The patient had no history of trauma, aneurysm or surgery. After abnormal non-invasive studies, a peripheral angiogram revealed significant disease and obstruction of bilateral superficial femoral arteries. Spontaneous bilateral femoral arteriovenous fistulas were also found incidentally. After discussing the risks and benefits of multiple treatment options with the patient, percutaneous intervention with orbital atherectomy, balloon angioplasty and covered stent placements were performed. This treatment improved the patient's symptoms significantly.

BACKGROUND

Arteriovenous fistulas (AVFs) are anomalous direct communications between the arterial and venous systems that bypass the normal anatomic capillary beds. Penetrating trauma and iatrogenic injury following percutaneous catheterisation are the main causes of AVFs in civilian as well as in military populations.¹ Postinflammatory SFAs have also been reported.² Spontaneous AVFs are extremely rare with very few case reports available in the literature.³ Although AVFs are rare, awareness of these fistulas is clinically extremely important because they can steal the blood flow, resulting in markedly decreased circulation and ischaemia in the arterial system below the communication. When this occurs, the patient can become significantly symptomatic with potentially detrimental sequelae.

We present a case of bilateral spontaneous AVFs of the superficial femoral vessel level diagnosed during work up for peripheral arterial disease.

CASE PRESENTATION

A 71-year-old man with a history of multiple sclerosis presented after 10 years of severe claudications, progressive discolouration and hypaesthesia of bilateral lower extremities, which had worsened insidiously over the past few months. A few days before his presentation, he started developing wounds likely of an ischaemic nature on his medial malleoli. In spite of his mobility, which was relatively restricted due to his neurological deficit from multiple sclerosis, he was suffering from severe

claudications (Rutherford category 3–4) even at rest. The patient denied any trauma or surgery of the legs. The patient had a remote history of tobacco abuse. The patient did not have diabetes, hypertension or dyslipidaemia. There was no evidence of family history of coronary or peripheral vascular disease. Examination revealed bluish to purple discolouration of bilateral feet (*figure 1*), with feeble but palpable pedal pulses.

INVESTIGATIONS

Bilateral venous ultrasound of the lower extremities ruled out any evidence of venous thrombi. Glycated haemoglobin was checked and noted to be within normal limits. In view of a high suspicion for peripheral arterial disease, Ankle-Brachial Index (ABI) accompanied by Doppler ultrasound was carried out. This is considered a good, non-invasive, accurate and reliable screening tool. The results were abnormal at 0.72 for the left leg and 0.84 for the right leg.

Owing to lifestyle limiting claudication and rest pain as well as abnormal ABI, a bilateral lower extremity peripheral angiogram was performed. It showed high-grade stenosis of the distal left superficial femoral artery (SFA) along with an AVF

Figure 1 Bilateral lower extremity oedema and bluish discolouration.

CrossMark

To cite: Hwang I, Bomb K, Askari R, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-213092

Figure 2 Peripheral angiogram demonstrating superficial femoral artery stenosis and fistula before (left panel) and after intervention (right panel).

communicating between the SFA and the femoral vein ([figure 2](#) left panel/[video 1](#)).

Almost identical findings were noted on the right lower extremity. It was thought that the high-grade stenoses combined with the AVFs were contributing to the patient's symptoms, likely due to a vascular steal phenomenon.

DIFFERENTIAL DIAGNOSIS

In view of this patient's presentation, the differential diagnosis initially included peripheral vascular disease, venous disease, small vessel diabetic vascular disease with peripheral neuropathy (of note, our patient did not have diabetes) or neuropathy secondary to multiple sclerosis.

TREATMENT

In such cases, the treatment options depend on the degree of patient symptoms. Options include surgical ligation or endovascular techniques either by transvenous embolisation or stent grafting.

Video 1 Peripheral angiogram demonstrating superficial femoral artery stenosis and fistula before intervention.

Video 2 Peripheral angiogram demonstrating improvement of flow and exclusion of fistula after intervention, and covered stent placement.

After consulting with the vascular surgeon and the interventional cardiologist as a team approach, as well as reviewing the risks and benefits of each treatment option at length with the patient and his family, it was decided that endovascular therapy for the bilateral high-grade stenoses and the AVFs would be the best initial viable approach to attempt. The benefit of this approach is its effectiveness as well as its minimally invasive technique compared to vascular surgery, whereas the adverse effects can potentially include rare but possible bleeding, infection and vessel. Orbital atherectomy, percutaneous balloon angioplasty and, finally, placement of covered stents (VIABAHN), were successfully performed bilaterally in a sequential pattern. Postintervention angiography showed successful angiographic results and complete coverage of the AV fistulas ([figure 2](#) right panel/[video 2](#)). After the procedure, we evaluated the success of the treatment by re-evaluating the patient's symptoms. We found that the patient's symptoms had improved significantly.

OUTCOME AND FOLLOW-UP

The patient presented for follow-up visits 3 and 6 months after successful procedures were performed. He has noted significant improvement in claudications, as well as healing of wounds.

DISCUSSION

Spontaneous AVF of the superficial femoral artery is a rare clinical entity. In the lower extremities, AVFs are commonly iatrogenic, primarily due to percutaneous groin access. Occasionally, AVFs are reported from penetrating injuries such as stab wounds or high velocity bullet wounds. AVFs may be congenital, as in Klippel-Trenaunay syndrome, or spontaneous, as presented in our case.

In their case report, Beecher *et al*⁴ postulated persistent arterial remnants as the cause of congenital AVF of the popliteal artery. Spontaneous unilateral AVF has been reported in association with atherosclerosis,⁵ and as an arteritis-related aneurysm from an HIV patient.² Our case is a rare presentation of an incidental discovery of bilateral spontaneous AVFs without any obvious leading aetiology.

Learning points

- ▶ A spontaneously occurring arteriovenous fistula of the superficial femoral vessels is rarely seen and its pathogenesis remains an enigma. Common leading causes such as trauma, surgery or aneurysm have to be ruled out prior to such conclusion.
- ▶ Recognition of this pathology, and prompt diagnosis and treatment, are crucial in preventing further possible potential detrimental complications leading to worsening ischaemia.
- ▶ The clinical presentation of arteriovenous fistulas varies considerably based on the anatomic location, duration and nature of the trauma. There is no noted immediate risk; however, chronic fistula may lead to limb discrepancy, venous hypertension, arterial insufficiency and high output cardiac failure, with serious consequences. Early diagnosis and efficient management of such cases is therefore warranted.
- ▶ Treatment options range from non-invasive ultrasound and guided compression for asymptomatic cases to surgical ligation or endovascular techniques either by transvenous embolisation or by stent grafting for symptomatic patients.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Davidovic L, Lotina S, Vojnovic B, *et al*. Post-traumatic AV fistulas and pseudoaneurysms. *J Cardiovasc Surg* 1997;38:645–51.
- 2 Nair R, Chetty R, Woolgar J, *et al*. Spontaneous arteriovenous fistula resulting from HIV arteritis. *J Vasc Surg* 2001;33:186–7.
- 3 Willaert WI, Scheltinga MR. A case of a spontaneous popliteal arteriovenous fistula. *Acta Chirurgica Belgica* 2007;107:208–10.
- 4 Beecher S, Alawy M, Elbakar A, *et al*. Incidental discovery of a long standing arteriovenous fistula after thrombectomy for acute lower limb ischaemia. *Int J Surg Case Rep* 2014;5:1031–4.
- 5 Vallance R, Quin RO, Forrest H. Arterio-venous shunting complicating occlusive atherosclerotic peripheral vascular disease. *Clin Radiol* 1986;37:389–93.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow