CASE REPORT

Aggressive eosinophilic granulomatosis with polyangiitis and transverse sinus thrombosis

Snegha Ananth,¹ Rajeswari Sankaralingam,² Manesh Manoj³

¹Kilpauk Medical College, Chennai, Tamil Nadu, India

²Institute of Rheumatology, Madras Medical College, Chennai, Tamil Nadu, India

³Department of Internal Medicine, Kilpauk Medical College, Nilesishwar, Kerala, India

Correspondence to

Dr Snegha Ananth, sneghaananth@gmail.com

Accepted 21 December 2015

SUMMARY

We report a case of a patient with eosinophilic granulomatosis with polyangiitis (EGPA), previously known as Churg-Strauss syndrome) who succumbed to respiratory failure complicated by sepsis, pneumonitis and transverse sinus thrombosis. In the absence of antineutrophilic cytoplasmic antibodies positivity, a clinical diagnosis was made based on The American College of Rheumatology–1990 criteria. Glucocorticoids, cyclophosphamide, rituximab and intravenous immunoglobulin were administered at different stages. The patient recovered momentarily after each cycle of treatment, but his vasculitis progressed and he eventually died. Cardiac or renal involvement, which is more commonly associated with an adverse outcome, was not present.

BACKGROUND

Arterial and venous thromboses are being increasingly associated with eosinophilic granulomatosis with polyangiitis (EGPA), and their pathogenesis is yet to be clearly elucidated. Cerebral venous thrombosis is a rare occurrence in EGPA, with only one case reported so far. The potential role of eosinophils as the primary cause of thrombosis is under evaluation. Hence, apart from the routine anti-inflammatory and anticoagulation measures, the role of antieosinophilic agents in the management and prophylaxis of thrombosis is to be explored.

CASE PRESENTATION

A 17-year-old boy presented with intermittent fever and breathlessness at rest for 4 days, with neither haemoptysis nor chest pain. He had similar episodes for the past 1 year and was a known asthmatic since childhood, controlled with daily inhaled formoterol and budesonide. On inspecting the skin, there were multiple erythematous macules and nodules over the upper and lower limbs (figure 1). On examination, the patient was dyspnoeic and tachypnoeic with bilateral wheeze. His respiratory rate was 24/min with SpO₂ of 94% on 60% oxygen with a flow rate of 15 L/min through a Venturi mask. The patient had bilateral cervical, axillary, epitrochlear and inguinal lymphadenopathy. Cardiovascular, nervous system and abdomen examinations were normal. High-resolution CT (HRCT) of the chest and sinuses (figure 2) showed bilateral basal ground glass opacities and left maxillary sinusitis, respectively. Complete blood count, renal and liver function tests, electrolytes, peripheral smear and urine analysis were normal except for eosinophilia in blood (12%) and raised IgE

levels (>2500 IU/mL). ECHO and ECG were also normal. Biopsy of cervical and inguinal lymph node showed extravascular eosinophils with inflammatory cell infiltration around blood vessel (figure 3). Skin biopsy from an erythematous macular rash on the forearm revealed leucocytoclastic vasculitis (figure 4). Antineutrophilic cytoplasmic antibodies (ANCA) and antinuclear antibody (ANA) were negative. EGPA was diagnosed based on five of six American College of Rheumatology Criteria: prodromal asthma, eosinophilia, tissue infiltration of eosinophils (perivascular in lymph nodes), sinusitis and transient pulmonary infiltrates (the basal opacities resolved with steroid therapy; figure 5). The patient was started on antibiotics and oral prednisolone (1 mg/kg/day) as first-line therapy as per 2015 EGPA Consensus Task force recommendations. His symptoms subsided and he was subsequently discharged. He was advised to take prednisolone (1 mg/kg/day) daily and appear for monthly follow-up for evaluation and tapering of dose. For the next few months, he had repeated admissions for fever and breathlessness. A repeat chest X-ray showed bilateral infiltrates and chest HRCT showed bilateral diffuse reticulonodular shadows with interlobar septal thickening (figure 6). Following premedication with intravenous dexchlorpheniramine (5 mg) and paracetamol (1 g), rituximab infusion at 50 mg/h was started. Less than 15 min later, the patient developed severe difficulty in breathing with wheezing. The infusion was stopped and salbutamol nebulisation restored his breathing to baseline. No attempt at desensitisation was made due to the patient's already compromised respiratory status. Thereafter, cyclophosphamide (15 mg/kg) pulse therapy was started along with prophylaxis against *Pneumocystis jiroveci*. There was good response after two pulses

Figure 1 Vasculitic skin lesions on leg.

CrossMark

To cite: Ananth S, Sankaralingam R, Manoj M. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-213099

Figure 2 CT of the head and paranasal sinuses showing left maxillary sinusitis with inferior turbinate hypertrophy.

(2 weeks apart) but later, the patient developed acute abdominal pain and melena. Upper gastrointestinal endoscopy showed healed duodenal ulceration and contrast CT abdomen showed signs of mesenteric lymphadenopathy. There was no vomiting, no constipation and no diarrhoea; and physical examination revealed no guarding or rigidity. A third pulse of cyclophosphamide was continued and remission was achieved. Two days later, the patient developed an episode of generalised tonic clonic seizure with no focal neurological deficits. CT of the brain showed transverse sinus thrombosis (figure 7). Intravenous immunoglobulin was initiated due to the rapid deterioration and continued dyspnoea. The patient then developed bilateral lower limb weakness with decreased sensation. Nerve conduction study revealed axonal sensorimotor neuropathy.

Two days later, while attempting to intubate him, the patient went into sudden cardiac arrest as his saturation was rapidly declining ($\text{SpO}_2 \leq 88\%$) and pCO_2 (65 mm Hg) was rising. He died despite maximum resuscitative efforts.

INVESTIGATIONS

Laboratory findings showed a slightly raised white cell count of $11.0 \times 10^9/\text{L}$, an elevated absolute eosinophil count of $6.8 \times 10^9/\text{L}$, elevated immunoglobulin E (IgE $>2500 \text{ IU/mL}$) and erythrocyte sedimentation rate of 60 mm/h. The patient had negative cytoplasmic-ANCA, perinuclear-ANCA, ANA, antidouble stranded DNA, anticyclic citrullinated peptide, rheumatoid factor, lupus anticoagulant and anticardiolipin antibodies. Antihepatitis C virus, hepatitis B surface antigen and anti-HIV were also negative.

Figure 3 Cervical lymph node biopsy depicting abundant eosinophils with neutrophilic infiltration around the blood vessel.

Direct and indirect Coombs tests were negative. Bone marrow aspiration cytology was normal. Endoscopic paranasal sinus biopsy showed diffuse inflammatory infiltrates, predominantly of eosinophils. Flow cytometry analysis showed natural killer cell deficiency.

Sputum and blood samples revealed no organisms in culture. Epstein-Barr virus, Brucella and TORCH (toxoplasmosis, rubella, cytomegalovirus, herpes) complex panel were all negative. Cardiac echo (ejection fraction: 65%) and ECG were normal. Mantoux test was negative and sputum smear examination showed no acid-fast bacilli.

DIFFERENTIAL DIAGNOSIS

At the patient's initial presentation, vasculitic syndromes such as Kawasaki's disease, Kikuchi's necrotising lymphadenitis and idiopathic hypereosinophilic syndrome were suspected in view of the lymphadenopathy. A bone marrow biopsy was unremarkable and did not show signs of haematological malignancy. The history of asthma and eosinophilia made EGPA a strong possibility despite absence of ANCA. The second CT of the chest revealed bilateral diffuse reticular nodular shadows, suggesting an interstitial lung disease pattern, which is more common in granulomatosis with polyangiitis (Wegener's granulomatosis) rather than in EGPA, which is usually associated with consolidatory findings and ground glass opacities.¹ Opportunistic infections such as miliary tuberculosis, *P. jiroveci* and allergic broncho pulmonary aspergillosis were also suspected due to

Figure 4 Skin biopsy shows extensive neutrophilic infiltration around a blood vessel.

immunosuppression, and necessary investigations were carried out.

TREATMENT

Prednisolone 50 mg/day was initiated followed by cyclophosphamide (15 mg/kg) pulse therapy, after an adverse event with rituximab. The pulse therapy was administered at 2-week intervals along with trimethoprim/sulfamethoxazole (800/160 mg on alternate days). There was good response at 3 months. Intravenous immunoglobulin (12 g/day for 5 days) was initiated following neurological complications. The patient was also covered under broad spectrum antibiotics and antituberculous treatment according to national guidelines. Low-molecular-weight heparin (enoxaparin) was given at 1.5 mg/kg subcutaneously (6th hourly) along with warfarin after the development of transverse sinus thrombosis. Thromboprophylaxis was not considered as there were no risk factors and because EGPA is a rare cause of thrombosis.

DISCUSSION

During the past decade, there have been increased descriptions of EGPA characterised by vascular occlusions such as peripheral venous thrombosis, mesenteric thrombosis, retinal vein thrombosis and coronary artery thrombosis, but only one case of cerebral venous thrombosis as an onset of EGPA has been described so far.² The prevalence of arterial occlusion has reportedly been between 3.1% and 18.7%, and venous occlusion between 5.8% and 30%, whereas a specific survey for venous thromboembolism in EGPA yielded a prevalence of 8.1%.³ Elevated plasma levels of prothrombin fragment 1+2 and D-dimer during active disease indicating a prothrombotic state has also been reported.⁴ However, recent studies revealed the potential role of eosinophils in thrombosis.⁵ The role of an antieosinophilic drug (mepolizumab) in the treatment of thrombosis along with routine anticoagulation and immunosuppression has proven to be beneficial in a case of hypereosinophilia syndrome with arterial thrombosis.⁶

Many cases of arterial and venous thrombosis have also been linked to the presence of antiphospholipid antibodies (APA) in EGPA, considering a possible overlap syndrome.⁷ Our patient tested negative for lupus anticoagulant and anticardiolipin antibody.

ANCA negativity is associated with an increased proportion of cardiac involvement, gastrointestinal involvement, pulmonary infiltrates and the outcome of a life-threatening event or death, when compared with ANCA-positive cases. ANCA negativity has been associated with a decreased proportion of peripheral neuropathy. These findings support the hypothesis that the presence or absence of autoantibodies influences the clinical expression and severity.⁸

Oka *et al* classified EGPA into two subtypes based on ANCA and neuropathy. Peripheral neuropathy, though less frequently found in ANCA-negative subtypes, was present in our patient, and the role of eosinophils in this subtype is evidenced by the large number of eosinophilic infiltrations in the epineurium against the classical necrotising vasculitis seen in ANCA-positive patients. Therapy targeting activated eosinophils may be a possible treatment for intractable neuropathy.⁹

Based on the Factor Five Score (FFS) by the French Vasculitis Study Group, cyclophosphamide in addition to glucocorticoids had a significant survival benefit over glucocorticoids alone in a retrospective analysis of high-risk patients (FFS $\geq$ 1).¹⁰ Despite the overall survival benefit of baseline FFS=0, 45% of the patients required an immunosuppressant in addition to

Figure 5 Contrast CT of the chest with no opacities after initiation of steroids.

steroids.¹¹ Before the occurrence of transverse sinus thrombosis, our patient scored FFS=0, yet cyclophosphamide pulse therapy was initiated due to his unresolving dyspnoea. Thus the choice and combination of agents must be personalised due to the varied chronology of presentations.

A meta-analysis of three randomised controlled trials concluded that pulsed cyclophosphamide, more so than continuous oral therapy, was likely to result in remission, with a lower risk of side effects. However, pulsed therapy may be associated with a higher risk of relapse, as occurred in our patient.¹²

Figure 6 Contrast CT of the chest revealing a bilateral diffuse reticular pattern predominant in the posterobasal segments.

Figure 7 Plain CT of the brain showing right transverse sinus thrombosis.

Rituximab has been reported to cause bronchospasm in two patients with EGPA, similar to our patient's experience.¹³ A possible mechanism could be the immediate release of proinflammatory cytokines.¹⁴ Though desensitisation was successful in many cases where rituximab was used in lymphoproliferative malignancies, it has to be carried out with caution in patients with EGPA, as they are more susceptible due to their frequent allergic manifestations.

Patient's father's perspective

Patient's father: It is very unfortunate for me and my family to lose my son to this rare disease at this very young age. We just thought it was seasonal asthma in his childhood and never imagined that it would one day progress and take him away from us. From his 6th birthday on, he was constantly taken to hospitals for asthma exacerbations and only two years back, when he had severe fever along with swellings and rashes all over his body, did we come to know that he had this rare disease. Our son has suffered a lot in these two years. He couldn't attend his school and always remained very thin and weak. In the last one year, he spent most of his time in the hospital, rather than anywhere else. We are uneducated and couldn't understand the complexity of this disease. All our family members are healthy and it makes us clueless on why our son should suffer from this disease. Doctors said his own immune system was attacking him and we thought maybe that's why he was not gaining weight and always falling sick. The powerful medications made him weak, but we all hoped he would return to normal. Dr Rajeswari took great efforts for our son and has performed all of this free of cost through government insurance, relieving the huge burden on our poor family. I now ask my relatives to be careful whenever their children are diagnosed of asthma and fever. I miss my son.

Learning points

- ▶ Severe hypereosinophilia with elevated IgE should be monitored for hypercoagulopathy and the role of prophylactic anticoagulation should be explored.
- ▶ Monitor every patient with suitable prognostic scales and initiate aggressive treatment for unresolving symptoms irrespective of predictive scores.
- ▶ Be cautious while administering rituximab to patients with eosinophilic granulomatosis with polyangiitis.

Acknowledgements The author expresses her sincere condolences to the patient's family who, despite their grief, patiently cooperated with the procedures and gave their consent for this publication. Thanks to Dr Shriksha, Rheumatologist, Apollo Hospitals, Chennai, for her critical view of this manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Szczeklik W, Sokolowska B, Mastalerz L, et al. Pulmonary findings in Churg–Strauss syndrome in chest X-rays and high resolution computed tomography at the time of initial diagnosis. *Clin Rheumatol* 2010;29:1127–34.
- 2 Teresa Sartori M, Briani C, Munari M, et al. Cerebral venous thrombosis as a rare onset of Churg–Strauss syndrome. *Thromb Haemost* 2006;96:90–2.
- 3 Ames PR, Margaglione M, Mackie S, et al. Eosinophilia and thrombophilia in Churg–Strauss syndrome: a clinical and pathogenetic overview. *Clin Appl Thromb Hemost* 2010;16:628–36.
- 4 Marzano AV, Tedeschi A, Rossio R, et al. Prothrombotic state in Churg–Strauss syndrome: a case report. *J Investig Allergol Clin Immunol* 2010;20:616–19.
- 5 Khoury P, Grayson PC, Klion AD. Eosinophils in vasculitis: characteristics and roles in pathogenesis. *Nat Rev Rheumatol* 2014;10:474–83.
- 6 Leon-Ferre RA, Weiler CR, Halfdanarson TR. Hypereosinophilic syndrome presenting as an unusual triad of eosinophilia, severe thrombocytopenia, and diffuse arterial thromboses, with good response to mepolizumab. *Clin Advan Hematol Oncol* 2013;11:317–19.
- 7 Paroli M, Polidoro A, Romano S, et al. Churg–Strauss syndrome associated with antiphospholipid antibodies in a patient with recurrent myocardial and cerebral ischemia. *Int Med Case Rep J* 2012;5:79–81.

- 8 Healy B, Bibby S, Steele R, *et al.* Antineutrophil cytoplasmic autoantibodies and myeloperoxidase autoantibodies in clinical expression of Churg-Strauss syndrome. *J Allergy Clin Immunol* 2013;131:571–6.
- 9 Oka N, Kawasaki T, Matsui M, *et al.* Two subtypes of Churg-Strauss syndrome with neuropathy: the roles of eosinophils and ANCA. *Mod Rheumatol* 2011;21: 290–5.
- 10 Pagnoux C. Churg-Strauss syndrome: evolving concepts. *Discov Med* 2010;9:243–52.
- 11 Samson M, Puéchal X, Devilliers H, *et al.* Eosinophilic granulomatosis with polyangiitis (Churg–Strauss)(EGPA), polyarteritis nodosa (PAN) and microscopic polyangiitis (MPA) with no initial Five-Factor Score-defined poor-prognosis factors (FFS= 0): Baseline factors associated with cytotoxic agent and immunomodulator prescription. *La Presse Médicale* 2013;42(4P2):666.
- 12 De Groot K, Adu D, Savage CO, EUVAS (European Vasculitis Study Group). The value of pulse cyclophosphamide in ANCA-associated vasculitis: meta-analysis and critical review. *Nephrol Dial Transplant* 2001;16:2018–27.
- 13 Bouldouyre MA, Cohen P, Guillemin L. Severe bronchospasm associated with rituximab for refractory Churg–Strauss syndrome. *Ann Rheum Dis* 2009;68:606.
- 14 Kulkarni HS, Kasi PM. Rituximab and cytokine release syndrome. *Case Rep Oncol* 2012;5:134–40.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow