Published in final edited form as:

Curr Opin Pharmacol. 2016 February ; 26: 96–101. doi:10.1016/j.coph.2015.10.009.

Neuro-immune Dysfunction During Brain Aging: New Insights in Microglial Cell Regulation

Stephanie M. Matt¹ and Rodney W. Johnson^{1,2,3}

¹Neuroscience Program, University of Illinois Urbana-Champaign

²Department of Animal Sciences, University of Illinois Urbana-Champaign

³Integrative Immunology and Behavior Program, University of Illinois Urbana-Champaign

Abstract

Microglia, the resident immune cells of the brain, are at the center of communication between the central nervous system and immune system. While these brain-immune interactions are balanced in healthy adulthood, the ability to maintain homeostasis during aging is impaired. Microglia develop a loss of integrated regulatory networks including aberrant signaling from other brain cells, immune sensors, and epigenetic modifiers. The low-grade chronic neuroinflammation associated with this dysfunctional activity likely contributes to cognitive deficits and susceptibility to age-related pathologies. A better understanding of the underlying mechanisms responsible for neuro-immune dysregulation with age is crucial for providing targeted therapeutic strategies to support brain repair and healthy aging.

Introduction

The central nervous system (CNS) and immune system are intimately coupled, and maintain homeostasis through sophisticated bidirectional crosstalk [1]. However, the aged brain's ability to preserve the balance between the protective and possibly detrimental effects of the innate immune system is profoundly impaired. The low-grade chronic neuroinflammation that accompanies normal aging likely contributes to cognitive deficits and susceptibility to a plethora of age-related pathologies such as autoimmune diseases, cancer, and neurodegenerative diseases. As microglia are the brain's innate immune effector cells, dysregulation of microglial behavior appears to be a critical component of this progression [2]. This review discusses several recent developments in the understanding of neuroimmune dysregulation with age by focusing on regulatory mechanisms that normally constrain microglia but become dysfunctional with old age. Understanding the underlying mechanisms responsible for the age-related increase in brain inflammation is critical for improving the likelihood for successful aging.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors declare no conflict of interest.

Microglia at the Nexus of Immune-Brain Communication

The brain was once thought to be an immune-privileged organ, devoid of lymphatic vasculature and considered pathological with immune cell entry. Now it is clear that interactions between the immune system and the brain are integral in eliciting behavioral, endocrine, and physiological modifications in the presence of peripheral inflammation. A key player in this interaction is the microglial cell, as this resident immune effector cell in the brain is responsive to signals emerging from the periphery [3,4]. The peripheral immune system is able to stimulate microglial cell activity through at least four major communication pathways: (1) diffusion of cytokines through leaky regions of the blood-brain barrier (e.g. choroid plexus and circumventricular organs), (2) passage of cytokines through energy-dependent, saturable transport systems on brain endothelium, (3) activation of cytokine receptors on endothelial cells that subsequently release cytokines or second messengers, and (4) transmission of cytokine signals via afferent nerve fibers, such as the vagus nerve. Further, with the discovery of functional lymphatic vessels lining the dural sinuses [5,6], there is potential for yet another way in which brain-resident immune cells could communicate with peripheral immune cells. The lymphatic vessels have molecular characteristics of lymphatic endothelial cells, and act as direct clearance routes for the brain and cerebrospinal fluid (CSF) into the deep cervical lymph nodes. These lymphatic vessels also absorb CSF from the adjacent subarachnoid space and brain interstitial fluid via the recently discovered glymphatic system [7]. Irrespective of the communication pathway employed, a common outcome appears to be stimulation of microglial cells and induction of a neuroinflammatory response that elicits behavioral, endocrine, and physiological responses [8]. The large body of work in this area established beyond doubt the relevance of neuroimmune interactions and highlighted the fundamental role of microglia in neuroinflammation. The new findings concerning the glymphatic and lymphatic systems may form the basis for innovative therapeutic strategies to regulate microglia, for example boosting growth of meningeal lymphatic vessels to help clear debris that becomes harder for microglia to phagocytize with age.

Microglia were first thought to be similar to macrophages, but it is now apparent that these cell types differ in ontogeny and functional capacity [9]. Unlike bone marrow-derived macrophages, microglia are derived from yolk sac progenitors and migrate to the CNS at approximately embryonic day 8.5 in mice [10]. Fate-mapping studies have also established that the transcription factor Myb, required for hematopoiesis, is not necessary for the generation of microglia [11], and that microglia specifically originate from CD45-c-kit+ erythromyeloid progenitor cells [12]. Therefore, microglia are not hematopoietically derived in the same way as macrophages. An additional unique feature is that in the adult brain, the microglial population is maintained by local self-renewal without significant contribution from bone marrow derived progenitors. Specifically, colony-stimulating factor 1 receptor (CSF-1R), but not colony-stimulating factor 1, is necessary for regulating the microglial lineage, because inhibition of CSF-1R in mice leads to depletion of 99% of the microglial population [13].

In addition to their unique ontological properties, microglia perform a number of vital functions in maintaining brain homeostasis (Fig. 1). During development, microglia regulate

neurons in at least two important ways. First, microglia act as “gate keepers” to monitor the number of neural precursor cells in the developing cerebral cortex. Cunningham et al. demonstrated that microglia selectively colonize the cortical proliferative zones in the developing neocortex, and phagocytize neural precursor cells in late stages of cortical neurogenesis [14]. By activating microglia with LPS or eliminating them with liposome clodronate, the neural precursor numbers were either reduced or increased, respectively, thus identifying a crucial role for microglia in controlling the size of the precursor population pool. Second, precursor cells that differentiate into neurons often make far more synapses than needed, and many become eliminated by microglia in a process called synaptic pruning. The complement system has been suggested to play a major role in mediating microglia-synapse interactions, and one study revealed that CR3/C3 signaling is specifically involved, as CR3 and C3 deficient mice have dramatic pruning defects [15].

In adulthood microglia are said to be “resting” or “quiescent”, but they are in fact active surveyors of the CNS parenchyma, continually scanning the surrounding extracellular space to communicate with other brain cells to maintain tissue homeostasis [16]. Using in vivo two-photon microscopy of fluorescent-labeled microglia, it was established that processes of microglia make contacts with neuronal synapses for a brief duration of about 5 minutes, occurring about once every hour [17]. Moreover, after cerebral ischemia, the duration of these microglia-synapse contacts are prolonged for over an hour, and frequently followed by synapse elimination, suggesting that microglia can restore synapse function or initiate future removal.

Lastly, microglia play a fundamental role in clearing amyloid- β (A β) aggregates, which is thought to regulate the development and progression of Alzheimer’s disease [18]. It is known that A β deposits have potent chemoattractant activity on microglia, but direct phagocytosis of A β by microglia has not been demonstrated in vivo. Although outside the scope of this review, it is important to highlight microglia as a key player in eliminating harmful debris within the brain parenchyma. As evidenced later, the aging of microglia progressively diminishes their normal regulatory capacity, potentially leading to insufficient phagocytic activity along with a host of other complications.

Microglial Dysfunction with Age

Despite the dynamic role of microglia in maintaining homeostasis, their long-lived nature and general inability to be replaced by circulating peripheral cells makes them particularly sensitive to oxidative stress, DNA damage, and a lifetime of inflammatory insults. Over time, these cells may become hypersensitive or primed in the healthy aged brain (Fig. 1) [19]. The traditional view holds that microglia in the aged (but otherwise healthy) brain are primed towards a pro-inflammatory M1 phenotype with diminished anti-inflammatory (M2) properties. Approximately 25% of microglia isolated from aged mice stain positive for major histocompatibility complex class II compared to less than 3% of microglia isolated from young adults [20]. Furthermore, aging reduces expression of IL-4 receptor by microglia, rendering them less responsive to the M2-promoting effects of IL-4 [21]. This results in exaggerated immune responsiveness typified by increased pro-inflammatory cytokine production and signaling including upregulation of interleukin (IL)-1 β , IL-6, and

tumor necrosis factor (TNF)- α both in whole brain and specifically in microglia [22,23]. Accordingly, when the immune system is challenged by a pro-inflammatory agent such as lipopolysaccharide (LPS), microglial activation is amplified and prolonged in the aged brain compared with adults, leading to exaggerated neuroinflammation, sickness behavior and cognitive deficits (Fig. 1) [20,24]. This age-related microglial activity can also be brain-region specific [25,26]. For example, release of pro-inflammatory cytokines from microglia impair hippocampal-dependent memory by way of blocking long-term potentiation [25], and can activate NF- κ B signaling in hypothalamic neurons to hinder neurogenesis and accelerate cognitive decline [26].

Given the complexity of microglial behavior, classifying them as only “damaging” M1- and “healing” M2-phenotypes fails to recognize the subtle nuances within the spectrum of phenotypes and activation patterns necessary to understand neuroinflammation in various environmental conditions [27]. An increasing number of transcriptomic studies using primary microglia have revealed gene signatures extending far beyond upregulation of pro-inflammatory genes with age [28-30]. Notably, Hickman et al. identified 100 genes specifically enriched in microglia that were collectively named the microglial sensome, thus creating a new way to discriminate between other CNS cells and immune cells [28]. Surprisingly, the most upregulated genes in aged microglia belonged to neuroprotective pathways, whereas genes in neurotoxic pathways were downregulated. Therefore, studies of highly purified microglia taking into account the transcriptional profile instead of a limited number of markers are rapidly changing the way microglia are viewed [31].

Mechanisms that Mediate Microglial Inflammation in Aging

Within the aged brain, control of immune-sensing and regulatory pathways begins to break down, producing microglial priming and heightened responsiveness to inflammatory stimuli. Considerable findings have suggested an instrumental role of immune sensors such as inflammasomes [32], as they are necessary for activating caspases, which then cleave the precursor forms of pro-inflammatory cytokines into their active forms. Of note, it was found that aged nucleotide-binding domain, leucine-rich-containing family, pyrin domain-containing-3 (NLRP3)^{-/-} mice were protected from age-related increases in astrogliosis, microglia activation, elevation of IL-1 β and TNF- α , and displayed cognitive improvement [33]. In addition, aging-associated increases in interferon and complement pathways were reduced in mice deficient in NLRP3 or its associated inflammasome adapter proteins.

To maintain control of microglial activation, neurons produce several immunomodulatory factors that interact with microglial receptors, including the most well known ligand-receptor pairs CX3CL1-CX3CR1 and CD200-CD200r. Aging is associated with reduced signaling of both of these pairs [34,35], and mice deficient in either CD200 or CX3CR1 exhibit an activated microglial phenotype [36,37]. In addition, triggering receptor expressed on myeloid cells 2 (TREM2) is a microglial surface receptor that belongs to the immunoglobulin superfamily. TREM2 is reduced in aged microglia [28], and is suspected to be required for removal of damaged myelin and support myelin regeneration by oligodendrocytes. TREM2 deficiency produced fewer microglia with an abnormal morphology in aged mice, particularly in the corpus callosum, which is rich in myelin and

may provide a major stimulus for TREM2 signaling [38]. Transforming growth factor β 1 (TGF- β 1), a protein that has protective properties in the brain, was demonstrated to be critical for microglia to populate the brain [39]. By transcriptional profiling of purified microglia, Butovsky et al. uncovered a unique gene signature present in both mouse and human primary microglia, but absent in cultured microglia, recruited monocytes, and other brain cells. This signature reemerged in cultured microglia after addition of macrophage colony-stimulating factor and TGF- β 1. To demonstrate an in vivo role for TGF- β 1, mice deficient for TGF- β 1 in the CNS had microglial loss, late onset motor dysfunction, and defects related to glutamate recycling and synaptic plasticity [39,40]. Additionally, inhibition of TGF β signaling in the brain produced exaggerated microglial activation and sickness behavior after LPS [41].

Unlike most other tissue macrophages, adult microglia express high levels of the fractalkine receptor CX₃CR1. Several recent studies have utilized unique mouse models that for the first time allow for specific genetic targeting of microglia in the adult brain, particularly the generation of *Cx3Cr1^{creER}* mice [42-44]. For example, the specific deletion of transforming growth factor- β -activated kinase 1 in microglia reduced disease course of experimental autoimmune encephalomyelitis [43], and conditional removal of brain-derived neurotrophic factor from microglia resulted in deficits in motor-learning tasks and learning-dependent synapse formation [44]. A method for generation of microglial cells derived from embryonic stem (ESdM) cells has also been described [45]. ESdM develop in a mixed neural culture through a yolk sac-like primitive macrophage intermediate, which mirrors the in vivo situation of microglial development. When these cells were genetically modified to express the neuroprotective neurotrophin 3, mice recovered from EAE. This suggests that microglia could be used as delivery vehicles or even as therapeutic targets themselves.

Epigenetics and Microglial Regulation

The cellular and physiological phenotypic trait variation evident in aged microglial cells is likely related to epigenetic changes such as DNA methylation and histone modifications. To date, only a handful of studies have used high-throughput sequencing to address the epigenetic landscape of primary microglia [46,47], and there is yet to be a study comparing young to old. Fortunately, a number of recent studies have investigated specific epigenetic regulators in microglia. Sirtuin 1 (SIRT1) is a member of class III histone deacetylases and is known to play a role in longevity [48]. SIRT1 deficiency in aged microglia is related to increases of IL-1 β transcription and decreased methylation of CpG sites within the IL-1 β proximal promoter [49]. As epigenetic regulation of other pro-inflammatory genes such as TNF- α and IL-6 has been demonstrated in other immune cell types [50,51], it will be pertinent to explore this possibility in microglia. It has also been observed that the suppressing histone H3K27me3 demethylase Jumonji domain containing 3 (Jmjd3) in the substantia nigra in vivo caused exaggerated microglial activation and provoked greater dopamine neuron death in a mouse model of Parkinson's disease [52]. Moreover, the Jmjd3 level was lower in the midbrain of aged mice, which was accompanied by an elevated level of H3K27me3.

Micro-RNAs (mi-RNAs) are small non-coding mRNAs that can negatively control their target gene expression post-translationally, and have been suggested to participate in epigenetic machinery [53]. Importantly, an increasing number of miRNAs are being associated with specific microglial phenotypes. Of note, mirR-155 is the most upregulated miRNA in primary microglia stimulated with LPS, and an emerging hypothesis is that miR-155 is required to skew microglia to a more pro-inflammatory M1-like phenotype [54]. Another report also indicates that insulin-like growth factor 1 and CX3CL1 are reduced in aged microglia and associated with increased levels of miR-29b [55]. Lastly, dietary factors such as resveratrol, luteolin, and curcumin regulate microglia [56], and have been shown to modify the epigenome [57].

Coda

The pathways shared between the CNS and peripheral immune system represent therapeutic targets in preventing neuro-immune dysregulation in aging. As microglia are at the center of communication between these two systems, uncovering ways to protect the balance between protective and detrimental effects of these cells is a high priority. Aging results in the loss of integrated regulatory networks needed to maintain appropriate microglial function, including aberrant signaling from other brain cells, immune sensors such as inflammasomes, and epigenetic modifiers such as histone deacetylases and miRNAs. Pharmacological manipulation of these regulators could direct microglia towards conditions that support brain repair and tissue homeostasis, through subtle adjustment of their continuum of phenotypes.

Acknowledgements

This work was supported by NIH R01 AG16710 and HD069899.

References

1. Jurgens HA, Johnson RW. Dysregulated neuronal-microglial cross-talk during aging, stress and inflammation. *Exp Neurol*. 2012; 233:40–48. [PubMed: 21110971]
2. Norden DM, Godbout JP. Review: microglia of the aged brain: primed to be activated and resistant to regulation. *Neuropathol Appl Neurobiol*. 2013; 39:19–34. [PubMed: 23039106]
3. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci*. 2008; 9:46–56. [PubMed: 18073775]
4. Quan N, Banks WA. Brain-immune communication pathways. *Brain Behav Immun*. 2007; 21:727–735. [PubMed: 17604598]
5. Louveau A, Smirnov I, Keyes TJ, Eccles JD, Rouhani SJ, Peske JD, Derecki NC, Castle D, Mandell JW, Lee KS, et al. Structural and functional features of central nervous system lymphatic vessels. *Nature*. 2015; 523:337–341. [PubMed: 26030524] [** This group was the first to discover lymphatic vasculature within the brain, revealing a novel pathway in which the brain can interact with the immune system.]
6. Aspelund A, Antila S, Proulx ST, Karlsson TV, Karaman S, Detmar M, Wiig H, Alitalo K. A dural lymphatic vascular system that drains brain interstitial fluid and macromolecules. *J Exp Med*. 2015; 212:991–999. [PubMed: 26077718]
7. Iliff JJ, Wang M, Liao Y, Plog BA, Peng W, Gundersen GA, Benveniste H, Vates GE, Deane R, Goldman SA, et al. A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta. *Sci Transl Med*. 2012; 4:147ra111.

8. Dilger RN, Johnson RW. Aging, microglial cell priming, and the discordant central inflammatory response to signals from the peripheral immune system. *J Leukoc Biol.* 2008; 84:932–939. [PubMed: 18495785]
9. Nayak D, Roth TL, McGavern DB. Microglia development and function. *Annu Rev Immunol.* 2014; 32:367–402. [PubMed: 24471431]
10. Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, Mehler MF, Conway SJ, Ng LG, Stanley ER, et al. Fate Mapping Analysis Reveals That Adult Microglia Derive from Primitive Macrophages. *Science.* 2010; 330:841–845. [PubMed: 20966214]
11. Schulz C, Gomez Perdiguero E, Chorro L, Szabo-Rogers H, Cagnard N, Kierdorf K, Prinz M, Wu B, Jacobsen SE, Pollard JW, et al. A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science.* 2012; 336:86–90. [PubMed: 22442384]
12. Kierdorf K, Erny D, Goldmann T, Sander V, Schulz C, Perdiguero EG, Wieghofer P, Heinrich A, Riemke P, Holscher C, et al. Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-dependent pathways. *Nat Neurosci.* 2013; 16:273–280. [PubMed: 23334579]
13. Elmore MR, Najafi AR, Koike MA, Dagher NN, Spangenberg EE, Rice RA, Kitazawa M, Matusow B, Nguyen H, West BL, et al. Colony-stimulating factor 1 receptor signaling is necessary for microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron.* 2014; 82:380–397. [PubMed: 24742461] [* This study demonstrated the crucial role of CSF1R in regulating the adult microglial population. Inhibition of CSF1R signaling depleted 99% of microglia from the brain, and they further found that microglia remarkably repopulate the CNS within 7 days. These findings identify a novel mechanism for maintaining microglial homeostasis.]
14. Cunningham CL, Martínez-Cerdeño V, Noctor SC. Microglia regulate the number of neural precursor cells in the developing cerebral cortex. *J Neurosci.* 2013; 33:4216–4233. [PubMed: 23467340] [* This study highlights the importance of microglia in regulating the number of precursor cells by phagocytosis during cortical development. Through pharmacological manipulation, deactivating or activating microglia increased or decreased the size of the neural precursor pool, respectively.]
15. Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, Ransohoff RM, Greenberg ME, Barres BA, Stevens B. Microglia Sculpt Postnatal Neural Circuits in an Activity and Complement-Dependent Manner. *Neuron.* 2012; 74:691–705. [PubMed: 22632727]
16. Nimmerjahn A, Kirchhoff F, Helmchen F. Resting Microglial Cells Are Highly Dynamic Surveillants of Brain Parenchyma in Vivo. *Science.* 2005; 308:1314–1318. [PubMed: 15831717]
17. Wake H, Moorhouse AJ, Jinno S, Kohsaka S, Nabekura J. Resting Microglia Directly Monitor the Functional State of Synapses In Vivo and Determine the Fate of Ischemic Terminals. *J Neurosci.* 2009; 29:3974–3980. [PubMed: 19339593]
18. Thériault P, ElAli A, Rivest S. The dynamics of monocytes and microglia in Alzheimer's disease. *Alzheimers Res Ther.* 2015; 7:41. [PubMed: 25878730]
19. Sparkman NL, Johnson RW. Neuroinflammation associated with aging sensitizes the brain to the effects of infection or stress. *Neuroimmunomodulation.* 2008; 15:323–330. [PubMed: 19047808]
20. Henry CJ, Huang Y, Wynne AM, Godbout JP. Peripheral lipopolysaccharide (LPS) challenge promotes microglial hyperactivity in aged mice that is associated with exaggerated induction of both pro-inflammatory IL-1 β and anti-inflammatory IL-10 cytokines. *Brain Behav Immun.* 2009; 23:309–317. [PubMed: 18814846]
21. Fenn AM, Henry CJ, Huang Y, Dugan A, Godbout JP. Lipopolysaccharide-induced interleukin (IL)-4 receptor- α expression and corresponding sensitivity to the M2 promoting effects of IL-4 are impaired in microglia of aged mice. *Brain Behav Immun.* 2012; 26:766–777. [PubMed: 22024136]
22. Sierra A, Gottfried-Blackmore AC, McEwen BS, Bulloch K. Microglia derived from aging mice exhibit an altered inflammatory profile. *Glia.* 2007; 55:412–424. [PubMed: 17203473]
23. Ye SM, Johnson RW. Regulation of interleukin-6 gene expression in brain of aged mice by nuclear factor kappaB. *J Neuroimmunol.* 2001; 117:87–96. [PubMed: 11431008]
24. Godbout JP, Chen J, Abraham J, Richwine AF, Berg BM, Kelley KW, Johnson RW. Exaggerated neuroinflammation and sickness behavior in aged mice after activation of the peripheral innate immune system. *The FASEB Journal.* 2005

25. Ojo JO, Rezaie P, Gabbott PL, Stewart MG. Impact of age-related neuroglial cell responses on hippocampal deterioration. *Front Aging Neurosci.* 2015; 7:57. [PubMed: 25972808]
26. Tang Y, Purkayastha S, Cai D. Hypothalamic microinflammation: a common basis of metabolic syndrome and aging. *Trends Neurosci.* 2015; 38:36–44. [PubMed: 25458920]
27. Prinz M, Priller J. Microglia and brain macrophages in the molecular age: from origin to neuropsychiatric disease. *Nat Rev Neurosci.* 2014; 15:300–312. [PubMed: 24713688]
28. Hickman SE, Kingery ND, Ohsumi TK, Borowsky ML, Wang LC, Means TK, El Khoury J. The microglial sensome revealed by direct RNA sequencing. *Nat Neurosci.* 2013; 16:1896–1905. [PubMed: 24162652] [** This group identified a novel phenotypic signature of microglia characterized by transcripts encoding proteins for sensing endogenous ligands and microbes. These “sensome” genes are downregulated with increasing age, and provide a new way to differentiate microglia from a range of activation states as well as other immune cells.]
29. Orre M, Kamphuis W, Osborn LM, Melief J, Kooijman L, Huitinga I, Klooster J, Bossers K, Hol EM. Acute isolation and transcriptome characterization of cortical astrocytes and microglia from young and aged mice. *Neurobiol Aging.* 2014; 35:1–14. [PubMed: 23954174]
30. Ma W, Cojocaru R, Gotoh N, Gieser L, Villasmil R, Cogliati T, Swaroop A, Wong WT. Gene expression changes in aging retinal microglia: relationship to microglial support functions and regulation of activation. *Neurobiol Aging.* 2013; 34:2310–2321. [PubMed: 23608111]
31. Wes PD, Holtman IR, Boddeke EW, Moller T, Eggen BJ. Next generation transcriptomics and genomics elucidate biological complexity of microglia in health and disease. *Glia.* 2015
32. Goldberg EL, Dixit VD. Drivers of age-related inflammation and strategies for healthspan extension. *Immunol Rev.* 2015; 265:63–74. [PubMed: 25879284]
33. Youm YH, Grant RW, McCabe LR, Albarado DC, Nguyen KY, Ravussin A, Pistell P, Newman S, Carter R, Laque A, et al. Canonical Nlrp3 inflammasome links systemic low-grade inflammation to functional decline in aging. *Cell Metab.* 2013; 18:519–532. [PubMed: 24093676]
34. Frank MG, Barrientos RM, Biedenkapp JC, Rudy JW, Watkins LR, Maier SF. mRNA up-regulation of MHC II and pivotal pro-inflammatory genes in normal brain aging. *Neurobiol Aging.* 2006; 27:717–722. [PubMed: 15890435]
35. Wynne AM, Henry CJ, Huang Y, Cleland A, Godbout JP. Protracted downregulation of CX3CR1 on microglia of aged mice after lipopolysaccharide challenge. *Brain Behav Immun.* 2010; 24:1190–1201. [PubMed: 20570721]
36. Hoek RM, Ruuls SR, Murphy CA, Wright GJ, Goddard R, Zurawski SM, Blom B, Homola ME, Streit WJ, Brown MH, et al. Down-regulation of the macrophage lineage through interaction with OX2 (CD200). *Science.* 2000; 290:1768–1771. [PubMed: 11099416]
37. Cardona AE, Pioro EP, Sasse ME, Kostenko V, Cardona SM, Dijkstra IM, Huang D, Kidd G, Dombrowski S, Dutta R, et al. Control of microglial neurotoxicity by the fractalkine receptor. *Nat Neurosci.* 2006; 9:917–924. [PubMed: 16732273]
38. Poliani PL, Wang Y, Fontana E, Robinette ML, Yamanishi Y, Gilfillan S, Colonna M. TREM2 sustains microglial expansion during aging and response to demyelination. *J Clin Invest.* 2015; 125:2161–2170. [PubMed: 25893602]
39. Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, Koeglspenger T, Dake B, Wu PM, Doykan CE, et al. Identification of a unique TGF-beta-dependent molecular and functional signature in microglia. *Nat Neurosci.* 2014; 17:131–143. [PubMed: 24316888] [** Transcriptomic analyses from this study revealed a unique primary microglial molecular signature, and for the first time demonstrated the requirement for TGF- β to promote this signature. Cultured microglia without TGF- β did not produce the same expression of markers and mice deficient for TGF- β 1 had an absence of microglia.]
40. Koeglspenger T, Li S, Brenneis C, Saulnier JL, Mayo L, Carrier Y, Selkoe DJ, Weiner HL. Impaired glutamate recycling and GluN2B-mediated neuronal calcium overload in mice lacking TGF-beta1 in the CNS. *Glia.* 2013; 61:985–1002. [PubMed: 23536313]
41. Norden DM, Fenn AM, Dugan A, Godbout JP. TGFbeta produced by IL-10 redirected astrocytes attenuates microglial activation. *Glia.* 2014; 62:881–895. [PubMed: 24616125]

42. Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, Strauss-Ayali D, Viukov S, Williams M, Misharin A, et al. Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity*. 2013; 38:79–91. [PubMed: 23273845]
43. Goldmann T, Wieghofer P, Muller PF, Wolf Y, Varol D, Yona S, Brendecke SM, Kierdorf K, Staszewski O, Datta M, et al. A new type of microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation. *Nat Neurosci*. 2013; 16:1618–1626. [PubMed: 24077561]
44. Parkhurst CN, Yang G, Ninan I, Savas JN, Yates JR 3rd, Lafaille JJ, Hempstead BL, Littman DR, Gan WB. Microglia promote learning-dependent synapse formation through brain-derived neurotrophic factor. *Cell*. 2013; 155:1596–1609. [PubMed: 24360280]
45. Beutner C, Lepperhof V, Dann A, Linnartz-Gerlach B, Litwak S, Napoli I, Prinz M, Neumann H. Engineered stem cell-derived microglia as therapeutic vehicle for experimental autoimmune encephalomyelitis. *Gene Ther*. 2013; 20:797–806. [PubMed: 23324824]
46. Gosselin D, Link VM, Romanoski CE, Fonseca GJ, Eichenfield DZ, Spann NJ, Stender JD, Chun HB, Garner H, Geissmann F, et al. Environment drives selection and function of enhancers controlling tissue-specific macrophage identities. *Cell*. 2014; 159:1327–1340. [PubMed: 25480297]
47. Lavin Y, Winter D, Blecher-Gonen R, David E, Keren-Shaul H, Merad M, Jung S, Amit I. Tissue-resident macrophage enhancer landscapes are shaped by the local microenvironment. *Cell*. 2014; 159:1312–1326. [PubMed: 25480296]
48. Satoh A, Brace CS, Rensing N, Cliften P, Wozniak DF, Herzog ED, Yamada KA, Imai S. Sirt1 extends life span and delays aging in mice through the regulation of Nk2 homeobox 1 in the DMH and LH. *Cell Metab*. 2013; 18:416–430. [PubMed: 24011076]
49. Cho SH, Chen JA, Sayed F, Ward ME, Gao F, Nguyen TA, Krabbe G, Sohn PD, Lo I, Minami S, et al. SIRT1 deficiency in microglia contributes to cognitive decline in aging and neurodegeneration via epigenetic regulation of IL-1 β . *J Neurosci*. 2015; 35:807–818. [PubMed: 25589773] [* This was the first study to demonstrate a role for DNA methylation in regulating IL-1 β expression in aged microglia. Hypomethylation of IL-1 β was strongly associated with age and increased IL-1 β mRNA expression. This identifies a novel mechanism in which microglial priming can occur during aging.]
50. Sullivan KE, Reddy ABM, Dietzmann K, Suriano AR, Kocieda VP, Stewart M, Bhatia M. Epigenetic Regulation of Tumor Necrosis Factor Alpha. *Mol Cell Biol*. 2007; 27:5147–5160. [PubMed: 17515611]
51. Eddy JL, Krukowski K, Janusek L, Mathews HL. Glucocorticoids regulate natural killer cell function epigenetically. *Cell Immunol*. 2014; 290:120–130. [PubMed: 24978612]
52. Tang Y, Li T, Li J, Yang J, Liu H, Zhang XJ, Le W. Jmjd3 is essential for the epigenetic modulation of microglia phenotypes in the immune pathogenesis of Parkinson's disease. *Cell Death Differ*. 2014; 21:369–380. [PubMed: 24212761]
53. Sato F, Tsuchiya S, Meltzer SJ, Shimizu K. MicroRNAs and epigenetics. *Febs J*. 2011; 278:1598–1609. [PubMed: 21395977]
54. Cardoso AL, Guedes JR, Pereira de Almeida L, Pedrosa de Lima MC. miR-155 modulates microglia-mediated immune response by down-regulating SOCS-1 and promoting cytokine and nitric oxide production. *Immunology*. 2012; 135:73–88. [PubMed: 22043967]
55. Fenn AM, Smith KM, Lovett-Racke AE, Guerau-de-Arellano M, Whitacre CC, Godbout JP. Increased miR-29b in the aged brain correlates with the reduction of IGF-1 and CX(3)CL1. *Neurobiol Aging*. 2013; 34:2748–2758. [PubMed: 23880139]
56. Johnson RW. Feeding the beast: Can microglia in the senescent brain be regulated by diet? *Brain Behav Immun*. 2015; 43:1–8. [PubMed: 25451610]
57. Szic KS, Declerck K, Vidakovi M, Vanden Berghe W. From inflammaging to healthy aging by dietary lifestyle choices: is epigenetics the key to personalized nutrition? *Clin Epigenetics*. 2015; 7:33. [PubMed: 25861393]

Highlights

- Brain microglia react to peripheral immune signals, inducing neuroinflammation
- Brain microglia become dysregulated with age and cause chronic neuroinflammation
- Understanding microglial cell regulation is important to promote healthy aging

Figure 1.

Age-related changes in microglia. Microglia participate in brain development by regulating the number of neuronal precursors and engaging in synaptic pruning. In adulthood they maintain tissue homeostasis through active surveillance of the brain parenchyma, effective phagocytosis to clear harmful debris such as A β aggregates, and facilitate synaptic plasticity by either supporting or removing synapses. Upon stimulation by signals emerging from the peripheral immune system, microglia produce pro-inflammatory cytokines to initiate adaptive behavioral and physiological responses. During aging however, microglia are primed to be more reactive to inflammatory stimuli, overproducing pro-inflammatory cytokines for a longer period of time resulting in maladaptive sickness responses. Age-related changes in neural-microglial communication and epigenetic modifications may underlie microglial cell hyperactivity.