Published in final edited form as:

Org Lett. 2015 February 20; 17(4): 856–859. doi:10.1021/ol5037024.

Intramolecular [4+2] Trapping of a Hexadehydro-Diels–Alder (HDDA) Benzyne by Tethered Arenes

Vedamayee D. Pogula, Tao Wang, and Thomas R. Hoyer*

Department of Chemistry, University of Minnesota, 207 Pleasant Street SE Minneapolis, Minnesota 55455, United States

Abstract

We report here the efficient, intramolecular trapping in a Diels–Alder (DA) sense of thermally generated benzyne by one of two pendant arene rings. A more electron rich ring (*p*-methoxyphenyl) reacted substantially faster than a simple phenyl ring, which was in turn, slightly more reactive vs. a 4-carbomethoxyphenyl ring. Photoinduced di- π -methane rearrangement of the initial DA adducts gave rise to unusual isomeric polycyclic adducts.

Abstract

Benzene (and its substituted derivatives) are generally unwilling partners in typical Diels–Alder (DA) [4+2] cycloaddition reactions. In fact, a solvent like benzene, toluene, xylene, or 1,2-dichlorobenzene is often used as an effectively inert solvent medium for DA reactions between other pairs of diene/dienophile reactants. The energetic penalty of dearomatization of the benzenoid ring is typically too costly to permit it to participate in cycloaddition. *o*-Benzyne (**1**) and its substituted derivatives comprise one of the very few classes of dienophiles with sufficient reactivity to engage a benzenoid as a diene. The first example of the reaction of 1,2-dehydrobenzene (**1**) with benzene produced benzobar-relene (**3a** or **3b** with X = H) in 12% yield (along with biphenyl and benzocyclooctatetraene).¹ This accentuates the inherent difficulty in DA reactions of arenes, even when a highly energetic benzyne is conscripted as the dienophile.

*Corresponding Author: hoye@umn.edu.

Supporting Information

Experimental procedures, characterization data, copies of ¹H- and ¹³C NMR spectra for all isolated compounds, ORTEP renderings, and DFT computational methods (146 pages). This material is available free of charge via the Internet at <http://pubs.acs.org>.

Author Contributions

V.P. performed all of the experiments; T.W. performed all of the computations. / The manuscript was written through contributions from all authors. / All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interests.

In 1977 Yoshida, Oda, and coworkers reported² the relative reactivity of **1** with several monosubstituted benzenoid derivatives (i.e., **2**) through sets of bimolecular competition experiments (Scheme 1). The reactivity pattern was that more electron-rich arenes reacted faster; however, the range of relative rates varied only from ca. 15 for the fastest (anisole) to 1 for the slowest (trifluoromethylbenzene). Anisole was also the only trapping agent that showed a propensity to favor addition across the C1/C4 positions of the arene ring (to give **3a**); the other derivatives favored formation of the C2/C5 adduct **3b**. In all of these experiments the aromatic trapping agents were used as the solvent; the sole isolated yield reported (for the anisole case) were 32% and 8% for **3a** (X = OMe) and **3b** (X = Me), respectively.

The cycloisomerization of triynes like **4** (and **7**) to give benzyne like **5** (and **8**) by a simple thermal process³ has considerable generality (Scheme 2).⁴ We have termed this cyclization a hexadehydro-Diels–Alder (HDDA) reaction. The reactive benzyne intermediates can be trapped in many useful ways, including by reaction in a [4+2] Diels–Alder sense with an aromatic benzenoid ring in either a bimolecular (e.g., **5** to **6**)⁴ or unimolecular (e.g., **8** to **9**)^{4,5} fashion. To understand in more detail the influence on reactivity of substituents on the aromatic ring acting in the role of the diene, we designed a substrate that would pit two aromatic rings in competition for the benzyne dienophile. Specifically, we have prepared and studied the reactivity of a number of substrates of general structure **10**, having two aryl groups that bear substituents of differing electronic character, yet otherwise were equally well-poised to engage in intramolecular [4+2] trapping of the HDDA-generated benzyne. The ratio of the constitutionally isomeric products **11** (where X = Y) formed upon heating **10** in an inert solvent would then provide a direct indication of the impact of substituents X vs. Y on the rate of the intramolecular Diels–Alder (IMDA) trapping reaction.

The substrates examined in this study were triynes **10a–f**. Their preparation by cross coupling of the diyne **12**⁶ and bromoalkynes **13** (derived from the known benzhydryl derivatives **14**) is outlined in Scheme 3 [see the Supporting Information (SI) for details].

The rate-limiting step in the conversion of **10** to **11** (Scheme 2c) is the initial HDDA cycloisomerization reaction; the subsequent Diels–Alder trapping event by the pendant arene is relatively rapid. We have reported that the half-life for the HDDA reaction of a triynone similar to that of **10** is ca. 4 h at 85 °C.⁴ Accordingly, we heated each of **10a–f** in an 80–85 °C bath in a sealed vial containing a solution of the triyne for ca. 48 hours in CDCl₃. This solvent has proven repeatedly to be inert toward HDDA-generated benzyne, underscoring the fact that these benzyne should not be considered to engage in reaction processes that are radical in character. Analysis of the ¹H NMR spectra of the reaction mixture itself was used to judge the product ratios in the cases where competitive reaction pathways were detected.

The diphenyl-containing triyne **10a** gave the trapped adduct **11a** (Figure 1.) in 70% yield following chromatographic purification.⁷ Hence, a simple phenyl ring is an efficient and competent trap for the benzyne intermediate (cf. **5**, R = CH₂OCHAr¹Ar²). The first substrate we studied with two different pendant arenes was the mono-*p*-methoxy benzhydryl ether **10b**. It produced a single isolable IMDA adduct—namely, **11b-maj** in 88% yield. We did

not definitively detect the isomeric, phenyl-trapped adduct **11b-min** (^1H NMR). Thus, the more electron-rich arene is considerably more reactive (cf. $k_{\text{anisole}}/k_{\text{benzene}} = 8$, Scheme 1).

We then studied the mono-carbomethoxy mono-phenyl ether **10c**. In this substrate an electron-poor arene is pitted against the parent phenyl ring. The major product in this case was the phenyl-trapped adduct **11c-maj**, isolated in 64% yield. The minor component **11c-min** was also isolated following careful separation by HPLC (product ratio = 3.2:1 by ^1H NMR analysis of the reaction mixture). An interesting behavior was observed for **11c-min** (as well as for the analogous **11d** and **11e**, which also contain a bridgehead carbomethoxy group). Several resonances in both the ^1H and ^{13}C NMR spectra were broadened when recorded at ambient temperature. We attributed this to slow rotation about the bridgehead to carbonyl carbons in these hindered esters. These resonances sharpened considerably when the ^1H spectrum was recorded at 100 °C (see SI), consistent with this hindered rotation hypothesis. Similarly, when the bisester-containing substrate **10d** (or the mono-ester **10e**) was heated (in the dark, see below), only the primary Diels-Alder adduct **11d** (or **11e**) was observed.

We also examined, in one instance, the effect of substitution pattern. Substrate **10f** contains both a *para*- and an *ortho*-methoxyphenyl substituent. When heated in 1,2-dichloroethane at 85 °C for 2 days, **10f** produced both **11f-maj** and the ketone **11f-min**⁸ (8.8:1 ratio from ^1H NMR analysis) in 89% and 5% isolated yield (MPLC separation). Oda et al. reported facile *in situ* hydrolysis of a primary enol ether product arising from anisole trapping to its ketone, the process we also presume gives rise to **11f-min**. They also observed a similar preference for formation of the bridgehead rather than the alkenyl methoxylated isomer (cf. Scheme 1).

We examined whether DFT calculations are effective at rationalizing the observed selectivities. Specifically, we have computed the transition state (TS) structures for the competing IMDA trapping reactions of the benzyne derived from **10b**, **10c**, and **10f**. These lead via **TS15** (Figure 2a) to the major and minor products of **11b**, **11c**, and **11f**. Also given in Figure 2a are the differences in computed free energy of activation ($\Delta G^\ddagger$) along with the associated computed and experimentally observed product ratios. DFT does a good job of accounting for the relative energies of the competing TS structures for the trapping of the benzyne. The geometries of those TSs all indicated a high degree of asynchronicity in these concerted IMDA reactions (Figure 2b).

We were also successful in using DFT to account for the sense of diastereoselectivity (i.e., anti) observed in the formation of **11f-min**⁸ (cf. Figure 1). The TS **TS15f-min-anti** was favored by 1.8 kcal•mol⁻¹ over the alternative **TS15f-min-syn** (Figure 2b), presumably reflecting the greater steric congestion in the latter, which possesses cis-(and endo-)oriented methoxy and aryl groups.

We observed that some of the IMDA products containing bridgehead substituents on their benzobicyclo[2.2.2]octatriene units underwent facile photo-induced rearrangement reactions (Figure 3.). Either ambient laboratory light or exposure of a chloroform solution to a 75 W incandescent light source was sufficient to induce these changes in these typically yellow fluorenone compounds. Specifically, the primary HDDA-IMDA products **11c-min**,

11d, and **11e** gave rise via di- π -methane rearrangement⁹ to the cyclopropane-containing products **16c**, **16d**, and **16e**, respectively. Irradiation of **11f** in CDCl₃ produced a structure whose ¹H NMR spectrum was consistent with the cyclopropane **16f**, which, upon chromatographic purification on silica gel, cleanly gave rise to the ketone **17f**. The structure assignments for **16c** and **17f** were confirmed by single crystal X-ray analysis. The hydrogen locations in the renderings in Figure 3 are idealized and some have been omitted for clarity. Interestingly, products of trapping by a parents phenyl ring, i.e. **11a** and **11c-maj**, did not show evidence of the di- π -methane rearrangement, likely reflecting the greater steric strain in those dibenzobar-relenes having a non-hydrogen bridgehead substituent.

Finally, a pendant naphthalene substituent was shown to efficiently capture the HDDA benzyne. Triyne **18** gave the Diels–Alder adduct **19** in high yield following chromatographic purification (Figure 4). The *in situ* ¹H NMR spectrum of this reaction mixture is indicative of the cleanliness of this transformation.

In conclusion, we have studied intramolecular Diels–Alder reactions between various pendent aromatic groups to trap thermally generated benzynes. The arene functions as the 4 π dienic component and the HDDA benzyne the 2 π dienophile. More electron rich arenes are more reactive dienes, consistent with the view of the benzyne as an electrophilic partner. DFT calculations were effective in accounting for the observed product ratios for three substrates containing two different and competing arenes. The transition state structures for these reactions show a substantial degree of asynchronicity (Figure 2b). Adducts containing a bridgehead carbomethoxy substituent show hindered rotation about the ester carbonyl to barrelene bridgehead C–C bond. Some of the initial Diels–Alder adducts—yellow, dibenzobarrelene, fluorenone-containing derivatives—underwent smooth di- π -methane rearrangement when exposed to ambient light. Finally, a pendant naphthalene substituent is a particularly effective diene. Collectively these results provide guidance about what types of substituents can be selected to enhance bimolecular Diels–Alder trapping of arynes by structurally more complex arenes, a topic that is under further study here.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

This investigation was supported by a grant awarded by the U.S. Department of Health and Human Services (National Institute of General Medical Sciences, GM-65597). Portions of this work were performed with hardware and software resources available through the University of Minnesota Supercomputing Institute (MSI). T. W. acknowledges the support of a Wayland E. Noland Fellowship and a University of Minnesota Graduate School Doctoral Dissertation Fellowship. We thank Victor G. Young, Jr. of the University of Minnesota for assistance with the X-ray diffraction analysis.

REFERENCES

1. Miller RG, Stiles M. J. Am Chem. Soc. 1963; 85:1798–1800.
2. Tabushi I, Yamada H, Yoshida Z, Oda R. Bull. Chem. Soc. Jpn. 1977; 50:285–290.
3. a Miyawaki K, Suzuki R, Kawano T, Ueda I. Tetrahedron Lett. 1997; 38:3943–3946. b Bradley AZ, Johnson RP. J. Am. Chem. Soc. 1997; 119:9917–9918.

4. Hoye TR, Baire B, Niu D, Willoughby PH, Woods BP. *Nature*. 2012; 490:208–212. [PubMed: 23060191]
5. Niu D, Wang T, Woods BP, Hoye TR. *Org. Lett.* 2014; 16:254–257. [PubMed: 24329102]
6. Suffert J, Abraham E, Raeppl S, Brückner R. *Liebigs Ann.* 1996:447–456.
7. We typically observed some amount of competitive cleavage of the benzhydryl ether to its corresponding ketone (or benzylic ether to aldehyde in the case of **10e**). For **10a** benzophenone itself was observed. Although this byproduct formation reduced the overall yield of IMDA products isolated from these experiments, it did not compromise our ability to assess the relative propensity of two different arenes to participate in the Diels–Alder trapping reactions. The IMDA products do not further fragment to give these benzophenone derivatives
8. The relative configuration of the two stereocenters in **11f-min** was assigned on the basis of NOE experiments (see Supporting Information)
9. Liu RSH, Krespan CG. *J. Org. Chem.* 1969; 34:1271–1278.

Figure 1.

Products from competitive IMDA trapping within benzyne intermediates **15**, derived from **10a-f**

Figure 2.

(a) DFT calculation [M06-2X/6-31G(d) with SMD (chloroform)] of the relative TS energies for the competitive IMDA reactions proceeding via **TS15** to **11-maj** vs. **11-min**. (b) DFT geometries and relative energies for the TSs accounting for the preferential formation of **11f-maj** as well as the observed anti relative configuration in **11f-min** from **10f**.

Figure 3. Di- π -methane rearrangement products from light-induced transformations of the primary HDDA-IMDA products **11c-min**, **11d**, **11e**, and **11f-maj**.

Figure 4.
 ^1H NMR spectrum of the product mixture produced directly upon heating **18** in CDCl_3 at 85°C .

Scheme 1.

Reported bimolecular Diels–Alder reactions between *o*-benzyne (**1**) as the dienophile and various monosubstituted benzene derivatives **2** as the diene partner to give [4+2] adducts **3a-b**

Scheme 2.

Examples of HDDA-generated benzyne reacting with arenes either (a) bimolecularly or (b) unimolecularly and (c) the substrates designed for the intramolecular competition reactions reported here

Scheme 3. General route used for the synthesis of HDDA triyne substrates **10**