Published in final edited form as:

J Mol Neurosci. 2016 January ; 58(1): 88–92. doi:10.1007/s12031-015-0647-x.

CSF β -amyloid 1-42 predicts progression to cognitive impairment in newly diagnosed Parkinson disease

Mark Terrelonge Jr., MPH^a, Karen S. Marder, MD, MPH^a, Daniel Weintraub, MD^b, and Roy N. Alcalay, MD, MS^a

^aColumbia University Department of Neurology, New York, NY 10032

^bDepartments of Psychiatry and Neurology, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania, USA; Philadelphia Veterans Affairs Medical Center, Philadelphia, Pennsylvania

Abstract

BACKGROUND—Low CSF β -amyloid 1-42 has been associated with cognitive decline in advanced Parkinson's Disease; data from a single cohort suggest β -amyloid 1-42 may be an early marker of cognitive impairment.

DESIGN/METHODS—Newly diagnosed Parkinson's participants (mean duration: 6.9 months) in the Parkinson's Progression Markers Initiative (n=341) were assessed at baseline (untreated state) and followed for two years. CSF β -amyloid 1-42, α -synuclein, total tau, and tau phosphorylated at threonine 181 were collected at baseline. Participants were classified as having CI if scores on two of six cognitive tests were 1.5 standard deviations below the standardized mean based on published norms in healthy controls. Multivariable regression analyses was used to determine the association between baseline CSF markers with cognitive impairment, defined by neuropsychological testing performance at two-year follow-up.

RESULTS—55 participants (16.1%) had CI at baseline and were not included in further analyses. 37 of the 286 participants without CI at baseline (12.9%) developed CI at two years. Participants with CI at two years had significantly lower mean baseline CSF β -amyloid 1-42 levels than non-CI

Corresponding Author: Roy Alcalay, 710 W 168th St, New York, NY 10032, (212)-305-5554, rna2104@columbia.edu.

Author Roles

Research project: Conception: MT, KM, RA; Organization: MT, RA; Execution: MT, RA

Statistical Analysis: Design: MT, RA, KM, DW; Execution: MT

Review and Critique: MT, KM, DW, RA

Manuscript Preparation: Writing of the first draft: MT, RA; Review and Critique: KM, DW

Financial Disclosures:

Dr. Marder reports grants from NIH [#NS036630 (PI), 1UL1 RR024156-01(Director PCIR), PO412196- G (Co-I), and PO412196-G (Co-I)], grants from steering committee for U01NS052592, grants from Parkinson Disease Foundation, and grants from MJ Fox Foundation, and research funds from Teva Pharmaceuticals outside the submitted work. Dr. Weintraub received research funding from Michael J. Fox Foundation for Parkinson's Research, National Institutes of Health, Novartis Pharmaceuticals, Department of Veterans Affairs, and Alzheimer's Disease Cooperative Study; honoraria from Teva Pharmaceuticals, Lundbeck Inc., Pfizer, Avanir Pharmaceuticals, Merck & Co., UCB, Bristol-Myers Squibb Company, Novartis Pharmaceuticals, Eli Lilly and Company, Clintrex LLC, Theravance, Medivation, CHDI Foundation, and Alzheimer's Disease Cooperative Study; license fee payments from the University of Pennsylvania for the QUIP and QUIP-RS; and fees for testifying in court case related to impulse controls disorders in Parkinson's disease (March 2013). Dr. Alcalay is supported by the Parkinson's Disease Foundation, the National Institutes of Health (K02NS080915, NS036630 and UL1 TR000040, formerly the National Center for Research Resources, Grant Number UL1 RR024156) and the Brookdale Foundation. He consulted for Genzyme/Sanofi and Prophase.

participants (343.8 vs. 380.4pg/mL, $p<0.01$); no significant difference was seen for α -synuclein, T-tau, or P-tau 181. In a regression model of 286 participants without baseline CI adjusted for age, gender, disease duration, education, motor severity, and depression status, lower baseline β -amyloid 1-42 levels were associated with higher odds of CI at two years. ($OR_{10pg/mL}=1.04$, 95% CI: 1.01–1.08, $p<0.05$).

CONCLUSION—CSF β -amyloid 1-42 level at disease onset is an independent predictor of cognitive impairment in early Parkinson's Disease.

Keywords

CSF β -amyloid 1-42; cognitive impairment; Parkinson's Disease; PPMI

Introduction

Up to 80% of Parkinson Disease (PD) patients develop dementia during the course of their illness.¹ Several baseline factors have been associated with more rapid cognitive decline in PD including advanced age at time of diagnosis, male gender, and education level.² The degenerative process may begin years before the first signs of cognitive impairment or dementia become apparent in both Alzheimer's disease (AD) and PD, and these changes may be reflected in CSF marker profile.^{3,4} Additionally, evidence has shown that lower CSF levels of β -amyloid predict faster cognitive decline among PD patients.^{5–7} A study from the Norwegian ParkWest project, a population based cohort study of newly diagnosed PD patients (mean disease duration of 2.2years), found that CSF β -amyloid 1-42 ($A\beta_{1-42}$) was associated with memory decline at the time of patient diagnosis.⁸ Later analysis from the same cohort found that low CSF $A\beta_{1-42}$ at baseline was associated with early dementia associated with PD (median time to dementia: 4.3 years).⁹

The Parkinson's Progression Markers Initiative (PPMI) is an international, multi-center prospective study following PD cases and controls with baseline CSF sampling and biannual neurological and cognitive evaluations.¹⁰ A previous cross-sectional design study using partial PPMI data showed slight but significantly lower levels of CSF $A\beta_{1-42}$, α -synuclein (α -syn), total tau (T-tau), and tau phosphorylated at threonine 181 (P-tau₁₈₁) in PD compared with healthy controls, with $A\beta_{1-42}$ and P-tau₁₈₁ remaining lower than healthy control data after controlling for age, gender, and education.¹¹ We examined whether CSF concentration of $A\beta_{1-42}$, α -syn, T-tau, and P-tau₁₈₁ at baseline predicts diagnosis of cognitive impairment at two-year follow up among newly diagnosed, drug-naïve PD patients without cognitive impairment.

Methods

Participants

Data were obtained from the PPMI database (www.ppmi-info.org/data) on August 10, 2015.¹⁰ Participants were included in this PD cohort if they enrolled in the study within two years of diagnosis and were not expected to require PD medication within 6 months of their baseline evaluation. Participants were excluded if they had a clinical diagnosis of dementia, were unable to participate in lumbar puncture, or had MRI evidence of another clinically

significant neurological disorder. Scans without evidence for dopaminergic deficit (SWEDD) participants and control participants were not included in this study.

Individuals from the PD cohort who had two years of follow-up were included in the analysis. At the time of data acquisition, 341 PD participants (of original 406 with baseline data) had completed two years of study follow-up. Written informed consent was obtained from all participants, and all PPMI sites received approval from their respective ethics committee on human experimentation prior to study initiation.

Assessments

The annual assessment included six cognitive tests, which can be divided into four cognitive domains: Memory (Hopkins Verbal Learning Test-Revised [HVLT-R] Recall, HVLT-R Recognition Discrimination),¹² Visuospatial (Judgment of Line Orientation [JOLO]),¹³ Working Memory- Executive Function (Letter Number Sequencing [LNS], Semantic Fluency),^{14,15} and Attention- Processing Speed (Symbol Digit Modalities Test [SDMT]).¹⁶ Cognitive impairment was defined as having at least 2 test scores (of six; irrespective of test domain) greater than 1.5 standard deviation below the age- and education-standardized mean score based on published norms in healthy controls.¹⁷

CSF Analysis

CSF samples were collected from all participants enrolled in the study at baseline including A β 1-42, α -syn, T-tau, and P-Tau₁₈₁. Additional information on how CSF samples were collected and analyzed was previously reported.¹¹

Statistical Analysis

Data were analyzed using SAS v 9.3. Participants with baseline cognitive impairment by the aforementioned definition were not included in future analyses. Baseline demographics (age, gender, PD duration from diagnosis, education, and 15-item Geriatric Depression Scale [GDS-15]), disease characteristics (Movement Disorder Society- Unified Parkinson's Disease Rating Scale [MDS- UPDRS] motor score Part 3, MDS-UPDRS Part 1 Cognitive Impairment self-report at baseline, and MDS-UPDRS Part 1 Cognitive Impairment self-report at 2 years), and baseline CSF marker levels and their ratios (α -syn/T-tau, T-tau/A β 1-42, P-Tau₁₈₁/A β 1-42, P-Tau₁₈₁/T-tau) were compared between participants with and without CI at two year follow up using either the Student t-test or the Chi-square test. Non-normal data were compared using the Wilcoxon rank sum test. Multivariable logistic regression was used to determine the association of baseline CSF measures and their ratios to CI at 2 years. The adjusted model included baseline age, gender, disease duration, education, MDS-UPDRS Part 3 motor score, and Geriatric Depression Scale score. Significance level for all non-ratio tests was set at $P < 0.05$ (2-tailed test). Adjustment for multiple comparisons was used for all ratio tests with a Bonferroni-Holm correction.

Results

Participant Characteristics

Two year follow-up data was available for 341 participants out of 406 assessed at baseline. There were no significant differences in baseline characteristics or biomarker profiles between those with two years of follow-up and those without two years of follow-up (Supplemental Table 1). 55 participants had CI at baseline (16.1%) and were not included in future analyses. Those with CI at baseline had greater motor impairment reflected by higher MDS-UPDRS motor scores (24.6 vs. 20.3, $p<0.01$), had higher scores on depression rating scale (5.7 vs. 5.2, $p=0.04$), and had a higher proportion of self-reported cognitive decline at baseline (43.6% vs. 23.4%, $p<0.01$) (Supplemental Table 2).

Of the 286 remaining participants without CI at baseline, 37 (12.9%) developed CI by year 2. Demographics and disease characteristics were similar between those with and without CI at two years (Table 1). Of note, in this cohort baseline self-reported cognitive changes on the MDS-UPDRS was not associated with later development of CI; however, at year-2 evaluation, those with CI were more likely to report cognitive changes during that evaluation (Table 1).

CSF markers as predictor of CI

Participants with CI at two years had significantly lower mean baseline CSF A β 1-42 levels than those who did not (343.8 vs. 380.4 pg/mL, $p<0.01$). No significant differences in CSF level were noted for α -syn, T-tau, or P-tau₁₈₁. Significant differences were not noted in the ratio of CSF T-tau/A β 1-42, T-tau/ α -syn, CSF P-tau₁₈₁/A β 1-42, or P-tau₁₈₁/T-tau (Table 1).

Logistic regression was used to test whether baseline CSF markers could predict CI at two years, adjusting for baseline demographic characteristics (Table 2). After adjustment for age, education, disease duration, gender, MDS-UPDRS motor score, and Geriatric Depression Score, CSF A β 1-42 still predicted CI at two years. α -syn, T-tau, and P-tau₁₈₁ were not predictive of CI at two years in the univariate or multivariable models. CSF ratios of α -syn/T-tau, T-tau/A β 1-42, Ptau-181/A β 1-42, and P tau-181/Total tau were also not associated with CI at two years in multivariable models.

Discussion

In the present study of a medication naïve PD cohort with two years follow-up, we found that lower baseline CSF A β 1-42 was associated with higher odds of having CI after two years of follow-up after adjusting for baseline demographic characteristics. To our knowledge, this is the first study to demonstrate an association between lower baseline CSF A β 1-42 levels in medication naïve newly diagnosed PD cases without cognitive impairment and later development of CI as defined by neuropsychological testing. This finding is consistent with findings from the Norwegian ParkWest study, which reported an association between lower levels of CSF A β 1-42 and cognitive deficits in the domain of memory,⁸ and later development of PD dementia.⁹ It is also consistent with studies of more advanced PD cohorts associating lower CSF A β 1-42 with development of dementia.^{4,7,18}

That lower CSF A β 1-42 among medication naïve early PD cases predicts CI two years later argues that the neuropathological process that leads to cognitive impairment in these cases may have started before or in parallel to motor symptom onset. However, the neuropathological correlates of lower CSF A β 1-42 are not clear. In AD, it is hypothesized that low CSF A β 1-42 may reflect higher levels of aggregation of A β protein in the cortex, signifying a more advanced neurodegenerative process.^{19–21} Whether those with low levels of CSF A β 1-42 may represent a subset of PD with coincident AD remains to be seen, and may be elucidated with further follow-up (including post-mortem neuropathological evaluation).^{22,23}

Strengths of the present study include finding a robust relationship between CSF markers and CI in a multicenter longitudinal study including a large sample size of participants with newly diagnosed PD without CI. Limitations include a definition of CI dependent only on neuropsychological test scores without assessment of participant's functional impairment,²⁴ which was not available at time of participant enrollment. Taken together, this study provides evidence that CSF A β 1-42 may play a role in prediction of early cognitive impairment among persons with newly diagnosed PD with mild motor symptoms. Early identification of markers of cognitive impairment is important for determining which patients may benefit most from therapies once they become available. Further longitudinal analysis following changes in CSF A β 1-42 as they relate to cognition can be the next step in exploring the role of CSF A β 1-42 in predicting cognitive impairment and dementia.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Sources: No funding sources required for this study. Data support provided by the Michael J Fox Foundation Parkinson's Progression Markers Initiative (PPMI) study.

PPMI is sponsored by the Michael J. Fox Foundation for Parkinson's Research (MJFF) and is co-funded by MJFF, Abbvie, Avid Radiopharmaceuticals, Biogen Idec, Bristol-Myers Squibb, Covance, Eli Lilly & Co., F. Hoffman-La Roche, Ltd., GE Healthcare, Genentech, GlaxoSmithKline, Lundbeck, Merck, MesoScale, Piramal, Pfizer and UCB.

References

1. Aarsland D, Andersen K, Larsen JP, Lolk A, Kragh-Sorensen P. Prevalence and characteristics of dementia in Parkinson disease: an 8-year prospective study. *Arch Neurol*. 2003; 60:387–92. [PubMed: 12633150]
2. Hughes TA, Ross HF, Musa S, et al. A 10-year study of the incidence of and factors predicting dementia in Parkinson's disease. *Neurology*. 2000; 54:1596–602. [PubMed: 10762499]
3. Mollenhauer B, Trenkwalder C, von Ahsen N, et al. Beta-amyloid 1-42 and tau-protein in cerebrospinal fluid of patients with Parkinson's disease dementia. *Dement Geriatr Cogn Disord*. 2006; 22:200–8. [PubMed: 16899997]
4. Parnetti L, Castrioto A, Chiasserini D, et al. Cerebrospinal fluid biomarkers in Parkinson disease. *Nat Rev Neurol*. 2013; 9:131–40. [PubMed: 23419373]
5. Siderowf A, Xie SX, Hurtig H, et al. CSF amyloid {beta} 1-42 predicts cognitive decline in Parkinson disease. *Neurology*. 2010; 75:1055–61. [PubMed: 20720189]

6. Montine TJ, Shi M, Quinn JF, et al. CSF Abeta(42) and tau in Parkinson's disease with cognitive impairment. *Mov Disord.* 2010; 25:2682–5. [PubMed: 20818673]
7. Parnetti L, Farotti L, Eusebi P, et al. Differential role of CSF alpha-synuclein species, tau, and Abeta42 in Parkinson's Disease. *Front Aging Neurosci.* 2014; 6:53. [PubMed: 24744728]
8. Alves G, Bronnick K, Aarsland D, et al. CSF amyloid-beta and tau proteins, and cognitive performance, in early and untreated Parkinson's disease: the Norwegian ParkWest study. *J Neurol Neurosurg Psychiatry.* 2010; 81:1080–6. [PubMed: 20547614]
9. Alves G, Lange J, Blennow K, et al. CSF Abeta42 predicts early-onset dementia in Parkinson disease. *Neurology.* 2014; 82:1784–90. [PubMed: 24748671]
10. Parkinson Progression Marker I. The Parkinson Progression Marker Initiative (PPMI). *Prog Neurobiol.* 2011; 95:629–35. [PubMed: 21930184]
11. Kang JH, Irwin DJ, Chen-Plotkin AS, et al. Association of cerebrospinal fluid beta-amyloid 1-42, T-tau, P-tau181, and alpha-synuclein levels with clinical features of drug-naïve patients with early Parkinson disease. *JAMA Neurol.* 2013; 70:1277–87. [PubMed: 23979011]
12. Vanderploeg RD, Schinka JA, Jones T, Small BJ, Graves AB, Mortimer JA. Elderly norms for the Hopkins Verbal Learning Test-Revised. *Clin Neuropsychol.* 2000; 14:318–24. [PubMed: 11262707]
13. Mitrushina, MN. Handbook of normative data for neuropsychological assessment. 2. New York: Oxford University Press; 2005.
14. Groth-Marnat, G. Handbook of psychological assessment. 5. Hoboken, N.J: John Wiley & Sons, Inc; 2009.
15. Tombaugh TN, Kozak J, Rees L. Normative data stratified by age and education for two measures of verbal fluency: FAS and animal naming. *Arch Clin Neuropsychol.* 1999; 14:167–77. [PubMed: 14590600]
16. Sheridan LK, Fitzgerald HE, Adams KM, et al. Normative Symbol Digit Modalities Test performance in a community-based sample. *Arch Clin Neuropsychol.* 2006; 21:23–8. [PubMed: 16139470]
17. Weintraub D, Simuni T, Caspell-Garcia C, Coffey C, Lasch S, Siderowf A. Cognitive Performance and Neuropsychiatric Symptoms in Early, Untreated Parkinson's Disease. *Mov Disord.* 2015 In press.
18. Compta Y, Pereira JB, Rios J, et al. Combined dementia-risk biomarkers in Parkinson's disease: a prospective longitudinal study. *Parkinsonism Relat Disord.* 2013; 19:717–24. [PubMed: 23643469]
19. Strozzyk D, Blennow K, White LR, Launer LJ. CSF Abeta 42 levels correlate with amyloid-neuropathology in a population-based autopsy study. *Neurology.* 2003; 60:652–6. [PubMed: 12601108]
20. Fagan AM, Mintun MA, Mach RH, et al. Inverse relation between in vivo amyloid imaging load and cerebrospinal fluid Abeta42 in humans. *Ann Neurol.* 2006; 59:512–9. [PubMed: 16372280]
21. Shaw LM, Vanderstichele H, Knapik-Czajka M, et al. Qualification of the analytical and clinical performance of CSF biomarker analyses in ADNI. *Acta Neuropathol.* 2011; 121:597–609. [PubMed: 21311900]
22. Burack MA, Hartlein J, Flores HP, Taylor-Reinwald L, Perlmuter JS, Cairns NJ. In vivo amyloid imaging in autopsy-confirmed Parkinson disease with dementia. *Neurology.* 2010; 74:77–84. [PubMed: 20038776]
23. Hyman BT, Phelps CH, Beach TG, et al. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimers Dement.* 2012; 8:1–13. [PubMed: 22265587]
24. Litvan I, Goldman JG, Troster AI, et al. Diagnostic criteria for mild cognitive impairment in Parkinson's disease: Movement Disorder Society Task Force guidelines. *Mov Disord.* 2012; 27:349–56. [PubMed: 22275317]

Table 1

Comparison of demographics and Parkinson's disease characteristics between participants with and without cognitive impairment at 2 years

Baseline (Year 0)			
Baseline characteristics	No Cognitive Impairment at 2 years (N=249)	Cognitive Impairment at 2 years (N=37)	P-value
Gender (% male)	161 (64.7%)	26 (70.3%)	P=0.44
Age (years)	60.6	61.8	p=0.48
Age at Disease Onset (years)	60.0	61.2	P=0.48
Disease Duration (months)	6.9	7.7	p=0.51
Education (years)	15.8	15.3	p=0.26
Movement Disorders Society-United Parkinson's Disease Rating Scale Motor Score	8.7	9.8	p=0.23
Geriatric Depression Scale Score	5.2	4.9	p=0.11
Self-reported Cognitive changes at baseline on the Movement Disorders Society –United Parkinson's Disease Rating Scale	59 (23.7%)	8 (21.6%)	p=0.96
CSF A β 1-42 levels pg/mL	380.4	343.8	p<0.01
CSF Alpha-synuclein pg/mL	1897.3	1689.2	p=0.14
CSF Total tau pg/ml	44.1	40.9	p=0.29
CSF Phosphorylated tau-181 pg/ml	16.2	14.9	p=0.31
CSF Alpha-synuclein/Total tau	44.9	42.3	p=0.28
CSF Total tau/A β 1-42	0.12	0.13	p=0.63
CSF Phosphorylated tau-181/A β 1-42	0.04	0.04	p=0.88
CSF Phosphorylated tau-181/Total tau	0.39	0.38	p=0.83
Year 2			
	No Cognitive Impairment at 2 years (N=249)	Cognitive Impairment at 2 years (N=37)	P-value
Self-reported cognitive changes at two year follow up on the Movement Disorders Society-United Parkinson's Disease Rating Scale	76 (30.5%)	19 (51.4%)	p<0.01

Values denote means or percentage (n) unless otherwise indicated. P-values computed using student t-test, Wilcoxon rank sum test, or χ^2 test as appropriate.

Table 2

The association between cognitive impairment (outcome) and CSF markers (predictors) in univariate and multivariable regression models. [OR for decrease in CSF marker levels]

CSF Markers	Univariable analysis			Multivariable analysis ^d		
	OR	95% CI	P value	OR	95% CI	P value
Aβ1-42 _{10pg/mL}	0.96	(0.92, 0.99)	p=0.03	0.96	(0.92, 0.99)	p=0.04
α-synuclein _{10pg/mL}	0.99	(0.99, 1.00)	p=0.13	0.99	(0.99, 1.00)	p=0.14
total tau _{10pg/mL}	0.89	(0.70, 1.11)	p=0.27	0.88	(0.69, 1.11)	p=0.27
p-tau _{10pg/mL}	0.88	(0.60, 1.27)	p=0.49	0.88	(0.61, 1.27)	p=0.49
CSF Alpha-synuclein/Total tau	0.99	(0.97, 1.01)	p=0.38	0.99	(0.96, 1.01)	p=0.39
Total tau/Aβ1-42	3.64	(0.02, 794.9)	p=0.64	4.46	(0.01, 999.9)	p=0.62
Phosphorylated tau-181/Aβ1-42	1.57	(0.01, 999.9)	p=0.92	0.97	(0.01, 999.9)	p=0.99
Phosphorylated tau-181/Total tau	0.88	(0.20, 3.94)	P=0.87	0.88	(0.19, 4.02)	P=0.86

^dModel adjusted for age, gender, education, duration, Movement Disorder Society –United Parkinson’s Disease Rating Scale Part 3 motor score, and Geriatric Depression Scale.