CORRESPONDENCE

De Novo Acute Heart Failure and Acutely Decompensated Chronic Heart Failure

Dr. med. Astrid Hummel, PD Dr. med. Klaus Empen, Prof. Dr. med. Marcus Dörr, Prof. Dr. med. Stephan B. Felix in issue 17/2015

Which Patients Will Benefit?

The special topic of cardiogenic shock associated with ST-elevation myocardial infarction (STEMI) requires a more detailed explanation.

In spite of the results of the SCHOCK-I study, which showed a clear survival advantage at 6 months and 1 year after emergency revascularization (1), hospital mortality on the basis of multi-organ failure remains high in this group of patients, at 50%, in spite of the brief therapeutic window recommended by guidelines (door-to-balloon-time) of below 90 minutes (2). It remains unclear which patients actually benefit from early revascularization.

For this reason, and independently of the etiology of the cardiogenic shock, the focus should primarily be on restoring adequate hemodynamics and thus peripheral organ perfusion. Once all conservative measures have been exhausted, the timely use of temporary mechanical circulation support is crucial. Veno-arterial extracorporeal membrane oxygenation (ECMO) is used routinely as bridge-to-decision therapy these days. In order to increase the likelihood of myocardial recovery it is important to additionally relieve the left ventricle of volume. To this end, either an intra-aortic balloon pump is implanted contralaterally or, in transthoracically applied ECMO, an additional drain is placed in the left ventricle. The ECMO treatment is continued for seven days. The decision about further therapeutic procedures can then be made depending on the patient's condition.

According to the insights gained from the SCHOCK-I study, the isolated use of an intra-aortic balloon pump does not affect patients' survival and is no longer recommended (3).

DOI: 10.3238/arztebl.2016.0039a

REFERENCES

1. Hochman JS, Sleeper LA, White HD, et al.: One-year survival following early revascularization for cardiogenic shock. *JAMA* 2001; 285: 190–2.
2. Meneses DS, Peterson ED, Wang Y, et al.: Door-to-balloon time and mortality among patients undergoing primary PCI. *N Engl J Med* 2013; 369: 901–9.
3. Thiele H, Zeymer U, Neumann FJ, et al.: Intraaortic Balloon support for myocardial infarction with cardiogenic shock. *N Engl J Med* 2012; 367: 1287–96.
4. Hummel A, Empen K, Dörr M, Felix SB: De novo acute heart failure and acutely decompensated chronic heart failure. *Dtsch Arztebl Int* 2015; 112: 298–310.

Dr. med. Sotirios Spiliopoulos

Prof. Dr. med. Dr. h.c. Reiner Körfer

Prof. Dr. med. Gero Tenderich

Evangelisches Krankenhaus Duisburg

Klinik für Herz- und Gefäßchirurgie

Abteilung für die chirurgische Therapie der terminalen Herzinsuffizienz und

Kunstherzversorgung

Sotirios.Spiliopoulos@evkdn.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

The Renaissance of Digitoxin Is Overdue

A 12-page review article on heart failure is very commendable (1). However, it is a source of irritation that in such a comprehensive article, a total of six lines is all that is given to the positive inotropic digitalis glycosides. A distinction between digoxin and digitoxin was not provided. Positive experiences have been reported for a long time for digitoxin (made from *Digitalis purpurea*). Some 50 years ago, digitoxin (oral, i.v.) and strophanthin (i.v.) were used in clinical practice on a daily basis. When clinical pharmacologists then claimed that digoxin (made from *Digitalis lanata*) was more easily controllable—its saturation dosage is lower and it is excreted more rapidly—digoxin was seen as more modern and displaced digitoxin almost completely (2). This however, meant that right from the beginning no attention was given to the fact that clinical pharmacologists' data came from young students, whereas usually heart failure occurs in old age. Renal function decreases physiologically in old persons. Digoxin is excreted via the kidneys, which means that in this setting there is a risk of cumulation with symptoms of toxicity, such as heart arrhythmia. By contrast, digitoxin is excreted via the liver and does not cumulate in older persons. In case of doubt, concentrations can easily be measured. As might have been expected, clinical experiences with digoxin were disappointing. The indications were reduced and are currently restricted to absolute tachyarrhythmia. However, study results are increasingly reporting higher lethality for digoxin for this indication too, with reported increases of up to 27% (3). A digitoxin study is being conducted at Hanover Medical School. I am certain that this will confirm the positive effect of digitoxin and thus confirm that the preference for digoxin is an aberration. A renaissance of digoxin is overdue, in the interest of the many affected patients.

DOI: 10.3238/arztebl.2016.0039b

REFERENCES

1. Hummel A, Empen K, Dörr M, Felix SB: De novo acute heart failure and acutely decompensated chronic heart failure. *Dtsch Arztebl Int* 2015; 112: 298–310.
2. Overbeck P: Vorhofflimmern: Digoxin gerät erneut unter Verdacht. *Ärzte Zeitung* 2014; 33: 10.
3. Overbeck P: Digitalis-Therapie bei Vorhofflimmern: Auf ewig unter Verdacht? *Ärzte Zeitung* 2015; 34: 2 und 8.

Prof. Dr. med. habil. Christian Tauchnitz

Mikrobiologie und Innere Medizin, Leipzig

tauchnitz.christian@web.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Rapid Application Is Crucial

In the section on mechanical circulatory support, the authors mention the option of using ventricular assist devices (VADs) or extracorporeal membrane oxygenation (ECMO/ECLS) only if the cause of the cardiogenic shock is potentially reversible (1).

The presence of a reversible cause is not crucial for the treatment with mechanical circulatory support (MCS) systems. On

the contrary, we wish to suggest making the decision in favor of implantation on the basis of a patient's prognosis once heart failure has been removed from the scenario. If cardiac function improves under MCS then the system can be explanted without any substitute. If the patient's general condition stabilizes but his/her heart function remains unsatisfactory (in a scenario of an otherwise acceptable prognosis!) a change to a permanent VAD (or even heart transplantation) is possible.

What is crucial is that mechanical support is administered as rapidly as possible, as the success rate of such systems decreases in tandem with the increasing extent of the shock (2).

Acutely decompensated chronic heart failure is a primary focus of the article. In this context the role of MCS should be explicitly mentioned. According to the ESC guidelines, VAD implantation is indicated if a patient who receives adequate medication recurrently decompensates and, in addition to left ventricular heart failure, is found to have impaired right ventricular function and secondary organ function impairments (reduced liver or kidney function) (3). We see this constellation often in our clinical practice, with patients not being informed about the option of VAD implantation. If the device is implanted only in the context of hemodynamic decompression, the chance of success is clearly lower (4).

DOI: 10.3238/arztebl.2016.0039c

REFERENCES

1. Hummel A, Empen K, Dörr M, Felix SB: De novo acute heart failure and acutely decompensated chronic heart failure. *Dtsch Arztebl Int* 2015; 112: 298–310.
2. Lebreton G, Pozzi M, Mastroianni C, Léger P, Pavie A, Leprince P: Extracorporeal life support as a bridge to bridge: a strategy to optimize ventricular assist device results. *Eur J Cardiothorac Surg* 2015; doi:10.1093/ejcts/ezu516.
3. McMurray JJ, Adamopoulos S, Anker SD, et al.: ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The task force for the diagnosis and treatment of acute and chronic heart failure 2012 of the European Society of Cardiology. developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2012; 33: 1787–847.
4. Kirklin JK, Naftel DC, Pagani FD, et al.: Sixth INTERMACS annual report: A 10 000-patient database. *J Heart Lung Transplant* 2014; 33: 555–64.

Dr. med. Felix Hennig

PD Dr. med. Evgenij V. Potapov

Prof. Dr. med. Volkmar Falk

Prof. Dr. med. Thomas Krabatsch

on behalf of the MCS-Team Deutsches Herzzentrum Berlin
fhennig@dhzb.de

Conflict of interest statement

Dr Potapov has been reimbursed conference delegate fees by HeartWare and travel and hotel expenses from HeartWare and Thoratec. He has received study support from Berlin Heart, HeartWare, and Thoratec for a research project that he himself initiated.

Prof. Falk has received consultancy fees from Berlin Heart and HeartWare. He has been reimbursed conference delegate fees and travel and hotel expenses from Berlin Heart, HeartWare, Edwards, Medtronic, and St Jude Medical. He has received study support from Philips and Siemens for a research project that he himself initiated. He has received funding from Berlin Heart and HeartWare for carrying out clinical studies.

Dr Hennig and Prof. Krabatsch declare that no conflict of interest exists.

In Reply:

For a long time, there have been indications of excess mortality associated with the use of digoxin (1), albeit when used in stable chronic heart failure. In patients in the acute phase of heart failure, digitalis preparations are given primarily not for their inotropic effects but because of their negative dromotropic

effects in absolute tachyarrhythmia, so their range of application in this setting has been restricted. In progressive chronic heart failure of NYHA stages III and IV, digitalis preparations do, however, have a positive effect on symptom improvement as well as in terms of reducing rehospitalizations and can be considered on an individual basis.

Further developments and increasing experience in using temporary mechanical circulation support systems allow their use in different decision scenarios in cardiogenic shock—for example, as bridge-to-diagnostics, bridge-to-bridge, bridge-to-recovery, and bridge-to-transplantation. We thank Spiliopoulos et al. and Hennig et al. for pointing out, firstly, the timely use in order to prevent irreversible multiorgan failure in cardiogenic shock, and, secondly, the need for circulatory support before, during and after coronary revascularization after infarct-associated cardiogenic shock. Established or promising areas of use for ECMO are, furthermore, fulminant myocarditis, shock after massive pulmonary embolism, therapy-refractory ventricular tachycardias or atrial fibrillation, and support to recovery after heart transplantations and after heart surgery. However, mortality in individual areas of application remains high, at 58% (2). Factors with a poor prognosis even when ECMO therapy is administered are chronic renal failure, pre-existing multiorgan failure including long periods of ventilation, congenital heart diseases, low-amplitude pulses, and low serum bicarbonate concentrations, as well as prior cardiopulmonary resuscitation. Scores have been suggested for the purpose of assessing the survival probability when using ECMO, but these require further evaluation (2). Similarly, large randomized controlled studies that characterize the optimal patient population for the use of ECMO are lacking, as are cost-benefit analyses of this intervention. In spite of mobile ECMO teams the systems can currently not be offered to all patients because they are not ubiquitously available in a timely fashion (3). Mechanical support systems are a promising solitary building block in the total concept of therapy of cardiogenic (infarct-associated) shock (4).

DOI: 10.3238/arztebl.2016.0040

REFERENCES

1. Vamos M, Erath JW, Hohnloser SH: Digoxin-associated mortality: a systematic review and meta-analysis of the literature. *Eur Heart J* 2015; 143. (Epub ahead of print).
2. Schmidt M, Burell A, Roberts L, et al.: Predicting survival after ECMO for refractory cardiogenic shock: the survival after veno-arterial-ECMO (SAVE)-score. *Eur Heart J* 2015; 194. (epub ahead of print)
3. Abrams D, Combes A, Brodie MD: Extracorporeal membrane oxygenation in cardiopulmonary disease in adults. *J Am Coll Cardiol* 2014; 63: 2769–78.
4. Pöss J, Vollert JO, Böhm M, et al.: Infarktbedingter kardiogener Schock. *Kardiologie* 2014; 8: 302–12.
5. Hummel A, Empen K, Dörr M, Felix SB: De novo acute heart failure and acutely decompensated chronic heart failure. *Dtsch Arztebl Int* 2015; 112: 298–310.

Prof. Dr. med. Stefan B. Felix, Dr. med. Astrid Hummel,

Dr. med. Klaus Empen; Prof. Dr. med. Marcus Dörr

Universitätsmedizin Greifswald, Klinik und Poliklinik für Innere Medizin B, Greifswald
innereb@uni-greifswald.de

Conflict of interest statement

Prof. Felix has served as a paid consultant for Novartis and Cardioventris. He has received reimbursement of congress participation fees and lecture honoraria relating to continuing medical education events from Novartis, Bayer, Berlin Chemie, Cardioventris, and Servier. He has received honoraria in a third-party account for carrying out clinical studies relating to the topic of this article, as well as financial support for a research project that he initiated, from Bayer, Novartis, and Medtronic.

Prof. Empen, Prof. Dörr, and Dr. Hummel state that they have no conflict of interest.