Published in final edited form as:

Psychol Med. 2016 March ; 46(4): 865–875. doi:10.1017/S0033291715002378.

Personality diatheses and Hurricane Sandy: effects on post-disaster depression

D. C. Kopala-Sibley^{1,*}, R. Kotov^{1,2}, E. J. Bromet², G. A. Carlson², A. P. Danzig¹, S. R. Black¹, and D. N. Klein¹

¹Psychology Department, Stony Brook University, Stony Brook, NY, USA

²Psychiatry Department, Stony Brook University, Stony Brook, NY, USA

Abstract

Background—According to diathesis–stress models, personality traits, such as negative emotionality (NE) and positive emotionality (PE), may moderate the effects of stressors on the development of depression. However, relatively little empirical research has directly examined whether NE and PE act as diatheses in the presence of stressful life events, and no research has examined whether they moderate the effect of disaster exposure on depressive symptoms. Hurricane Sandy, the second costliest hurricane in US history, offers a unique opportunity to address these gaps.

Method—A total of 318 women completed measures of NE and PE 5 years prior to Hurricane Sandy. They were also assessed for lifetime depressive disorders on two occasions, the latter occurring an average of 1 year before the hurricane. Approximately 8 weeks after the disaster (mean = 8.40, *s.d.* = 1.48 weeks), participants completed a hurricane stress exposure questionnaire and a measure of current depressive symptoms.

Results—Adjusting for lifetime history of depressive disorders, higher levels of stress from Hurricane Sandy predicted elevated levels of depressive symptoms, but only in participants with high levels of NE or low levels of PE.

Conclusions—These findings support the role of personality in the development of depression and suggest that personality traits can be useful in identifying those most vulnerable to major stressors, including natural disasters.

Keywords

Depression; natural disasters; negative emotionality; personality; positive emotionality

*Address for correspondence: D. C. Kopala-Sibley, Ph.D., Stony Brook University, Stony Brook, NY, USA. (daniel.kopala-sibley@mail.mcgill.ca).

Declaration of Interest
None.

Introduction

According to diathesis–stress models, personality may moderate the effects of stressors on the development of psychopathology (Monroe & Simons, 1991; Blatt & Zuroff, 1992; Ingram & Price, 2010). Hurricane Sandy, the second costliest hurricane in US history, hit Long Island, New York as a category 1 hurricane with a record 14-foot (4.25 m) storm surge on 29 October 2012, destroying 100 000 homes. Natural disasters, such as Sandy, offer a unique opportunity to test diathesis–stress models. Studies of mental disorders after hurricanes, including Hurricane Sandy, find increased levels of depressive symptoms post-hurricane (e.g. Norris *et al.* 1999, 2002; Kessler *et al.* 2006; Bonanno *et al.* 2010; Neria & Shultz, 2012; Boscarino *et al.* 2013; North & Pfefferbaum, 2013). However, it is unknown why some individuals and not others became symptomatic.

Much literature has suggested that individual differences in personality traits, such as negative emotionality (NE) or the very closely related construct of neuroticism, and positive emotionality (PE) or the similar construct of extraversion, may respectively confer vulnerability to, or resilience against, depression (e.g. Bagby *et al.* 1997; De Fruyt *et al.* 2006; Kotov *et al.* 2010; Klein *et al.* 2011; Ormel *et al.* 2013; Wilson *et al.* 2014). NE/neuroticism is characterized by a general tendency to experience fear, anger and sadness, and a susceptibility to the effects of stress on mood. PE/extraversion is characterized by a general tendency to experience positive emotions (e.g. joy, exuberance), and being gregarious and engaged with the environment (Watson *et al.* 2006). Both NE/neuroticism and PE/extraversion are higher-order factors in all Big Three (Eysenck, 1991; Watson *et al.* 1999) and Big Five (McCrae & Costa, 1999) personality models.

Despite high NE/neuroticism and low PE/extraversion being conceptualized as vulnerabilities to psychopathology, only a few longitudinal studies have examined whether these traits act as diatheses that increase sensitivity to life stress (Ormel *et al.* 2001; Kendler *et al.* 2004; Hutchinson & Williams, 2007; Brown & Rosellini, 2011; Vinkers *et al.* 2014), and no research of which we are aware has tested whether they moderate the impact of natural disasters on depressive symptoms. The current study examined whether the personality traits of NE and PE predicted the occurrence of depressive symptoms following Hurricane Sandy and moderated the effects of exposure to stress from Sandy on subsequent depressive symptoms.

Impact of life stress and natural disasters on depression

There is considerable evidence that stress plays a causal role in the onset of depression (for reviews, see Kessler, 1997; Tennant, 2002; Hammen, 2005; Stroud *et al.* 2008; Monroe *et al.* 2014). Much of this research has focused on relatively common hassles (e.g. arguments with a loved one) or major events (e.g. the death of a loved one). In parallel, a substantial literature shows that exposure to natural disasters, including hurricanes, confers risk for depressive symptoms (e.g. Norris *et al.* 1999, 2002; Acierno *et al.* 2006, 2007; Weisler *et al.* 2006; Bonanno *et al.* 2010; North & Pfefferbaum, 2013).

However, regardless of the nature of the stressor, most persons do not experience a significant increase in symptoms, with stress typically accounting for only a modest

percentage of variance in depressive symptoms (Monroe, *et al.* 1986; Coyne & Downey, 1991; Norris *et al.* 2002; Tennant, 2002; Hammen, 2005). The considerable heterogeneity in individual responses to stress suggests that personal characteristics may moderate the effect of stress on depression. These findings have prompted research into how personality or cognitive individual differences confer vulnerability to depression in the presence of stressful life events (e.g. Hammen *et al.* 1985; Blatt & Zuroff, 1992; Segal *et al.* 1992; Ingram *et al.* 1998; Clark & Beck, 1999).

Personality diatheses and vulnerability to depression

Much literature has supported the possibility that personality or cognitive characteristics, such as self-criticism and dependency (Blatt & Zuroff, 1992; Kopala-Sibley & Zuroff, 2010, 2014), sociotropy and autonomy (Beck *et al.* 1983; Clark & Beck, 1999), hopelessness/helplessness (Rose & Abramson, 1992; Iacoviello *et al.* 2013), and perfectionism (Hewitt & Flett, 1991), increase the effects of stress on depression. Other evidence has shown that the personality trait of hardiness buffers the impact of disaster stress on psychiatric symptoms (Bonanno, 2004). Similarly, the personality traits of NE/ neuroticism and PE/extraversion (see Watson *et al.* 2005) have received substantial attention as risk for, or protective factors against, depression, with high levels of NE/neuroticism and low levels of PE/extraversion being hypothesized as key vulnerabilities (e.g. Enns & Cox, 1997; Gershuny & Sher, 1998; Bienvenu *et al.* 2004; Watson *et al.* 2005; Brown, 2007; Verstraeten *et al.* 2009; Kotov *et al.* 2010; Klein *et al.* 2011).

However, the vast majority of studies of PE/extraversion and NE/neuroticism have examined only their main effects on the onset or course of depressive symptoms or diagnoses (for reviews, see Kotov *et al.* 2010; Klein *et al.* 2011; Ormel *et al.* 2013). We are aware of only a handful of studies to date which have tested whether NE/neuroticism exacerbates the effects of stress on depression, and even fewer have examined whether PE/extraversion buffers the impact of stress. Across periods of several weeks to 1 year, evidence generally suggests that the effect of stress on subsequent depressive symptoms is greater in individuals with high levels of NE/neuroticism (Ormel *et al.* 2001; Kendler *et al.* 2004; Hutchinson & Williams, 2007; Brown & Rosellini, 2011; Vinkers *et al.* 2014), although some evidence has been less supportive (Spinhoven *et al.* 2011). Regarding PE/extraversion, the only study (Spinhoven *et al.* 2011) of which we are aware did not find that extraversion buffered the effect of stress on depression. However, other evidence has found that positive emotional states, as opposed to trait-levels of PE, may buffer the effect of stress on negative affect or depressive symptoms (Fredrickson, 2001; Compas *et al.* 2004; Wichers *et al.* 2007).

All studies to date have examined stressors that are relatively common and have been limited to intervals of 1 year or less between the assessments of personality and stress. As personality is conceptualized as an enduring diathesis but has not been tested as such over an interval of longer than 1 year, measuring traits well in advance of the stressor would provide a more appropriate and conservative test of personality vulnerability models of depression. In addition, examining a natural disaster as the form of stress is particularly advantageous as its occurrence is independent of personality.

Researchers have examined several factors that may moderate the effect of disasters on psychopathology, with evidence suggesting that more severe exposure, female gender, middle age, ethnic minority status, prior psychiatric problems, and weak or deteriorating psychosocial resources increase the likelihood of adverse outcomes following a natural disaster (Norris *et al.* 2002; Bonanno *et al.* 2010; North & Pfefferbaum, 2013). However, no research has attempted to integrate the personality diathesis literature with the literature on natural disasters and psychopathological outcomes.

Overview and hypotheses

This paper examined whether the personality traits of NE/neuroticism and PE/extraversion predict levels of depressive symptoms following Hurricane Sandy and tested whether they moderate the effects of hurricane-related stress on depressive symptoms shortly following the disaster. Given the prior literature demonstrating the effects of various demographic and psychiatric factors on outcomes following natural disasters, all analyses adjusted for participants' age, race/ethnicity, marital status, education, household income, and lifetime history of depressive disorders. We hypothesized that higher levels of stress, higher levels of NE/ neuroticism and lower levels of PE/extraversion would be related to higher levels of depressive symptoms after the hurricane. We also hypothesized that the effect of hurricane-related stress on depressive symptoms would be stronger for those high in NE/neuroticism and weaker for those high in PE/extraversion.

Method

Procedure and participants

Our effective sample comprised 318 mothers. Between October 2004 and June 2007, 559 families with a 3-year-old child were recruited to participate in a longitudinal study of temperament. Specifically, lists of families with 3-year-old children who lived within a 20-mile (32 km) radius of Stony Brook University in Long Island, New York, were purchased from a commercial mailing list broker. Commercial mailing lists are available to those who wish to conduct marketing campaigns via direct mail or, in our case, research projects. The sample included families living in a contiguous 20-mile radius of Stony Brook University. Potential participants were screened to select families with at least one English-speaking biological parent and a 3-year old child with no significant medical problems or developmental disabilities. At wave 2, conducted when the children were 6 years old, 50 families were added to increase the demographic diversity of the sample, yielding a total sample of 609 pairs. This time point is our baseline assessment for the current study. Wave 3 was conducted between September 2010 and April 2013, when the children were 9 years old. In all, 446 mothers participated at this time point, and had completed their assessments prior to 29 October 2012, when Hurricane Sandy struck.

At 6 weeks after the hurricane, the 446 mothers were invited to complete an on-line survey about the disaster. A total of 362 of the 446 mothers (81.2%) participated. For the current analysis, we excluded 15 mothers who had moved to other parts of the country, and were therefore outside the hurricane zone as well as 29 mothers with incomplete data on the key measures used in the analysis. Thus, the final analysis sample included 318 mothers, 96.0%

of whom lived in Federal Emergency Management Agency (FEMA)-declared disaster zones (see AP Danzig *et al.* unpublished observations).

Most mothers (96.5%) completed the on-line survey 7–11 weeks after Hurricane Sandy (mean = 8.4, *s.d.* = 1.48 weeks). There were no significant differences between the Hurricane Sandy participants and either non-participants or the full sample on the measures included in this report (all *p*'s > 0.20). Demographically, the 318 mothers had a mean age of 36.1 (*s.d.* = 4.5) years, an average income of about \$100 000, and some college education (90.9%). Most were married (>85%) and non-Hispanic Caucasian (83.02%).

The study was approved by the Stony Brook University Institutional Review Board.

Materials

Exposure to Hurricane Sandy stress—Mothers were asked to indicate to what extent their family was affected by 13 Hurricane Sandy stressors/ exposures (Table 1; for details, see AP Danzig *et al.* unpublished observations). Participants were asked to complete this measure at their earliest, and did so an average of 8.4 (*s.d.* = 1.48) weeks post-Sandy. The 13 items were drawn primarily from questionnaires administered in studies of Hurricane Ike (Norris *et al.* 2010) and Hurricane Katrina (Galea *et al.* 2007). Eight exposures were rated on a 1 (not at all affected) to 5 (extremely affected) scale: life disrupted by the hurricane, children fearing for their own or their family's safety, difficulty finding food or warmth, difficulty finding gasoline, children complaining more than usual, damage to home or possessions, family's safety threatened, and financial hardship. These items were dichotomized into present *v.* absent for ease of interpretation and analysis: the impact typically had to be rated as 'quite a bit' or 'extremely' for the event to be coded as present. Exceptions to this were made for items that were very rare (i.e. financial hardship, where 'moderate' ratings were included) and highly endorsed (i.e. difficulty finding gasoline, where only 'extreme' ratings were included) to capture an appropriate threshold for exposure. Two items, time school was closed and time without power, were rated on a five-point duration scale ranging from 0 days to 2 weeks or more, and the stressor was considered 'present' when lasting for 7 or more days. Three stressors were rated yes/no: having to evacuate the home; applying for FEMA assistance; and members of the immediate family, other relatives or friends and/or pets being either injured, victimized, robbed or lost. A continuous 'total stressors' variable was computed by summing the 13 dichotomous stressors (see Table 1).

Depressive symptoms post-Sandy—Depressive symptoms following Hurricane Sandy were assessed with the Diagnostic Inventory for Depression (DID) (Zimmerman *et al.* 2004), a 22-item measure designed to assess Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) criterion symptoms of major depressive disorder (MDD) occurring in the past week. The DID correlates well with diagnoses of MDD based on the Structured Clinical Interview for the DSM-IV (SCID), discriminates the likelihood of MDD *v.* other disorders, and is sensitive to clinical change in depression levels (Zimmerman *et al.* 2004). Each question contains five response options, with higher scores indicating greater levels of each symptom. Scores from each item are summed, with higher total scores

indicating higher levels of depressive symptoms. An example item includes ‘During the past week, have you been feeling sad or depressed?’ with response options ranging from 0 (‘no, not at all’) to 4 (‘yes, I have been extremely sad or depressed nearly all the time’). In the current study, coefficient α for the DID was 0.86, and scores ranged from 0 to 47.

PE and NE—Mothers completed the General Temperament Survey (GTS; Clark & Watson, 1992) during the wave 2 assessment when children were 6 years old. The GTS is a 43-item factor-analytically derived measure of the Big Three personality dimensions, namely, PE and NE and disinhibition. The current paper focuses on NE and PE, which are vulnerability factors for depression. Conversely, disinhibition was not included because it is generally not considered as a vulnerability factor for depressive symptoms (Klein *et al.* 2011). NE and PE are clearly distinct from each other (Clark & Watson, 1992; Watson *et al.* 2005), and are strongly related in expected directions with measures of state positive and negative affect (Watson & Clark, 1992). NE correlates highly with measures of neuroticism, and PE correlates highly with measures of extra-version (Watson *et al.* 2005). High NE and low PE are robustly related to a range of psychopathological symptoms, including depression (Watson *et al.* 2005). Both scales also show excellent internal consistency (Watson & Clark, 1992). In the current study, internal consistencies were 0.81 and 0.82 for NE and PE, respectively.

Lifetime history of depressive disorders—History of depressive disorders (MDD and dysthymic disorder) was determined from the non-patient version of the SCID (Williams *et al.* 1992; First *et al.* 1996). The SCID was administered at two time points: initial assessment at wave 1 (lifetime version) and interval assessment at wave 3 for the prior 6 years. Lifetime depressive disorder was considered positive if a diagnosis was present at either assessment point. The SCID is among the most widely used diagnostic interviews, and its inter-rater reliability and validity have been well documented (Williams *et al.* 1992). The SCID interviews were conducted by master’s-level clinicians and clinical psychology graduate students. Inter-rater reliability was excellent ($\kappa = 0.93$ at wave 1 and 0.91 at wave 3). In the current sample, 117 (36.8%) participants met criteria for a lifetime depressive disorder.

Data analyses

Data were analysed in SAS 9.3 (USA). Primary analyses used multiple linear regression. Given the moderate correlation between NE and PE ($r = -0.39$, $p < 0.001$), we were concerned that their mutual effects on depressive symptoms might impair statistical power or result in suppression effects. We therefore tested two separate regression models in which either NE or PE moderated the effect of Hurricane Sandy stress on post-Sandy depressive symptoms. All analyses adjusted for participants’ age, race/ethnicity (Caucasian and non-Hispanic *v.* non-Caucasian and/or Hispanic), marital status (married *v.* not), level of education (graduated 4-year college *v.* not), household income, and lifetime history of depressive disorders. In each model, the main effect of lifetime depressive disorder history was entered first, followed by the demographic covariates listed above, followed by the main effects of exposure and one personality variable (NE or PE). The interaction between exposure and personality was subsequently included. Non-categorical predictor variables

were centered (Aiken & West, 1991). Interactions were interpreted by comparing simple slopes at high and low levels (± 1 s.d.) of the moderator variable. Confidence intervals (CIs) are presented for all significant results.

Results

Descriptive statistics and bivariate correlations

Rates of exposures in the analysis sample are presented in Table 1. The most commonly reported exposures were prolonged school closing, losing electric power, life disruption, and children fearing for their own or their family's safety. The average number of stressors (out of 13) was 2.20 (s.d. = 2.19).

Correlations and descriptive statistics are presented in Table 2. Having had a pre-Sandy depressive episode predicted greater depressive symptoms post-Sandy. Consistent with past research, higher levels of NE and lower levels of PE were related to greater depressive symptoms post-Sandy. Higher levels of exposure to stress from Hurricane Sandy were also related to greater depressive symptoms post-Sandy.

Regression analyses predicting post-Sandy depressive symptoms

Regression results are presented in Table 3. Our first model tested the effects of NE, stress and their interaction on post-Sandy depressive symptoms. A lifetime history of depressive disorder, lower household income, and higher NE predicted greater depressive symptoms post-Sandy. In addition, there was a significant interaction between NE and Sandy-related stress (Fig. 1). The effect of exposure on depressive symptoms was significant at high levels of NE ($\beta = 1.21$, $t = 2.79$, $p < 0.01$, 95% CI 0.15–1.82), but not at low levels ($\beta = -0.21$, $t = -0.49$, $p = 0.63$, 95% CI –1.23 to 0.97). The full model explained 25.12% of variance in post-Sandy depressive symptoms.

In our second model, a lifetime history of depressive disorders, lower income, lower PE and greater hurricane-related stress predicted higher depressive symptoms post-Sandy. Additionally, there was a significant interaction between PE and exposure (Fig. 2). The effect of stress on depressive symptoms was significant at low ($\beta = 1.75$, $t = 3.85$, $p < 0.001$, 95% CI 0.86–2.64), but not high PE ($\beta > 0.34$, $t = 0.86$, $p = 0.39$, 95% CI –0.43 to 1.11). The full model explained 23.27% of variance in post-Sandy depressive symptoms^{1†}.

Discussion

We conducted a novel test of diathesis–stress models by examining whether NE and PE exacerbated or buffered the effect of stress caused by a natural disaster, Hurricane Sandy, on subsequent depressive symptoms. Adjusting for participants' history of depressive disorders, as well as age, race/ethnicity, marital status, education and household income, we found that both PE and NE, assessed 3–5 years prior to Hurricane Sandy, moderated the effects of disaster exposure on subsequent depressive symptoms. Specifically, greater stress

¹Analyses were repeated using a continuously scored stress measure (i.e. item scores were summed, rather than dichotomized and counted). The overall pattern of results remained similar, with a significant interaction between PE and stress ($p = 0.05$), and a trend towards an interaction between NE and stress ($p = 0.08$).

significantly predicted greater depressive symptoms in those with low levels of PE, but not in those who showed high levels of PE. In addition, the effect of hurricane-related stress was significant for those high in NE, but not for those low in NE. Taken together, these results support the role of PE as a protective factor against, and NE as a vulnerability factor for, depressive symptoms following exposure to a natural disaster. Such personality diathesis x stress effects have been frequently assumed but rarely tested.

NE/neuroticism, disaster stress and depressive symptoms

The significant main effect of NE supports the contention that NE/neuroticism contributes to the development of depressive symptoms (Watson *et al.* 2005; Kotov *et al.* 2010; Klein *et al.* 2011; Ormel *et al.* 2013). The interaction of NE and stress is also consistent with most of the limited literature available on this topic (Ormel *et al.* 2001; Kendler *et al.* 2004; Hutchinson & Williams, 2007; Brown & Rosellini, 2011; Vinkers *et al.* 2014). However, our findings extend this literature by demonstrating that NE moderates the effects of stressors that occur up to 5 years later, and that this personality characteristic interacts not only with common stressors such as job loss or interpersonal conflict, but also with stress ensuing from a natural disaster.

The reasons for why or how NE/neuroticism moderates the effects of disaster stress on depressive symptoms were not examined in the current study, but may be a fruitful area for future research. For instance, the emotional lability that is characteristic of high NE/neuroticism may deplete individuals' cognitive resources, resulting in difficulties with executive functioning and self-regulation in the face of major stressors (Gomez *et al.* 1999). Individuals high on NE/neuroticism may also view negative events as more stressful or more hopeless than they really are (Chioqueta & Stiles, 2005), employ less problem-focused and more avoidant coping methods (Gomez *et al.* 1999; Compas *et al.* 2004), and be less likely to seek out or receive instrumental social support (Borja *et al.* 2009), all of which may ultimately result in greater depressive symptoms post-disaster.

PE/extroversion, disaster stress and depressive symptoms

Results regarding PE support the established relationship between higher levels of PE/extraversion and lower levels of depressive symptoms (Watson *et al.* 2005; Kotov *et al.* 2010; Klein *et al.* 2011). In contrast to the effects of NE, PE protected against the effects of Hurricane Sandy exposure in that there was no effect on depressive symptoms in individuals high in this trait. There is a relative paucity of literature examining PE/extraversion from a diathesis–stress perspective. The one study of which we are aware was not supportive (Spathov *et al.* 2011), although this may be due to its use of a clinical, rather than community, sample. Our results highlight the adaptive nature of PE by showing that it mitigates the effect of disaster-related stress on depressive symptoms.

As with NE/neuroticism, the mechanisms underlying the protective effect of PE/extraversion were not examined in this study. Individuals high in PE/extraversion may benefit from a greater ability to make decisions and plans under stressful situations given their greater cognitive resources (Gomez *et al.* 1999). They may also view difficult situations more optimistically (Duberstein *et al.* 2001; Chioqueta & Stiles, 2005) and be more likely to

solicit more instrumental social support. Broaden-and-build models of positive affect (e.g. Fredrickson, 2001) suggest that the elevated level of positive affect characteristic of individuals high in PE/extraversion may facilitate thinking of novel and adaptive ways to cope with stress from a natural disaster. Future research will probably benefit from examining these and other potential mediators of the stress-exacerbating effects of NE/neuroticism and stress-mitigating effects of PE/extraversion.

Our results are also consistent with prior evidence of increased depressive symptoms in mothers following natural disasters (e.g. Adams *et al.* 2011; Goto *et al.* 2015), as well as evidence that mothers, relative to women without children, are particularly at risk for depression following a natural disaster (Havenaar *et al.* 1997). Our results therefore emphasize the importance of interventions geared towards mothers with high NE/neuroticism or low PE/extraversion who are caring for children, as they may be particularly at risk in the context of major stressors, such as natural disasters.

Finally, our results support vulnerability models of personality (Clark & Beck, 1999; Compas *et al.* 2004; Klein *et al.* 2011) in which traits place individuals at risk for psychopathology. However, our results cannot speak to other models of the personality–depression relationship, such as stress generation (e.g. Hammen, 2006), scar, or pathoplasticity (see Compas *et al.* 2004; Klein *et al.* 2011), although our findings of trait x stress interactions are difficult to reconcile with spectrum/continuum models (Clark & Beck, 1999; Klein *et al.* 2011).

Above all, however, the current results support the role of these relatively stable personality traits as factors which moderate the effect of stress on depression, even when measured 3 to 5 years prior to the stress. These results support the utility of assessing the traits of PE/extraversion and NE/neuroticism in order to identify those most vulnerable to major stressors, including natural disasters.

Limitations and future directions

Although this study had several notable strengths, including a 5-year longitudinal design and assessment of personality and lifetime history of depressive disorders prior to Hurricane Sandy, several limitations should be noted. Perhaps the most substantial concern is that depressive symptoms were assessed at the same time as stress from Hurricane Sandy. As such, it is possible that individuals with elevated levels of depressive symptoms were more likely to report greater stress from Sandy (e.g. recall bias). Our assessments of stress and depressive symptoms were also both self-report; it is not clear whether our results would extend to clinicians' ratings of depressive symptomology or diagnoses, or whether interviews with the mothers and other informants would have provided a more accurate picture of participants' experiences during Hurricane Sandy. However, the DID converges well with clinician ratings of depressive symptoms severity (Zimmerman *et al.* 2004). Moreover, although these biases may inflate personality–depression associations, they are less likely to account for the interactions between personality and stress.

Another concern is that our measure of stress primarily reflected events that happened to participants' families; we cannot be sure if events which were specific to the women in this

study, as opposed to their family, would show similar interactions with personality. To the extent that this concern is a factor, it probably reduced effects of exposure, making our estimates conservative. Similarly, because we aimed to assess exposure to stress from Sandy as quickly as possible after the hurricane, we were unable to determine whether there were other stressors in participants' lives that were either unrelated, secondary to, or ongoing from Sandy, but may have compounded its effects or resulted in additional enduring stress post-Sandy (e.g. Galea *et al.* 2007; Kessler *et al.* 2008). However, it is unlikely that very many major unrelated stressors occurred in the relatively short time period between Sandy and our assessment. As well, our measure of stress from Sandy included stressors that were likely to be ongoing for a significant period of time (e.g. damage to home, evacuation, financial hardship).

It is also unclear how long the effects will persist after Hurricane Sandy. In addition, the most commonly reported stressors from Hurricane Sandy could arguably be considered to be hassles, rather than major stressors; that is, we cannot know from these data whether similar effects would be found in an even more serious disaster in which, for instance, most participants lost their homes, or there was a high rate of death and injury. If anything, the effects of such disasters are likely to be larger. Furthermore, while adjusting for participants' lifetime depressive history is quite conservative, due to the impromptu nature of natural disasters, we were unable to assess symptoms immediately prior to Sandy. Finally, due to the time-sensitive nature of these data, we were unable to assess other psychiatric symptoms, such as post-traumatic stress disorder (PTSD) or anxiety, and do not know if the results would extend to other symptoms, or if adjusting for other symptoms would affect our results, given some evidence that neuroticism moderates the effect of stress on PTSD symptoms (Breslau & Schultz, 2013).

Conclusion

This study provided a novel test of diathesis–stress models by showing that the personality traits of PE/extraversion conferred resilience, while NE/neuroticism conferred vulnerability, to the effects of exposure to a natural disaster. Results further our understanding of the role of personality in vulnerability to depression, shedding light on the origins of the robust links between these variables observed in cross-sectional studies. Such an understanding is probably critical in future efforts to identify those at risk for psychopathology prior to disorder onset as they can be targeted for additional support and preventive interventions.

Acknowledgements

The present study was supported by National Institute of Mental Health (NIMH) grant RO1 MH45757 and NIMH grant R01 MH093479.

References

- Acierno R, Ruggiero KJ, Galea S, Resnick HS, Koenen K, Roitzsch J, de Arellano M, Boyle J, Kilpatrick DG. Psychological sequelae resulting from the 2004 Florida hurricanes: implications for postdisaster intervention. *American Journal of Public Health*. 2007; 97(Suppl. 1):S103–S108. [PubMed: 17413067]

- Acierio R, Ruggiero KJ, Kilpatrick DG, Resnick HS, Galea S. Risk and protective factors for psychopathology among older *versus* younger adults after the 2004 Florida hurricanes. *American Journal of Geriatric Psychiatry*. 2006; 14:1051–1059. [PubMed: 17035356]
- Adams RE, Guey LT, Gluzman SF, Bromet EJ. Psychological well-being and risk perceptions of mothers in Kyiv, Ukraine, 19 years after the Chernobyl disaster. *International Journal of Social Psychiatry*. 2011; 57:637–645. [PubMed: 21813484]
- Aiken, LS.; West, SG. *Multiple Regression: Testing and Interpreting Interactions*. Newbury Park, CA: Sage; 1991.
- Bagby RM, Bindseil KD, Schuller DR, Rector NA, Young LT, Cooke RG, Seeman MV, McCay EA, Joffe RT. Relationship between the five-factor model of personality and unipolar, bipolar and schizophrenic patients. *Psychiatry Research*. 1997; 70:83–94. [PubMed: 9194202]
- Beck, AT.; Epstein, N.; Harrison, RP.; Emery, G. *Development of the Sociotropy–Autonomy Scale: A Measure of Personality Factors in Depression*. Philadelphia, PA: University of Pennsylvania; 1983.
- Bienvenu OJ, Samuels JF, Costa PT, Reti IM, Eaton WW, Nestadt G. Anxiety and depressive disorders and the five-factor model of personality: a higher- and lower-order personality trait investigation in a community sample. *Depression and Anxiety*. 2004; 20:92–97. [PubMed: 15390211]
- Blatt SJ, Zuroff DC. Interpersonal relatedness and self-definition: two prototypes for depression. *Clinical Psychology Review*. 1992; 12:527–562.
- Bonanno GA. Loss, trauma, and human resilience: have we underestimated the human capacity to thrive after extremely aversive events? *American Psychologist*. 2004; 59:20–28. [PubMed: 14736317]
- Bonanno GA, Brewin CR, Kaniasty K, La Greca AM. Weighing the costs of disaster consequences, risks, and resilience in individuals, families, and communities. *Psychological Science in the Public Interest*. 2010; 11:1–49. [PubMed: 26168411]
- Borja SE, Callahan JL, Rambo PL. Understanding negative outcomes following traumatic exposure: the roles of neuroticism and social support. *Psychological Trauma: Theory, Research, Practice, and Policy*. 2009; 1:118.
- Boscarino JA, Hoffman SN, Kirchner HL, Erlich PM, Adams RE, Figley CR, Solkhah R. Mental health outcomes at the Jersey shore after hurricane Sandy. *International Journal of Emergency Mental Health*. 2013; 15:147–158. [PubMed: 24558743]
- Breslau N, Schultz L. Neuroticism and post-traumatic stress disorder: a prospective investigation. *Psychological Medicine*. 2013; 43:1697–1702. [PubMed: 23199934]
- Brown TA. Temporal course and structural relationships among dimensions of temperament and *DSM-IV* anxiety and mood disorder constructs. *Journal of Abnormal Psychology*. 2007; 116:313–328. [PubMed: 17516764]
- Brown TA, Rosellini AJ. The direct and interactive effects of neuroticism and life stress on the severity and longitudinal course of depressive symptoms. *Journal of Abnormal Psychology*. 2011; 120:844–856. [PubMed: 21381799]
- Chioqueta AP, Stiles TC. Personality traits and the development of depression, hopelessness, and suicide ideation. *Personality and Individual Differences*. 2005; 38:1283–1291.
- Clark, DA.; Beck, AT. *Scientific Foundations of Cognitive Theory and Therapy of Depression*. Wiley: New York; 1999.
- Compas BE, Connor-Smith J, Jaser SS. Temperament, stress reactivity, and coping: implications for depression in childhood and adolescence. *Journal of Clinical Child and Adolescent Psychology*. 2004; 33:21–31. [PubMed: 15028538]
- Coyne JC, Downey G. Social factors and psychopathology: stress, social support, and coping processes. *Annual Review of Psychology*. 1991; 42:401–425.
- De Fruyt F, Van Leeuwen K, Bagby RM, Rolland JP, Rouillon F. Assessing and interpreting personality change and continuity in patients treated for major depression. *Psychological Assessment*. 2006; 18:71–80. [PubMed: 16594814]
- Duberstein PR, Laurent J, Conner KR, Conwell Y, Cox C. Personality correlates of hopelessness in depressed inpatients 50 years of age and older. *Journal of Personality Assessment*. 2001; 77:380–390. [PubMed: 11693865]

- Enns MW, Cox BJ. Personality dimensions and depression: review and commentary. *Canadian Journal of Psychiatry*. 1997; 42:274–284. [PubMed: 9114943]
- Eysenck HJ. Dimensions of personality: 16, 5 or 3? -Criteria for a taxonomic paradigm. *Personality and Individual Differences*. 1991; 12:773–790.
- First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. *The Structured Clinical Interview for DSM-IV Axis I Disorders – Non-Patient Edition*. New York: Biometrics Research Department, New York State Psychiatric Institute; 1996.
- Fredrickson BL. The role of positive emotions in positive psychology. The broaden-and-build theory of positive emotions. *American Psychologist*. 2001; 56:218–226. [PubMed: 11315248]
- Galea S, Brewin CR, Gruber M, Jones RT, King DW, King LA, McNally RJ, Ursano RJ, Petukhova M, Kessler RC. Exposure to hurricane-related stressors and mental illness after Hurricane Katrina. *Archives of General Psychiatry*. 2007; 64:1427–1434. [PubMed: 18056551]
- Gershuny BS, Sher KJ. The relation between personality and anxiety: findings from a 3-year prospective study. *Journal of Abnormal Psychology*. 1998; 107:252–262. [PubMed: 9604554]
- Gomez R, Holmberg K, Bounds J, Fullarton C, Gomez A. Neuroticism and extraversion as predictors of coping styles during early adolescence. *Personality and Individual Differences*. 1999; 27:3–17.
- Goto A, Bromet EJ, Fujimori K. Immediate effects of the Fukushima nuclear power plant disaster on depressive symptoms among mothers with infants: a prefectural-wide cross-sectional study from the Fukushima Health Management Survey. *BMC Psychiatry*. 2015; 15:59. [PubMed: 25885267]
- Hammen C. Stress and depression. *Annual Reviews of Clinical Psychology*. 2005; 1:293–319.
- Hammen C. Stress generation in depression: reflections on origins, research, and future directions. *Journal of Clinical Psychology*. 2006; 62:1065–1082. [PubMed: 16810666]
- Hammen C, Marks T, Mayol A, DeMayo R. Depressive self-schemas, life stress, and vulnerability to depression. *Journal of Abnormal Psychology*. 1985; 94:308–319. [PubMed: 4031228]
- Havenaar JM, Rumyantzeva GM, van den Brink W, Poelijoe NW, van den Bout J, van Engeland H, Koeter MW. Long-term mental health effects of the Chernobyl disaster: an epidemiologic survey in two former Soviet regions. *American Journal of Psychiatry*. 1997; 154:1605–1607. [PubMed: 9356574]
- Hewitt PL, Flett GL. Perfectionism in the self and social contexts: conceptualization, assessment, and association with psychopathology. *Journal of Personality and Social Psychology*. 1991; 60:456–470. [PubMed: 2027080]
- Hutchinson JG, Williams PG. Neuroticism, daily hassles, and depressive symptoms: an examination of moderating and mediating effects. *Personality and Individual Differences*. 2007; 42:1367–1378.
- Iacoviello BM, Alloy LB, Abramson LY, Choi JY, Morgan JE. Patterns of symptom onset and remission in episodes of hopelessness depression. *Depression and Anxiety*. 2013; 30:564–573. [PubMed: 23495016]
- Ingram, RE.; Miranda, J.; Segal, ZV. *Cognitive Vulnerability to Depression*. New York: Guilford Press; 1998.
- Ingram, RE.; Price, JM. Understanding psychopathology: the role of vulnerability. In: Ingram, RE.; Price, JM., editors. *Vulnerability to Psychopathology: Risk Across the Lifespan*. 2nd. New York: Guilford Press; 2010. p. 3–17.
- Kendler KS, Kuhn J, Prescott CA. The interrelationship of neuroticism, sex, and stressful life events in the prediction of episodes of major depression. *American Journal of Psychiatry*. 2004; 161:631–636. [PubMed: 15056508]
- Kessler RC. The effects of stressful life events on depression. *Annual Reviews of Psychology*. 1997; 48:191–214.
- Kessler RC, Galea S, Gruber MJ, Sampson NA, Ursano RJ, Wessely S. Trends in mental illness and suicidality after Hurricane Katrina. *Molecular Psychiatry*. 2008; 13:374–384. [PubMed: 18180768]
- Kessler RC, Galea S, Jones RT, Parker HA. Mental illness and suicidality after Hurricane Katrina. *Bulletin of the World Health Organization*. 2006; 84:930–939. [PubMed: 17242828]
- Klein DN, Kotov R, Bufferd SJ. Personality and depression: explanatory models and review of the evidence. *Annual Review of Clinical Psychology*. 2011; 7:269–295.

- Kopala-Sibley DC, Zuroff DC. Proximal predictors of depressive symptomatology: perceived losses in self-worth and interpersonal domains and introjective and anaclitic mood states. *Cognitive Behaviour Therapy*. 2010; 39:270–282. [PubMed: 20936547]
- Kopala-Sibley DC, Zuroff DC. The developmental origins of personality factors from the self-definitional and relatedness domains: a review of theory and research. *Review of General Psychology*. 2014; 18:137–155.
- Kotov R, Gamez W, Schmidt F, Watson D. Linking “big” personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychological Bulletin*. 2010; 136:768–821. [PubMed: 20804236]
- McCrae, RR.; Costa, PT, Jr. A five-factor theory of personality. In: John, OP.; Robins, RW., editors. *Handbook of Personality: Theory and Research*. New York: Guilford Press; 1999. p. 159-181.
- Monroe SM, Bromet EJ, Connell MM, Steiner SC. Social support, life events, and depressive symptoms: a 1-year prospective study. *Journal of Consulting and Clinical Psychology*. 1986; 54:424–431. [PubMed: 3745594]
- Monroe SM, Simons AD. Diathesis–stress theories in the context of life stress research: implications for the depressive disorders. *Psychological Bulletin*. 1991; 110:406–425. [PubMed: 1758917]
- Monroe, SM.; Slavich, GM.; Georgiades, K. The social environment and depression: the roles of life stress. In: Gotlib, IH.; Hammen, CL., editors. *Handbook of Depression and Its Treatment*. 3rd. New York: Guilford Press; 2014. p. 296-314.
- Neria Y, Shultz JM. Mental health effects of Hurricane Sandy: characteristics, potential aftermath, and response. *Journal of the American Medical Association*. 2012; 308:2571–2572. [PubMed: 23160987]
- Norris FH, Friedman MJ, Watson PJ. 60 000 disaster victims speak: Part II. Summary and implications of the disaster mental health research. *Psychiatry: Interpersonal and Biological Processes*. 2002; 65:240–260.
- Norris FH, Perilla JL, Riad JK, Kaniasty K, Lavizzo EA. Stability and change in stress, resources, and psychological distress following natural disaster: findings from Hurricane Andrew. *Anxiety, Stress and Coping*. 1999; 12:363–396.
- Norris FH, Sherrieb K, Galea S. Prevalence and consequences of disaster related illness and injury from Hurricane Ike. *Rehabilitation Psychology*. 2010; 55:221–230. [PubMed: 20804265]
- North CS, Pfefferbaum B. Mental health response to community disasters: a systematic review. *Journal of the American Medical Association*. 2013; 310:507–518. [PubMed: 23925621]
- Ormel J, Jeronimus JF, Kotov R, Riese H, Bos EE, Hankin B, Rosmalen JGM. Neuroticism and common mental disorders: meaning and utility of a complex relationship. *Clinical Psychology Review*. 2013; 5:686–697. [PubMed: 23702592]
- Ormel J, Oldehinkel AJ, Brilman EI. The interplay and etiological continuity of neuroticism, difficulties, and life events in the etiology of major and subsyndromal, first, and recurrent depressive episodes in later life. *American journal of Psychiatry*. 2001; 158:885–891. [PubMed: 11384895]
- Rose, D.; Abramson, L. Developmental predictors of depressive cognitive style: research and theory. In: Cicchetti, D.; Toth, S., editors. *Developmental Perspectives on Depression*. Rochester, NY: University of Rochester Press; 1992. p. 323-349.
- Segal ZV, Shaw BF, Vella DD, Katz R. Cognitive and life stress predictors of relapse in remitted unipolar depressed patients: test of the congruency hypothesis. *Journal of Abnormal Psychology*. 1992; 101:26–36. [PubMed: 1537969]
- Spinhoven P, Elzinga BM, Hovens JGFM, Roelofs K, van Oppen P, Zitman FG, Penninx BWJH. Positive and negative life events and personality traits in predicting course of depression and anxiety. *Acta Psychiatrica Scandinavica*. 2011; 124:462–473. [PubMed: 21848705]
- Stroud CB, Davila J, Moyer A. The relationship between stress and depression in first onsets *versus* recurrences: a meta-analytic review. *Journal of Abnormal Psychology*. 2008; 117:206–213. [PubMed: 18266498]
- Tennant C. Life events, stress and depression: a review of recent findings. *Australian and New Zealand Journal of Psychiatry*. 2002; 36:173–182. [PubMed: 11982537]

- Verstraeten K, Vasey MW, Raes F, Bijttebier P. Temperament and risk for depressive symptoms in adolescence: mediation by rumination and moderation by effortful control. *Journal of Abnormal Child Psychology*. 2009; 37:349–361. [PubMed: 19107592]
- Vinkers CH, Joëls M, Milaneschi Y, Kahn RS, Penninx BW, Boks MP. Stress exposure across the lifespan cumulatively increases depression risk and moderated by neuroticism. *Depression and Anxiety*. 2014; 31:737–745. [PubMed: 24753162]
- Watson D, Clark LA. On traits and temperament: general and specific factors of emotional experience and their relation to the five-factor model. *Journal of Personality*. 1992; 60:441–476. [PubMed: 1635050]
- Watson D, Gamez W, Simms LJ. Basic dimensions of temperament and their relation to anxiety and depression: a symptom-based perspective. *Journal of Research in Personality*. 2005; 39:46–66.
- Watson, D.; Kotov, R.; Gamez, W. Basic dimensions of temperament in relation to personality and psychopathology. In: Krueger, RF.; Tackett, JL., editors. *Personality and Psychopathology*. New York: Guilford Press; 2006. p. 7-38.
- Watson D, Wiese D, Vaidya J, Tellegen A. The two general activation systems of affect: structural findings, evolutionary considerations, and psychobiological evidence. *Journal of Personality and Social Psychology*. 1999; 76:820–838.
- Weisler RH, Barbee JG, Townsend MH. Mental health and recovery in the Gulf Coast after Hurricanes Katrina and Rita. *Journal of the American Medical Association*. 2006; 296:585–588. [PubMed: 16882968]
- Wichers MC, Myin-Germeys I, Jacobs N, Peeters F, Kenis G, Derom C, Vlietinck R, Delespaul P, van Os J. Evidence that moment-to-moment variation in positive emotions buffer genetic risk for depression: a momentary assessment twin study. *Acta Psychiatrica Scandinavica*. 2007; 115:451–457. [PubMed: 17498156]
- Williams JB, Gibbon M, First MB, Spitzer RL, Davies M, Borus J, Howes MJ, Kane J, Pope HG, Rounsaville B, Wittchen HU. The Structured Clinical Interview for DSM-III-R (SCID): II. Multisite test–retest reliability. *Archives of General Psychiatry*. 1992; 49:630–636. [PubMed: 1637253]
- Wilson S, DiRago AC, Iacono WG. Prospective inter-relationships between late adolescent personality and major depressive disorder in early adulthood. *Psychological Medicine*. 2014; 44:567–577. [PubMed: 23689064]
- Zimmerman M, Sheeran T, Young D. The Diagnostic Inventory for Depression: a self-report scale to diagnose DSM-IV major depressive disorder. *Journal of Clinical Psychology*. 2004; 60:87–110. [PubMed: 14692011]

Fig. 1. Relationship between stress from Hurricane Sandy and post-Sandy depressive symptoms at high and low levels of baseline negative emotionality. Only the slope at high negative emotionality is significant. Error bars represent the standard error of the slope.

Fig. 2. Relationship between stress from Hurricane Sandy and post-Sandy depressive symptoms at high and low levels of positive emotionality. Only the slope at low positive emotionality is significant. Error bars represent the standard error of the slope.

Table 1Frequency of participant family exposure to Hurricane Sandy stressors ($n = 318$)^a

Item	Number (%) of participants endorsing item
1. Damage to home or possessions	35 (11.01)
2. Own or family's safety threatened	38 (12.50)
3. Financial hardships	40 (13.16)
4. Children fear for their own or their family's safety	68 (22.30)
5. Life disrupted by Hurricane Sandy	72 (23.68)
6. Difficulty finding gasoline	48 (15.79)
7. Difficulty getting food, water or warmth	59 (19.34)
8. Children quarreling or complaining more than usual	47 (15.46)
9. Length of time without power	108 (35.41)
10. Length of time children's school closed	147 (48.36)
11. Self, friends, family, pets or co-workers robbed, injured or got lost	9 (2.83)
12. Apply to FEMA, government aid, or Red Cross or other aid agency	14 (4.40)
13. Evacuated home	12 (3.77)
Average number of stressors reported	2.20

FEMA, Federal Emergency Management Agency.

^aItem scores dichotomized as present or not present.

Table 2

Descriptive statistics and bivariate correlations (*n* = 318)

	1	2	3	4	5
1. Lifetime depressive diagnosis ^a	-	0.28**	-0.19**	0.03	0.28**
2. Negative emotionality		-	-0.39**	0.14**	0.42**
3. Positive emotionality			-	0.09	-0.34**
4. Hurricane Sandy exposure				-	0.12*
5. Depressive symptoms post-Sandy					-
Mean	Yes = 117 (36.79%)	4.28	9.22	2.20	5.56
Standard deviation		3.58	3.10	2.19	6.02

^aFor lifetime depressive diagnosis, 0 = no, 1 = yes.

* *p* <0.05,

** *p* <0.01.

Table 3Results of regression models predicting post-Sandy depressive symptoms^a

	β	s.e.	<i>t</i>	R^2	95% CI
Model 1: NE				0.25**	
Age 9 years lifetime depressive history	1.87**	0.64	3.00		0.73 to 3.25
Age	0.04	0.07	0.54		-0.10 to 0.18
Marital status	0.46	1.03	0.46		-1.54 to 2.51
Education	0.25	0.70	0.22		-1.23 to 1.53
Household income	-1.00**	0.34	-2.92		-1.63 to -0.30
Race/ethnicity	0.18	0.83	0.22	0.12	-1.03 to 2.43
NE	1.96**	0.31	6.33	0.11	1.38 to 2.60
Stress	0.42	0.30	1.42	0.006	-0.09 to 1.09
NE x stress	0.62*	0.31	2.02	0.013	0.10 to 1.32
Model 2: PE				0.23**	
Age 9 years lifetime depressive history	2.32**	0.63	3.67		1.08 to 3.56
Age	-0.01	0.07	-0.10		-0.15 to 0.13
Marital status	0.78	1.04	0.76		-1.27 to 2.83
Education	0.20	0.71	0.28		-1.20 to 1.60
Household income	-1.06**	0.34	-3.11		-1.73 to -0.39
Race/ethnicity	0.01	0.89	0.01	0.12	-1.74 to 1.76
PE	-1.78*	0.30	-5.87	0.076	-2.37 to -1.19
Stress	1.04**	0.30	3.45	0.025	0.45 to 1.63
PE x stress	-0.71*	0.30	-2.37	0.014	-1.13 to -0.12

s.e., Standard error; CI, confidence interval; NE, negative emotionality; PE, Positive emotionality; stress, exposure to hurricane Sandy-related stress.

^aMarital status coded as 1 = married, 0 = not married; education coded as 1 = 4-year college degree or more, 0 = less than 4-year college degree; ethnicity coded as 0 = Caucasian/non-Hispanic, 1 = non-Caucasian and/or Hispanic.* $p < 0.05$,** $p < 0.01$.