Design, Synthesis, and Immunological Evaluation of Benzyloxyalkyl-Substituted 1,2,3-Triazolyl α -GalCer AnaloguesYogesh Kumar Verma,[†] Bonam Srinivasa Reddy,^{†,‡} Mithun S. Pawar,^{†,‡} Debabrata Bhunia,[†] and Halmuthur M. Sampath Kumar^{*,†,‡}[†]Vaccine Immunology Laboratory, Natural Products Chemistry Division, Indian Institute of Chemical Technology, Hyderabad 500007, India[‡]Academy of Scientific and Innovative Research, CSIR–Indian Institute of Chemical Technology, Hyderabad 500007, India

Supporting Information

ABSTRACT: Replacement of the amide moiety in the structure of α -GalCer with a 1,2,3-triazole linker is known to elicit a response skewed toward Th2 immunity, and glycolipids containing an aromatic ring in the terminus of their acyl or phytosphingosine structural component exhibit an enhanced Th1 immune response. In the current study, synthesis and immunological screening of a focused library of benzyloxyalkyl-substituted 1,2,3-triazolyl α -GalCer analogues are reported. The novel α -GalCer analogues activate invariant natural killer T (iNKT) cells via CD1d mediated presentation, which was confirmed by *in vitro* tests performed on iNKT hybridomas incubated with CD1d proteins. When tested on isolated murine splenocytes, the T1204B and T1206B compounds stimulated higher levels of both IFN- γ and IL-4 cytokine expression *in vitro* compared to that of α -GalCer.

KEYWORDS: benzyloxyalkyl α -GalCer, Th1/Th2, IL-4, IFN- γ , kinetic release, iNKT hybridoma

α -Galactosylceramide (α -GalCer) was isolated from the extracts of the Japanese marine sponge *Agelas mauritanus* from the Okinawan Sea.^{1,2} In the last two decades, α -GalCer and its derivatives have drawn considerable interest due to their versatile utility as iNKT activators, immunomodulators,³ and adjuvants in many diseases including malaria,⁴ HIV,⁵ tuberculosis,⁶ and tumor immunotherapy.⁷ Natural killer T (NKT) cells are subsets of T lymphocytes that share common features of both NK cells and conventional T cells. NKT cells that express semi-invariant surface receptor (iNKT cells) specifically recognize α -GalCer presented by an MHC class-I like molecule (i.e., CD1d). Activation of iNKT cells stimulates production of both IFN- γ and IL-4.^{8–10} Studies have revealed that production of Th1 cytokines may correlate with antiviral, antibacterial, antitumor, and adjuvant activities. However, Th2 cytokine production may subdue autoimmune diseases. Therefore, the identification of compounds that are capable of inducing varying degrees of Th1/Th2 polarization as reflected by their cytokine responses is desirable. In the search for more effective glycolipids, several analogues have been synthesized through modifications at suitable positions on the α -GalCer structure, and as a result, more potent analogues, such as OCH, C-GalCer, 7DW8–5, RCal-56, Nu- α -GalCer, SMC-124, and EF77, were prepared.^{11–15} Among the large number of α -GalCer analogues that have been synthesized over the past several years, some analogues with a fascinating immune stimulatory nature have attracted our attention. The first variety

of these analogues are α -GalCer derivatives that include a 1,2,3-triazole moiety as a replacement for the amide linkage in the structure.^{16–18} These compounds exhibited comparable iNKT stimulatory effect with Th2 bias on mice splenocytes.¹⁶ The triazole moiety served as a rigid linker that is more stable to hydrolysis and oxidative/reductive conditions in biological systems. In addition, two additional hydrogen bonds were reported between the Thr154 residue and the triazole moiety.¹⁶ α -GalCer derivatives with different aryl moieties at the acyl chain terminus have exhibited remarkable activity.^{19,20} These analogues exhibited strong iNKT stimulation with Th1 bias.^{20,21} According to Wu et al., their docking models revealed additional hydrogen bonding between the phenyl/aryl ring of the fatty acyl chain and the aromatic amino acid residues present on the wall of the A' pocket in the CD1d hydrophobic groove.²² One such analogue (7DW8–5) from the same structural class exhibited superior adjuvant activity compared to that of α -GalCer in HIV and malaria vaccines in mice.^{20,23}

To achieve a better stimulatory effect with an enhanced Th1 and Th2 response, we have designed novel α -GalCer analogues that combine both the previously discussed properties. A benzyloxy group is utilized to serve as an aryl terminus of the alkyl chain,

Received: August 20, 2015

Accepted: December 10, 2015

Published: December 10, 2015

which was attached to the azido-galactosylceramide moiety through a triazole linker. Additionally, this molecular design would enable the evaluation of the possible role of the benzyl and triazolyl moieties present in the same structural framework in modulating the immune response. To explore the immunomodulatory property of these glycolipid entities, we prepared a focused library of α -GalCer analogues where a 1,2,3-triazole replaced the amide linkage of α -GalCer along with lipid chains of varying lengths bearing a terminal benzyl group with some intervening oxygen atoms. These structural modifications may enhance the hydrogen bonding involved during the iNKT cell activation cascade, which may result in a more pronounced Th1 and Th2 response. In continuation of our ongoing research program to develop novel immunomodulators, we present the synthesis and immunopharmacological studies of novel benzyloxyalkyl-substituted 1,2,3-triazolyl α -GalCer analogues.

The synthesis of α -GalCer analogues has been accomplished by following literature procedures with some minor modifications. To achieve the synthesis of the designed analogues, two key fragments (i.e., azido α -GalCer and acetylenic lipid spacers of varying chain lengths) were prepared, and their syntheses are briefly discussed below. The synthesis of mesylate **1** was accomplished by starting from D-galactose pentaacetate according to previously published methods (Scheme 1).^{24,25}

Scheme 1. Synthesis of Azido α -GalCer^a

^aReagents and conditions: (i) NaN_3 , DMF, 100 °C, 6 h, 65%. (ii) TBSPSCl, imidazole, DCM, rt, 88%. (iii) *p*-methoxybenzoyl chloride, Et_3N , rt, 1 h, 95%. (iv) TBAF, THF, -10 °C, 3 h, 75%. (v) TBAI, DIPEA, DCM, 4 Å MS, rt, 4 h, 85%. (vi) NaOMe, MeOH, 20 min, 94%.

Mesylate **1** was subjected to $\text{S}_{\text{N}}2$ substitution to achieve azido phytosphingosine **2**. Compound **2** was selectively protected as a silyl ether followed by the orthogonal protection of the secondary hydroxyls with *p*-methoxybenzoyl (MBz) protection to afford compound **4**, which was treated with a TBAF solution for silyl deprotection to prepare intermediate **5**. A *p*-methoxybenzoyl group was employed to minimize the benzoyl migration to the primary hydroxyl group under TBAF deprotection conditions, which has been reported as a hurdle encountered with conventional benzoyl protection strategies.²⁶ The glycosyl acceptor **5** was subjected to glycosidation with persilylated galactose under TBAI/DIPEA catalysis, as described by Hague et al. to afford protected GalCer **6** with a selective α configuration.²⁷ The high yields (>80%) of the glycosidation reaction with *p*-methoxybenzoyl protection are impressive because glycosidation yields are reported to be substantially reduced with changes in the orthogonal protection

in the phytosphingosine acceptor.²⁸ Subsequent treatment of compound **6** with NaOMe/MeOH was performed to remove the *p*-methoxybenzoyl protection, and azido α -GalCer intermediate **7** was obtained as a white solid.

The terminal benzylated acetylenic lipid intermediates with varying chain lengths that are required for click chemistry were generated by following standard aliphatic condensation protocols starting from the corresponding diols with different chain lengths (Scheme 2I). The monobenzoylation of diols was

Scheme 2. Synthesis of Acetylenic Lipid Chains^a

^aReagents and conditions: (a) BnBr, NaH, THF, rt, 12 h, >75%. (b) Propargyl bromide, NaH, THF, rt, 2 h, 80%. (c) Propargyl bromide, NaH, THF, rt, 2 h, 86%. (d) TsCl, DCM, Et_3N , rt, 1 h, 90%. (e) NaH, THF, reflux, 12 h, $\geq 65\%$.

carried out using benzyl bromide/NaH in THF at rt to afford alcohols **8a–f**, which afforded propargylated intermediates **9a–f** upon condensation with propargyl bromide in the presence of sodium hydride at rt. Alternatively, propargyl intermediates **11a–c** with longer chain length lipids have been prepared by monopropargylation of 1,12-dodecanediol followed by tosylation to afford acetylenic tosylate **10**, which yielded the required benzylated acetylenic lipids **11a–c** (Scheme 2II) upon substitution with benzylated alcohols **8a–c**.²⁹

The synthesis of the final constructs was accomplished by applying a copper-mediated “click chemistry” protocol using sodium ascorbate in a tBuOH /water mixture (Scheme 3).¹⁶ All of the products were purified by column chromatography with CHCl_3 /methanol as eluents, and the products were obtained in quantitative yields.

Scheme 3. “Click Chemistry” Assembly of 1,2,3-Triazolyl α -GalCer Analogues

CD1d–GalCer engagement with a T cell receptor of iNKT cells activates their costimulatory molecules followed by activation of bystander cells. This activation leads to the production of a large amount of various cytokines and chemokines, leading to Th1 and Th2 immune responses.³⁰ Based on previous results, the synthesized analogues were screened at three different concentrations (1000, 100, and 10 ng/mL).³¹ All of the α -GalCer analogues exhibited good solubility in DMSO. The α -GalCer and its T2B, T4B, T6B,

T8B, T10B, T12B, T1202B, T1204B, and T1206B analogues were examined for their splenocyte proliferation potential and determine to be nontoxic up to 1000 ng/mL. In comparison to the untreated splenocytes, some analogues, such as T8B, T1204B, and T1206B, exhibited satisfactory proliferative effect when incubated *in vitro* with splenocytes isolated from the spleen of BALB/c mice (Figure 1).

Figure 1. Effect of α -GalCer and its analogues on splenocyte proliferation. Splenocytes were treated with α -GalCer and its analogues for 48 h, and proliferation was estimated using the MTT method.

The effect of α -GalCer and its analogues on IL-2, IL-4, and IFN- γ expression in mouse splenocytes was determined *in vitro* by ELISA. Cytokine estimation after 48 h treatment with glycolipids revealed that all of the analogues exhibited T-cell stimulation in a dose-dependent manner.³² All of the GalCer analogues exhibited uniform IL-2 expression with α -GalCer except for T2B bearing the shortest acyl lipid chain. However, the analogues with longer acyl chains, such as T8B, T12B, T1202B, T1204B, and T1206B, exhibited a significant increase in IL-2 (Figure 2a) production. Further analysis revealed that compounds T1204B and T1206B exhibited a notable increase in both IL-4 and IFN- γ (Figure 2b,c) at all three concentrations compared to that of α -GalCer, and T6B and T1202B exhibited a slight increase in IFN- γ expression at 1000 ng/mL. The high level of IL-4 and IFN- γ expressed by T1204B and T1206B may influence Th1 and Th2 cells to modulate cellular immunity.

Earlier studies demonstrated that α -GalCer induced kinetic release of cytokines by iNKT cells in a signature fashion at longer periods of time after treatment with the glycolipid, but shorter periods of time indicate dominance of IL-4 followed by high IFN- γ production after a long period of time. The kinetic release of IL-4 and IFN- γ was studied over the second, 12th and 24th hour after *in vitro* treatment of mice splenocytes with glycolipids at a 1000 ng/mL concentration. IL-4 estimation (Figure 3a) from cell culture supernatants was analyzed, and the T1204B and T1206B analogues induced a substantial increase in IL-4 at 2 h with a gradual decrease up to 24 h compared to α -GalCer. Study of kinetic release of IFN- γ (Figure 3b) revealed no significant changes in the cytokine expression up to 12 h. However, a remarkable increase in the IFN- γ levels was observed at 24 h compared to that of α -GalCer. Therefore, compounds T1204B and T1206B exhibited a good stimulatory effect expressing high levels of both the IL-4 and IFN- γ cytokines.

The test compounds (i.e., T1204B and T1206B) exhibited cytokine expression polarized toward Th1 after 48 h (Figures

Figure 2. Cytokine secretion by mouse splenocytes when stimulated with α -GalCer and its analogues. IL-2, IFN- γ , and IL-4 production was measured after 48 h treatment.

2b,c). However, the cytokine levels in kinetic release studies at 2, 12, and 24 h confirmed a bias toward Th2 behavior. Further studies on iNKT stimulation and *in vivo* kinetic release of test compounds were carried out with direct comparison of Kim's triazolyl derivative (i.e., α -GalCer-analogue-8, which is referred to as GC-8) under similar conditions.

To further confirm that the triazole analogues are activators of iNKT cells, iNKT hybridoma cells were treated with T1204B, T1206B, GC-8, and α -GalCer, incubated with CD1d protein for 16 h and assayed for IL-2 production (Figure 4). The level of cytokine IL-2 that was expressed in the *in vitro* study revealed that all three compounds were efficient stimulators of iNKT cells at a concentration of 100 ng/mL even though slightly less compared to that of α -GalCer. The T1206B analogue slightly enhanced IL-2 stimulation compared to the GC-8 compound (Figure 4).

In addition, *in vivo* experiments were performed with all four compounds individually administered to BALB/c mice intravenously at a concentration of 1 μ g, and both analogues (i.e.,

Figure 3. Kinetic release of cell-secreted cytokines (IL-4 and IFN- γ) when stimulated with α -GalCer and its analogues.

Figure 4. IL-2 expression on compound treatment with CD1d protein with Va14 V β 8.2 DN3A4–1.2 (1.2) iNKT hybridoma cells.

T1204B and T1206B) stimulated low levels of IFN- γ (Figure 5a) and IL-4 (Figure 5b) expression compared to that of α -GalCer or GC-8. Interestingly, these derivatives also exhibit signature trends of expression of both representative cytokines, as observed for α -GalCer.

In conclusion, the syntheses and immunological results of a focused library of novel benzyloxyalkyl-substituted triazolyl α -GalCer analogues that are useful immunomodulators have been reported. The results from the bioassays indicated that T1204B and T1206B exhibited significantly improved Th1 as well as Th2 cytokine expression, which may be due to the presence of directing moieties in their chemical structure. However, similar levels of expression were not observed when these tests were performed *in vitro* on isolated iNKT cells or *in vivo* in an animal model. The observed immune response may be due to the dose-dependent stimulation of other heterogeneous innate cell populations present in spleen cells by the test compounds in the *in vivo* environment. Overall, it is important to note that the additive effect observed in the current study will aid in designing analogues that induce higher levels of either Th1 or Th2 immunity and direct the immune response of interest.

Figure 5. IFN- γ and IL-4 production by BALB/c mice on *in vivo* treatment of compounds at a concentration of 1 μ g.

Modulation of Th1/Th2 activation by these α -GalCer analogues makes them promising candidates for their possible application as vaccine adjuvants and ligands to treat autoimmune disorders.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acsmedchemlett.5b00340.

Detailed experimental procedures including biology and chemistry, analytical data of all compounds, ^1H NMR, ^{13}C NMR, and HRMS spectra for selected compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*Tel: +91-40-27191824. E-mail: sampath@iict.res.in.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors wish to thank CSIR, New Delhi for financial support as part of XII Five Year plan programme under title DENOVA (CSC-0205). The authors also wish to thank Dr. Mitchell Kronenberg, La Jolla Institute for Allergy & Immunology, USA for providing the iNKT hybridoma cells. Y.K.V. wishes to thank CSIR for a fellowship.

■ ABBREVIATIONS

IFN- γ , interferon gamma; IL, interleukin; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; ELISA, enzyme linked immunosorbent assay; HIV, human immunodeficiency virus; OD, optical density; TBAF, tetrabutylammonium fluoride; MS, molecular sieves; THF, tetrahydrofuran

REFERENCES

- (1) Morita, M.; Motoki, K.; Akimoto, K.; Natori, T.; Sakai, T.; Sawa, E.; Yamaji, K.; Koezuka, Y.; Kobayashi, E.; Fukushima, H. Structure-Activity Relationship of α -Galactosylceramides against B16-Bearing Mice. *J. Med. Chem.* **1995**, *38*, 2176–2187.
- (2) Natori, T.; Koezuka, Y.; Higa, T. Agelasphins, novel α -galactosylceramides from the marine sponge *Agelas mauritanus*. *Tetrahedron Lett.* **1993**, *34*, 5591–5592.
- (3) Hayakawa, Y.; Godfrey, D. I.; Smyth, M. J.; Hayakawa, Y. α -Galactosylceramide: potential immunomodulatory activity and future application. *Curr. Med. Chem.* **2004**, *11*, 241–252.
- (4) G-Aseguinolaza, G.; Kaer, V.; Bergmann, C.; Wilson, M.; Schmieg, J.; Kronenberg, M.; Nakayama, K.; Taniguchi, M.; Koezuka, Y.; Tsuji, M. Natural Killer T Cell Ligand α -Galactosylceramide Enhances Protective Immunity Induced by Malaria Vaccines. *J. Exp. Med.* **2002**, *195*, 617–624.
- (5) Huang, Y.; Chen, A.; Li, X.; Chen, Z.; Zhang, W.; Song, Y.; Gurner, D.; Gardiner, D.; Basu, S.; Ho, D.; Tsuji, M. Enhancement of HIV DNA vaccine immunogenicity by the NKT cell ligand, α -galactosylceramide. *Vaccine* **2008**, *26*, 1807–1816.
- (6) Sada-Ovalle, I.; Sköld, M.; Tian, T.; Besra, G. S.; Behar, S. M. α -Galactosylceramide as a Therapeutic Agent for Pulmonary *Mycobacterium tuberculosis* Infection. *Am. J. Respir. Crit. Care Med.* **2010**, *182*, 841–847.
- (7) Giaccone, G.; Punt, C. J. A.; Ando, Y.; Ruijter, R.; Nishi, N.; Peters, M.; von Blomberg, M. E.; Scheper, R. J.; van der Vliet, H. J. J.; van den Eertwegh, A. J. M.; Roelvink, M.; Beijnen, J.; Zwierzina, H.; Pinedo, H. M. A phase I study of the natural killer T-cell ligand α -galactosylceramide (KR7000) in patients with solid tumors. *Clin. Cancer Res.* **2002**, *8*, 3702–3709.
- (8) Matsuda, J. L.; Mallevaey, T.; Scott-Browne, J.; Gapin, L. CD1d-restricted iNKT cells, the ‘Swiss-Army knife’ of the immune system. *Curr. Opin. Immunol.* **2008**, *20*, 358–368.
- (9) Godfrey, D. I.; MacDonald, H. R.; Kronenberg, M.; Smyth, M. J.; Van Kaer, L. NKT cells: what’s in a name? *Nat. Rev. Immunol.* **2004**, *4*, 231–237.
- (10) Godfrey, D. I.; Hammond, K. J. L.; Poulton, L. D.; Smyth, M. J.; Baxter, A. G. NKT cells: facts, functions and fallacies. *Immunol. Today* **2000**, *21*, 573–583.
- (11) Miyamoto, K.; Miyake, S.; Yamamura, T. A synthetic glycolipid prevents autoimmune encephalomyelitis by inducing TH2 bias of natural killer T cells. *Nature* **2001**, *413*, 531–534.
- (12) Franck, R. W. C-Galactosylceramide: Synthesis and Immunology. *C. R. Chim.* **2012**, *15*, 46–56.
- (13) Aspeslagh, S.; Li, Y.; Yu, E. D.; Pauwels, N.; Trappeniers, M.; Girardi, E.; Decruy, T.; van Beneden, K.; Venken, K.; Drennan, M. Galactose-modified iNKT cell agonists stabilized by an induced fit of CD1d prevent tumour metastasis. *EMBO J.* **2011**, *30*, 2294–2305.
- (14) Banchet-Cadédu, A.; Hénon, E.; Dauchez, M.; Renault, J.-H.; Monneaux, F.; Haudrechy, A. The stimulating adventure of KR7000. *Org. Biomol. Chem.* **2011**, *9*, 3080–3104.
- (15) Tyznik, A.; Farber, E.; Girardi, E.; Birkholz, A.; Li, Y.; Chitale, S.; So, R.; Arora, P.; Khurana, A.; Wang, J.; Porcelli, S.; Zajonc, D.; Kronenberg, M.; Howell, A. Glycolipids that Elicit IFN- γ -Biased Responses from Natural Killer T Cells. *Chem. Biol.* **2011**, *18*, 1620–1630.
- (16) Lee, T.; Cho, M.; Ko, S. Y.; Youn, H. J.; Baek, D. J.; Jea, Cho, W.; Kang, C. Y.; Kim, S. Synthesis and Evaluation of 1,2,3-Triazole Containing Analogues of the Immunostimulant α -GalCer. *J. Med. Chem.* **2007**, *50*, 585–589.
- (17) Tashiro, T.; Mori, K.; Shiozaki, M.; Taniguchi, M.; Watarai, H. Novel Synthetic Glycolipid and Use Thereof. US patent 2013, US2013/0005669A1.
- (18) Pilgrim, W.; O’Reilly, C.; Murphy, P. V. Synthesis of α -O- and α -S-Glycosphingolipids Related to *Sphingomonas* cell Wall Antigens Using Anomerisation. *Molecules* **2013**, *18*, 11198–11218.
- (19) Chang, Y.-J.; Huang, J.-R.; Tsai, Y.-C.; Hung, J.-T.; Wu, D.; Fujio, M.; Wong, C.-H.; Yu, A. L. Potent immune-modulating and anticancer effects of NKT cell stimulatory glycolipids. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 10299–10304.
- (20) Li, X.; Fujio, M.; Imamura, M.; Wu, D.; Vasan, S.; Wong, C. H.; Ho, D. D.; Tsuji, M. Design of a potent CD1d-binding NKT cell ligand as a vaccine adjuvant. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 13010–13015.
- (21) Chang, Y. J.; Huang, J. R.; Tsai, Y. C.; Hung, J. T.; Wu, D.; Fujio, M.; Wong, C. H.; Yu, A. L. Potent immune-modulating and anticancer effects of NKT cell stimulatory glycolipids. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 10299–10304.
- (22) Wu, D.; Zajonc, D. M.; Fujio, M.; Sullivan, B. A.; Kinjo, Y.; Kronenberg, M.; Wilson, I. A.; Wong, C.-H. Design of natural killer T cell activators: Structure and function of a microbial glycosphingolipid bound to mouse CD1d. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 3972–3977.
- (23) Padte, N. N.; Li, X.; Tsuji, M.; Vasan, S. Clinical Development of a Novel CD1d-binding NKT Cell Ligand as a Vaccine Adjuvant. *Clin. Immunol.* **2011**, *140*, 142–151.
- (24) León, F.; Brouard, I.; Rivera, A.; Torres, F.; Rubio, S.; Quintana, J.; Estévez, F.; Bermejo, J. Isolation, Structure Elucidation, Total Synthesis, and Evaluation of New Natural and Synthetic Ceramides on Human SK-MEL-1 Melanoma Cells. *J. Med. Chem.* **2006**, *49*, 5830–5839.
- (25) Asai, N.; Fusetani, N.; Matsunaga, S. Sex Pheromones of the Hair Crab *Erimacrus isenbeckii*. II. Synthesis of Ceramides. *J. Nat. Prod.* **2001**, *64*, 1210–1215.
- (26) Long, D. E.; Karmakar, P.; Wall, K. A.; Suchek, S. J. Synthesis of α -L-rhamnosyl ceramide and evaluation of its binding with anti-rhamnose antibodies. *Bioorg. Med. Chem.* **2014**, *22*, 5279–5289.
- (27) Schombs, M.; Park, F. E.; Du, W.; Kulkarni, S. S.; Gervay-Hague, J. One-Pot Syntheses of Immunostimulatory Glycolipids. *J. Org. Chem.* **2010**, *75*, 4891–4898.
- (28) Du, W.; Gervay-Hague, J. Efficient Synthesis of α -Galactosyl Ceramide Analogues Using Glycosyl Iodide Donors. *Org. Lett.* **2005**, *7*, 2063–2065.
- (29) Ouchi, M.; Inoue, Y.; Kanzaki, T.; Hakushi, T. Molecular design of crown ethers. 1. Effects of methylene chain length: 15- to 17-crown-5 and 18- to 22-crown-6. *J. Org. Chem.* **1984**, *49*, 1408–1412.
- (30) Van Kaer, L. α -Galactosylceramide Therapy for Autoimmune Diseases: Prospects and Obstacles. *Nat. Rev. Immunol.* **2005**, *5*, 31–42.
- (31) Yu, K. O. A.; Im, J. S.; Molano, A.; Dutronc, Y.; Illarionov, P. A.; Forestier, C.; Fujiwara, N.; Arias, I.; Miyake, S.; Yamamura, T.; Chang, Y.-T.; Besra, G. S.; Porcelli, S. A. Modulation of CD1d-restricted NKT cell responses by using N-acyl variants of α -galactosylceramides. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 3383–3388.
- (32) Boyman, O.; Sprent, J. The role of interleukin-2 during homeostasis and activation of the immune system. *Nat. Rev. Immunol.* **2012**, *12*, 180–190.