Published in final edited form as:

Cardiol Young. 2015 August ; 25(0 2): 58–66. doi:10.1017/S1047951115000840.

Stem Cell Therapy For Congenital Heart Disease: Toward Translation

Brody Wehman, Osama T. Siddiqui, Rachana Mishra, Sudhish Sharma, and Sunjay Kaushal

Division of Cardiac Surgery, University of Maryland School of Medicine, 110 S. Paca St., 7th Floor, Baltimore, MD, USA, 21201

Abstract

Stem cell therapy has the optimistic goal of regenerating the myocardium as defined by regrowth of lost or destroyed myocardium. As applied to patients with heart failure, many confuse or limit the regenerative definition to just improving myocardial function and/or decreasing myocardial scar formation which may not be the most important clinical outcome to achieve in this promising field of molecular medicine. Many different stem cell-based therapies have been tested and have demonstrated a safe and feasible profile in adult patients with heart failure but with varied efficacious endpoints reported. While not been achieved as of yet, the encompassing goal to regenerate the heart is still believed to be within reach using these cell-based therapies in adult patients with heart failure as the first generation therapies are now being tested in different phases of clinical trials. Similar efforts to foster the translation of stem cell therapy to children with heart failure have, however, been limited. With this review, we aim to summarize the findings from preclinical models and clinical experience to date that have focused on the evaluation of stem cell therapy in children with heart failure. Finally, we present methodological considerations pertinent to the design of a stem cell-based trial for children with heart failure since they represent a population of patients with very different sets of issues when compared to adult patients. As has been taught by many learned clinicians, children are not small adults!

Keywords

congenital heart disease; stem cell; hypoplastic left heart syndrome; single ventricle

Growing Epidemic Of Pediatric Heart Failure

The epidemic of congestive heart failure is a growing, worldwide concern and is expected to become even more prevalent. Among adults, there are approximately 15 million people worldwide and 5 million patients in the United States who have congestive heart failure,

Address correspondence to: Sunjay Kaushal, MD, PhD, Division of Cardiac Surgery, University of Maryland Medical Center, 110 S. Paca St., 7th Floor, Baltimore, MD, USA, 21201; skaushal@smail.umaryland.edu; Phone: 410-328-5842; Fax: 410-328-2750.

Conflicts of Interest

None.

Ethical Standards

Not applicable.

with 400,000 new cases per year in the United States alone. An equally alarming trend of this disease is being witnessed among children with an increase in prevalence observed over the last decade. There were almost 3000 more congestive heart failure-related hospitalizations among children in 2006 than in 2000 with over a billion dollars more in total charges to hospitals¹. This cost in simply managing children with congestive heart failure is not far from the amount spent for management of all cancer-related diagnoses (\$2.24 billion) and myocardial infarction (\$3.18 billion) combined. These numbers in children will only increase during the next decade as patients living with congenital heart disease (CHD) now outnumber patients being born with CHD. Current therapies do not adequately address the spectrum of pathophysiology that encumbers patients with CHD and cardiomyopathy, which includes pressure and volume overload, ischemia and dysrhythmias².

Considerable optimism has developed over the last decade for stem cell therapy as an option to recover dysfunctional myocardium due to encouraging preclinical studies and early clinical trials in human patients. In tandem with advancements in adult patients with ischemic heart disease, there has been a steadily emerging effort to determine the potential of these stem cells for patients with CHD. Here, we review relevant preclinical studies and summarize the clinical experience to date with stem cell therapy for patients with CHD, including a discussion of the recently published first clinical trial to evaluate the safety and feasibility of stem cell therapy in patients with hypoplastic left heart syndrome. Additionally, we discuss methodological considerations for delivery of stem cells and the design of clinical trials as it pertains to patients with CHD. Finally, we provide an overview of important unanswered translational questions and upcoming clinical trials.

Building Upon Experience With Ischemic Heart Disease

Stem cell therapy has shifted to the forefront of translational and clinical research in cardiovascular medicine. A series of advances in the laboratory has driven the implementation of numerous early clinical trials, employing a wide range of types of stem cells, methods of delivery and indications for therapy. Emerging from this assortment of preliminary clinical studies has been a handful of stem cell types that have produced reliable, safe and favorable results in both the preclinical setting and early clinical trials. These include, among others, bone marrow-derived mononuclear and mesenchymal stem cells, cardiosphere-derived cells and c-kit⁺ cardiac stem cells.

Bone marrow-derived mononuclear cells represent a heterogeneous population of precursor cells that have the ability to differentiate into multiple cardiac lineages, secrete paracrine factors and recover left ventricular function in animal models of myocardial infarction.^{3–5} Initially, two promising clinical trials, BOOST trial (bone marrow transfer to enhance ST elevation infarct regeneration) and REPAIR-AMI (the reinfusion of enriched progenitor cells and infarct remodeling in acute myocardial infarction), gathered much enthusiasm after an increase of 6.7% and 5.5% in left ventricular function, respectively, was seen after myocardial infarction in adults.^{6–8} Thereafter, many clinical trials attempted to refine the optimal timing of intracoronary delivery and dosage of these cells after myocardial infarction. However, these multiple clinical trials failed to reproduce the previously reported

improvements of left ventricular function seen in adult ischemic patients as were initially reported in the BOOST and REPAIR-MI studies.^{9–12} Even patients with chronic ischemic cardiomyopathy failed to show improvements in left ventricular function after intracoronary injection of these cells. Explanations for this discrepancy are still unclear but the percentage of CD34⁺ and CD133⁺ cells within the bone marrow-derived preparation appear crucial for their regenerative abilities. Patient characteristics, like aging, can negatively impact the regenerative abilities of these autologous stem cells and need further clarification.

Regardless of left ventricular function improvement, the clinical trials using bone marrow mononuclear cells have demonstrated an extremely safe profile in humans, an advantage to be considered when developing therapies for children with heart failure^{6–12} Furthermore, the ease of harvest and delivery of these cells add to that advantage in the pediatric population. What is still unclear is the optimal timing of these cells in adults suffering from a myocardial infarction. This unknown may also directly impact children with heart failure due to possible implications that the regenerative efficacy of these cells may be influenced by the progression of myocardial disease. Finally, these bone marrow derived cells have not been shown to routinely improve myocardial function and clinical symptoms in adults. A multicenter, randomized controlled phase III trial is now in progress to more definitively answer this question, the BAMI trial (effect of intracoronary reinfusion of bone marrow-derived mononuclear cells on all cause mortality in acute myocardial infarction).⁵ This trial is powered to determine whether infusion of mononuclear cells can effectively reduce all cause mortality in patients with an ejection fraction less than or equal to 45% when compared to a control group receiving optimal medical therapy. Results from this clinical trial will directly impact research focused on children with heart failure as demonstrating efficacy in adults is the critical first step before the application of this cell preparation in the more vulnerable pediatric population.

Another source of bone marrow-derived progenitor cells are mesenchymal stem cells. These cells have shown safety, feasibility and preliminary efficacy in improving regional contractility and quality of life, and decreasing scar formation in early phase clinical trials.^{13–15} Distinct advantages of mesenchymal stem cells include the capacity to self-replicate and differentiate into various tissue lineages, as well as the potential for allogeneic use. Mesenchymal stem cells have the unique property of being immunologically more inert due to a reduced expression of major histocompatibility complex class-I molecule, and lack of major histocompatibility complex class-II and co-stimulatory molecules CD80 (B7-1), CD86(B7-2), and CD40.^{16, 17} This has been verified extensively in preclinical animal models and recently tested in phase I, double-blind randomized clinical trials. In the initial clinical trial, intravenous infusion of allogeneic mesenchymal stem cells were delivered to patients after an acute myocardial infarction and showed that the mesenchymal stem cells did not trigger an immune response. Additionally, the mesenchymal stem cells promoted improvements in pulmonary function, left ventricular function, and symptomatic global assessment with a decrease of cardiac arrhythmias.¹⁸ Subsequently, a phase I/II randomized trial, the POSEIDON trial (the percutaneous stem cell injection delivery effects on neomyogenesis), was designed to compare the use of allogeneic and autologous mesenchymal stem cells in patients with chronic ischemic cardiomyopathy. The investigators found that allogeneic MSCs caused no relative increase in serious adverse

events and did not stimulate significant alloimmune reactions.¹⁴ Additionally, both autologous and allogeneic stem cell injections in this study reduced the size of the infarct by approximately 33%, reduced left-ventricular sphericity index, improved physical functional capacity, and improved quality of life. The exact mechanism of action underlying the recovery of myocardial function is still unknown, but may include differentiation into mature cardiomyocytes, decreased inflammation and scar formation, decreased apoptosis of cardiomyocytes, secretion of paracrine factors and stimulation of the resident c-kit⁺ cardiac stem cells.

Another area of intense focus has been the therapeutic potential of two types of resident cardiac stem cells: c-kit⁺ resident cardiac stem cells and cardiosphere-derived cells. The c-kit⁺ cardiac stem cells are defined by their multipotent, self-renewing and clonogenic properties and have been clinically studied in the SCIPIO trial (administration of cardiac stem cells in patients with ischemic cardiomyopathy) which was reported as a randomized, open-label phase I trial in patients with ischemic heart disease having undergone coronary bypass revascularization.¹⁹ With the intracoronary delivery of c-kit⁺ cardiac stem cells there were no serious adverse events reported and, despite a small number of patients, the cell treated patients showed a reduction in scar formation and an average improvement in the left ventricular ejection fraction by 12.3% during the first year of cellular transplantation compared to baseline. Another cell type extensively studied is the cardiosphere-derived cells, which are a heterogeneous population of CD105⁺/CD45⁻ mononuclear cells. The CADUCEUS trial (cardiosphere-derived autologous stem cells to reverse ventricular dysfunction and autologous cardiosphere-derived cells) was a prospective, randomized, phase Ib safety trial in adult patients with post-infarction left ventricular dysfunction. The trial demonstrated that intracoronary delivery of endocardial biopsy-derived cardiosphere-derived cells resulted in no arrhythmias, formation of tumors, myocardial infarction or other serious adverse cardiac events. Although there were no significant improvements in cardiac function, there was significant reduction in scar mass, increased viable heart mass, regional contractility and systolic wall thickening. With these encouraging results a phase 1–2 trial using allogeneic cardiosphere-derived cells rather than the autologous cells are being studied in adult patients with ischemic heart disease.²⁰

Evaluation Of Stem Cell Therapy In Animal Models Relevant to Congenital Heart Disease

With the majority of stem cell research aimed at the regeneration or recovery of ischemic myocardium, less attention has been given to the evaluation of stem cell therapy under pathophysiologic conditions relevant to patients with CHD. Although no animal model can truly replicate all of the salient features of complex CHD, reproducible models of pressure or volume overload can mimic some of the cardinal features of these diseases. However, use of these models to determine the safety and efficacy of stem cell therapy has been scarce. In a notable exception, Davies et al used a model of right ventricular pressure overload in neonatal sheep to determine the effects of human cord blood stem cells on right ventricular function²¹. In these experiments, immunosuppressed neonatal sheep underwent banding of the pulmonary artery followed by epicardial injection of human-derived cord blood stem

cells into the right ventricle and were compared to post-banding parameters of right ventricular function as well as to animals receiving injection of a placebo solution. At four weeks post-injection, marked improvement was observed in load-independent indices of systolic and diastolic function of the right ventricle in the cell-treated sheep compared to placebo-treated sheep.

To further evaluate the safety profile of the injection of cord blood stem cells to the right ventricle, a recent double-blinded randomized study injected cord blood stem cells or a placebo into the right ventricle of healthy juvenile swine²². The animals were rigorously monitored for three months, including electrophysiological assessments using implanted event recorders. The investigators observed no major adverse events, including systemic infection, tumor formation, myocardial necrosis, ventricular dysfunction or dysrhythmias. Although the number of animals was relatively small at six per group, this study established the first evidence of a favorable safety profile for injection of cord blood stem cells to the right ventricle in a juvenile model.

In a rat model of chronic pressure-overload of the right ventricle, Hoashi et al demonstrated that sheets of skeletal myoblasts transplanted four weeks after banding of the pulmonary artery improved diastolic function, reduced fibrosis of the right ventricle and increased capillary density in the myocardium compared to controls²³. Additionally, real time polymerase chain reaction of explanted specimens from the right ventricle revealed increased expression of genes encoding the angiogenic cytokines, hepatocyte growth factor and vascular endothelial growth factor, in rats that received transplantation of skeletal myoblasts compared to controls, suggesting a possible mechanism for the improved angiogenesis observed in the treatment group.

Clinical Experience With Stem Cell Therapy In Patients With Congenital Heart Disease

Case Reports and Series

The majority of experience with stem cell therapy in human pediatric patients has been limited to isolated case reports and small case series. In total, over twenty reported pediatric patients have received a stem cell therapy product for CHD or end-stage cardiomyopathy (reviewed in Tarui et al).²⁴ Among these patients, dilated cardiomyopathy has been the most common etiology of heart failure, with improvements in left ventricular ejection fraction of 20–23% from measured values at baseline in patients ranging from four months to seventeen years of age. A similar increase in left ventricular ejection fraction was reported in a nine year-old girl with congestive heart failure after an anterior myocardial infarction, which was presumed to be due to Takayasu arteritis.²⁵ After initial treatment with medical therapy and placement of a left main stent, the patient was discharged from the hospital with an ejection fraction of 28.5% by cardiac magnetic resonance imaging (MRI). She re-presented one year later with severe fatigue and dyspnea and was found to have an ejection fraction of 19%. She underwent a bone marrow biopsy and was successfully treated with autologous bone-marrow derived progenitor cells via transcatheter injection to the left anterior descending

artery. At twelve weeks of follow-up a remarkable improvement in her symptoms and ejection fraction was observed, which had improved to 47.8% as assessed by cardiac MRI.

Case reports that describe delivery of stem cells to patients with hypoplastic left heart syndrome have shown similar benefits to the performance of the single right ventricle. One patient developed failure of the single right ventricle following a hybrid stage I procedure that was complicated by obstruction of the arterial duct.²⁶ This patient was treated with intracoronary delivery of bone marrow cells to the proximal and distal right coronary artery. Twelve months after administration of stem cells, the ejection fraction of the systemic right ventricle improved from 22% to 44% and the serum level of brain natriuretic peptide was reduced from 2200 picograms/milliliter to 132 picograms/milliliter.

More recently, Burkhardt et al published the first reported intramyocardial delivery of autologous umbilical cord blood stem cells to patients with hypoplastic left heart syndrome during stage II surgical palliation.²⁷ This patient suffered no sustained dysrhythmias, myocardial necrosis or systemic infection related to cell injections and the ejection fraction of the right ventricle improved from 30–35% prior to stage II to 50% at three months post-injection by transthoracic echocardiography. Despite the heterogeneity in age, indication, type of cell and method of delivery, a commonality between all reported cases in children has been the robust response to cellular therapy. Until recently, however, no clinical trial had been completed to formally evaluate the safety, feasibility and preliminary efficacy of stem cell therapy for children with complex forms of CHD.

A Clinical Trial For Patients With Hypoplastic Left Heart Syndrome

Prior to the development of palliative strategies for hypoplastic left heart syndrome, this lesion was universally lethal in infancy. With development over the last few decades of staged surgical treatments, outcomes of surgical palliation for hypoplastic left heart syndrome have steadily improved. Typically, HLHS patients undergo three-staged surgical procedures: stage I Norwood palliative operation in the neonatal period, stage II palliative bidirectional cavopulmonary connection operation at ~4 months of age, and the stage III palliative total cavopulmonary (Fontan) operation at ~three years of age. Despite these advancements in surgical technique, the five-year mortality rate still remains unacceptably high at 50–60%, with cardiac transplantation remaining as the only alternative for patients with failing circulatory systems.^{28, 29} While the etiology of this attrition is multifactorial, RV dysfunction plays an important role³⁰. In a report by Altmann et al., those patients who presented with depressed RV function had an 18-month survival of 35% compared to 70% for those with normal function.³¹ Furthermore, as an increasing number of these patients survive the neonatal period, there is a growing population of patients who present with dysfunction of the single right ventricle, whose cellular and gross architecture is not structurally or genetically designed to chronically support the systemic circulation.^{32–35} Interventions aimed at improving the function of the systemic right ventricle which avoid or defer transplantation in patients with hypoplastic left heart syndrome are therefore urgently needed.

The TICAP trial (transcoronary infusion of cardiac progenitor cells in patients with single ventricle physiology) was the first completed clinical trial to address these needs in patients

with hypoplastic left heart syndrome using a cell-based therapy³⁶. In this study, autologous cardiosphere-derived cells were isolated and expanded from biopsies of the right atrial appendage obtained from each patient, which were then administered via intracoronary delivery four to five weeks following stage II or stage III surgical palliation. The primary objectives of the study were to evaluate the safety and feasibility of infusion of cardiosphere-derived cells, with a secondary aim to assess preliminary evidence of efficacy to enhance performance of the single ventricle. Seven patients were included in each arm of the study, which was not randomized. Function of the single ventricle was serially measured using cardiac MRI, transthoracic echocardiography and single-photon emission computed tomography.

No adverse events were reported in the form of perioperative complications during delivery of the cells, formation of new tumors, or other major adverse cardiac events. Other favorable findings included a reduced incidence of readmission to the hospital and end-organ dysfunction in the cell-treated group compared to controls. Interestingly, cell-treated patients demonstrated an improvement in right ventricular ejection fraction from $46.9 \pm 4.6\%$ to $54.0 \pm 2.8\%$ and a significant reduction in the diameter of the tricuspid valve annulus at 18 months of follow-up, while control patients showed little improvement in right ventricular ejection fraction, going from $46.7 \pm 4.4\%$ to $48.7 \pm 6.7\%$, and no change in diameter of the tricuspid valve annulus. At 18 months, there were observed reductions in free wall mass and indexed end-systolic and end-diastolic volumes of the right ventricle in cell-treated patients. The authors also tracked somatic growth of enrolled patients, and found that cell-treated patients had significantly improved z-scores for height and weight at 18 months compared to values at baseline, while there was no significant change in growth in the group of control patients. Additionally, a reduction in heart failure status, levels of brain natriuretic peptide and number of coil interventions was found in the cell-treated group compared to controls. While the numbers in this initial study were small, and the study was not randomized, the authors effectively demonstrated safety of intracoronary delivery of cardiosphere-derived cells to patients with hypoplastic left heart syndrome. Furthermore, this study showed the potential efficacy of a cell-based therapy to enhance the ventricular function of patients with complex CHD.

Methodologic Considerations

The potential of stem cell therapy for patients with CHD presents a different set of challenges not seen in adults with ischemic heart disease. Given the diversity of the types of stem cells and approaches to delivery of stem cells that have been described, investigators are faced with designing clinical trials to the best of their ability using existing literature. The optimal type of stem cell for therapeutic use in patients with CHD is not known, and current choices for use in clinical trials may be based on experience of the investigators, cost, regulatory issues and feasibility. An additional consideration is the use of an allogeneic, rather than autologous, cell therapy product, which has emerged as a potential option for some types of stem cells. Clear advantages to this approach include “off-the-shelf” availability of a pre-screened and expanded stem cell product for clinical use, as well as the avoidance of an additional invasive procedure, such as a bone marrow biopsy¹⁷. A recent phase I/II randomized trial comparing autologous to allogeneic mesenchymal stem

cells demonstrated allogeneic cells were safe and equally effective in the recovery of ischemic myocardium¹⁴. Additionally, an allogeneic preparation of cardiosphere-derived cells has been recently shown not to elicit an immune response and to have equal efficacy as allogeneic cells in the recovery of function of the left ventricle in rodent and swine models of myocardial infarction.^{37, 38} These findings support the basis of the ALL-STAR trial (allogeneic heart stem cells to achieve myocardial regeneration) which is a phase I/II trial to determine the safety and efficacy of allogeneic cardiosphere-derived cells administered to patients with a history of an anterior myocardial infarction.²⁰

The optimal method for delivery of stem cells remains an active area of investigation. The ideal method would ultimately be tailored to the needs of the age of the patient, diagnosis and whether or not a surgical procedure is planned to occur at the time of administration of cells to provide direct exposure of cardiac structures. The existing delivery strategies include catheter-based intracoronary delivery, intramyocardial injection or transendocardial injection. A novel method under investigation in our animal laboratory is the potential for cardiopulmonary bypass-assisted intracoronary infusion of cardiac stem cells via the root of the aorta, which is anticipated to facilitate a global distribution of the cell product as an adjunct to a planned cardiac surgical procedure (Figure 1). In a population of neonatal and infant patients, especially those with complex cardiac malformations and a high association of coronary anomalies, a strategy of intramyocardial injection or bypass-assisted coronary infusion at the time of surgical reconstruction may have reduced risk than a catheter-based coronary intervention.^{39, 40} Furthermore, the repeated interruption of coronary blood flow that occurs with the commonly used stop-reflow technique during intracoronary delivery may pose additional risks to patients with complex CHD or end-stage cardiomyopathy. Use of this technique has been safe thus far in clinical trials in adults where cells are delivered to a focal area of ischemic or scarred myocardium, usually at the mid to distal left anterior descending artery.^{19, 41} There are key differences, however, in the approach to patients with CHD who have global, non-ischemic ventricular dysfunction, and the clinical implication of intracoronary delivery to these patients has yet to be fully determined. In the TICAP trial, for example, operators were required to engage and repeatedly occlude proximal coronary blood flow to each of the major coronary arteries supplying a univentricular heart, one month out from surgical palliation.³⁶ The authors reported no complications experienced with the procedure and only transient ST-segment elevations using this technique, but concern for the possibility of future complications is warranted. In the design of future trials, intracoronary infusion without balloon inflation may be a safer approach, and is the strategy employed in an upcoming trial designed for adult patients with non-ischemic cardiomyopathy. However, whether cells are as effectively delivered to the myocardium as with the stop-reflow technique has yet to be verified.

One of the key biological questions is the mechanism of action of stem cells in patients with CHD. We have shown that the main action of recovery of the cardiosphere-derived cells obtained from patients with CHD is the more potent cytokine release by the younger derived cardiosphere-derived cells when compared to older derived cardiosphere-derived cells.⁴² We are now testing these younger derived cardiosphere-derived cells in other non-ischemic ventricular dysfunctional rodent models, which may have more relevance to patients with CHD. In addition to the impact of age of the donor on stem cell functionality,

characterization of the ideal cell product may also be determined using quantifiable growth properties such as telomere length, telomerase activity and population doubling time.⁴³ As recent work from ours and others continues to show the central importance of a high functioning secretome in the repair capabilities of stem cells⁴⁴, it is likely that the stem cell type and donor with the most potent secretome will emerge as the most efficacious stem cell product for repair of injured myocardium.⁴⁵

Finally, the indication of stem cell therapy for patients with CHD remains to be defined. For instance, a subset of patients with hypoplastic left heart syndrome and right ventricular dysfunction may have the greatest clinical benefit from stem cell treatment. The answers to these translational questions will be paramount to achieve the greatest efficacy in the treatment of these challenging patients with CHD.

Future Clinical Trials

A number of planned or ongoing clinical trials have been designed to evaluate the safety, feasibility and preliminary efficacy of stem cell therapy as a novel treatment for CHD. Interestingly, patients with hypoplastic left heart syndrome are the population of interest for each of the planned or ongoing clinical trials, a testament to the severe unmet medical needs of this population.

Randomized phase I clinical trials at Duke University and Mayo Clinic are both aimed at evaluating the safety and feasibility of administration of autologous cord blood stem cells to patients with hypoplastic left heart syndrome. At Mayo Clinic, cord blood stem cells are administered intra-operatively via intramyocardial injection at the time of the stage II palliation, with a secondary objective to evaluate the preliminary efficacy to enhance performance of the right ventricle.²⁷ The trial at Duke University, by contrast, utilizes intravenous delivery of cord blood stem cells with a focus on improvement of neurologic outcomes.²⁴ Also recruiting patients are the same investigators of the TICAP trial for its successor, a phase II, PERSEUS trial (cardiac progenitor cell infusion to treat univentricular heart disease).²⁴ This is a randomized study aimed at assessing the efficacy of treatment with cardiosphere-derived cells for patients with single ventricle lesions of either the left or right ventricle.

Our institution will soon begin enrollment in a phase I trial to intra-operatively administer allogeneic mesenchymal stem cells to patients with hypoplastic left heart syndrome undergoing stage II palliation (Figure 2)⁴⁶. Mesenchymal stem cells have a long and proven safety record in adult clinical trials. In addition, the regenerative efficacy of mesenchymal stem cells is not affected when used as an allogeneic cell product. This is a unique feature of mesenchymal stem cells, and allows the potential to eliminate many of the variables present in an autologous cell product.¹⁴ Mesenchymal stem cells have also been shown in a large animal study to recruit and activate the endogenous pool of c-kit⁺ cardiac stem cells to areas of myocardial injury.⁴⁷ This has significant implications for the application of mesenchymal stem cell therapy in children with CHD, as our laboratory has demonstrated an age-dependent decline in the number and regenerative efficacy of resident c-kit⁺ cardiac stem cells present in pediatric versus adult myocardium.⁴² In our planned trial, we chose the stage

II palliative operation as the time of stem cell injection for several reasons. First, there is a well established risk for inter-stage mortality following the stage I operation, as high as 12% according to follow-up of 426 patients from the Single Ventricular Reconstruction trial.⁴⁸ Thus, the high early mortality rate after the stage I operation could potentially mask the safety endpoints of a phase I study. Secondly, an intervention at the stage II operation will allow for parallel study of the natural history of the systemic right ventricle following the bidirectional cavopulmonary connection operation. Specifically, we will have the opportunity to assess the effects of volume unloading on systolic and diastolic function over an extended period of follow-up using cardiac MRI, in addition to the evaluation of mesenchymal stem cell therapy on right ventricular function. Lastly, restricting enrollment to patients undergoing the stage II operation will create a more homogenous population of patients with hypoplastic left heart syndrome and eliminate patient selection variables in the final analysis of clinical results.

In summary, stem cell therapy has promise as a novel therapy for patients with CHD and could potentially offer a new paradigm for the management of these patients. Several pioneering trials are underway, with a resolute focus on the treatment of hypoplastic left heart syndrome, which will establish the foundation for stem cell therapy in patients with CHD. As preclinical and clinical research continues to advance, regular discourse between basic science and clinical investigators is essential for the effective translation of this therapy from bench to bedside in the safest and most efficacious manner.

Acknowledgments

None.

Financial Support

Dr. Kaushal is supported by the National Institutes of Health Grant R01HL118491 and the Children's Heart Foundation.

References

1. Rossano JW, Kim JJ, Decker JA, Price JF, Zafar F, Graves DE, Morales DL, Heinle JS, Bozkurt B, Towbin JA, Denfield SW, Dreyer WJ, Jefferies JL. Prevalence, morbidity, and mortality of heart failure-related hospitalizations in children in the united states: A population-based study. *J Card Fail.* 2012; 18:459–470. [PubMed: 22633303]
2. Rossano JW, Shaddy RE. Heart failure in children: Etiology and treatment. *J Pediatr.* 2014; 165:228–233. [PubMed: 24928699]
3. Karantalis V, Balkan W, Schulman IH, Hatzistergos KE, Hare JM. Cell-based therapy for prevention and reversal of myocardial remodeling. *Am J Physiol Heart Circ Physiol.* 2012; 303:H256–H270. [PubMed: 22636682]
4. Williams AR, Hare JM. Mesenchymal stem cells: Biology, pathophysiology, translational findings, and therapeutic implications for cardiac disease. *Circulation research.* 2011; 109:923–940. [PubMed: 21960725]
5. Telukuntla KS, Suncion VY, Schulman IH, Hare JM. The advancing field of cell-based therapy: Insights and lessons from clinical trials. *J Am Heart Assoc.* 2013; 2:e000338. [PubMed: 24113326]
6. Wollert KC, Meyer GP, Lotz J, Ringes-Lichtenberg S, Lippolt P, Breidenbach C, Fichtner S, Korte T, Hornig B, Messinger D, Arseniev L, Hertenstein B, Ganser A, Drexler H. Intracoronary autologous bone-marrow cell transfer after myocardial infarction: The boost randomised controlled clinical trial. *Lancet.* 2004; 364:141–148. [PubMed: 15246726]

7. Schachinger V, Erbs S, Elsasser A, Haberbosch W, Hambrecht R, Holschermann H, Yu J, Corti R, Mathey DG, Hamm CW, Suselbeck T, Assmus B, Tonn T, Dimmeler S, Zeiher AM. Investigators R-A. Intracoronary bone marrow-derived progenitor cells in acute myocardial infarction. *N Engl J Med*. 2006; 355:1210–1221. [PubMed: 16990384]
8. Schachinger V, Erbs S, Elsasser A, Haberbosch W, Hambrecht R, Holschermann H, Yu J, Corti R, Mathey DG, Hamm CW, Suselbeck T, Werner N, Haase J, Neuzner J, Germing A, Mark B, Assmus B, Tonn T, Dimmeler S, Zeiher AM. Investigators R-A. Improved clinical outcome after intracoronary administration of bone-marrow-derived progenitor cells in acute myocardial infarction: Final 1-year results of the repair-ami trial. *European heart journal*. 2006; 27:2775–2783. [PubMed: 17098754]
9. Traverse JH, Henry TD, Pepine CJ, Willerson JT, Zhao DX, Ellis SG, Forder JR, Anderson RD, Hatzopoulos AK, Penn MS, Perin EC, Chambers J, Baran KW, Raveendran G, Lambert C, Lerman A, Simon DI, Vaughan DE, Lai D, Gee AP, Taylor DA, Cogle CR, Thomas JD, Olson RE, Bowman S, Francescon J, Geither C, Handberg E, Kappenman C, Westbrook L, Piller LB, Simpson LM, Baraniuk S, Loghin C, Aguilar D, Richman S, Zierold C, Spoon DB, Bettencourt J, Sayre SL, Vojvodic RW, Skarlatos SI, Gordon DJ, Ebert RF, Kwak M, Moye LA, Simari RD. Cardiovascular Cell Therapy Research N. Effect of the use and timing of bone marrow mononuclear cell delivery on left ventricular function after acute myocardial infarction: The time randomized trial. *JAMA*. 2012; 308:2380–2389. [PubMed: 23129008]
10. Traverse JH, Henry TD, Ellis SG, Pepine CJ, Willerson JT, Zhao DX, Forder JR, Byrne BJ, Hatzopoulos AK, Penn MS, Perin EC, Baran KW, Chambers J, Lambert C, Raveendran G, Simon DI, Vaughan DE, Simpson LM, Gee AP, Taylor DA, Cogle CR, Thomas JD, Silva GV, Jorgenson BC, Olson RE, Bowman S, Francescon J, Geither C, Handberg E, Smith DX, Baraniuk S, Piller LB, Loghin C, Aguilar D, Richman S, Zierold C, Bettencourt J, Sayre SL, Vojvodic RW, Skarlatos SI, Gordon DJ, Ebert RF, Kwak M, Moye LA, Simari RD. Cardiovascular Cell Therapy R. Effect of intracoronary delivery of autologous bone marrow mononuclear cells 2 to 3 weeks following acute myocardial infarction on left ventricular function: The latetime randomized trial. *JAMA*. 2011; 306:2110–2119. [PubMed: 22084195]
11. Sürder D, Manka R, Lo Cicero V, Moccetti T, Rufibach K, Soncin S, Turchetto L, Radrizzani M, Astori G, Schwitter J, Erne P, Zuber M, Auf der Maur C, Jamshidi P, Gaemperli O, Windecker S, Moschovitis A, Wahl A, Bühler I, Wyss C, Kozerke S, Landmesser U, Lüscher TF, Corti R. Intracoronary injection of bone marrow derived mononuclear cells, early or late after acute myocardial infarction: Effects on global left ventricular function four months results of the swiss-ami trial. *Circulation*. 2013
12. Perin EC, Willerson JT, Pepine CJ, Henry TD, Ellis SG, Zhao DX, Silva GV, Lai D, Thomas JD, Kronenberg MW, Martin AD, Anderson RD, Traverse JH, Penn MS, Anwaruddin S, Hatzopoulos AK, Gee AP, Taylor DA, Cogle CR, Smith D, Westbrook L, Chen J, Handberg E, Olson RE, Geither C, Bowman S, Francescon J, Baraniuk S, Piller LB, Simpson LM, Loghin C, Aguilar D, Richman S, Zierold C, Bettencourt J, Sayre SL, Vojvodic RW, Skarlatos SI, Gordon DJ, Ebert RF, Kwak M, Moye LA, Simari RD. Cardiovascular Cell Therapy Research N. Effect of transendocardial delivery of autologous bone marrow mononuclear cells on functional capacity, left ventricular function, and perfusion in chronic heart failure: The focus-cctn trial. *JAMA*. 2012; 307:1717–1726. [PubMed: 22447880]
13. Karantalis V, DiFede DL, Gerstenblith G, Pham S, Symes J, Zambrano JP, Fishman J, Pattany P, McNiece I, Conte J, Schulman S, Wu K, Shah A, Breton E, Davis-Sproul J, Schwarz R, Feigenbaum G, Mushtaq M, Suncion VY, Lardo AC, Borrello I, Mendizabal A, Karas TZ, Byrnes J, Lowery M, Heldman AW, Hare JM. Autologous mesenchymal stem cells produce concordant improvements in regional function, tissue perfusion, and fibrotic burden when administered to patients undergoing coronary artery bypass grafting: The prospective randomized study of mesenchymal stem cell therapy in patients undergoing cardiac surgery (prometheus) trial. *Circulation research*. 2014; 114:1302–1310. [PubMed: 24565698]
14. Hare JM, Fishman JE, Gerstenblith G, DiFede Velazquez DL, Zambrano JP, Suncion VY, Tracy M, Ghersin E, Johnston PV, Brinker JA, Breton E, Davis-Sproul J, Schulman IH, Byrnes J, Mendizabal AM, Lowery MH, Rouy D, Altman P, Wong Po Foo C, Ruiz P, Amador A, Da Silva J, McNiece IK, Heldman AW, George R, Lardo A. Comparison of allogeneic vs autologous bone marrow-derived mesenchymal stem cells delivered by transendocardial injection in patients with

ischemic cardiomyopathy: The poseidon randomized trial. *JAMA*. 2012; 308:2369–2379. [PubMed: 23117550]

15. Heldman AW, DiFede DL, Fishman JE, Zambrano JP, Trachtenberg BH, Karantalis V, Mushtaq M, Williams AR, Suncion VY, McNiece IK, Ghersin E, Soto V, Lopera G, Miki R, Willens H, Hendel R, Mitrani R, Pattany P, Feigenbaum G, Oskoue B, Byrnes J, Lowery MH, Sierra J, Pujol MV, Delgado C, Gonzalez PJ, Rodriguez JE, Bagno LL, Rouy D, Altman P, Foo CW, da Silva J, Anderson E, Schwarz R, Mendizabal A, Hare JM. Transendocardial mesenchymal stem cells and mononuclear bone marrow cells for ischemic cardiomyopathy: The tac-hft randomized trial. *JAMA*. 2014; 311:62–73. [PubMed: 24247587]
16. Selem SM, Kaushal S, Hare JM. Stem cell therapy for pediatric dilated cardiomyopathy. *Current cardiology reports*. 2013; 15:369. [PubMed: 23666883]
17. Karantalis V, Schulman IH, Balkan W, Hare JM. Allogeneic cell therapy: A new paradigm in therapeutics. *Circulation research*. 2015; 116:12–15. [PubMed: 25552688]
18. Hare JM, Traverse JH, Henry TD, Dib N, Strumpf RK, Schulman SP, Gerstenblith G, DeMaria AN, Denktas AE, Gammon RS, Hermiller JB Jr, Reisman MA, Schaer GL, Sherman W. A randomized, double-blind, placebo-controlled, dose-escalation study of intravenous adult human mesenchymal stem cells (prochymal) after acute myocardial infarction. *Journal of the American College of Cardiology*. 2009; 54:2277–2286. [PubMed: 19958962]
19. Bolli R, Chugh AR, D'Amario D, Loughran JH, Stoddard MF, Ikram S, Beache GM, Wagner SG, Leri A, Hosoda T, Sanada F, Elmore JB, Goichberg P, Cappetta D, Solankhi NK, Fahsah I, Rokosh DG, Slaughter MS, Kajstura J, Anversa P. Cardiac stem cells in patients with ischaemic cardiomyopathy (scipio): Initial results of a randomised phase 1 trial. *The Lancet*. 2011; 378:1847–1857.
20. Makkar R, Schatz R, Traverse J, Hamer A, Beattie K, Smith RR, Kivel F, Marbán L, Marbán E, Henry TD. Abstract 20536: Allogeneic heart stem cells to achieve myocardial regeneration (allstar): The one year phase i results. *Circulation*. 2014; 130:A20536.
21. Davies B, Elwood NJ, Li S, Cullinane F, Edwards GA, Newgreen DF, Brizard CP. Human cord blood stem cells enhance neonatal right ventricular function in an ovine model of right ventricular training. *The Annals of thoracic surgery*. 2010; 89:585–593. 593, e581–e584. [PubMed: 20103347]
22. Cantero Peral S, Burkhart HM, Oommen S, Yamada S, Nyberg SL, Li X, O'Leary PW, Terzic A, Cannon BC, Nelson TJ. Wanek Program Porcine Pipeline G, Wanek Program Porcine Pipeline G. Safety and feasibility for pediatric cardiac regeneration using epicardial delivery of autologous umbilical cord blood-derived mononuclear cells established in a porcine model system. *Stem cells translational medicine*. 2015
23. Hoashi T, Matsumiya G, Miyagawa S, Ichikawa H, Ueno T, Ono M, Saito A, Shimizu T, Okano T, Kawaguchi N, Matsuura N, Sawa Y. Skeletal myoblast sheet transplantation improves the diastolic function of a pressure-overloaded right heart. *J Thorac Cardiovasc Surg*. 2009; 138:460–467. [PubMed: 19619796]
24. Tarui S, Sano S, Oh H. Stem cell therapies in patients with single ventricle physiology. *Methodist Debakey Cardiovasc J*. 2014; 10:77–81. [PubMed: 25114758]
25. Limsuwan A, Pienvichit P, Limpijankit T, Khowsathit P, Hongeng S, Pornkul R, Siripornpitak S, Boonbaichaiyapruk S. Transcoronary bone marrow-derived progenitor cells in a child with myocardial infarction: First pediatric experience. *Clin Cardiol*. 2010; 33:E7–E12. [PubMed: 20632394]
26. Rupp S, Zeiher AM, Dimmeler S, Tonn T, Bauer J, Jux C, Akintuerk H, Schranz D. A regenerative strategy for heart failure in hypoplastic left heart syndrome: Intracoronary administration of autologous bone marrow-derived progenitor cells. *J Heart Lung Transplant*. 2010; 29:574–577. [PubMed: 20044280]
27. Burkhart HM, Qureshi MY, Peral SC, O'Leary PW, Olson TM, Cetta F, Nelson TJ. the Wanek Program Clinical Pipeline G. Regenerative therapy for hypoplastic left heart syndrome: First report of intraoperative intramyocardial injection of autologous umbilical-cord blood-derived cells. *J Thorac Cardiovasc Surg*. 2014
28. Newburger JW, Sleeper LA, Frommelt PC, Pearson GD, Mahle WT, Chen S, Dunbar-Masterson C, Mital S, Williams IA, Ghanayem NS, Goldberg CS, Jacobs JP, Krawczeski CD, Lewis AB,

- Pasquali SK, Pizarro C, Gruber PJ, Atz AM, Khaikin S, Gaynor JW, Ohye RG. Pediatric Heart Network I. Transplantation-free survival and interventions at 3 years in the single ventricle reconstruction trial. *Circulation*. 2014; 129:2013–2020. [PubMed: 24705119]
29. Feinstein JA, Benson DW, Dubin AM, Cohen MS, Maxey DM, Mahle WT, Pahl E, Villafane J, Bhatt AB, Peng LF, Johnson BA, Marsden AL, Daniels CJ, Rudd NA, Caldarone CA, Mussatto KA, Morales DL, Ivy DD, Gaynor JW, Tweddell JS, Deal BJ, Furck AK, Rosenthal GL, Ohye RG, Ghanayem NS, Cheatham JP, Tworetzky W, Martin GR. Hypoplastic left heart syndrome: Current considerations and expectations. *Journal of the American College of Cardiology*. 2012; 59:S1–S42. [PubMed: 22192720]
 30. Lee TM, Aiyagari R, Hirsch JC, Ohye RG, Bove EL, Devaney EJ. Risk factor analysis for second-stage palliation of single ventricle anatomy. *The Annals of thoracic surgery*. 2012; 93:614–618. discussion 619. [PubMed: 22197533]
 31. Altmann K, Printz BF, Solowiejczyk DE, Gersony WM, Quaegebeur J, Apfel HD. Two-dimensional echocardiographic assessment of right ventricular function as a predictor of outcome in hypoplastic left heart syndrome. *The American journal of cardiology*. 2000; 86:964–968. [PubMed: 11053708]
 32. Kaneko S, Khoo NS, Smallhorn JF, Tham EB. Single right ventricles have impaired systolic and diastolic function compared to those of left ventricular morphology. *Journal of the American Society of Echocardiography : official publication of the American Society of Echocardiography*. 2012; 25:1222–1230. [PubMed: 22975791]
 33. d'Udekem Y, Xu MY, Galati JC, Lu S, Iyengar AJ, Konstantinov IE, Wheaton GR, Ramsay JM, Grigg LE, Millar J, Cheung MM, Brizard CP. Predictors of survival after single-ventricle palliation: The impact of right ventricular dominance. *Journal of the American College of Cardiology*. 2012; 59:1178–1185. [PubMed: 22440217]
 34. Walsh MA, McCrindle BW, Dipchand A, Manlhiot C, Hickey E, Caldarone CA, Van Arsdel GS, Schwartz SM. Left ventricular morphology influences mortality after the norwood operation. *Heart*. 2009; 95:1238–1244. [PubMed: 19457871]
 35. Voelkel NF, Quaife RA, Leinwand LA, Barst RJ, McGoon MD, Meldrum DR, Dupuis J, Long CS, Rubin LJ, Smart FW, Suzuki YJ, Gladwin M, Denholm EM, Gail DB. National Heart L, Blood Institute Working Group on C, Molecular Mechanisms of Right Heart F. Right ventricular function and failure: Report of a national heart, lung, and blood institute working group on cellular and molecular mechanisms of right heart failure. *Circulation*. 2006; 114:1883–1891. [PubMed: 17060398]
 36. Ishigami S, Ohtsuki S, Tarui S, Ousaka D, Eitoku T, Kondo M, Okuyama M, Kobayashi J, Baba K, Arai S, Kawabata T, Yoshizumi K, Tateishi A, Kuroko Y, Iwasaki T, Sato S, Kasahara S, Sano S, Oh H. Intracoronary autologous cardiac progenitor cell transfer in patients with hypoplastic left heart syndrome (ticap): A prospective phase 1 controlled trial. *Circulation research*. 2014
 37. Ibrahim AG, Cheng K, Marban E. Exosomes as critical agents of cardiac regeneration triggered by cell therapy. *Stem Cell Reports*. 2014; 2:606–619. [PubMed: 24936449]
 38. Suzuki G, Weil BR, Leiker MM, Ribbeck AE, Young RF, Cimato TR, Canty JM Jr. Global intracoronary infusion of allogeneic cardiosphere-derived cells improves ventricular function and stimulates endogenous myocyte regeneration throughout the heart in swine with hibernating myocardium. *PloS one*. 2014; 9:e113009. [PubMed: 25402428]
 39. Bergersen L, Marshall A, Gauvreau K, Beekman R, Hirsch R, Foerster S, Balzer D, Vincent J, Hellenbrand W, Holzer R, Cheatham J, Moore J, Lock J, Jenkins K. Adverse event rates in congenital cardiac catheterization - a multi-center experience. *Catheter Cardiovasc Interv*. 2010; 75:389–400. [PubMed: 19885913]
 40. Lin CH, Hegde S, Marshall AC, Porras D, Gauvreau K, Balzer DT, Beekman RH 3rd, Torres A, Vincent JA, Moore JW, Holzer R, Armsby L, Bergersen L. Incidence and management of life-threatening adverse events during cardiac catheterization for congenital heart disease. *Pediatr Cardiol*. 2014; 35:140–148. [PubMed: 23900744]
 41. Makkar RR, Smith RR, Cheng K, Malliaras K, Thomson LE, Berman D, Czer LS, Marbán L, Mendizabal A, Johnston PV. Intracoronary cardiosphere-derived cells for heart regeneration after myocardial infarction (caduceus): A prospective, randomised phase 1 trial. *The Lancet*. 2012; 379:895–904.

42. Simpson DL, Mishra R, Sharma S, Goh SK, Deshmukh S, Kaushal S. A strong regenerative ability of cardiac stem cells derived from neonatal hearts. *Circulation*. 2012; 126:S46–S53. [PubMed: 22965993]
43. D'Amario D, Leone AM, Iaconelli A, Luciani N, Gaudino M, Kannappan R, Manchi M, Severino A, Shin SH, Graziani F, Biasillo G, Macchione A, Smaldone C, De Maria GL, Cellini C, Siracusano A, Ottaviani L, Massetti M, Goichberg P, Leri A, Anversa P, Crea F. Growth properties of cardiac stem cells are a novel biomarker of patients' outcome after coronary bypass surgery. *Circulation*. 2014; 129:157–172. [PubMed: 24249720]
44. Li TS, Cheng K, Malliaras K, Smith RR, Zhang Y, Sun B, Matsushita N, Blusztajn A, Terrovitis J, Kusuoka H, Marban L, Marban E. Direct comparison of different stem cell types and subpopulations reveals superior paracrine potency and myocardial repair efficacy with cardiosphere-derived cells. *Journal of the American College of Cardiology*. 2012; 59:942–953. [PubMed: 22381431]
45. Mirotso M, Jayawardena TM, Schmeckpeper J, Gnechi M, Dzau VJ. Paracrine mechanisms of stem cell reparative and regenerative actions in the heart. *Journal of molecular and cellular cardiology*. 2011; 50:280–289. [PubMed: 20727900]
46. Wehman B, Kaushal S. The emergence of stem cell therapy for patients with congenital heart disease. *Circulation research*. 2015; 116:566–569. [PubMed: 25677515]
47. Hatzistergos KE, Quevedo H, Oskouei BN, Hu Q, Feigenbaum GS, Margitich IS, Mazhari R, Boyle AJ, Zambrano JP, Rodriguez JE, Dulce R, Pattany PM, Valdes D, Revilla C, Heldman AW, McNiece I, Hare JM. Bone marrow mesenchymal stem cells stimulate cardiac stem cell proliferation and differentiation. *Circulation research*. 2010; 107:913–922. [PubMed: 20671238]
48. Ghanayem NS, Allen KR, Tabbutt S, Atz AM, Clabby ML, Cooper DS, Eghtesady P, Frommelt PC, Gruber PJ, Hill KD, Kaltman JR, Laussen PC, Lewis AB, Lurito KJ, Minich LL, Ohye RG, Schonbeck JV, Schwartz SM, Singh RK, Goldberg CS. Pediatric Heart Network I. Interstage mortality after the norwood procedure: Results of the multicenter single ventricle reconstruction trial. *J Thorac Cardiovasc Surg*. 2012; 144:896–906. [PubMed: 22795436]

Figure 1. Schematic of cardiopulmonary bypass-assisted intracoronary infusion of a cell-based therapy via the aortic root during cardiac surgery

Figure 2. Schematic of planned phase I clinical trial at the University of Maryland Medical Center for patients with hypoplastic left heart syndrome

Allogeneic mesenchymal stem cells will be delivered via intramyocardial injections at the time of stage II surgical palliation, after take down of the previously placed RVPA (Sano) shunt. RV=right ventricle; RPA=right pulmonary artery; SVC=superior caval vein; RVPA=right ventricle to pulmonary artery shunt.

Planned or ongoing clinical trials to evaluate safety, feasibility and preliminary efficacy of stem cell therapy for patients with congenital heart disease.

Table

Institution	Year Initiated	Phase	Disease	Estimated Enrollment	Stem Cell	Source	Delivery Method
Okayama University	2013	2	Single ventricle (Left or Right)	34	CDCs	Autologous	IC
Mayo Clinic	2013	1	HLHS	10	CBSCs	Autologous	IM
Duke University	2011	1	HLHS	20	CBSCs	Autologous	IV
University of Maryland Medical Center	2015	1	HLHS	30	MSCs	Allogeneic	IM

CDCs = cardiosphere-derived cells; IC=intracoronary; HLHS=hypoplastic left heart syndrome; CBSCs=umbilical cord blood stem cells; IM=intramyocardial; IV=intravenous