Published in final edited form as:

Med Sci Sports Exerc. 2016 March ; 48(3): 472–480. doi:10.1249/MSS.0000000000000789.

Muscle Characteristics and Substrate Energetics in Lifelong Endurance Athletes

John J. Dubé¹, Nicholas T. Broskey^{1,2}, Alex A. Despines¹, Maja Stefanovic-Racic¹, Frederico G.S. Toledo¹, Bret H. Goodpaster¹, and Francesca Amati^{1,2}

¹Department of Medicine, Division of Endocrinology and Metabolism, School of Medicine, University of Pittsburgh, Pittsburgh, PA ²Department of Physiology & Institute of Sport Sciences, School of Biology and Medicine, University of Lausanne, Lausanne, Switzerland

Abstract

Purpose—The goal of this study was to explore the effect of lifelong aerobic exercise (i.e. chronic training) on skeletal muscle substrate stores (intramyocellular triglyceride [IMTG] and glycogen), skeletal muscle phenotypes, and oxidative capacity (ox), in older endurance-trained master athletes (OA) compared to non-competitive recreational younger (YA) athletes matched by frequency and mode of training.

Methods—Thirteen OA (64.8±4.9 yo) exercising 5 times/week were compared to 14 YA (27.8±4.9 yo) males and females. IMTG, glycogen, fiber types, succinate dehydrogenase (SDH) and capillarization were measured by immunohistochemistry in *vastus lateralis* biopsies. Fat-ox and carbohydrate (CHO)-ox were measured by indirect calorimetry before and after an insulin clamp and during a cycle ergometer graded maximal test.

Results— $\dot{V}O_{2peak}$ was lower in OA than YA. OA had greater IMTG in all fiber types and lower glycogen stores than YA. This was reflected in greater proportion of type I and less type II fibers in OA. Type I fibers were similar in size, while type II fibers were smaller in OA compared to YA. Both groups had similar SDH content. Numbers of capillaries per fiber were reduced in OA but with a higher number of capillaries per area. Metabolic flexibility and insulin sensitivity were similar in both groups. Exercise metabolic efficiency was higher in OA. At moderate exercise intensities, CHO-ox was lower in OA but with similar Fatox.

Conclusion—Lifelong exercise is associated with higher IMTG content in all muscle fibers and higher metabolic efficiency during exercise that are not explained by differences in muscle fibers

Corresponding author: Francesca Amati, University of Lausanne, Bugnon 7, Lausanne 1005, Switzerland, Francesca.amati@unil.ch, Phone: + 41-21-692-5552, Fax: + 41-21-692-5505.

The authors declare no conflict of interest. The results in the present study do not constitute endorsement by ACSM.

AUTHORS CONTRIBUTIONS

J.J.D. researched data, contributed to the study concept, design and wrote the manuscript. N.T.B and A.D. researched data. F.G.S.T and M.S.R. performed biopsies. B.H.G. contributed to the study concept, interpretation of the data and edited the manuscript. F.A. researched data, contributed to the study concept, design, analysis, and interpretation of the data; and wrote the manuscript.

CONFLICT OF INTEREST

The authors declare no conflict of interest. The results in the present study do not constitute endorsement by ACSM.

types and other muscle characteristics when comparing older to younger athletes matched by exercise mode and frequency.

Keywords

Aging; chronic exercise; muscle fibers; capillary density; energy expenditure; substrate oxidation; insulin sensitivity

INTRODUCTION

Aging is associated with a decline in physical capacity and modifications of muscle phenotype (34) leading to increased overall morbidity and risk for development of cardiometabolic diseases. Aerobic training interventions suggest that aged skeletal muscle remains malleable to sustain the functional and metabolic demands of exercise (6) demonstrated by a shift towards higher content of type I fibers and relative decrease in type IIx fibers (29), increased fiber cross sectional area (22), enhanced oxidative capacity (39), capillary angiogenesis (35) and elevated glycogen stores (33). Further, we have previously demonstrated that chronic aerobic training in older adults increases intramyocellular triglyceride (IMTG) stores (9) and reliance on fat metabolism (2) during exercise.

Despite the growing body of literature demonstrating alterations in skeletal muscle substrate content and capacity for oxidation in previously sedentary subjects, few studies have compared chronic aerobic training adaptations in young and old athletes. Current evidence supports the notion that being physically active throughout a person's life (lifelong) protects oxidative fiber number and size, as well as mitochondrial function when compared to younger trained (39) and older sedentary (1, 45) subjects. These retained muscle adaptations to exercise seem to provide functional benefits such as improved balance, gait speed and ability to get up from a chair (45), which in turn are likely to improve quality of life and reduce risk of falling. Yet, the impact of lifelong aerobic training on skeletal muscle metabolism within the context of whole-body substrate oxidation and insulin sensitivity is still largely unknown.

The primary goal of this study was to determine skeletal muscle substrate storage and capacity for oxidation, as well as exercise metabolic efficiency in older masters athletes and younger subjects matched by frequency and mode of training. A secondary goal was to determine if differences in skeletal muscle substrate storage was associated with differences in substrate oxidation under different physiological conditions. We hypothesized that despite lower peak aerobic capacity in older master athletes, lifelong aerobic training in this group would result in similar skeletal muscle substrate storage compared to the younger athletes matched by exercise mode and frequency, as well as similar oxidative capacity, metabolic efficiency and substrate oxidation under same physiological conditions.

METHODS

Subjects

Fourteen younger (age 18-39) and 13 older (age 60-75) endurance-trained athletes were recruited for this cross-sectional comparison. To be included, older women and men were

training 5 or more structured aerobic exercise sessions per week either in running, cycling, swimming, or aerobic dancing (fitness classes). Younger athletes were non-competitive recreational athletes matched by frequency and mode of training with at least 3 years of uninterrupted (>3 months) training. Habitual physical activity was self-reported and discussed during the screening visit medical interview, including exercise mode, frequency and training years. All subjects were in general good health, non-smokers, weight stable and training stable for the last 6 months. The University of Pittsburgh Institutional review board approved the protocol and all volunteers gave written consent.

Body composition

Total body fat-free (FFM), fat mass (FM) and percent body fat were measured by dual-emission X-ray absorptiometry (Lunar Prodigy; GE Healthcare, Milwaukee, MI).

Physical fitness

$\dot{V}O_2$ peak was assessed by a graded exercise test on an electronically braked cycle ergometer (Excalibur, Lode B.V., Groningen, The Netherlands) in conjunction with indirect calorimetry (Moxus, AEI Technologies, Pittsburgh, PA). The initial workload was set depending on the sex and age of the individual (50 W for younger and older women, 75 W for older men, 100 W for younger men) for the first 2 minutes and then increased by 50 W (men) or 25 W (women) every 2 minutes thereafter until volitional exhaustion or one of the established criteria for $\dot{V}O_2$ peak had been reached (38). Heart rate, blood pressure, and ECG were recorded before, during and immediately after this test.

Skeletal muscle biopsies

Percutaneous muscle biopsies were obtained from the *vastus lateralis* as described previously (1). Subjects were asked to refrain from exercise in the last 48 hours before the biopsy. Subjects were admitted to the Clinical and Translational Research Center (CTRC) in the evening and received a standard dinner (7.5 kcal·kg⁻¹ of body weight, 50% carbohydrate, 30% fat and 20% protein). The biopsy was performed the following morning at 7 AM after an overnight fast. Samples were trimmed of all visible adipose tissue with a dissecting microscope (Leica EZ4, Leica Microsystems, Wetzlar, Germany) and blotted dry. The muscle specimen was mounted on a small piece of cork with mounting medium, placed in liquid nitrogen cooled isopentane and then placed into liquid nitrogen. All samples were stored at -80 degrees Celsius until analysis.

Immunohistochemistry

Histochemical analyses were performed on 10 µm serial sections using methods previously described (9). IMTG content was determined by Oil Red O (ORO) and fiber type costain (1) allowing fiber specific IMTG measurements and cross sectional area. Succinate dehydrogenase (SDH, complex II of the electron transport chain) staining was used as a marker of oxidative capacity (40). Glycogen content was measured using a standard Shiffs reagent protocol (23). Capillary density was determined as previously described (9). Capillary density was computed as total number of capillaries per cross sectional area of

tissue (capillaries/area). The number of fibers in the cross sectional area of tissue is reported as the ratio fiber/area and the number of capillaries per fiber as the ratio capillaries/fiber.

Whole body substrate oxidation and exercise efficiency

Indirect calorimetry was used to measure $\dot{V}O_2$ and $\dot{V}CO_2$ under three physiological conditions: 1) in the fasted state between 6 and 7 AM (prior to the biopsy described above), 2) in the post-prandial state at the end of an hyperinsulinemic euglycemic clamp, and 3) during the graded exercise test described above. Systemic rates of fat oxidation (Fat-ox) and carbohydrate (CHO-ox) were calculated using the adapted stoichiometric equations of Frayn (13):

$$\text{Fat-ox (mg/min)} = 1.67 \dot{V}O_{2(\text{ml/min})} - 1.67 \dot{V}CO_{2(\text{ml/min})}$$

$$\text{CHO-ox (mg/min)} = 4.55 \dot{V}CO_{2(\text{ml/min})} - 3.21 \dot{V}O_{2(\text{ml/min})}$$

To compute the proportion of energy expended from carbohydrates or fat, Fat-ox and CHO-ox were transformed in kilocalories per minute and expressed as a proportion of resting energy derived from fat or carbohydrates as used previously (2). For substrate oxidation during the graded exercise test, only points corresponding to a respiratory exchange ratio <1 were used to account for possible changes in the size of the bicarbonate pool during maximal exercise (26). Protein oxidation rates were not included based on our laboratory's prior work demonstrating that rates of urinary nitrogen excretion were similar in different body phenotypes during resting conditions (19) and on the assumption that the amount of protein oxidized, as well as other metabolic processes, such as gluconeogenesis from protein, ketone body formation, and lipogenesis during exercise, are quantitatively negligible compared with glucose and fatty acid oxidation (37).

To account for possible aging and sex biases, all physiological data were normalized to FFM. Glucose uptake (glucose oxidase, [YSI, Yellow Springs, Colorado]) and plasma insulin (ELIZA, [Millipore, Billerica, MA]) were used to calculate insulin sensitivity ($\text{mg} \cdot \text{kgFFM}^{-1} \cdot \text{min}^{-1} \cdot \text{unit insulin}^{-1}$) during the steady state of the clamp.

During the graded exercise test, metabolic efficiency was measured as delta efficiency in percent for each consecutive stages as the difference in watts divided by the difference in $\dot{V}O_2$ (14). This was performed for each submaximal stage using the average $\dot{V}O_2$ for the last 30 seconds of each stage. Further, to obtain overall delta efficiency (η), linear regressions were drawn for each subject using all the submaximal stages. The average slopes and intercepts for each group were used to define the relationship $\dot{V}O_2 = bW + a$, where b is the slope and a the intercept. The inverse of the slope $1/b = \dot{W} / \dot{V}O_2$ is η (12).

Statistical Procedures

Subject characteristics are presented as means $\pm$ SD, all other data are presented as means $\pm$ SEM. After checking normality and equality of variance, two tailed independent t -tests were

performed to examine group differences. If the equality of variance assumption was not met, comparisons between groups were performed with the Welch corrected *t*-test. If the normality assumption was not met, comparisons between groups were performed with the non-parametric Median test. For substrate oxidation comparisons in fasted and fed conditions, 2×2 mixed MANOVA were performed. For substrate use during the graded exercise test, repeated mixed MANOVA were used with group X time. When needed pair-wise post hoc analyses were used to identify the significant difference.

RESULTS

Subject characteristics

Subject characteristics are presented in Table 1. Training years were between ~ 35-40 years for the older masters athletes and 5-13 years for younger subjects. FFM, FM and percent body fat were not different between age groups. Younger athletes had a higher $\text{VO}_{2\text{peak}}$ than older athletes with a magnitude of ~25% when expressed relative to FFM. Self reported activities were on average 6 sessions/week with running as the most common physical activity (62%), followed by biking (23%), brisk walking and aerobic fitness classes (both 8%). In addition of their main exercise mode, cross-training and seasonal activities included skiing, golfing and swimming.

Skeletal muscle lipid storage is greater in older compared to younger endurance-trained athletes

Chronic aerobic training increases skeletal muscle substrate storage in young and old previously sedentary subjects. Yet the effects of lifelong aerobic training on skeletal muscle adaptations are largely unknown. Older athletes had higher content of IMTG in each fiber type measured (Figure 1, Panel A), as well as overall greater content of IMTG. Glycogen content (Figure 1, Panel B) was higher in young athletes compared to old, while no differences in SDH (Figure 1, Panel C) were noted.

Oxidative fibers are higher in older compared to younger endurance-trained athletes

Older athletes had higher proportion of type I fibers and lower type IIa fibers than younger athletes (Figure 2, Panel A). The proportion of type IIx fibers was not different between groups. Mean area of type I fibers was similar in both groups, while younger athletes had larger IIa and IIx fiber area (Figure 2, Panel B). These data suggest that lifelong physical activity may not prevent the proposed age related decline in type II fiber area (31).

Capillary density is lower in older compared to younger endurance-trained athletes

As skeletal muscle capillary density is affected by aging and type 2 diabetes (21) and is associated with oxidative capacity (9), we next determined if capillary density was associated with the observed differences in oxidative fibers. While the number of capillaries per fiber was higher in the younger (Figure 3, Panel A) athletes, capillary density relative to muscle area was higher in the older athletes (Figure 3, Panel B). These data suggest that the decline in capillary density associated with sedentary aging (21) is attenuated with lifelong aerobic exercise.

Metabolic flexibility and insulin sensitivity are similar in older compared to younger endurance-trained athletes

Given the observed differences in skeletal muscle substrate composition and capacity for oxidation, we next examined whether or not these differences translated into changes in whole-body substrate oxidation and insulin sensitivity. Older athletes had higher resting energy expenditure in fasting condition, while younger athletes had higher energy expenditure in postprandial condition (Figure 4, Panel A, significant interaction $P=0.01$). The proportion of substrate use during both states was comparable in both groups (Figure 4, panel B). Metabolic flexibility, originally defined by the overall change in RQ from fasting to postprandial (28) was similar in both groups (Figure 4, Panel C, insulin effect $P<0.0001$). Insulin stimulated glucose uptake was similar in younger and older athletes (Figure 4, Panel D), with no differences in nonoxidative and oxidative disposal. Together these data suggest that lifelong endurance training protects older adults from declines in metabolic flexibility and insulin sensitivity. Moreover, relative fat- and carbohydrate-oxidation rates for basal and insulin-stimulated substrate use under non-exercising conditions are maintained throughout the lifespan with aerobic exercise.

Exercise metabolic efficiency is enhanced in older compared to younger endurance-trained athletes

We previously demonstrated that exercise training resulted in improved skeletal muscle oxidative capacity (9) and exercise efficiency (2) in previously sedentary older adults. Based on the differences in peak aerobic capacity and substrate storage in older athletes, we next calculated exercise metabolic efficiency during a graded exercise test. Older athletes had higher exercise metabolic efficiency compared to younger athletes (η of 9.03 ± 0.32 and $8.03\pm0.26\%$, $P=0.02$). Regression curves for each group, including slope and intercept are presented in Figure 5, Panel A ($P=0.02$ [older] and $P=0.13$ [younger]). Stage by stage delta efficiency is presented in Figure 5, Panel B (2×5 MANOVA not significant, point by point independent T tests are presented in the figure).

Moderate intensity exercise carbohydrate oxidation rates are lower in older compared to younger endurance-trained athletes

At moderate relative intensities, younger athletes had greater rates of carbohydrate oxidation compared to older (Figure 5, Panel C). Fat oxidation was higher in the younger at very low intensities (Figure 5, Panel C). It is important to note that the removal of time points with $RER>1$, to avoid underestimation of fat oxidation and overestimation of carbohydrate oxidation due to a potential difference in the bicarbonate buffering system and excess non oxidative CO_2 exhaled, reduces greatly the number of subjects included in the higher intensities measurements. Nevertheless, these data suggest that the observed increase in IMTG and oxidative fibers may contribute to the enhanced exercise metabolic efficiency.

DISCUSSION

The overall goal of this study was to investigate chronic aerobic exercise training on skeletal muscle substrate adaptations, as well as systemic oxidation in young and older endurance trained subjects. To achieve this goal we examined skeletal muscle phenotypes, as well as

whole-body substrate utilization using indirect calorimetry in two cohorts of subjects with similar endurance training regimens. We found that, despite lower peak aerobic capacity, lifelong master athletes have higher intramyocellular triglyceride (IMTG) and proportion of oxidative fibers compared to younger athletes. These differences were reflected in enhanced exercise metabolic efficiency with lower reliance on carbohydrate oxidation during exercise in the older subjects (at higher intensities). Together the data suggest that lifelong aerobic exercise, not only attenuates the age associated decreases in muscle oxidative potential, but also provides older endurance-trained subjects with an enhanced capacity for fatty acid oxidation.

Age-induced increases in intramyocellular lipids have been observed in previous human studies. Under sedentary conditions, this phenomenon is associated with a decline in muscle mass and strength (8, 16), as well as decreased insulin action (36). While decreases in muscle mass, fiber cross sectional area, and shifts in fiber type composition may explain, in part, intramyocellular lipid deposition in sedentary conditions (8, 18), this is not the case for the chronically trained older individuals in the current study. We have previously exposed that the “athlete's paradox” observed in younger endurance trained athletes (17) was also present in older endurance trained athletes compared to sedentary controls (1). A key novel finding in the present study is that older endurance trained athletes have greater lipid, yet lower carbohydrate stores, compared to younger athletes with similar training regimens. While aging *per se* has been associated with increased lipid uptake (44), chronic exercise training increases factors associated with IMTG turnover (i.e. storage and lipolysis) (1). We hypothesize that the combination of these age- and exercise-related alterations in IMTG turnover likely mediates, in part, the increased IMTG in this cohort. Proteins involved in IMTG storage are elevated in exercise-trained muscle (1, 4, 10) (amati, diabetes, 2011; dube, diabetologia 2011, Bergman, JAP, 2010). Additional studies are needed to investigate whether these, or other mechanisms for the increased IMTG storage, are altered in older endurance athletes.

In contrast to higher IMTG levels, older subjects demonstrated lower muscle glycogen stores compared to younger subjects. Although controversial, there is a suggestion that glycolytic activity (5), as well as type II fiber proportion and size (discussed below) may be reduced with aging. However, aerobic exercise training in previously sedentary older adults has been demonstrated to increase muscle glycogen content (9). Possible explanations to the lower glycogen content in older trained subjects is that younger endurance athletes may engage in relatively more frequent high-intensities and/or that younger athletes may have altered post-exercise carbohydrate consumption relative to older athletes, thus providing the necessary stimulus for enhanced glycogen storage (25). Nevertheless, lower glycogen content in our older athletes did not contribute to alterations in basal or insulin-stimulated rates of substrate oxidation. Rather, the functional relevance was only observed at maximal intensity exercise. These data support the notion that lifelong endurance training may better position older athletes for moderate intensity activities with relative higher fat oxidation, while young athletes may be positioned for high intensity exercise (i.e. higher glycogen). Thus the capacity for moderate high fat oxidation activity may be enhanced with lifelong endurance training.

Based on our novel demonstration of increased lipid stores with lifelong exercise training, we next examined the potential mechanisms associated with this phenomenon. While several studies have suggested that aging results in the atrophy of type II fibers (20, 39), with a relative increase of the area occupied by type I fibers (30), this is not without controversy. Our data suggest that lifelong exercise training is accompanied by a shift toward greater slow oxidative fibers with no change in the overall size of these fibers (45). Interestingly, not only was the relative percentage of glycolytic fibers decreased in older trained subjects, the mean area was also decreased. These data suggest that if an aging decrease in glycolytic fibers occurs, perhaps exercise training promotes a compensatory increase in oxidative fibers. This new harmony between type I and type II fibers observed in the aging and trained muscle may explain, at least in part, the distinction in substrate stores between older and younger muscle of endurance trained athletes witnessed in this study.

Previous studies have demonstrated that, while master athletes have significantly higher peak fitness levels compared to sedentary age-matched controls (41), the age-related decline in fitness persists despite continuous training. Thus, as expected, $\dot{V}O_{2peak}$, both absolute and adjusted to fat free mass, was higher in younger than older athletes. Peak fitness may be limited by two key peripheral factors, capillarization (3, 24) and mitochondrial capacity (3). While capillary density, relative to the number of fibers, was lower in older trained subjects, adjusting the data to the lower number and cross sectional area of glycolytic fibers suggests that capillary density is not different between the cohorts (7). This interpretation is in accord with previous studies that found similar adaptations in capillarization between older and younger adults undergoing an exercise intervention (15, 35). With respect to mitochondria, it has been reported that mitochondrial respiration (21), mitochondrial biogenesis (32), and perhaps oxidative capacity and energy production decline with aging. However, it's generally accepted that aerobic exercise training, in both older (9) and younger (11) previously sedentary subjects, results in enhanced mitochondrial oxidative capacity. In agreement with data from Proctor et al. (39), we did not observe any differences in mitochondrial capacity between the cohorts in this study. Thus, the difference in $\dot{V}O_{2peak}$ observed in our younger and older athletes seems to be explained mostly by the central component. This is in agreement with previous studies suggesting that peripheral factors play an important role in the elderly in the response to endurance exercise training (33). Together our data suggest that while lifelong exercise training may not prevent the age-associated loss of skeletal muscle capillarization, the overall capacity for substrate oxidation, as well as overall fitness is enhanced relative to sedentary subjects regardless of age (1).

Based on our demonstration of enhanced lipid stores and similar capacity for oxidation, we next examined whole-body substrate utilization under different physiological conditions. Previous studies have reported age-related declines in the capacity of skeletal muscle to oxidize fat in the fasting state and during exercise (42, 44). In this study, higher energy expenditure at rest was not associated with differences in substrate selection in the older athletes. These data are in stark contrast to previous reports from sedentary subjects (27) demonstrating a significant reduction in resting energy expenditure in older subjects adjusted for fat free mass. We speculate that the increased basal energy expenditure may be due to the modest but not significant BMI and gender difference between the groups (see below). Nevertheless, our data clearly indicate the lifelong training preserves basal energy

expenditure, as well as rates of both fat and carbohydrate oxidation in the basal and insulin-stimulated conditions. Thus, lifelong exercise training preserves metabolic flexibility and substrate selection with aging.

During moderate intensity exercise, older athletes used fewer carbohydrates for energy. These data are in agreement with our demonstration of greater muscle glycogen content in younger subjects. At maximal intensity, no conclusions can be drawn from our indirect calorimetry data due to the limitations of stoichiometric computations in anaerobic conditions. Intervention studies have concluded that previously sedentary older subjects undergoing endurance exercise interventions of 16 weeks were able to improve their reliance of fat during a one hour submaximal exercise (2, 43), thus our data may be explained by the maintenance of substrate oxidation in older athletes as well as by the shift towards type I fibers. Interestingly in our cohort, the higher muscle efficiency observed in the older athletes during the graded exercise test cannot be explained by different substrate use during exercise, but may be influenced by the greater number of capillaries per fibers and the higher proportion of type I fibers (2). Together these data suggest that lifelong aerobic exercise preserves, or perhaps enhances, resting exercise expenditure, as well as metabolic flexibility and substrate oxidation under physiological conditions.

This study is not without limitations. First, training regimens (frequency, mode) were self-reported. However, our data are in accord with previous reports of overall fitness and body composition in older and younger athletes (7, 39). Although we attempted to include equal numbers of males and females, males represent 50% in the younger group and 69% in the older group. While the chi-square test for sampling distribution was not significant, this discrepancy may influence some of the results. We believe that if so, this would have been in disfavor of the older group as women have relative lower exercise capacity and higher insulin sensitivity than men and thus, if the gender balance was important, we would have probably seen unequal insulin sensitivity and markers of oxidative capacity between our two groups.

In summary, the results of the present study demonstrate that lifelong endurance training results in increased skeletal muscle lipid stores and shift toward greater numbers of oxidative fibers. While exercise metabolic efficiency was enhanced, older endurance trained subjects had lower glycogen and glycolytic fiber content, as well as a lower reliance on carbohydrate oxidation at moderate intensities. We conclude that these physiological adaptations to chronic aerobic training in older subjects may place them in an optimal position for moderate high-fat oxidation activity. Moreover, these data provide further evidence against triglyceride-mediated impairments in metabolic function. Conversely, the demonstration of higher muscle glycogen content in younger subjects supports the notion of a higher capacity for high-intensity training, supported by enhanced carbohydrate oxidation observed in this study. Our studies raise further questions on lifelong adaptations to exercise in terms of increased efficiency without modifying the balance between sources of substrate oxidation. Additionally, these data further emphasize the importance of chronic exercise throughout life to attenuate the deleterious effects of aging and sedentary lifestyle.

ACKNOWLEDGMENTS

We appreciate the cooperation of our research volunteers, Steve Anthony, Nicole Helbling and the nursing staff of the Clinical and Translational Research Center of the University of Pittsburgh. This study was supported by the National Institutes of Health (NIH) Grants (K01-DK-084213-01 to J.J.D. and R01-AG20128 to B.H.G.), and University of Pittsburgh Student and Faculty Grants (to F.A.), and to the University of Pittsburgh CTRC (M01-RR-00056).

REFERENCES

- Amati F, Dube JJ, Alvarez-Carnero E, et al. Skeletal muscle triglycerides, diacylglycerols, and ceramides in insulin resistance: another paradox in endurance-trained athletes? *Diabetes*. 2011; 60(10):2588–97. [PubMed: 21873552]
- Amati F, Dube JJ, Shay C, Goodpaster BH. Separate and combined effects of exercise training and weight loss on exercise efficiency and substrate oxidation. *Journal of Applied Physiology*. 2008; 105(3):825–31. [PubMed: 18617627]
- Bassett DR Jr, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *MedSciSportsExerc*. 2000; 32(1):70–84.
- Bergman BC, Perreault L, Hunerdosse DM, Koehler MC, Samek AM, Eckel RH. Increased intramuscular lipid synthesis and low saturation relate to insulin sensitivity in endurance-trained athletes. *Journal of applied physiology* (Bethesda, Md.: 1985). 2010; 108(5):1134–41.
- Coggan AR, Spina RJ, King DS, et al. Histochemical and enzymatic comparison of the gastrocnemius muscle of young and elderly men and women. *Journal of gerontology*. 1992; 47(3):B71–6. [PubMed: 1573181]
- Coggan AR, Spina RJ, King DS, et al. Skeletal muscle adaptations to endurance training in 60- to 70-yr-old men and women. *J Appl Physiol*. 1992; 72(5):1780–6. [PubMed: 1601786]
- Coggan AR, Spina RJ, Rogers MA, et al. Histochemical and enzymatic characteristics of skeletal muscle in master athletes. *J Appl Physiol*. 1990; 68(5):1896–901. [PubMed: 2361892]
- Cree MG, Newcomer BR, Katsanos CS, et al. Intramuscular and liver triglycerides are increased in the elderly. *J Clin Endocrinol Metab*. 2004; 89(8):3864–71. [PubMed: 15292319]
- Dube JJ, Amati F, Stefanovic-Racic M, Toledo FGS, Sauers SE, Goodpaster BH. Exercise-induced alterations in intramyocellular lipids and insulin resistance: the athlete's paradox revisited. *American Journal of Physiology-Endocrinology and Metabolism*. 2008; 294(5):E882–E8. [PubMed: 18319352]
- Dube JJ, Amati F, Toledo FG, et al. Effects of weight loss and exercise on insulin resistance, and intramyocellular triacylglycerol, diacylglycerol and ceramide. *Diabetologia*. 2011; 54(5):1147–56. [PubMed: 21327867]
- Dube JJ, Coen PM, DiStefano G, et al. Effects of acute lipid overload on skeletal muscle insulin resistance, metabolic flexibility, and mitochondrial performance. *Am J Physiol EndocrinolMetab*. 2014; 307(12):E1117–24.
- Francescato MP, Girardis M, di Prampero PE. Oxygen cost of internal work during cycling. *European journal of applied physiology and occupational physiology*. 1995; 72(1-2):51–7. [PubMed: 8789570]
- Frayn KN. Calculation of substrate oxidation rates in vivo from gaseous exchange. *Journal of applied physiology: respiratory, environmental and exercise physiology*. 1983; 55(2):628–34.
- Gaesser GA, Brooks GA. Muscular efficiency during steady-rate exercise: effects of speed and work rate. *J Appl Physiol*. 1975; 38(6):1132–9. [PubMed: 1141128]
- Gavin TP, Ruster RS, Carrithers JA, et al. No difference in the skeletal muscle angiogenic response to aerobic exercise training between young and aged men. *The Journal of Physiology*. 2007; 585(Pt 1):231–9. [PubMed: 17884919]
- Goodpaster BH, Carlson CL, Visser M, et al. Attenuation of skeletal muscle and strength in the elderly: The Health ABC Study. *JApplPhysiol*. 2001; 90(6):2157–65.

17. Goodpaster BH, He J, Watkins S, Kelley DE. Skeletal muscle lipid content and insulin resistance: evidence for a paradox in endurance-trained athletes. *J Clin Endocrinol Metab.* 2001; 86(12): 5755–61. [PubMed: 11739435]
18. Goodpaster BH, Park SW, Harris TB, et al. The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. *J Gerontol A BiolSci Med Sci.* 2006; 61(10):1059–64.
19. Goodpaster BH, Wolfe RR, Kelley DE. Effects of obesity on substrate utilization during exercise. *Obesity research.* 2002; 10(7):575–84. [PubMed: 12105277]
20. Grimby G, Danneskiold-Samsøe B, Hvid K, Saltin B. Morphology and enzymatic capacity in arm and leg muscles in 78-81 year old men and women. *Acta Physiol Scand.* 1982; 115(1):125–34. [PubMed: 7136797]
21. Groen BB, Hamer HM, Snijders T, et al. Skeletal muscle capillary density and microvascular function are compromised with aging and type 2 diabetes. *Journal of applied physiology* (Bethesda, Md.: 1985). 2014; 116(8):998–1005.
22. Harber MP, Konopka AR, Undem MK. Aerobic exercise training induces skeletal muscle hypertrophy and age-dependent adaptations in myofiber function in young and older men. *JApplPhysiol.* 2012; 113(9):1495–504.
23. He J, Kelley DE. Muscle glycogen content in type 2 diabetes mellitus. *Am J Physiol EndocrinolMetab.* 2004; 287(5):E1002–7.
24. Honig CR, Connett RJ, Gayeski TE. O₂ transport and its interaction with metabolism; a systems view of aerobic capacity. *Med Sci Sports Exerc.* 1992; 24(1):47–53. [PubMed: 1548995]
25. Jensen J, Tantiwong P, Stuenkel JT, et al. Effect of acute exercise on glycogen synthase in muscle from obese and diabetic subjects. *Am J Physiol Endocrinol Metab.* 2012; 303(1):E82–9. [PubMed: 22510711]
26. Jeukendrup AE, Wallis GA. Measurement of substrate oxidation during exercise by means of gas exchange measurements. *International journal of sports medicine.* 2005; 26(Suppl 1):S28–37.
27. Johannsen DL, DeLany JP, Frisard MI, et al. Physical activity in aging: comparison among young, aged, and nonagenarian individuals. *Journal of applied physiology* (Bethesda, Md.: 1985). 2008; 105(2):495–501.
28. Kelley DE, Mandarino LJ. Fuel selection in human skeletal muscle in insulin resistance: a reexamination. *Diabetes.* 2000; 49(5):677–83. [PubMed: 10905472]
29. Konopka AR, Trappe TA, Jemiolo B, Trappe SW, Harber MP. Myosin heavy chain plasticity in aging skeletal muscle with aerobic exercise training. *J Gerontol A Biol Sci Med Sci.* 2011; 66(8): 835–41. [PubMed: 21659340]
30. Korhonen MT, Cristea A, Alen M, et al. Aging, muscle fiber type, and contractile function in sprint-trained athletes. *J Appl Physiol.* 2006; 101(3):906–17. [PubMed: 16690791]
31. Lee WS, Cheung WH, Qin L, Tang N, Leung KS. Age-associated decrease of type IIA/B human skeletal muscle fibers. *Clinical orthopaedics and related research.* 2006; 450:231–7. [PubMed: 16691139]
32. Lopez-Lluch G, Irusta PM, Navas P, de Cabo R. Mitochondrial biogenesis and healthy aging. *Experimental gerontology.* 2008; 43(9):813–9. [PubMed: 18662766]
33. Meredith CN, Frontera WR, Fisher EC, et al. Peripheral effects of endurance training in young and old subjects. *J Appl Physiol.* 1989; 66(6):2844–9. [PubMed: 2745349]
34. Miller MS, Callahan DM, Toth MJ. Skeletal muscle myofilament adaptations to aging, disease, and disuse and their effects on whole muscle performance in older adult humans. *Frontiers in physiology.* 2014; 5:369. [PubMed: 25309456]
35. Murias JM, Kowalchuk JM, Ritchie D, Hepple RT, Doherty TJ, Paterson DH. Adaptations in capillarization and citrate synthase activity in response to endurance training in older and young men. *J Gerontol A Biol Sci Med Sci.* 2011; 66(9):957–64. [PubMed: 21715648]
36. Pan DA, Lillioja S, Kriketos AD, et al. Skeletal muscle triglyceride levels are inversely related to insulin action. *Diabetes.* 1997; 46(6):983–8. [PubMed: 9166669]
37. Peronnet F, Massicotte D. Table of nonprotein respiratory quotient: an update. *Canadian journal of sport sciences = Journal canadien des sciences du sport.* 1991; 16(1):23–9. [PubMed: 1645211]

38. Pescatello, LS. ACSM's guidelines for exercise testing and prescription. 9th ed.. Vol. xxiv. Wolters Kluwer/Lippincott Williams & Wilkins Health; Philadelphia: 2014. American College of Sports Medicine.; p. 456
39. Proctor DN, Sinning WE, Walro JM, Sieck GC, Lemon PW. Oxidative capacity of human muscle fiber types: effects of age and training status. *J Appl Physiol.* 1995; 78(6):2033–8. [PubMed: 7665396]
40. Pruchnic R, Katsiaras A, He J, Kelley DE, Winters C, Goodpaster BH. Exercise training increases intramyocellular lipid and oxidative capacity in older adults. *Am J Physiol Endocrinol Metab.* 2004; 287(5):E857–62. [PubMed: 15226098]
41. Rogers MA, Evans WJ. Changes in skeletal muscle with aging: effects of exercise training. *Exerc Sport Sci Rev.* 1993; 21:65–102. [PubMed: 8504850]
42. Sial S, Coggan AR, Carroll R, Goodwin J, Klein S. Fat and carbohydrate metabolism during exercise in elderly and young subjects. *Am J Physiol.* 1996; 271(6 Pt 1):E983–9. [PubMed: 8997215]
43. Sial S, Coggan AR, Hickner RC, Klein S. Training-induced alterations in fat and carbohydrate metabolism during exercise in elderly subjects. *Am J Physiol.* 1998; 274(5 Pt 1):E785–90. [PubMed: 9612234]
44. Tucker MZ, Turcotte LP. Impaired fatty acid oxidation in muscle of aging rats perfused under basal conditions. *Am J Physiol Endocrinol Metab.* 2002; 282(5):E1102–9. [PubMed: 11934676]
45. Zampieri S, Pietrangelo L, Loeffler S, et al. Lifelong physical exercise delays age-associated skeletal muscle decline. *J Gerontol A Biol Sci Med Sci.* 2015; 70(2):163–73. [PubMed: 24550352]

Figure 1. Skeletal muscle fiber type proportion (panel A) and cross sectional area (panel B) in younger and older athletes. MHC= myosin heavy chain. * $P < 0.05$ ** $P < 0.001$ two tailed independent *t*-test.

Figure 2. Intramyocellular triglycerides (panel A), glycogen (panel B) and SDH content (panel C) in younger and older athletes. MHC= myosin heavy chain, A.U.= arbitrary units. * $P < 0.05$ ** $P < 0.001$ two tailed independent t -test.

Figure 3. Skeletal muscle capillary density: Number of capillaries per fiber (panel A), number of fibers per area and capillaries per area (panel B). **P<0.001 two tailed independent *t*-test, §<0.05 non parametric Median test.

Figure 4.

Energy expenditure (panel A) and substrate use at rest in the fasted and post-prandial phase (panel B), metabolic flexibility (panel C) and insulin-stimulated glucose uptake (panel D). FFM=fat free mass, RQ=respiratory quotient, CHO=carbohydrate. *Significant interaction effect, **Significant effect of time in 2x2 mixed MANOVA.

Figure 5.

Delta efficiency and substrate use during graded exercise test in older and younger endurance trained athletes. Panel A represents the regression lines defining oxygen uptake as a function of power output. The insert is the magnification of the origin of the axis (box).

*Significant difference on the slope but not on the intercept. Panel B is delta efficiency between consecutive stages. * $P < 0.05$, #= 0.09 two tailed independent t -test. Panel C is

carbohydrate and fat oxidation as a function of relative intensity of peak oxygen consumption. CHO=carbohydrate. * $P < 0.05$ two tailed independent t -test.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Subjects' characteristics

	Younger (7 M / 7 F)	Older (9 M / 4 F)	P-Value (two tailed)
Age (years)	27.8 ± 4.9	64.8 ± 4.9	<0.0001
Body weight (BW, kg)	65.66 ± 10.78	68.15 ± 9.97	0.541
BMI (kg/m ²)	22.12 ± 1.74	23.76 ± 2.50	0.058
Fat free mass (FFM, kg)	53.29 ± 10.66	54.00 ± 9.73	0.858
Fat mass (kg)	12.03 ± 4.17	14.15 ± 7.79	0.395
Body fat (%)	18.68 ± 6.47	20.63 ± 10.76	0.578
VO ₂ peak (l/min)	3.816 ± 1.088	3.031 ± 0.763	0.041
VO ₂ peak (ml/min/kgFFM)	70.53 ± 9.08	55.82 ± 7.22	<0.0001
VO ₂ peak (ml/min/kgBW)	57.41 ± 2.56	44.62 ± 2.86	0.003
Peak power output (watts)	285.7 ± 20.3	201.9 ± 13.4	0.002

Mean ± SD