Published in final edited form as:

J Infect. 2016 March ; 72(3): 302–308. doi:10.1016/j.jinf.2015.11.002.

Increased *In Vitro* Phenol-Soluble Modulin Production is Associated with Soft Tissue Infection Source in Clinical Isolates of Methicillin-Susceptible *Staphylococcus aureus*

Robert Qi¹, Hwang-Soo Joo², Batu Sharma-Kuinkel¹, Nicholas R. Berlon¹, Lawrence Park¹, Chih-lung Fu², Julia A. Messina^{1,3}, Joshua T. Thaden¹, Qin Yan¹, Felicia Ruffin¹, Stacey Maskarinec¹, Bobby Warren¹, Vivian H. Chu¹, Claudio Q. Fortes⁴, Efthymia Giannitsioti⁵, Emanuele Durante-Mangoni⁶, Zeina A. Kanafani⁷, Michael Otto², and Vance G. Fowler Jr^{1,3}

¹Division of Infectious Diseases, Department of Medicine, Duke University Medical Center, Durham, NC, USA

²Pathogen Molecular Genetics Section, Laboratory of Bacteriology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA

³Duke Clinical Research Institute, Duke University Medical Center, Durham, NC, USA

⁴Hospital Universitario Clementino Fraga Filho/UFRJ, Rio de Janeiro, Brazil

⁵Attikon University General Hospital, Athens, Greece

⁶Internal Medicine Section, Department of Cardiothoracic Sciences, and Division of Infectious and Transplant Medicine, Second University of Naples at Monaldi Hospital, Napoli, Italy

⁷Division of Infectious Diseases, Department of Internal Medicine, American University of Beirut Medical Center, Beirut, Lebanon

Abstract

Background—Phenol-soluble modulins (PSM) are amphipathic proteins produced by *Staphylococcus aureus* that promote virulence, inflammatory response, and biofilm formation. We previously showed that MRSA isolates from soft tissue infection (SSTI) produced significantly

*CORRESPONDING AUTHOR: Vance G. Fowler Jr., MD, MHS, Department of Medicine, Division of Infectious Diseases, Duke University Medical Center, Box 102359 Medical Center, Durham, NC 27710, Tel: (919) 613-5678, Fax: (919) 684-8902, fowle003@mc.duke.edu.

CONFLICTS OF INTEREST

Mr. Qi reports grant from the Eugene A. Stead Research Scholarship from Duke University School of Medicine. Dr. Otto reports grants from NIAID, during the conduct of the study; in addition, Dr. Otto has a patent 8,211,445 issued. Dr. Fowler reports grants from National Institutes of Health (K24-AI093969 and R01-AI068804), during the conduct of the study; personal fees from Scientific Advisory Board for Merck V710, personal fees from Consultancy from Pfizer, Novartis, Galderma, Novadigm, Durata, Debiopharm, Genentech, Achaogen, Affinium, Medicines Co., Cerexa, Tetrphase, Trius, MedImmune, Bayer, Theravance, Cubist, Basilea, Affinergy; personal fees from Duke University Medical Center, grants from Institutional Grants/grants pending NIH, MedImmune, Forest/Cerexa, Pfizer, Merck, Advanced Liquid Logics, Theravance, Novartis, and Cubist, personal fees from Royalties (UpTo- Date), personal fees from Payment for development of educational presentations (Green Cross, Cubist, Cerexa, Durata, Theravance), outside the submitted work. In addition, Dr. Fowler has a patent pending for rapid diagnostics.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

higher levels of PSM than MRSA isolates from hospital-acquired pneumonia (HAP) or infective endocarditis (IE). In this investigation, we sought to validate this finding in methicillin-susceptible *S. aureus* (MSSA) isolates.

Methods—MSSA isolates (n=162) from patients with SSTI, HAP, and IE were matched 1:1:1 based on geographic origin of the infection to form 54 triplets (North America n=27, Europe n=25, Australia n=2). All isolates underwent spa-typing and were classified using eGenomics. In vitro PSM production was quantified by high performance liquid chromatography/mass spectrometry. Fischer's Exact Test and the Kruskal-Wallis test were used for statistical analysis.

Results—Spa1 was more common in SSTI (14.81% SSTI, 3.70% HAP, 1.85% IE) ($p<0.03$). Spa2 was more common in HAP (0% SSTI, 12.96% HAP, 3.70% IE) ($p<0.01$). Levels of PSM α 1-4 all differed significantly among the three clinical groups, with SSTI isolates producing the highest levels and IE producing the lowest levels of PSM α 1-4. Spa1 isolates produced significantly more delta-toxin ($p<0.03$) than non-Spa1 isolates. No associations between PSM levels and clinical outcome of SSTI, HAP, or IE were identified.

Conclusion—Production of PSM α 1-4 is highest in SSTI MSSA isolates, supporting the hypothesis that these peptides are important for SSTI pathogenesis. These findings are similar to those described in MRSA, and demonstrate that associations between PSM levels and type of infection are independent of the methicillin resistance status of the isolate.

Keywords

phenol-soluble modulins; Methicillin-susceptible *Staphylococcus aureus*; skin and soft tissue infection; pneumonia; endocarditis

INTRODUCTION

Staphylococcus aureus produces many virulence factors that affect its clinical manifestations (1, 2). One recently discovered family of virulence factors are phenol-soluble modulins (PSM). Members of this group of amphipathic, alpha-helical peptides may influence inflammation induction, biofilm production, erythrocyte lysis, neutrophil lysis after *S. aureus* has been phagocytosed, and even antimicrobial activity (3). *S. aureus* is currently known to produce eight PSMs. α -type PSMs are 20–25 amino-acids long and are subdivided into PSM α 1-4, delta-toxin, and PSM-mec. β -type PSMs are 44 amino-acids long and include PSM β 1-2 (3). PSM α 1-4, PSM β 1-2, and delta-toxin are encoded on the core genome while PSM-mec is encoded on the mobile genetic element that confers methicillin resistance to *S. aureus* (4). All PSM α family proteins are encoded on the *psmA* operon, while PSM β 1-2 are both encoded on the *psm β* operons, facilitating co-transcription of PSM α family proteins and of PSM β 1-2 (5).

We previously showed that MRSA isolates from patients with SSTI produced significantly more PSMs *in vitro* than geographically-matched MRSA isolates from patients with either HAP or IE (6). However, our results could have been confounded by the presence of traits that were associated with the methicillin-resistance phenotype. For example, PSM-mec is present on the SCCmec element of MRSA isolates and decreases the production of PSM α peptides by inhibiting the positive regulator of PSM α -type production (7). MRSA and

MSSA isolates have also been shown to produce different amounts of enterotoxin A, alpha toxin, and Toxic Shock Syndrome Toxin (8, 9). Thus, potential associations between increased PSM production and specific infection types needs to be confirmed in MSSA isolates.

In the present investigation, we sought to confirm any such association. Our *a priori* hypothesis was that levels of *in vitro* PSM are higher in MSSA isolates from SSTI. To test this hypothesis, we utilized a unique, globally representative collection of *S. aureus* from a variety of clinical syndromes.

MATERIALS AND METHODS

Definitions

SSTI was defined as the presence of at least 1 of the following conditions: major abscess requiring surgical drainage, cellulitis, infected wound or ulcer, or infected burn plus purulent drainage/collection or 3 of the following conditions: heat, localized warmth, fluctuance, erythema, swelling, induration, pain/tenderness to palpation, temperature $\geq 38^{\circ}\text{C}$, WBC $10,000\text{ cells/mm}^3$, or $\geq 15\%$ bands (10). HAP was defined as: 1) presence of new or progressive infiltrates or consolidation, with or without pleural effusion on chest radiograph or computed tomography; 2) adequate respiratory specimen for culture and Gram stain and identification of an organism consistent with a respiratory pathogen isolated from respiratory tract or blood, or ≥ 2 of the following: cough, purulent sputum, auscultation findings, dyspnea, tachypnea, or hypoxemia acquired after 48 h in an inpatient acute or chronic care facility or that developed within 7 days after being discharged. In addition, patients had ≥ 2 of the following: temperature $\geq 38^{\circ}\text{C}$ or rectal/core temperature $\geq 35^{\circ}\text{C}$, respiratory rate ≥ 30 breathes per minute, pulse ≥ 120 beats per minute, altered mental status, mechanical ventilation, WBC $>10,000\text{ cells/mm}^3$, $<4500\text{ cells/mm}^3$, or $>15\%$ band neutrophils (11). IE was defined using the modified Duke criteria (12).

Isolates

SSTI isolates were acquired from two double-blinded, randomized, parallel group, identical method Phase-III trials (10, 13). HAP isolates were acquired from two double-blinded, randomized, parallel group, identical method Phase-III trials (11). IE isolates were acquired from the Microbiologic Repository of the International Collaboration on Endocarditis – Prospective Cohort Study (ICE-PCS). A total of 58 sites in 25 countries enrolled 2781 patients ≥ 18 years old with definite IE (14, 15). All clinical studies were approved by their respective Institutional Review Boards.

Matching of Bacterial Isolates According to Region of Origin

Isolates from SSTI, HAP, and IE were matched on a 1:1:1 basis to form a “triplet” of three MSSA isolates (one each from SSTI, HAP, and IE) that originated in the same geographic region. The total number of triplets per continent was determined by the clinical category with the fewest isolates. For studies with more isolates per continent than the number of isolates present in the numerically smallest category, we used every Xth isolate so that all

total number of isolates selected were equivalent, with $X = (\text{number of isolates in larger category}) / (\text{number of isolates in smallest category})$.

Growth and Supernatant Collection

Bacterial isolates originally frozen in tryptic soy broth (TSB)/glycerol stock were grown overnight at 37°C on trypticase soy agar (TSA) petri dishes. Approximately 1 µL of bacterial growth was inoculated into 10ml liquid TSB stock in 50mL conical tubes, then incubated overnight at 37°C at 200 RPM. The OD₆₀₀ of each sample was measured and the amount of TSB-bacteria needed to reach start OD₆₀₀ equal to 0.1 was added to 10mL liquid TSB in 50mL conical tubes. The new stocks were grown for 8 hours at 37°C and 200 RPM to reach early stationary phase with limited growth variation among different isolates as previously reported (26), and then placed on ice to arrest growth. Samples were centrifuged at 4°C and 4000 RPM for 10 minutes. 4 × 2mL supernatant (8mL total) were collected from each sample and stored at -80°C.

High-Performance Liquid Chromatography (HPLC) / Mass Spectrometry

Reversed phase HPLC / electrospray ionization mass spectrometry (RP-HPLC/ESI-MS) was used to measure PSM production in supernatant as previously described (5). Briefly, 50 µL of culture supernatant was injected into LC/MS (Agilent Technologies, Santa Clara, CA) equipped with C8 column (2.1 × 5 mm, 2.7 µm; PerkinElmer, Waltham, MA). PSMs were separated in flow rate of 0.5mL/min with the following gradient program: eluent A, 0.1% trifluoroacetic acid (TFA) in 100% water; eluent B, 0.1% TFA in 100% acetonitrile; 10% eluent B for 0.5 minutes; a linear gradient from 25% to 100% for 3 minutes; 100% for 2.5 minutes; 0% for 2.5 minutes. Parameters for operating mass spectrometer were as recommended by the manufacturer. Two major ions of each PSM with different charge states were used to generate extracted ion chromatogram for quantification by integration. We did not test for PSM-mec production because it is genetically located on the methicillin-resistance cassette, which MSSA lacks. Each sample was assayed three times and the results were averaged for each isolate and PSM peptide.

Spa Genotyping

Spa genotyping was performed as previously described (6, 17). The eGenomics online platform (<http://tools.egenomics.com>) was used to define *spa* type.

Statistical Methods

Spa type distribution among SSTI, HAP, and IE was evaluated with contingency tables and Fisher's Exact Test. The level of *in vitro* production of each PSM type for a particular MSSA isolate was defined as the average of three separate assays of the same sample. Differences in production across the three clinical categories and by *spa* types were compared using means and the Kruskal-Wallis test. P<0.05 was threshold for statistical significance in all tests.

Institutional Review

This study was approved by the Duke University Medical Center Institutional Review Board.

RESULTS

A total of 162 MSSA isolates from patients with SSTI, HAP, and IE were divided 1:1:1 into 54 geographically-matched triplets. Of the 54 triplets, 27 (50%) came from North America, 25 (46.3%) came from Europe, and 2 (3.7%) came from Australia.

Spa Genotyping According to Infectious Source

Among the 162 isolates, 65 known *spa* types were identified. The most common *spa* types were *spa1* (n=11, 6.8%), *spa2* (n=9, 5.6%), *spa33* (n=8, 4.9%), and *spa73* (n=8, 4.9%). Collectively, these 4 *spa* types accounted for 22.2% (n=36) of all isolates. The 61 other known *spa* types each was found in 6 or fewer isolates, and constituted 60.5% of isolates. New *spa* types (n=28) comprised the remaining 17.3% of isolates. *Spa1* was significantly more common in SSTI (SSTI 8/54 [14.81%], HAP 2/54 [3.70%], IE 1/54 [1.85%]) (p=0.0301), while *Spa2* was significantly more common in HAP (SSTI 0/54 [0%], HAP 7/54 [12.96%], IE 2/54 [3.70%]) (p=0.0111) (Table 1).

Association between Infectious Source and *in vitro* PSM production

Among the SSTI, HAP, and IE isolates, significant differences existed in levels of *in vitro* production of PSM α 1 (p=0.0262), PSM α 2 (p=0.0164), PSM α 3 (p=0.0021), and PSM α 4 (p=0.0016). SSTI isolates produced the highest levels of all four types of PSM α while IE isolates produced the lowest. There were no significant differences in the production of PSM β 1, PSM β 2, or delta-toxin among the three clinical groups (Figure 1).

Because *Spa1* was the most common single genotype in this study, we considered the possibility that its predominance in the SSTI cohort might confound results. PSM production patterns between *Spa1* and non-*Spa1* isolates were similar. *Spa1* did not produce significantly different quantities of PSM α 1-4 and PSM β 1-2 compared with non-*spa1* isolates. After excluding *spa1* isolates from analysis, statistically significant in-group differences remained in PSM α 1 (p=0.0293), PSM α 2 (p=0.0224), PSM α 3 (p=0.0050), and PSM α 4 (p=0.0042) (data not shown).

Association between clinical outcome and *in vitro* PSM production

Next, we explored the possible association between PSM level in MSSA isolates and the clinical outcome of the patients they infected. No statistically significant associations were identified between the clinical outcomes of patients with SSTI, HAP, or IE and the *in vitro* PSM level of the MSSA isolates causing the infection (Table 2).

DISCUSSION

The role of PSMs in the pathogenesis of *S. aureus* infections is increasingly recognized. Previously, we demonstrated an association between levels of *in vitro* PSM production of

MRSA isolates and the clinical source from those isolates were obtained. The current investigation confirmed this association, and showed that levels of *in vitro* PSM α 1-4 production were higher in MSSA from patients with SSTI than similar isolates from geographically-matched patients with HAP or IE.

The results of this study add to a growing body of evidence linking PSM production with specific manifestations of staphylococcal disease. Experimentally, PSM α -deficient MRSA isolates form smaller soft tissue infections in rabbit and murine models of SSTI than MRSA with normal PSM α production (5, 18). PSM α peptides have also been shown to promote bacterial spread across surfaces (19), destroy neutrophils (5) and erythrocytes (20), and inhibit dendritic cell activity (21) that could potentiate detection by immune response on the skin. Clinically, MRSA isolates from patients with SSTI exhibited higher levels of PSM than MRSA from other distinct clinical syndromes (6). The current study extends this observation into MSSA isolates as well, and provides further evidence for a key role for PSM in the pathogenesis of SSTI.

MSSA IE isolates produced less PSM α 1-4 than SSTI isolates did, while HAP isolates made intermediate amounts. These patterns are consistent with those in MRSA isolates from the same infection types (6). Nonetheless, the role of PSM in bloodstream infections remains unclear. Mice infected by PSM α -deletion or delta-toxin-deletion strain *S. aureus* showed decreased TNF- α inflammation and improved septicemia survival rates (5). PSM production is positively controlled by the *agr* regulatory system (22). *S. aureus agr* knockout mutants have diminished ability to induce experimental IE at low concentrations and adhere to valve vegetation (23). In addition, a unique PSM α 3 variant that elicits an attenuated cytolytic and inflammatory host response is associated with increased hematogenous spreading among the Clonal Complex 30 strain of *S. aureus* (24), which is associated with IE (25). Finally, both the *agr* regulatory system and PSMs themselves are neutralized by LDL and VLDL lipoproteins at physiologic conditions (26, 27), potentially decreasing the impact of PSMs secreted by *S. aureus* as it enters the host bloodstream. Collectively, these data suggest that attenuated PSM production by specific strains of *S. aureus* could reduce host recognition and response to *S. aureus* bacteremia (28). The data also provides a rationale for the hypothesis that MSSA isolates from HAP and IE produce less PSM α 1-4 to evade detection from and prevent destruction by the host immune system. Overall, more research is necessary to elucidate how PSMs affect pathogenicity of *S. aureus* within different host body sites.

The role of PSM β 1-2 in *S. aureus* infection remains unclear. PSM β peptides are far less cytolytic than the PSM α family (5), do not affect *S. aureus* ability to escape lysosomes (29), and are not bactericidal against other pathogens (3). In the current study and in MRSA (6), levels of PSM β 1-2 were lower than those of either PSM α or delta-toxin, and were not significantly associated with clinical types of infection. Taken together, these results suggest that PSM β production may be less important in the pathogenesis of SSTI than other forms of PSM peptides, such as PSM α .

The absence of PSM-mec in MSSA isolates may affect the total production levels of PSMs. PSM-mec is located on a regulatory RNA that inhibits *agrA* positive regulation of PSM α ,

PSM β , and delta-toxin (7). MRSA strains with their PSM-mec deleted produced more *agrA* and PSM α family peptides, and had increased virulence in mice models (7). Berlon *et al* showed that decreased PSM-mec production is associated with significantly increased PSM α , β , delta-toxin in MRSA SSTI isolates compared with HAP and IE (6). Moreover, PSM-mec vector-transformed *S. epidermidis* and *S. haemolyticus* had a 30% reduction in their combined PSM α and delta-toxin output compared with their null-vector transformed counterparts, suggesting that the PSM-mec downregulation of PSM production is conserved in the *Staphylococcus* family (30). Given that MSSA lacks the SCCmec cassette and by extension PSM-mec, total MSSA PSM production may be higher than total MRSA PSM production, even though the relative proportions of PSMs produced may have similar profiles. Thus, while the PSM production patterns appear independent of methicillin-resistance, the total amount of PSM produced may be influenced by methicillin-resistance acquisition.

Our study is limited by its *in vitro* nature, as *in vivo* factors, such as the presence of LDL and VLDL in blood, could alter PSM production. Although we geographically matched the SSTI, HAP, and IE isolates by continent of origin, this strategy may be insufficiently precise, as specific regions and even individual hospitals or hospital wards may often be affected by a handful of dominant bacterial clones. Furthermore, we were unable to investigate MSSA from regions other than North America, Europe, or Australia. Finally, our results confirm an association between *in vitro* levels of PSM and SSTI versus HAP or IE but do not establish the molecular basis for this association.

In conclusion, the results of this investigation confirm that *S. aureus* isolates from patients with SSTI exhibit significantly higher levels of *in vitro* PSM production than similar isolates from patients with either HAP or IE. This finding supports the hypothesis that PSMs play a significant role in SSTI pathogenesis. Future studies of this type are needed to begin to bridge the gap between the bench and the bedside by providing a better understanding of the determinants of virulence in *S. aureus*.

Acknowledgments

Funding: This work was supported by the Duke University School of Medicine Eugene A. Stead Student Research Scholarship [to RQ], grants provided by the National Institute of Health [R01-AI068804 and K24-AI093969 to VGF], and the Intramural Research Program of the National Institute of Allergy and Infectious Diseases [to MO].

References

1. Gordon RJ, Lowy FD. Pathogenesis of methicillin-resistant *Staphylococcus aureus* infection. Clin Infect Dis. 2008; 46:S350–359. [PubMed: 18462090]
2. Liu G. Molecular pathogenesis of *Staphylococcus aureus* infection. Pediatr Res. 2009; 65:71–77.
3. Otto M. Phenol-soluble modulins. Int J Med Microbiol. 2014; 304:164–169. [PubMed: 24447915]
4. Queck SY, Khan BA, Wang R, Bach TH, Kretschmer D, Chen L, Kreiswirth BN, Peschel A, DeLeo FR, Otto M. Mobile genetic element-encoded cytolytic connects virulence to methicillin resistance in MRSA. PLoS Pathog. 2009; 5:e1000533. [PubMed: 19649313]
5. Wang R, Braughton KR, Kretschmer D, Bach TH, Queck SY, Li M, Kennedy AD, Dorward DW, Klebanoff SJ, Peschel A, DeLeo FR, Otto M. Identification of novel cytolytic peptides as key virulence determinants for community-associated MRSA. Nat Med. 2007; 13:1510–1514. [PubMed: 17994102]

6. Berlon NR, Qi R, Sharma-Kuinkel BK, Joo HS, Park LP, George D, Thaden JT, Messina JA, Maskarinec SA, Mueller-Premru M, Athan E, Tattevin P, Pericas JM, Woods CW, Otto M, Fowler VG Jr. Clinical MRSA isolates from skin and soft tissue infections show increased in vitro production of phenol soluble modulins. *J Infection*. 2015 Epub ahead of print.
7. Kaito C, Saito Y, Ikuo M, Omae Y, Mao H, Nagano G, Fujiyuki T, Numata S, Han X, Obata K, Hasegawa S, Yamaguchi H, Inokuchi K, Ito T, Hiramatsu K, Sekimizu K. Mobile genetic element SCCmec-encoded psm-mec RNA suppresses translation of *agrA* and attenuates MRSA virulence. *PLoS Pathog*. 2013; 9:e1003269. [PubMed: 23592990]
8. Jordens JZ, Duckworth GJ, Williams RJ. Production of “virulence factors” by “epidemic” methicillin-resistant *Staphylococcus aureus* in vitro. *J Med Microbiol*. 1989; 30:245–252. [PubMed: 2600958]
9. Schlievert PM, Strandberg KL, Lin YC, Peterson ML, Leung DY. Secreted virulence factor comparison between methicillin resistant and methicillin-sensitive *Staphylococcus aureus*, and its relevance to atopic dermatitis. *J Allergy Clin Immunol*. 2010; 125:39–49. [PubMed: 20109735]
10. Stryjewski ME, Graham DR, Wilson SE, O’Riordan W, Young D, Lentnek A, Ross DP, Fowler VG, Hopkins A, Friedland HD, Barriere SL, Kitt MM, Corey GR. Assessment of Telavancin in Complicated Skin and Skin-Structure Infections Study. Telavancin versus vancomycin for the treatment of complicated skin and skin-structure infections caused by gram-positive organisms. *Clin Infect Dis*. 2008; 46:1683–1693. [PubMed: 18444791]
11. Rubinstein E, Lalani T, Corey GR, Kanafani ZA, Nannini EC, Rocha MG, Rahav G, Niederman MS, Kollef MH, Shorr AF, Lee PC, Lentnek AL, Luna CM, Fagon JY, Torres A, Kitt MM, Genter FC, Barriere SL, Friedland HD, Stryjewski ME. ATTAIn Study Group. Telavancin versus vancomycin for hospital-acquired pneumonia due to gram-positive pathogens. *Clin Infect Dis*. 2011; 52:31–40. [PubMed: 21148517]
12. Li JS, Sexton DJ, Mick N, Nettles R, Fowler VG Jr, Ryan T, Bashore T, Corey GR. Proposed modifications to the Duke criteria for the diagnosis of infective endocarditis. *Clin Infect Dis*. 2000; 30:633–638. [PubMed: 10770721]
13. Bae IG, Tonthat GT, Stryjewski ME, Rude TH, Reilly LF, Barriere SL, Genter FC, Corey GR, Fowler VG Jr. Presence of genes encoding the Pantone-Valentine Leukocidin exotoxin is not the primary determinant of outcome in patients with complicated skin and skin structure infections due to methicillin-resistant *Staphylococcus aureus*: results of a multinational trial. *J Clin Microbiol*. 2009; 47:3952–3957. [PubMed: 19846653]
14. Fowler VG Jr, Miro JM, Hoen B, Cabell CH, Abrutyn E, Rubinstein E, Corey GR, Spelman D, Bradley SF, Barsic B, Pappas PA, Anstrom KJ, Wray D, Fortes CQ, Anguera I, Athan E, Jones P, van der Meer JT, Elliott TS, Levine DP, Bayer AS. ICE Investigators. *Staphylococcus aureus* endocarditis: a consequence of medical progress. *JAMA*. 2005; 293:3012–3021. [PubMed: 15972563]
15. Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG, Bayer AS, Karchmer DW, Olaison L, Pappas PA, Moreillon P, Chambers ST, Chu VH, Falcó V, Holland DJ, Jones P, Klein JL, Raymond NJ, Read KM, Tripodi MF, Utili R, Wang A, Woods CW, Cabell CH. The ICE Investigators. Clinical presentation, etiology and outcome of infective endocarditis in the 21st century: The International Collaboration on Endocarditis-Pro prospective Cohort Study. *Arch Intern Med*. 2009; 169:463–473. [PubMed: 19273776]
16. Harris LG, Foster SJ, Richards RG. An introduction to *Staphylococcus aureus*, and techniques for identifying and quantifying *S. aureus* adhesins in relation to adhesion to biomaterials: review. *Eur Cells Mater*. 2002; 4:39–60.
17. Mathema B, Mediavilla J, Kreiswirth BN. Sequence analysis of the variable number tandem repeat in *Staphylococcus aureus* protein A gene: spa typing. *Methods Mol Biol*. 2008; 431:285–305. [PubMed: 18287764]
18. Kobayashi SD, Malachowa N, Whitney AR, Braughton KR, Gardner DJ, Long D, Bubeck Wardenburg J, Schneewind O, Otto M, Deleo FR. Comparative analysis of USA300 virulence determinants in a rabbit model of skin and soft tissue infection. *J Infect Dis*. 2011; 204:937–941. [PubMed: 21849291]

19. Tsompanidou E, Denham EL, Becher D, Jong A, Buist G, Oosten M, Manson WL, Back JW, Dijk JM, Dreisbach A. Distinct roles of phenol-soluble modulins in spreading of *Staphylococcus aureus* on wet surfaces. *Appl Environ Microb*. 2013; 79:886–895.
20. Cheung GY, Duong AC, Otto M. Direct and synergistic hemolysis caused by *Staphylococcus* phenol-soluble modulins: implications for diagnosis and pathogenesis. *Microbes Infect*. 2012; 14:380–386. [PubMed: 22178792]
21. Schreiner JKD, Klenk J, Otto M, Buhring HJ, Stevanovic S, Wang JM, Beer-Hammer S, Peschel A, Autenrieth SE. *Staphylococcus aureus* phenol-soluble modulin peptides modulate dendritic cell functions and increase in vitro priming of regulatory T cells. *J Immunol*. 2013; 190:3417–3426. [PubMed: 23460735]
22. Queck SY, Jameson-Lee M, Villaruz AE, Bach TH, Khan BA, Sturdevant DE, Ricklefs SM, Li M, Otto M. RNAIII-independent target gene control by the *agr* quorum-sensing system: insight into the evolution of virulence regulation in *Staphylococcus aureus*. *Mol Cell*. 2008; 32:150–158. [PubMed: 18851841]
23. Cheung AL, Eberhardt KJ, Chung E, Yeaman MR, Sullam PM, Ramos M, Bayer AS. Diminished virulence of a *sar-/agr-* mutant of *Staphylococcus aureus* in the rabbit model of endocarditis. *J Clin Invest*. 1994; 94:1815–1822. [PubMed: 7962526]
24. Cheung GY, Kretschmer D, Duong AC, Yeh AJ, Ho TV, Chen Y, Joo HS, Kreiswirth BN, Peschel A, Otto M. Production of an attenuated phenol-soluble modulin variant unique to the MRSA clonal complex 30 increases severity of bloodstream infection. *PLoS Pathog*. 2014; 10:e1004298. [PubMed: 25144687]
25. Nienaber JJ, Sharma Kuinkel BK, Clarke-Pearson M, Lamlertthong S, Park L, Rude TH, Barriere S, Woods CW, Chu VH, Marín M, Bukovski S, Garcia P, Corey GR, Korman T, Doco-Lecompte T, Murdoch DR, Reller LB, Fowler VG Jr. International Collaboration on Endocarditis-Microbiology Investigators. Methicillin-susceptible *Staphylococcus aureus* endocarditis isolates are associated with clonal complex 30 genotype and a distinct repertoire of enterotoxins and adhesins. *J Infect Dis*. 2011; 204:704–713. [PubMed: 21844296]
26. Peterson MM, Mack JL, Hall PR, Alsup AA, Alexander SM, Sully EK, Sawires YS, Cheung AL, Otto M, Gresham HD. Apolipoprotein B is an innate barrier against invasive *Staphylococcus aureus* infection. *Cell Host Microbe*. 2008; 4:555–566. [PubMed: 19064256]
27. Peschel A, Otto M. Phenol-soluble modulins and staphylococcal infection. *Nat Rev Microbiol*. 2013; 11:667–673. [PubMed: 24018382]
28. Spaan AN, Surewaard BG, Nijland R, van Strijp JA. Neutrophils versus *Staphylococcus aureus*: a biological tug of war. *Annu Rev Microbiol*. 2013; 67:629–650. [PubMed: 23834243]
29. Grosz M, Kolter J, Paprotka K, Winkler AC, Schäfer D, Chatterjee SS, Geiger T, Wolz C, Ohlsen K, Otto M, Rudel T, Sinha B, Fraunholz M. Cytoplasmic replication of *Staphylococcus aureus* upon phagosomal escape triggered by phenol-soluble modulin α . *Cell Microbiol*. 2014; 16:451–465. [PubMed: 24164701]
30. Ikuo M, Nagano G, Saito Y, Mao H, Sekimizu K, Kaito C. Inhibition of exotoxin production by mobile genetic element SCCmec-encoded psm-mec RNA is conserved in staphylococcal species. *PLoS One*. 2014; 9:e100260. [PubMed: 24926994]

Appendix

ICE Micro Investigator Index 2012: **Australia:** Eugene Athan MD, Owen Harris MBBS (*Barwon Health*); Tony M. Korman, MD (*Monash Medical Centre*), Despina Kotsanas, BS (*Southern Health*); Phillip Jones MD, Porl Reinbott, Suzanne Ryan, MHS (*The University of New South Wales*); **Brazil:** Claudio Querido Fortes, MD (*Hospital Universitario Clementino Fraga Filho/UFRJ*); **Chile:** Patricia Garcia, MD, Sandra Braun Jones, MD (*Hosp. Clínico Pont. Universidad Católica de Chile*); **Croatia** Bruno Barsic, MD, PhD, Suzana Bukovski, MD (*Univ. Hospital for Infectious Diseases*); **France:** Christine Selton-Suty, MD, Neijla Aissa, MD, Thanh Doco-Lecompte, MD (*CHU Nancy-Brabois*); Francois Delahaye, MD,

PhD, Francois Vandenesch, MD (*Hopital Louis Pradel*); Pierre Tattevin, MD, PhD (*Pontchaillou University*); Bruno Hoen, MD, PhD, Patrick Plesiat, MD (*University Medical Center of Besançon*); **Greece**: Helen Giamarellou MD PhD, Efthymia Giannitsioti, MD, Ekaterini Tarpazi, MD (*Attikon University General Hospital*); **Italy**: Marie Françoise Tripodi, MD, (*Università degli Studi di Salerno*), Emanuele Durante-Mangoni, MD, PhD, Riccardo Utili MD, PhD, Roberta Casillo, MD, PhD, Susanna Cuccurullo, MSc (*II Università di Napoli*), Pierre Yves Donnio, PhD, Annibale Raglio, MD, Fredy Suter, MD (*Ospedali Riuniti di Bergamo*); **Lebanon**: Tania Baban, MD, Zeina Kanafani, MD, MS, Souha S. Kanj, MD, Jad Sfeir, MD, Mohamad Yasmine, MD (American University of Beirut Medical Center); **New Zealand**: Arthur Morris, MD (*Diagnostic Medlab*), David R. Murdoch, MD MSc, DTM&H (*University of Otago*); **Slovenia**: Manica Mueller Premru, MD, PhD, Tatjana Lejko-Zupanc, MD, PhD (*Medical Center Ljubljana*); **Spain**: Manuel Almela, MD, Yolanda Armero, MD, Manuel Azqueta, MD, Ximena Castañeda, MD, Carlos Cervera, MD, Carlos Falces, MD, Cristina Garcia- de- la- Maria, PhD, Jose M. Gatell, MD, PhD, Jaume Llopis, MD, PhD, Francesc Marco, MD, PhD, Carlos A. Mestres, MD, PhD, José M. Miró, MD, PhD, Asuncion Moreno, MD, PhD, Salvador Ninot, MD, Carlos Paré, MD, PhD, Juan M Pericas, MD, Eduard Quintana, MD, Jose Ramirez, MD, PhD, Marta Sitges, MD (*Hospital Clinic – IDIBAPS. University of Barcelona*), Emilio Bouza, MD, PhD, Marta Rodríguez-Créixems, MD PhD, Victor Ramallo, MD, (*Hospital General Universitario Gregorio Marañón*); **USA**: Suzanne Bradley, MD (*Ann Arbor VA Medical Center*); Dannah Wray, MD, Lisa Steed, PhD, MHS, Robert Cantey, MD, (*Medical University of South Carolina*); Gail Peterson MD, Amy Stancoven, MD (*UT-Southwestern Medical Center*); Christopher Woods, MD, MPH, G. Ralph Corey, MD, L. Barth Reller, MD, Vance G. Fowler Jr., MD, MHS, Vivian H Chu, MD, MHS, (*Duke University Medical Center*)

ICE Coordinating Center: Khaula Baloch, MPH, Vivian H. Chu, MD, MHS, G. Ralph Corey, MD, Christy C. Dixon, Vance G. Fowler, Jr., MD, MHS, Tina Harding, RN, BSN, Marian Jones- Richmond, Paul Pappas, MS, Lawrence P. Park, PhD, Thomas Redick, MPH, Judy Stafford, MS ICE Publications Committee: Kevin Anstrom, PhD, Eugene Athan, MD, Arnold S. Bayer, MD, Christopher H. Cabell, MD, MHS, Vivian H. Chu, MD, MHS, G. Ralph Corey, MD, Vance G. Fowler, Jr., MD, MHS, Bruno Hoen, MD, PhD, A.W. Karchmer, MD, José M. Miró, MD, PhD, David R. Murdoch, MD, MSc, DTM&H, Daniel J. Sexton MD, Andrew Wang, MD

ICE Steering Committee: Arnold S. Bayer, MD, Christopher H Cabell, MD, MHS, Vivian Chu MD, MHS, G. Ralph Corey, MD, David T. Durack, MD, D Phil, Susannah Eykyn, MD, Vance G. Fowler, Jr., MD, MHS, Bruno Hoen, MD, PhD, José M. Miró, MD, PhD, Phillipe Moreillon, MD, PhD, Lars Olaison, MD, PhD, Didier Raoult, MD, PhD, Ethan Rubinstein MD, LLB, Daniel J, Sexton, MD

Key Point

In vitro production of most phenol-soluble modulin proteins is significantly higher among methicillin-susceptible *S. aureus* isolates obtained from skin and soft tissue infection when compared to *S. aureus* isolates obtained from patients with hospital acquired pneumonia or infective endocarditis.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- 162 MSSA isolates from SSTI, HAP, and IE were geographically matched and *spa*-typed
- PSM production was quantitated by liquid chromatography/mass spectrometry.
- SSTI isolates produced the most PSM α 1-4, supporting a PSM role in SSTI pathogenesis
- PSM production patterns are similar in MRSA and MSSA

Figure 1. Mean Levels of *In Vitro* Phenol-Soluble Modulin Production by Clinical Source of Isolates of Methicillin-susceptible *S. aureus*

Mean Levels of *In Vitro* Phenol-Soluble Modulin Production by Clinical Source of MSSA Isolate. PSM production in culture supernatant from SSTI, HAP, and IE isolates were quantified and averaged in technical triplicate (n=54 per group). Error bars show SEM. Kruskal-Wallis analysis was used to calculate differences in production. There were significant in-group differences in the production of PSMα1-4, with SSTI isolates producing the highest levels in all four PSMα subtypes.

Table 1

Spa Type Distribution Among Methicillin Susceptible *S. aureus* Isolates from Skin/Soft Tissue Infection, Hospital Acquired Pneumonia, and Infective Endocarditis.

Spa Type	Skin and Soft Tissue% (n)	Pneumonia % (n)	Endocarditis % (n)	p-value
1	14.8% (8)	3.7% (2)	1.9% (1)	0.030
2	0.0% (0)	13.0% (7)	3.7% (2)	0.011
33	1.9% (1)	9.3% (5)	3.7% (2)	0.278
73	7.4% (4)	3.7% (2)	3.7% (2)	0.732
NEW	18.5% (10)	18.5% (10)	14.8% (8)	0.900
OTHER	57.4% (31)	51.9% (28)	72.2% (39)	0.089

Spa typing was performed on 162 MSSA isolates from SSTI, HAP, and IE. All *spa* types with n = 8 are listed. *Spa*-type NEW are isolates with genomic *spa* sequences that did not match a known *spa* type on eGenomics. *Spa*-type OTHER consist of all isolates with known *spa* types but with total n < 8. Fischer's Exact Test showed that SSTI isolates are significantly more likely to be *spa* type 1 (p=0.030) and significantly less likely to be *spa* type 2 (p=0.011).

Table 2

Associations between Patient Outcome and Average Levels of *In Vitro* Phenol-Soluble Modulin Production from methicillin susceptible *S. aureus* isolates causing Skin and Soft Tissue Infection, Hospital-Acquired Pneumonia, and Infective Endocarditis.

Phenol Soluble Modulin Subtype	Average PSM production in Skin and Soft Tissue Infection				Average PSM production in Hospital-Acquired Pneumonia				Average PSM production in Infective Endocarditis*			
	Cure n=41	Failure n=9	Unknown n=4	p-value	Alive at 28 days n=34	Dead at 28 days n=7	Unknown n=13	p-value	Alive at Discharge n=40	In-hospital Mortality n=11	p-value	
PSMα1	1259.0	999.0	1168.4	0.77	1151.1	709.6	910.4	0.53	800.4	700.1	0.96	
PSMα2	616.0	481.6	523.5	0.58	525.0	338.1	407.9	0.58	402.9	361.5	0.89	
PSMα3	493.8	427.3	403.3	0.78	379.7	209.4	263.8	0.56	271.8	226.6	0.93	
PSMα4	1745.6	1736.0	1267.6	0.65	1453.3	1012.0	1177.3	0.57	1083.1	828.0	0.61	
PSMβ1	222.9	162.6	214.3	0.64	189.1	125.4	196.9	0.58	172.2	173.5	0.71	
PSMβ2	49.0	38.4	44.3	0.90	40.6	29.5	45.7	0.85	34.6	38.3	0.79	
δ-toxin	1752.6	2100.0	1375.7	0.73	1653.0	1367.0	1420.3	0.78	1308.4	1362.4	0.77	

Averaged PSM production classified by clinical outcomes (SSTI = treatment cure or failure; HAP = alive or death at 28 days; IE = alive at discharge or in-hospital mortality). Kruskal-Wallis analysis was used to calculate differences in production. There were no statistically significant differences. Values are in millions of units.

* Information for 3 subjects not available.