Published in final edited form as:

Commun Stat Theory Methods. 2015 ; 44(10): 2107–2135. doi:10.1080/03610926.2013.775304.

An Investigation of Quantile Function Estimators Relative to Quantile Confidence Interval Coverage

Lai Wei¹, Dongliang Wang², and Alan D. Hutson¹

¹Department of Biostatistics, State University of New York at Buffalo, Buffalo, New York, USA

²Department of Public Health and Preventive Medicine, State University of New York Upstate Medical University, Syracuse, New York, USA

Abstract

In this article, we investigate the limitations of traditional quantile function estimators and introduce a new class of quantile function estimators, namely, the semi-parametric tail-extrapolated quantile estimators, which has excellent performance for estimating the extreme tails with finite sample sizes. The smoothed bootstrap and direct density estimation via the characteristic function methods are developed for the estimation of confidence intervals. Through a comprehensive simulation study to compare the confidence interval estimations of various quantile estimators, we discuss the preferred quantile estimator in conjunction with the confidence interval estimation method to use under different circumstances. Data examples are given to illustrate the superiority of the semi-parametric tail-extrapolated quantile estimators. The new class of quantile estimators is obtained by slight modification of traditional quantile estimators, and therefore, should be specifically appealing to researchers in estimating the extreme tails.

Keywords

Characteristic function; Direct density estimation; Inversion theorem; Smoothed bootstrap; Tail extrapolation

1. Introduction

The estimators $\hat{Q}(u)$ of the population quantiles, $0 < u < 1$, are of interest in a broad spectrum of theories, methods, and applications of parametric, robust, and exploratory statistical analyses. Their importance is emphasized in the following quote from where Parzen (1993, p. 7) wishes to “... emphasize to applied statisticians our opinion that the first step in data analysis should be to form the sample quantile function, $Q^{(n)}(u)$, $0 < u < 1$, ...” Moreover, we will show that the quantile estimator in conjunction with the bootstrap method can overcome the shortcoming of the standard bootstrap method in finite sample size.

There have been several well-developed quantile estimators, from the simplest step function quantile estimator to the kernel quantile estimator. Let $X_{1:n}$ $X_{2:n}$ $\cdots$ $X_{n:n}$ denote the

order statistics from an i.i.d. sample of size n from a continuous distribution F with support over the entire real line. The quantile function $Q(u)$ is defined to be the left-continuous inverse of F , given by $Q(u) = \inf\{x : F(x) \geq u\}$ ($0 < u < 1$). Furthermore, if $F(\cdot)$ is strictly monotone and continuous, then $Q(u) = F^{-1}(u)$. The traditional step function estimator of $Q(u)$ can be denoted as

$$\hat{Q}_S(u) = X_{(\lfloor nu \rfloor + 1):n}, \quad (1.1)$$

where $\lfloor \cdot \rfloor$ denotes the floor function.

One of the most commonly used quantile estimator in applications is based on a linear combination of successive order statistics, and can be written as

$$\hat{Q}_L(u) = (1 - \varepsilon)X_{\lfloor n'u \rfloor : n} + \varepsilon X_{(\lfloor n'u \rfloor + 1):n}, \quad \frac{1}{n'} \leq u \leq \frac{n}{n'} \quad (1.2)$$

where $n' = n + 1$ and $\varepsilon = n'u - \lfloor n'u \rfloor$.

More recently, in order to overcome the lack of smoothness and its non differentiability of the estimators mentioned above, Mudholkar et al. (2001) constructed a smooth Hermitian quadratic estimator for $1.5/n' - u < (n - 0.5)/n'$, which is given by

$$\hat{Q}_H(u) = w_1 X_{\lfloor n'u - 1/2 \rfloor : n} + w_2 X_{(\lfloor n'u - 1/2 \rfloor + 1):n} + w_3 X_{(\lfloor n'u - 1/2 \rfloor + 2):n}, \quad (1.3)$$

where

$$w_1 = \frac{1}{2} - \tau + \frac{\tau^2}{2},$$

$$w_2 = \frac{1}{2} + \tau - \tau^2, \quad (1.4)$$

$$w_3 = 1 - w_1 - w_2$$

and

$$\tau = (n'u - 1/2 - \lfloor n'u - 1/2 \rfloor). \quad (1.5)$$

In order to reduce the variability, an L -estimator for the quantile function was proposed by forming a weighted average of all order statistics with different weight functions. Harrell and Davis (1982) and Kaigh and Lachenbruch (1982) developed the Harrell–Davis estimator, which is given by

$$\hat{Q}_{HD}(u) = \sum_{i=1}^n w_{i(u)} X_{i:n}, \quad (1.6)$$

with

$$w_{i(u)} = B_{n'u, n'(1-u)}\left(\frac{i}{n}\right) - B_{n'u, n'(1-u)}\left(\frac{i-1}{n}\right), \quad (1.7)$$

where $B_{p,q}(x)$ is the cumulative distribution function for a beta function with shape parameters p, q . In other words, it is the exact bootstrap estimator of $E(X_{n'u:n})$, where $n' = n + 1$ and $X_{n'u:n}$ denotes the fractional order statistics; see Harrell and Davis (1982) for the original derivation of this result.

Another popular class of the L -estimators is the kernel quantile estimator, which can be defined as

$$\hat{Q}_K(u) = \sum_{i=1}^n \left[\int_{(i-1)/n}^{i/n} K_h(t-u) dt \right] X_{i:n}, \quad (1.8)$$

where $K_h(\cdot) = K(\cdot/h)/h$ and K is a density function symmetric about 0 and that $h \rightarrow 0$ as $n \rightarrow \infty$; see Sheather and Marron (1990) and Cheng and Parzen (1997) for a detailed study of the kernel quantile function estimators. Here $h = h_n > 0$ is the smoothing parameter or bandwidth as it controls the amount of smoothness in the quantile estimator. Yang (1985) provided the asymptotic normality property and the mean squared consistency of this estimation. Falk (1984) showed that the asymptotic performance of $\hat{Q}_K(u)$ is better than that of the empirical sample quantile $\hat{Q}_S(u)$ in terms of relative deficiency for appropriately chosen kernels and sufficiently smooth distribution functions. Note that the Harrell–Davis estimator in (1.6) is a special case of the kernel quantile estimator with $K(\cdot)$ being the density function of a beta distribution. In our study, we will choose the Gaussian kernel and further develop the standardized version of $\hat{Q}_K(u)$ in (1.8) as

$$\hat{Q}_K(u) = \frac{\sum_{i=1}^n \left[\Phi\left(\frac{i/n-u}{h}\right) - \Phi\left(\frac{(i-1)/n-u}{h}\right) \right] X_{i:n}}{\Phi\left(\frac{1-u}{h}\right) - \Phi\left(\frac{0-u}{h}\right)}. \quad (1.9)$$

The construction of confidence intervals for the quantiles are of great importance. The traditional method is the well-known standard bootstrap resampling approach for a statistical functional $T(F(X))$ based upon the B bootstrap replicates, where $F(\cdot)$ denotes the population c.d.f. having support over the real line, in conjunction with the bootstrap percentile confidence interval method. Researchers have focused their efforts on using different estimators of F from which to resample in order to compensate for the discreteness of the classic empirical estimator F_n in small samples. For example, Silverman and Young (1987) suggested modifying the resampling procedure by employing a smoothed version of the empirical distribution function $\hat{F}_n$. Gamero et al. (1998) suggested eliminating “outlier” bootstrap samples in the resampling procedure in order to minimize their impact on the

estimation procedure. Hutson (1999) employed the linear interpolation of successive order statistics to improve the coverage probabilities of analytically calculated bootstrap percentile confidence intervals for quantiles. We will show in Sec. 3 that the standard bootstrap replicates can be generated through $X_i = Q(\hat{U}_i)$, where $U_i \sim U(0, 1)$ and $i = 1, 2, \dots, n$.

Lee (1994) has proposed combining parametric and non parametric bootstrapping procedures by defining an estimator of the form $T_n = \varepsilon \alpha(F_n) + (1 - \varepsilon) \alpha(F_0)$, where T_n is a convex combination of a non parametric and parametric estimator. In practice, Lee (1994) suggested that the optimal ε can be estimated from the data. Even though Lee's approach has been shown to be asymptotically optimal based on an MSE criteria, it is oftentimes complicated to carry out in terms of a resampling method. It is however an idea worth further exploration. More recently, Ho and Lee (2005) have developed theory with respect to determining the optimal bandwidth for kernel smoothed bootstrap procedures related to confidence intervals for quantiles, which are based on kernel estimators of the empirical distribution function.

Hutson (2002) developed a new semi-parametric tail-extrapolated quantile function estimation approach for generating bootstrap confidence intervals that had improved finite sample coverage probabilities as compared to simple percentile based intervals. This estimation is described in detail in Sec. 2. The notion of using the quantile function leads to an obvious question along the lines of Silverman and Young (1987): Can we improve the bootstrap resampling method in small samples through the use of smoothed kernel quantile estimators such as those studied in Sheather and Marron (1990) and having the form

$\sum_{i=1}^n X_{i:n} \left[\int_{(i-1)/n}^{i/n} K_h(t-u) dt \right]$, We argue that there is no real advantage to simply smoothing the quantile function estimator without considering the tail behavior of the estimator. In fact, simple kernel smoothing typically makes the coverage errors for percentile intervals worse, since the range of the bootstrap replicated statistics are typically reduced after smoothing, i.e., the bias plays a critical role in the bootstrap process.

Apart from utilizing the bootstrap method to obtain the confidence intervals for quantile estimators, Wang et al. (2012) proposed a kernel density estimation method for directly estimating the probability and cumulative distribution function of the L -estimators from a single population based on the characteristic function in Knight (1985) in conjunction with the classic inversion theory. This method provides us an exact estimation of the distribution function of quantile estimators and thus eliminates the error arising from the bootstrap resampling.

In this note, we follow the idea of Hutson (2002) with that of Silverman and Young (1987) and correct defects with respect to kernel smoothing by developing a tail-extrapolated kernel quantile function estimator for the purpose of improving quantile function estimation. We also show how to extrapolate the tail for other traditional quantile estimators defined in Sec. 1 to form a new class of semi-parametric quantile estimators. Confidence intervals are obtained through the smoothed bootstrap and the direct density estimation methods. We illustrate the improved coverage probabilities with respect to the percentile confidence interval calculations via an extensive simulation study. Data examples of the flood data and

Buffalo snowfall data are given to show the superiority of tail-extrapolated quantile estimators in estimating the extreme tails and improved coverage probabilities for bootstrap procedures.

2. A New Class of Semi-Parametric Tail-Extrapolated Quantile Estimators

Our new class of semi-parametric tail-extrapolated quantile estimators is to modify the traditional quantile estimators to obtain better estimated quantiles that are less than $\hat{Q}(1/n')$ and larger than $\hat{Q}(n/n')$, given a minimal set of assumptions about the extreme quantiles. The modified estimator should converge to $Q(u)$ as $n \rightarrow \infty$ and behave well in small finite samples, the latter being important with respect to bootstrap resampling.

To build our estimator we propose starting with an estimate of the derivative of $Q'(u) = 1/f(Q(u))$ at $1/n'$ based upon simple spacings. By our definition the derivative is given by the spacing $Q'(1/(n+1)) = n'(X_{2:n} - X_{1:n})$. Similarly, the estimate of $Q'(u)$ at n/n' is given by the spacing $n'(X_{n:n} - X_{n-1:n})$. Note also that given the assumption that X is absolutely continuous and has support over the real line, $\lim_{u \rightarrow 0^+} f(Q(u)) = 0$ and $\lim_{u \rightarrow 1^-} f(Q(u)) = 0$ for the density-quantile function. If we assume no other additional information is available pertaining to the tails of the density of X then a simple linear interpolation between the values of the density-quantile function 0 and $1/[n'(X_{2:n} - X_{1:n})]$ at the knots 0 and $1/n'$ and the values of the density quantile function $1/[n'(X_{n:n} - X_{n-1:n})]$ and 0 at the knots n/n' and 1 is possibly a reasonable approach. Note that between $i/n' \leq u < (i+1)/n'$, $i = 1, 2, \dots, n-1$, the values of $Q'(u)$ have been traditionally assumed to be constant given no other information about the density. Linearly interpolating the tail region makes use of the information regarding support of the distribution and leads to the tail extrapolation quantile function estimator based on the linear combination of order statistics defined for $0 < u < 1$ as follows:

$$\hat{Q}_{T_i}(u) = \begin{cases} \hat{Q}_i(1/n') + (X_{2:n} - X_{1:n}) \log(n'u), & 0 < u \leq 1/n', \\ \hat{Q}_i(u), & 1/n' < u < n/n', \\ \hat{Q}_i(n/n') - (X_{n:n} - X_{n-1:n}) \log(n'(1-u)), & n/n' \leq u < 1, \end{cases} \quad (2.1)$$

where $\hat{Q}_i(u)$ can be one of the traditional quantile estimators defined in Sec. 1. Note that the constant of integration was chosen such that the quantiles align with $\hat{Q}_i(u)$ at $1/n'$ and n/n' , respectively. The subscript i , for instance, can be L, H, D , or K as the linear combination of order statistics estimator at (1.2), the Hermitian estimator at (1.3), the Harrell–Davis estimator at (1.6), or the kernel quantile estimator at (1.9), respectively.

The properties and robustness of $\hat{Q}_{T_i}(u)$ will be examined in the next section. However, it is obvious that as $n \rightarrow \infty$ the “influence” of the extrapolation portion of the estimator diminishes at a rate $O(n^{-1})$.

In many instances, if the support of the data is assumed sufficiently near 0 and we know from some physical properties that $X > 0$. Then a slight modification of (2.1) yields the estimator:

$$\hat{Q}_{T_i}(u) = \begin{cases} \varepsilon \hat{Q}_i(1/n'), & 0 < u \leq 1/n', \\ \hat{Q}_i(u), & 1/n' < u < n/n', \\ \hat{Q}_i(n/n') - (X_{n:n} - X_{n-1:n}) \log(n'(1-u)), & n/n' \leq u < 1, \end{cases} \quad (2.2)$$

where $\varepsilon = n'u$.

3. Generating the Smoothed Bootstrap Samples based on $\hat{Q}_T(u)$

Given that the sample data $\mathbf{X} = (X_1, X_2, \dots, X_n)$ is held fixed, the traditional non parametric bootstrap replications are generated by taking a sample of size n with replacement from $\mathbf{X}$. Hutson (2000) have illustrated how this is equivalent to generating a random sample of size n from a uniform(0, 1) distribution, with corresponding order statistic $\mathbf{U} = (U_{1:n}, U_{2:n}, \dots, U_{n:n})$, and applying the sample quantile function estimators to the n uniform variates, i.e., a non parametric version of the probability integral transform. The resampling procedure by Hutson (2000) is described as follows:

1. *Step 1.* Generate a random sample of size n from a uniform (0,1) distribution denoted by $U^* = (U_1^*, U_2^*, \dots, U_n^*)$.
2. *Step 2.* Apply the quantile function estimator to each element of U^* to obtain one bootstrap replication, $X^* = (X_1^*, X_2^*, \dots, X_n^*)$, e.g., $X_1^* = \hat{Q}(U_1^*)$.
3. *Step 3.* Calculate the bootstrap replication of the statistics of interest $T_i(X^*)$.
4. *Step 4.* Repeat Steps 1–3 for B times in order to obtain the empirical bootstrap distribution for $T(X)$ from which various quantities such as variance and percentile estimators can be obtained.

Note that if the empirical version of the functional of interest, $T(F(X))$ consists of a function of order statistics, one might wish to use the algorithm developed by Schucany (1972) in conjunction with the new estimator for the purpose of programming efficiency, since it obviates the need to sort the observations after each resample.

4. Direct Density Estimation for L -estimators via Characteristic Functions

The goal of this section is to estimate the density function of the quantile estimators via a characteristic function approach utilizing the methodology derived in Wang et al. (2012). To eliminate the bootstrap resampling error, Hutson and Ernst (2000) developed the exact analytic expressions for the bootstrap mean and variance of any L -estimator, which are given as

Lemma 4.1

The exact bootstrap estimate of $\mu_{k(r)}$, $\sigma_{(r)}^2$, and $\sigma_{(rs)}$ $1 \leq r < s \leq n$, are

$$\hat{\mu}_{r:n}^k = \sum_{j=1}^n w_{j(r)} X_{j:n}^k, \quad (4.1)$$

$$\hat{\sigma}_{r:n}^2 = \sum_{j=1}^n w_{j(r)} (X_{j:n} - \mu_{r:n})^2, \quad (4.2)$$

$$\hat{\sigma}_{rs:n} = \sum_{j=2}^n \sum_{i=1}^{j-1} w_{ij(rs)} (X_{i:n} - \hat{\mu}_{r:n}) (X_{j:n} - \hat{\mu}_{s:n}) + \sum_{j=1}^n v_{j(rs)} (X_{j:n} - \hat{\mu}_{r:n}) (X_{j:n} - \hat{\mu}_{s:n}), \quad (4.3)$$

where

$$w_{j(r)} = {}_nC_r \cdot \int_{\frac{j}{n}}^{\frac{j-1}{n}} t^{r-1} (1-t)^{n-r} dt, \quad (4.4)$$

$$w_{ij(rs)} = \int_{(j-1)/n}^{j/n} \int_{(i-1)/n}^{i/n} f_{rs}(u_r, u_s) du_r du_s, \quad (4.5)$$

$$v_{j(rs)} = \int_{(j-1)/n}^{j/n} \int_{(j-1)/n}^{u_s} f_{rs}(u_r, u_s) du_r du_s, \quad (4.6)$$

and

$$f_{rs}(u_r, u_s) = {}_nC_{rs} u^{r-1} (u_s - u_r)^{s-r-1} (1 - u_s)^{n-s}, \quad (4.7)$$

is the joint distribution of two uniform order statistics $U_{r:n}$ and $U_{s:n}$ with ${}_nC_{rs} = n!/((r-1)!(s-r-1)!(n-s)!)$. The estimators for the moments of order statistics in Lemma 4.1 can get us to density estimation of L-estimators.

Direct application of Definition 3.1 in Wang et al. (2012) yields the following:

Definition 4.2

An empirical “plug-in” estimator for the c.g.f. of $Q(u)$ given an observed data set $X_1 = x_1, X_2 = x_2, \dots, X_n = x_n$ is defined as

$$\begin{aligned} \hat{\phi}_{\hat{Q}(u)}(t) = & \exp \left(\frac{1}{2} (2it \mathbf{w}' \hat{\boldsymbol{\mu}} - t^2 \mathbf{w}' \hat{\boldsymbol{\Sigma}} \mathbf{w}) \right) \\ & \times \left(1 + \frac{1}{6} \sum_{j=1}^n (it)^3 w_{j(u)}^3 \hat{\kappa}_{3j} + \frac{1}{24} \sum_{j=1}^n (it)^4 w_{j(u)}^4 \hat{\kappa}_{4j} \right. \\ & \left. + \frac{1}{72} \sum_{i=1}^n \sum_{j=1}^n (it)^6 w_{i(u)}^3 w_{j(u)}^3 \hat{\kappa}_{3i} \hat{\kappa}_{3j} \right), \end{aligned} \quad (4.8)$$

where $\mathbf{w} = (w_{1(u)}, w_{2(u)}, \dots, w_{n(u)})'$, $\hat{\boldsymbol{\mu}}$ and $\hat{\boldsymbol{\Sigma}}$ are the exact bootstrap estimators of mean vector and variance–covariance matrix of the order statistics and the estimated cumulants can be obtained by

$$\hat{\kappa}_{3j} = 2\hat{\mu}_{1,j:n}^3 - 3\hat{\mu}_{2,j:n}\hat{\mu}_{1,j:n} + \hat{\mu}_{3,j:n}, \quad (4.9)$$

$$\hat{\kappa}_{4j} = -6\hat{\mu}_{1,j:n}^4 + 12\hat{\mu}_{2,j:n}\hat{\mu}_{1,j:n}^2 - 3\hat{\mu}_{2,j:n}^2 - 4\hat{\mu}_{3,j:n}\hat{\mu}_{1,j:n} + \hat{\mu}_{4,j:n}. \quad (4.10)$$

Theorem 4.3

The inversion theorem based kernel estimators for the p.d.f. and c.d.f. for the kernel quantile function estimate $\hat{Q}(u)$ given the observed data $X_1 = x_1, X_2 = x_2, \dots, X_n = x_n$ are defined as

$$\hat{f}_{\hat{Q}(u)}(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp(-itx) \hat{\phi}_{\hat{Q}(u)}(t) dt, \quad (4.11)$$

$$\hat{F}_{\hat{Q}(u)}(x) = \frac{1}{2} + \frac{1}{2\pi} \int_0^{\infty} \frac{\exp(itx) \hat{\phi}_{\hat{Q}(u)}(-t) - \exp(-itx) \hat{\phi}_{\hat{Q}(u)}(t)}{it} dt. \quad (4.12)$$

The expressions for the p.d.f. and c.d.f. estimators of $\hat{Q}(u)$ at (4.11) and (4.12) follow from substituting $\hat{\phi}_{\hat{Q}(u)}(t)$ at (4.8) into the classic inversion theorem, respectively. Direct integration of these expressions is allowed by commonly used symbolic software packages, such as Maple and Mathematica.

Percentile Intervals—The $(1 - \alpha) \times 100$ percentile-type confidence interval for the u th population $Q(u)$ based on our new approach is defined as

$$\left(\hat{F}_{\hat{Q}(u)}^{-1}(\alpha/2), \hat{F}_{\hat{Q}(u)}^{-1}(1 - \alpha/2) \right), \quad (4.13)$$

where $\hat{F}_{\hat{Q}(u)}^{-1}(\cdot)$ is the inverse of $\hat{F}_{\hat{Q}(u)}(\cdot)$ defined at (4.12) above. Even though the calculation of the percentile interval requires numerical inversion of the c.d.f., it is straightforward using common statistical software packages, such as Maple and Mathematica or we can calculate it numerically using R or Matlab.

5. Bandwidth Selection Methods for Kernel Quantile Estimators

Sheather and Marron (1990) proposed a data-based choice of bandwidth h . The asymptotically optimal bandwidth for all p , apart from $p = .5$ when F is symmetric is

$$\hat{h}_{SM} = \alpha(K) \times \hat{\beta} \times n^{-1/3}, \quad (5.1)$$

where

$$\alpha(K) = \left[2 \int_{-\infty}^{\infty} u K(u) K^{(-1)}(u) du / \{ u^2 K(u) du \}^2 \right]^{1/3}, \quad (5.2)$$

$$\hat{\beta} = [\hat{Q}'_m(p)/\hat{Q}''_m(p)]^{2/3}, \quad (5.3)$$

and $\hat{Q}'_m(p)$ and $\hat{Q}''_m(p)$ are obtained through Theorem 3 in Sheather and Marron (1990). In our article, we choose $K(\cdot)$ and $K_*(\cdot)$ to be the normal kernel which is of order 2, so that

$$\int_{-\infty}^{\infty} uK(u)K^{(-1)}(u)du = 1/(2\sqrt{\pi}) \text{ and consequently, the } \alpha(K) = \pi^{-1/6}.$$

For an alternative bandwidth selection method, we will use the risk or mean integrated squared error (MISE) under the squared error loss as given by Miecznikowski et al. (2010),

$$\widehat{MISE}(Q, \hat{Q}) = \int_0^1 \left(\hat{\sigma}_{\hat{Q}(u;h)}^2 + (\hat{\mu}_{\hat{Q}(u;h)} - \hat{Q}(u))^2 \right) du, \quad (5.4)$$

where the $\hat{\sigma}_{\hat{Q}(u;h)}^2$ and $\hat{\mu}_{\hat{Q}(u;h)}$ can be developed by applying the approach described in Theorem 3.2 by Wang et al. (2012) as follows:

$$\hat{\mu}_{\hat{Q}(u;h)} = \mathbf{w}'\hat{\boldsymbol{\mu}}, \quad (5.5)$$

$$\hat{\sigma}_{\hat{Q}(u;h)}^2 = \mathbf{w}'\hat{\boldsymbol{\Sigma}}\mathbf{w}. \quad (5.6)$$

6. Properties of $\hat{Q}_{TK}(u)$

In this section, we illustrate how $\hat{Q}_{TK}(u)$ performs using the case of $\hat{Q}_{TK}(u)$ for finite samples and the large sample. The other quantile estimators in the class of $\hat{Q}_{Ti}(u)$ will follow similar patterns. The overall spirit of this section derives from Young (1994, p. 385) who notes that the “Attractiveness of the bootstrap approach is somewhat dimmed by the realization that favorable asymptotic of a particular bootstrap procedure is no guarantee of good small-sample behavior.”

In practice, it could be argued that it is more important to know how an estimator behaves for the small to moderate fixed n . A tool for graphically examining the behavior of $\hat{Q}_{TK}(u)$ in this context is the notion of sample profiles or “ideal samples,” e.g., see Mudholkar et al. (1991). The sample profile is given simply by the values $p_i = Q((i - 1/2)/n)$ for $i = 1, 2, \dots, n$. It can be shown that this particular profile is optimum among size n profiles of F . We used a slightly modified profile of $p_i = Q(i/n') = E(X_{i:n}) + O(n^{-1})$ since it provides a symmetric profile as compared to the traditionally defined profile. In Figs. 1–6, we plot the quantile function estimators $\hat{Q}_S(u)$, $\hat{Q}_K(u)$, and $\hat{Q}_{TK}(u)$ using the sample profiles for $n = 10$ and $n = 25$ for normal, Laplace, and exponential distributions, overlaid with the population quantile function. The bold solid lines represent the true quantile function estimator while the solid line, the dash line with the “x” sign, and the long dash line represent the step function $\hat{Q}_S$, the traditional kernel quantile estimator $\hat{Q}_K(u)$, and tail-extrapolated kernel quantile estimator $\hat{Q}_{TK}$, respectively. We also used the modified version of $\hat{Q}_{TK}(u)$ at (2.2) for the exponential distribution.

We see from the figures that $\hat{Q}_{TK}(u)$ fits well to the ideal sample for all distributions as small as $n = 10$, the exception being the Laplace distribution. The limitation of our method appears to be the inability to completely capture the tail behavior of heavy-tailed distributions such as Laplace as seen in Figs. 3 and 4. Note however, given the non parametric nature of the estimation, we believe that the $\hat{Q}_{TK}(u)$ provides a fairly flexible model and is still an improvement over other quantile function estimators, especially in terms of bootstrap resampling. The figures give some indication why $\hat{Q}_{TK}(u)$ might provide more accurate bootstrap samples in the extremes as compared to traditional quantile estimators.

The large-sample properties of $\hat{Q}_{TK}$ follow straightforwardly from well-known results pertaining to kernel quantile function estimators for interior points ($1/n' < u < n/n'$), e.g., see Sheather and Marron (1990) for detailed calculations. We refer the reader to the aforementioned reference for technical details.

Theorem 6.1

For large samples and $1/n' < u < n/n'$, $\hat{Q}_{TK}(u)$ is asymptotically normal with expectation and variance given as

$$E[\hat{Q}_{TK}(u)] = Q(u) + O(n^{-1}), \quad (6.1)$$

$$\text{Var}[\hat{Q}_{TK}(u)] = u(1-u)Q'(u)^2/n + Q(n^{-2}). \quad (6.2)$$

Proof—The details follow the results in Mudholkar and Hutson (1997) and Hutson (2002). Note that the interval $(1/n', n/n')$ will converge to $(0, 1)$ as n goes to infinity. Therefore, we conclude Equations (6.1) and (6.2) hold in $(0, 1)$ given n is sufficiently large.

7. Simulation Study

To compare the performance of traditional quantile estimators with the proposed semi-parametric tail-extrapolated quantile estimators, simulation studies were carried out to calculate the coverage probability with confidence interval lengths based on significant level of $\alpha = 0.05$. We ran 1,000 simulations and 1,000 smoothed bootstrap resampling per simulation for the following distributions: normal, logistic, Laplace, and exponential with sample sizes of 10, 25, and 50.

The quantile estimators we considered are $\hat{Q}_S, \hat{Q}_L, \hat{Q}_H, \hat{Q}_{HD}, \hat{Q}_K, \hat{Q}_{TL}, \hat{Q}_{THD}, \hat{Q}_{TK}$ defined at (1.1) to (1.6), (1.9), and (2.1) with $i = L, HD$, and K . The smoothed bootstrap and the direct density estimation via characteristic methods introduced in Secs. 3 and 4 were applied to obtain coverage probabilities for various quantile estimators. For simplicity, we used the data-based bandwidth selection method by Sheather and Marron (1990). When F is symmetric and $u = 0.5$, we chose the commonly used asymptotic bandwidth $\hat{h} = n^{-1/2}$ for kernel quantile estimators.

In general, as we can see from Tables 1 to 12, for small sample sizes ($n = 10, 25$), the commonly used quantile estimator $\hat{Q}_L$ and the Hermitian estimator $\hat{Q}_H$ are not obtainable for

extreme u 's and the coverage probabilities are poor for the rest of traditional quantile estimators. Our semi-parametric tail-extrapolated quantile estimators improve the coverage probability significantly for all kinds of distributions and parameter settings. The tail-extrapolated estimators with the smoothed bootstrap have outstanding performance over the traditional estimators, especially for the extreme tails, see Table 1 as an example. In comparison, the direct density estimation method is more robust with respect to u and the confidence intervals is narrower. This is especially obvious for heavy-tailed distributions, for example, in Tables 4 and 7. As sample size gets larger, the coverage probabilities for all kinds of quantile estimators become closer to nominal level.

Through the extensive simulation studies, we have several important findings which further demonstrated the performance of the proposed class of tail-extrapolated quantile estimators. First, the simulation results of the Q_{Ti} ($i = L, HD, K$) by the direct density estimation method are the same for $u = 0.95$ and $n = 10$ case (see Tables 1, 4, 7, and 10). Since $u = 0.95$ is within the tail region of the proposed tail-extrapolated quantile estimator, we are in fact using the same weighting scheme of order statistics.

Furthermore, the simulation results for Q_i and Q_{Ti} ($i = L, HD, K$) by the direct density method are identical for $u = 0.5, 0.75$ (and $u = 0.95$ when $n = 25, 50$), see Tables 2 and 3 as examples. The reason is under the above-mentioned simulation settings, the weights of order statistics, which determine the characteristic function of the quantile estimator, and ultimately the quantile estimators and the confidence intervals, are identical for the traditional estimators and the tail-extrapolated estimators.

Another pattern we observed is when we estimate $u = 0.5, 0.75$ in heavy-tail distributions, e.g., logistic distribution and Laplace distribution, the coverage probabilities and confidence interval lengths of Harrel–Davis estimator and the tail-extrapolated Harrel–Davis estimator by the smoothed bootstrap method yield the same results for large sample sizes as shown in Tables 6 and 9. This is due to the definition of Harrel–Davis estimator, which in heavy-tail distributions gives small weights to the order statistics along the two tails.

Lastly, it is also interesting to notice, for the same distribution, a decrease in the coverage probability of the tail-extrapolated quantile estimators when sample size increased from 10 to 25/50 for $u = 0.95$ case (see Tables 1–3 as an example). When sample size is 10, $u = 0.95$ is in the tail region of the proposed tail-extrapolated quantile estimators while for other sample sizes, $u = 0.95$ is within the region of $(1/n', n/n')$, which means the tail-extrapolated quantile estimators take the same form as the traditional ones. This phenomenon leads us to think there could be a further improved tail-extrapolated quantile estimator for which the defined region would vary by sample size accordingly.

8. Data Examples on Estimation of Extreme Quantiles

We used the flood data and the Buffalo snowfall amounts to compare the performance of the simple, Harrel–Davis, and kernel quantile function estimator with the corresponding tail-extrapolated quantile estimators in practical settings. The bandwidths are selected by minimizing MISE as described in Sec. 5. We did not include the direct density estimation method to calculate the confidence intervals based on our conclusion in Sec. 7 that smoothed

bootstrap method has better performance in estimating the extreme tails. Therefore, the 95% smoothed bootstrap confidence intervals are obtained from 10,000 resamples.

8.1. Flood Data

The annual maximum flood data for the Feather River at Oroville, CA, for a period of 59 years is given in Table 13; and that for the Blackstone River at Woonsocket, RI, for a period of 37 years appears in Table 14; for more details see Pericchi and Rodriguez-Iturbe (1985). The histograms of the standardized data with standard normal reference lines are shown in Figs. 7 and 9. The distributions of the data are right-skewed from the histograms. As it is well-known that the y th year flood amounts to estimating the u th quantile at $u = 1 - 1/y$, we use $u = 0.98$ and $u = 0.999$, respectively, to estimate the 50 and 1,000 year floods. The point estimates with 95% smoothed bootstrap confidence intervals are given for the Feather River and Blackstone River in Tables 17 and 18. We can easily see the limitations of the standard estimation procedures in terms of providing an upper confidence interval based-bound. This is particularly striking in the case of the 1,000 year flood estimation for both rivers.

Plots for the simple and kernel quantile estimators along with the tail-extrapolated kernel quantile estimators per river are provided in Figs. 8 and 10, respectively. We can see the significant improvement in potentially capturing the true underlying distributions for tail-extrapolated version quantile estimators through the examination of these plots.

8.2. Buffalo Snowfall Data

The 24 hour maximum snowfall amounts for 63 years from 1950 to 2012 and the annual snowfall amounts from 102 winters from 1910/11 to 2011/12 at Buffalo, NY, are given in Tables 15 and 16. The histograms of the standardized data with standard normal reference lines are shown in Figs. 11 and 13, respectively. While the 24 hour maximum snowfall amounts are skewed to the right, the annual snowfall amounts follow a normal distribution as discussed in Parzen (1979). The point estimates with 95% smoothed bootstrap confidence intervals for $u = 0.95, 0.975$, and 0.99 are given in Tables 19 and 20, respectively.

For Buffalo 24 hour maximum snowfall amount data, we get wider confidence intervals for the tail-extrapolated quantile estimators. We can see the limitations of the traditional quantile estimators in Table 19 in terms of providing reasonable upper confidence limits. For Table 20, the quantile estimators as well as the confidence intervals are very close since the sample size is large ($n = 102$) and the data follows the normal distribution with no significant skewness or long tails. Plots for the simple and kernel quantile estimators along with the tail-extrapolated kernel quantile estimators are provided in Figs. 12 and 14. Examination of these plots provides some insight into the quantile estimation and confidence interval calculations with respect to tail behavior estimation. The fitted line of the tail-extrapolated kernel quantile estimator is closer to the observations while the traditional kernel quantile estimator has certain limitation in estimating the extreme tails.

9. Summary and Discussion

For quantile estimation, the coverage probabilities obtained by the smoothed bootstrap and the direct density estimation methods through the use of the semi-parametric tail-

extrapolated quantile estimations improve upon the classic approach, which is based on the traditional quantile estimators such as the empirical sample quantile estimator. In general, the tail-extrapolated quantile estimators with the smoothed bootstrap method give the best coverage probabilities, especially for extreme-tails. For symmetric heavy tail distribution with small sample sizes, although we have decent coverage probabilities, it is important to note that much wider confidence lengths are obtained by applying the smoothed bootstrap method. In this case, we suggest the direct density estimation via characteristic function method if narrower confidence length is preferred. For large sample sizes ($n > 50$), the coverage probabilities obtained by these two methods are comparable.

Acknowledgments

We are thankful to the reviewers for their detailed reviews and helpful suggestions, which significantly improved our article.

References

- Cheng C, Parzen E. Unified estimators of smooth quantile and quantile density functions. *J. Stat. Plan. Inf.* 1997; 59(2):291–307.
- Falk M. Relative deficiency of kernel type estimators of quantiles. *Ann. Stat.* 1984; 12(1):261–268.
- Gamero J, Muñoz García J, Muñoz Reyes A. Bootstrapping statistical functionals. *Stat. Probab. Lett.* 1998; 39(3):229–236.
- Harrell F, Davis C. A new distribution-free quantile estimator. *Biometrika.* 1982; 69(3):635.
- Ho Y, Lee S. Iterated smoothed bootstrap confidence intervals for population quantiles. *Ann. Stat.* 2005; 33(1):437–462.
- Hutson A. Calculating non parametric confidence intervals for quantiles using fractional order statistics. *J. Appl. Stat.* 1999; 26(3):343–353.
- Hutson A. A composite quantile function estimator with applications in bootstrapping. *J. Appl. Stat.* 2000; 27(5):567–577.
- Hutson A. A semi-parametric quantile function estimator for use in bootstrap estimation procedures. *Stat. Comput.* 2002; 12(4):331–338.
- Hutson AD, Ernst MD. The exact bootstrap mean and variance of an l -estimator. *J. Royal Stat. Soc. Ser. B.* 2000; 62(1):89–94.
- Kaigh W, Lachenbruch P. A generalized quantile estimator. *Commun. Stat.-Theory Methods.* 1982; 11(19):2217–2238.
- Knight J. The joint characteristic function of linear and quadratic forms of non-normal variables. *Sankhy : Ind. J. Stat. Ser. A.* 1985; 47(2):231–238.
- Lee S. Optimal choice between parametric and non-parametric bootstrap estimates. *Math. Proc. Cambridge Philosophical Soc.* 1994; 115(2):335–364.
- Miecznikowski J, Wang D, Hutson A. Bootstrap mise estimators to obtain bandwidth for kernel density estimation. *Commun. Stat. - Simul. Comput.* 2010; 39(7):1455–1469.
- Mudholkar G, Hutson A. Asymmetric quasimedians: Remarks on an anomaly. *Stat. Probab. Lett.* 1997; 32(3):261–268.
- Mudholkar G, Hutson A, Marchetti C. A simple hermitian estimator of the quantile function. *J. Nonparametric Stat.* 2001; 13(2):229–243.
- Mudholkar G, Kollia G, Lin C, Patel K. A graphical procedure for comparing goodness-of-fit tests. *J. Royal Stat. Soc. Ser. B (Methodological).* 1991; 53(1):221–232.
- Parzen E. Nonparametric statistical data modeling. *J. Am. Stat. Assoc.* 1979; 74(365):105–121.
- Parzen E. Change pp plot and continuous sample quantile function. *Optimization.* 1993; 22(12):3287–3304.

- Pericchi, L.; Rodriguez-Iturbe, I. On the statistical analysis of floods. In: Atkinson, AC.; Fienberg, SE., editors. A Celebration of Statistics: The ISI Centenary Volume A Volume to Celebrate the Founding of the International Statistical Institute in 1885. New York: Springer; 1985. p. 511-541.
- Schucany W. Order statistics in simulation. *Journal of Statistical Computation and Simulation*. 1972; 1(3):281–286.
- Sheather S, Marron J. Kernel quantile estimators. *J. Am. Stat. Assoc.* 1990; 85(410):410–416.
- Silverman B, Young G. The bootstrap: to smooth or not to smooth? *Biometrika*. 1987; 74(3):469.
- Wang D, Miecznikowski J, Hutson A. Direct density estimate of L-estimates via characteristic functions with applications. *J. Stat. Plan. Inf.* 2012; 142(8):567–578.
- Yang S. A smooth nonparametric estimator of a quantile function. *J. Am. Stat. Assoc.* 1985; 80(392): 1004–1011.
- Young G. Bootstrap: more than a stab in the dark. *Stat. Sci.* 1994; 9(3):382–395.

Figure 1.
Profile plot for $N(0,1)$, $n = 10$.

Figure 2.
Profile plot for $N(0,1)$, $n = 25$.

Figure 3.
Profile plot for Laplace(0,1), $n = 10$.

Figure 4.
Profile plot for Laplace(0,1), $n = 25$.

Figure 5.
Profile plot for $\text{Exp}(1)$, $n = 10$.

Figure 6.
Profile plot for $\text{Exp}(1)$, $n = 25$.

Figure 7.
Histogram of standardized Feather River flood data for 59 years.

Figure 8.
Feather River estimated quantile plot.

Figure 9.
Histogram of standardized Blackstone River flood data for 37 years.

Figure 10.
Blackstone River estimated quantile plot.

Figure 11.
Histogram of standardized Buffalo 24 hour max snowfall amount for 63 years.

Figure 12.
Buffalo 24 hour maximum snowfall amount estimated quantile plot.

Figure 13.
Histogram of standardized Buffalo snowfall amount for 102 years.

Figure 14.
Buffalo annual snowfall amount estimated quantile plot.

Table 1Coverage probability with length for $N(0, 1)$ kernel with $n = 10$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.959(1.853)	0.951(1.976)	0.745(1.773)
$Q_S^{\hat{}}Boot$	0.934(1.642)	0.914(1.647)	0.393(1.067)
$Q_L^{\hat{}}Chara$	0.944(1.553)	0.938(1.692)	n/a
$Q_L^{\hat{}}Boot$	0.959(1.543)	0.905(1.422)	n/a
$Q_H^{\hat{}}Chara$	0.944(1.553)	0.931(1.644)	n/a
$Q_H^{\hat{}}Boot$	0.959(1.543)	0.904(1.405)	n/a
$Q_{HD}^{\hat{}}Chara$	0.926(1.370)	0.901(1.375)	0.692(1.585)
$Q_{HD}^{\hat{}}Boot$	0.941(1.388)	0.874(1.262)	0.382(0.999)
$Q_K^{\hat{}}Chara$	0.911(1.219)	0.899(1.556)	0.566(1.417)
$Q_K^{\hat{}}Boot$	0.659(0.636)	0.877(1.351)	0.348(1.013)
$Q_{TL}^{\hat{}}Chara$	0.944(1.553)	0.938(1.692)	0.873(2.521)
$Q_{TL}^{\hat{}}Boot$	0.958(1.637)	0.917(1.794)	0.888(3.886)
$Q_{THD}^{\hat{}}Chara$	0.926(1.370)	0.901(1.375)	0.873(2.521)
$Q_{THD}^{\hat{}}Boot$	0.944(1.424)	0.914(1.615)	0.897(3.923)
$Q_{TK}^{\hat{}}Chara$	0.911(1.219)	0.899(1.556)	0.873(2.521)
$Q_{TK}^{\hat{}}Boot$	0.866(1.085)	0.923(1.656)	0.896(3.995)

Table 2Coverage probability with length for $N(0, 1)$ kernel with $n = 25$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.974(1.154)	0.962(1.278)	0.860(1.623)
$Q_S^{\hat{}}Boot$	0.949(0.983)	0.918(1.039)	0.711(1.149)
$Q_L^{\hat{}}Chara$	0.974(1.154)	0.959(1.093)	0.862(1.273)
$Q_L^{\hat{}}Boot$	0.972(0.985)	0.967(1.083)	0.692(0.866)
$Q_H^{\hat{}}Chara$	0.966(1.001)	0.959(1.093)	n/a
$Q_H^{\hat{}}Boot$	0.972(0.969)	0.967(1.083)	n/a
$Q_{HD}^{\hat{}}Chara$	0.955(0.888)	0.946(0.975)	0.776(1.115)
$Q_{HD}^{\hat{}}Boot$	0.971(0.900)	0.958(0.992)	0.668(0.885)
$Q_K^{\hat{}}Chara$	0.955(0.816)	0.914(0.953)	0.778(1.179)
$Q_K^{\hat{}}Boot$	0.931(0.704)	0.945(0.981)	0.582(0.876)
$Q_{TL}^{\hat{}}Chara$	0.974(1.154)	0.959(1.093)	0.862(1.273)
$Q_{TL}^{\hat{}}Boot$	0.960(0.994)	0.959(1.120)	0.899(2.088)
$Q_{THD}^{\hat{}}Chara$	0.955(0.888)	0.946(0.975)	0.776(1.115)
$Q_{THD}^{\hat{}}Boot$	0.971(0.900)	0.959(1.002)	0.882(1.800)
$Q_{TK}^{\hat{}}Chara$	0.955(0.816)	0.914(0.953)	0.778(1.179)
$Q_{TK}^{\hat{}}Boot$	0.939(0.725)	0.954(1.007)	0.855(1.737)

Table 3Coverage probability with length for $N(0, 1)$ kernel with $n = 50$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_{\hat{S}}.Chara$	0.972(0.892)	0.966(0.947)	0.948(1.486)
$Q_{\hat{S}}.Boot$	0.938(0.688)	0.933(0.739)	0.911(1.159)
$Q_{\hat{L}}.Chara$	0.957(0.705)	0.954(0.796)	0.930(1.181)
$Q_{\hat{L}}.Boot$	0.968(0.689)	0.960(0.763)	0.877(0.964)
$Q_{\hat{H}}.Chara$	0.957(0.705)	0.949(0.779)	0.930(1.183)
$Q_{\hat{H}}.Boot$	0.968(0.689)	0.959(0.762)	0.877(0.964)
$Q_{\hat{HD}}.Chara$	0.942(0.648)	0.941(0.704)	0.886(0.988)
$Q_{\hat{HD}}.Boot$	0.958(0.650)	0.949(0.714)	0.846(0.860)
$Q_{\hat{K}}.Chara$	0.946(0.608)	0.898(0.652)	0.871(1.001)
$Q_{\hat{K}}.Boot$	0.964(0.622)	0.960(0.778)	0.769(0.770)
$Q_{\hat{TL}}.Chara$	0.957(0.705)	0.954(0.796)	0.930(1.181)
$Q_{\hat{TL}}.Boot$	0.950(0.692)	0.945(0.774)	0.917(1.338)
$Q_{\hat{THD}}.Chara$	0.942(0.648)	0.941(0.704)	0.886(0.988)
$Q_{\hat{THD}}.Boot$	0.958(0.650)	0.949(0.714)	0.911(1.213)
$Q_{\hat{TK}}.Chara$	0.946(0.608)	0.898(0.652)	0.871(1.001)
$Q_{\hat{TK}}.Boot$	0.964(0.623)	0.965(0.780)	0.903(1.187)

Table 4Coverage probability with length for $Logistic(0, 1)$ kernel with $n = 10$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.963(3.023)	0.951(3.379)	0.683(3.225)
$Q_S^{\hat{}}Boot$	0.932(2.753)	0.928(3.058)	0.407(2.127)
$Q_L^{\hat{}}Chara$	0.953(2.578)	0.944(3.061)	n/a
$Q_L^{\hat{}}Boot$	0.968(2.665)	0.909(2.634)	n/a
$Q_H^{\hat{}}Chara$	0.953(2.578)	0.939(3.000)	n/a
$Q_H^{\hat{}}Boot$	0.968(2.665)	0.906(2.603)	n/a
$Q_{HD}^{\hat{}}Chara$	0.940(2.314)	0.905(2.530)	0.618(2.914)
$Q_{HD}^{\hat{}}Boot$	0.951(2.426)	0.872(2.317)	0.387(1.978)
$Q_K^{\hat{}}Chara$	0.926(2.134)	0.906(2.866)	0.564(2.800)
$Q_K^{\hat{}}Boot$	0.664(1.115)	0.869(2.414)	0.369(2.021)
$Q_{TL}^{\hat{}}Chara$	0.953(2.578)	0.944(3.061)	0.830(4.496)
$Q_{TL}^{\hat{}}Boot$	0.968(2.756)	0.940(3.447)	0.892(8.500)
$Q_{THD}^{\hat{}}Chara$	0.940(2.314)	0.905(2.530)	0.830(4.496)
$Q_{THD}^{\hat{}}Boot$	0.955(2.505)	0.925(3.120)	0.898(8.527)
$Q_{TK}^{\hat{}}Chara$	0.926(2.134)	0.906(2.866)	0.830(4.496)
$Q_{TK}^{\hat{}}Boot$	0.852(2.044)	0.929(3.109)	0.902(8.684)

Table 5Coverage probability with length for $Logistic(0, 1)$ kernel with $n = 25$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.965(1.797)	0.961(2.104)	0.845(3.225)
$Q_S^{\hat{}}Boot$	0.946(1.612)	0.925(1.789)	0.729(2.458)
$Q_L^{\hat{}}Chara$	0.965(1.797)	0.956(1.910)	0.797(2.623)
$Q_L^{\hat{}}Boot$	0.966(1.632)	0.968(1.945)	0.691(1.887)
$Q_H^{\hat{}}Chara$	0.954(1.646)	0.956(1.910)	n/a
$Q_H^{\hat{}}Boot$	0.964(1.607)	0.968(1.945)	n/a
$Q_{HD}^{\hat{}}Chara$	0.936(1.472)	0.938(1.718)	0.757(2.411)
$Q_{HD}^{\hat{}}Boot$	0.954(1.505)	0.957(1.790)	0.682(1.900)
$Q_K^{\hat{}}Chara$	0.943(1.383)	0.917(1.690)	0.765(2.590)
$Q_K^{\hat{}}Boot$	0.906(1.220)	0.945(1.715)	0.635(1.935)
$Q_{TL}^{\hat{}}Chara$	0.965(1.797)	0.956(1.910)	0.797(2.623)
$Q_{TL}^{\hat{}}Boot$	0.954(1.636)	0.959(1.988)	0.861(4.826)
$Q_{THD}^{\hat{}}Chara$	0.936(1.472)	0.938(1.718)	0.757(2.411)
$Q_{THD}^{\hat{}}Boot$	0.954(1.505)	0.958(1.815)	0.838(4.088)
$Q_{TK}^{\hat{}}Chara$	0.943(1.383)	0.917(1.690)	0.765(2.590)
$Q_{TK}^{\hat{}}Boot$	0.917(1.269)	0.952(1.771)	0.833(3.936)

Table 6Coverage probability with length for *Logistic*(0, 1) kernel with $n = 50$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.955(1.244)	0.965(1.454)	0.936(2.880)
$Q_S^{\hat{}}Boot$	0.929(1.117)	0.941(1.248)	0.914(2.556)
$Q_L^{\hat{}}Chara$	0.938(1.132)	0.956(1.330)	0.907(2.602)
$Q_L^{\hat{}}Boot$	0.945(1.115)	0.961(1.315)	0.870(2.123)
$Q_H^{\hat{}}Chara$	0.938(1.132)	0.955(1.322)	0.907(2.599)
$Q_H^{\hat{}}Boot$	0.945(1.115)	0.961(1.312)	0.870(2.122)
$Q_{HD}^{\hat{}}Chara$	0.925(1.047)	0.941(1.206)	0.880(2.143)
$Q_{HD}^{\hat{}}Boot$	0.941(1.056)	0.951(1.237)	0.840(1.874)
$Q_K^{\hat{}}Chara$	0.935(0.994)	0.898(1.137)	0.879(2.236)
$Q_K^{\hat{}}Boot$	0.958(1.050)	0.969(1.318)	0.789(1.750)
$Q_{TL}^{\hat{}}Chara$	0.938(1.132)	0.956(1.330)	0.907(2.602)
$Q_{TL}^{\hat{}}Boot$	0.940(1.119)	0.958(1.315)	0.929(3.030)
$Q_{THD}^{\hat{}}Chara$	0.925(1.047)	0.941(1.206)	0.880(2.143)
$Q_{THD}^{\hat{}}Boot$	0.941(1.056)	0.951(1.237)	0.908(2.724)
$Q_{TK}^{\hat{}}Chara$	0.935(0.994)	0.898(1.137)	0.879(2.236)
$Q_{TK}^{\hat{}}Boot$	0.958(1.051)	0.977(1.329)	0.914(2.689)

Table 7Coverage probability with length for $Laplace(0, 1)$ kernel with $n = 10$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.973(2.265)	0.957(2.781)	0.663(2.787)
$Q_S^{\hat{}}Boot$	0.932(1.923)	0.928(2.390)	0.407(1.790)
$Q_L^{\hat{}}Chara$	0.969(1.803)	0.939(2.437)	n/a
$Q_L^{\hat{}}Boot$	0.971(1.936)	0.908(2.052)	n/a
$Q_H^{\hat{}}Chara$	0.969(1.803)	0.933(2.378)	n/a
$Q_H^{\hat{}}Boot$	0.971(1.936)	0.902(2.028)	n/a
$Q_{HD}^{\hat{}}Chara$	0.958(1.635)	0.899(1.970)	0.603(2.515)
$Q_{HD}^{\hat{}}Boot$	0.947(1.792)	0.870(1.793)	0.391(1.640)
$Q_K^{\hat{}}Chara$	0.933(1.586)	0.917(2.356)	0.559(2.430)
$Q_K^{\hat{}}Boot$	0.657(0.827)	0.878(1.823)	0.374(1.702)
$Q_{TL}^{\hat{}}Chara$	0.969(1.803)	0.939(2.437)	0.805(4.049)
$Q_{TL}^{\hat{}}Boot$	0.968(1.937)	0.939(2.777)	0.877(7.567)
$Q_{THD}^{\hat{}}Chara$	0.958(1.635)	0.899(1.970)	0.805(4.049)
$Q_{THD}^{\hat{}}Boot$	0.953(1.864)	0.916(2.523)	0.881(7.553)
$Q_{TK}^{\hat{}}Chara$	0.933(1.586)	0.917(2.356)	0.805(4.049)
$Q_{TK}^{\hat{}}Boot$	0.836(1.647)	0.928(2.452)	0.885(7.707)

Table 8Coverage probability with length for $Laplace(0, 1)$ kernel with $n = 25$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.970(1.183)	0.967(1.697)	0.844(2.986)
$Q_S^{\hat{}}Boot$	0.940(1.005)	0.908(1.326)	0.720(2.242)
$Q_L^{\hat{}}Chara$	0.970(1.183)	0.956(1.487)	0.820(2.450)
$Q_L^{\hat{}}Boot$	0.963(1.050)	0.960(1.511)	0.691(1.737)
$Q_H^{\hat{}}Chara$	0.960(1.021)	0.956(1.487)	n/a
$Q_H^{\hat{}}Boot$	0.961(1.035)	0.960(1.511)	n/a
$Q_{HD}^{\hat{}}Chara$	0.952(0.930)	0.943(1.322)	0.763(2.224)
$Q_{HD}^{\hat{}}Boot$	0.955(0.997)	0.943(1.392)	0.678(1.730)
$Q_K^{\hat{}}Chara$	0.945(0.947)	0.934(1.317)	0.779(2.433)
$Q_K^{\hat{}}Boot$	0.911(0.884)	0.943(1.281)	0.656(1.831)
$Q_{TL}^{\hat{}}Chara$	0.970(1.183)	0.956(1.487)	0.820(2.450)
$Q_{TL}^{\hat{}}Boot$	0.952(1.015)	0.952(1.536)	0.849(4.555)
$Q_{THD}^{\hat{}}Chara$	0.952(0.930)	0.943(1.322)	0.763(2.224)
$Q_{THD}^{\hat{}}Boot$	0.955(0.997)	0.944(1.416)	0.846(3.818)
$Q_{TK}^{\hat{}}Chara$	0.945(0.947)	0.934(1.317)	0.779(2.433)
$Q_{TK}^{\hat{}}Boot$	0.928(0.932)	0.951(1.330)	0.842(3.683)

Table 9Coverage probability with length for $Laplace(0, 1)$ kernel with $n = 50$

$Q(\hat{u})$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$Q_S^{\hat{}}Chara$	0.971(0.856)	0.959(1.162)	0.946(2.836)
$Q_S^{\hat{}}Boot$	0.947(0.663)	0.936(0.930)	0.928(2.520)
$Q_L^{\hat{}}Chara$	0.961(0.669)	0.955(1.023)	0.927(2.593)
$Q_L^{\hat{}}Boot$	0.961(0.680)	0.966(1.001)	0.891(2.099)
$Q_H^{\hat{}}Chara$	0.961(0.669)	0.954(1.005)	0.927(2.592)
$Q_H^{\hat{}}Boot$	0.961(0.680)	0.966(0.999)	0.891(2.099)
$Q_{HD}^{\hat{}}Chara$	0.953(0.634)	0.937(0.913)	0.901(2.113)
$Q_{HD}^{\hat{}}Boot$	0.948(0.656)	0.958(0.946)	0.865(1.834)
$Q_K^{\hat{}}Chara$	0.947(0.636)	0.928(0.868)	0.900(2.277)
$Q_K^{\hat{}}Boot$	0.962(0.729)	0.966(0.958)	0.830(1.806)
$Q_{TL}^{\hat{}}Chara$	0.961(0.669)	0.955(1.023)	0.927(2.593)
$Q_{TL}^{\hat{}}Boot$	0.954(0.666)	0.951(0.998)	0.929(3.080)
$Q_{THD}^{\hat{}}Chara$	0.953(0.634)	0.937(0.913)	0.901(2.113)
$Q_{THD}^{\hat{}}Boot$	0.948(0.656)	0.958(0.946)	0.914(2.749)
$Q_{TK}^{\hat{}}Chara$	0.947(0.636)	0.928(0.868)	0.900(2.277)
$Q_{TK}^{\hat{}}Boot$	0.962(0.730)	0.976(0.983)	0.925(2.740)

Table 10Coverage probability with length for $Exp(1)$ kernel with $u = 10$

$\hat{Q}(u)$	$u = 0.05$	$u = 0.25$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$\hat{Q}_S Chara$	0.933(1.157)	0.965(1.232)	0.966(1.964)	0.940(2.634)	0.664(2.693)
$\hat{Q}_S Boot$	0.410(0.345)	0.923(0.752)	0.922(1.532)	0.918(2.246)	0.392(1.696)
$\hat{Q}_L Chara$	n/a	0.959(1.001)	0.953(1.371)	0.924(2.315)	n/a
$\hat{Q}_L Boot$	n/a	0.905(0.710)	0.957(1.321)	0.893(1.878)	n/a
$\hat{Q}_H Chara$	n/a	0.945(0.928)	0.953(1.371)	0.920(2.255)	n/a
$\hat{Q}_H Boot$	n/a	0.902(0.703)	0.957(1.321)	0.893(1.856)	n/a
$\hat{Q}_{HP} Chara$	0.908(0.957)	0.922(0.717)	0.931(1.188)	0.882(1.835)	0.575(2.421)
$\hat{Q}_{HP} Boot$	0.357(0.364)	0.862(0.691)	0.926(1.210)	0.863(1.600)	0.38(1.55)
$\hat{Q}_K Chara$	0.735(0.708)	0.848(0.811)	0.953(1.347)	0.913(2.231)	0.492(2.23)
$\hat{Q}_K Boot$	0.370(0.817)	0.845(0.857)	0.928(1.206)	0.870(1.632)	0.331(1.55)
$\hat{Q}_{TK} Chara$	0.971(1.260)	0.959(1.001)	0.953(1.371)	0.924(2.315)	0.786(3.801)
$\hat{Q}_{TK} Boot$	0.961(0.942)	0.935(0.750)	0.958(1.373)	0.924(2.607)	0.871(7.077)
$\hat{Q}_{THP} Chara$	0.971(1.260)	0.922(0.717)	0.931(1.188)	0.882(1.835)	0.786(3.801)
$\hat{Q}_{THP} Boot$	0.971(0.993)	0.926(0.764)	0.930(1.250)	0.894(2.285)	0.87(7.051)
$\hat{Q}_{TK} Chara$	0.971(1.260)	0.848(0.811)	0.953(1.347)	0.913(2.231)	0.786(3.801)
$\hat{Q}_{TK} Boot$	0.992(1.341)	0.916(0.961)	0.928(1.220)	0.918(2.202)	0.874(7.19)

Table 11Coverage probability with length for $Exp(1)$ kernel with $u = 25$

$\hat{Q}(u)$	$u = 0.05$	$u = 0.25$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$\hat{Q}_S Chara$	0.958(1.282)	0.964(0.889)	0.962(1.100)	0.965(1.708)	0.836(2.932)
$\hat{Q}_S Boot$	0.740(0.216)	0.922(0.478)	0.937(0.823)	0.936(1.326)	0.710(2.188)
$\hat{Q}_L Chara$	0.940(1.899)	0.959(0.581)	0.962(1.100)	0.965(1.488)	0.794(2.411)
$\hat{Q}_L Boot$	0.723(0.146)	0.963(0.448)	0.959(0.832)	0.967(1.505)	0.680(1.701)
$\hat{Q}_H Chara$	n/a	0.959(0.581)	0.947(0.881)	0.965(1.488)	n/a
$\hat{Q}_H Boot$	n/a	0.963(0.448)	0.955(0.819)	0.967(1.505)	n/a
$\hat{Q}_{HP} Chara$	0.876(1.855)	0.944(0.527)	0.932(0.756)	0.949(1.317)	0.760(2.177)
$\hat{Q}_{HP} Boot$	0.670(0.168)	0.926(0.423)	0.939(0.771)	0.957(1.379)	0.669(1.694)
$\hat{Q}_K Chara$	0.696(1.319)	0.805(0.502)	0.920(0.767)	0.946(1.325)	0.749(2.300)
$\hat{Q}_K Boot$	0.210(0.609)	0.902(0.643)	0.943(0.800)	0.952(1.230)	0.584(1.663)
$\hat{Q}_{TK} Chara$	0.940(1.899)	0.959(0.581)	0.962(1.100)	0.965(1.488)	0.794(2.411)
$\hat{Q}_{TK} Boot$	0.910(0.245)	0.962(0.451)	0.946(0.832)	0.964(1.545)	0.863(4.448)
$\hat{Q}_{THD} Chara$	0.876(1.855)	0.944(0.527)	0.932(0.756)	0.949(1.317)	0.760(2.177)
$\hat{Q}_{THD} Boot$	0.905(0.252)	0.930(0.424)	0.939(0.771)	0.957(1.403)	0.846(3.723)
$\hat{Q}_{TK} Chara$	0.696(1.319)	0.805(0.502)	0.920(0.767)	0.946(1.325)	0.749(2.300)
$\hat{Q}_{TK} Boot$	0.795(0.769)	0.904(0.653)	0.946(0.813)	0.953(1.275)	0.821(3.567)

Table 12Coverage probability with length for $Exp(1)$ kernel with $u = 50$

$\hat{Q}(u)$	$u = 0.05$	$u = 0.25$	$u = 0.5$	$u = 0.75$	$u = 0.95$
$\hat{Q}_S Chara$	0.969(2.511)	0.968(0.884)	0.969(0.869)	0.959(1.162)	0.936(2.830)
$\hat{Q}_S Boot$	0.930(0.136)	0.940(0.320)	0.949(0.575)	0.925(0.925)	0.910(2.492)
$\hat{Q}_L Chara$	0.964(3.193)	0.970(0.713)	0.955(0.588)	0.940(1.018)	0.913(2.576)
$\hat{Q}_L Boot$	0.907(0.114)	0.970(0.320)	0.961(0.561)	0.959(0.994)	0.880(2.080)
$\hat{Q}_H Chara$	0.965(3.205)	0.968(0.663)	0.955(0.588)	0.936(1.002)	0.912(2.576)
$\hat{Q}_H Boot$	0.907(0.114)	0.969(0.319)	0.961(0.561)	0.959(0.991)	0.880(2.079)
$\hat{Q}_{HP} Chara$	0.940(3.384)	0.952(0.574)	0.943(0.535)	0.924(0.907)	0.875(2.106)
$\hat{Q}_{HP} Boot$	0.850(0.113)	0.956(0.302)	0.951(0.531)	0.952(0.938)	0.847(1.821)
$\hat{Q}_K Chara$	0.713(2.564)	0.792(0.550)	0.924(0.536)	0.911(0.854)	0.880(2.159)
$\hat{Q}_K Boot$	0.175(0.439)	0.895(0.520)	0.965(0.579)	0.956(0.888)	0.783(1.625)
$\hat{Q}_{TK} Chara$	0.964(3.193)	0.970(0.713)	0.955(0.588)	0.940(1.018)	0.913(2.576)
$\hat{Q}_{TK} Boot$	0.937(0.126)	0.961(0.320)	0.958(0.565)	0.941(0.993)	0.917(3.022)
$\hat{Q}_{THD} Chara$	0.940(3.384)	0.952(0.574)	0.943(0.535)	0.924(0.907)	0.875(2.106)
$\hat{Q}_{THD} Boot$	0.935(0.130)	0.956(0.302)	0.951(0.531)	0.952(0.938)	0.905(2.692)
$\hat{Q}_{TK} Chara$	0.713(2.564)	0.792(0.550)	0.924(0.536)	0.911(0.854)	0.880(2.159)
$\hat{Q}_{TK} Boot$	0.380(0.523)	0.905(0.516)	0.965(0.586)	0.959(0.920)	0.908(2.618)

Table 13

The consecutive annual flood discharge ranges of the Feather River for 59 years at Oroville, CA

Year	Flood Discharge in (ft ³ /s)
1902–1909	42,000, 102,000, 118,000, 81,000, 128,000, 230,000, 16,300, 140,000
1910–1917	31,000, 75,400, 16,400, 16,800, 122,000, 81,400, 42,400, 80,400
1918–1925	28,200, 65,900, 23,400, 62,300, 36,400, 22,400, 42,400, 64,300
1926–1933	55,700, 94,000, 185,000, 14,000, 80,100, 11,600, 22,600, 88,60
1934–1941	20,300, 58,600, 85,400, 19,200, 185,000, 8,080, 152,000, 84,200
1942–1949	110,000, 108,000, 24,900, 60,100, 54,400, 45,600, 36,700, 16,800
1950–1957	46,400, 92,100, 59,200, 13,000, 113,000, 54,800, 203,000, 83,100
1958–1960	102,000, 34,500, 135,000

Table 14

The consecutive annual flood discharge rates of the Blackstone River for 37 years at Woonsocket, RI

Year	Flood Discharge in (ft ³ /s)
1929–1938	4,570, 1,970, 8,220, 4,530, 5,780, 6,560, 7,500, 15,000, 6,340, 15,100
1939–1948	3,840, 5,860, 4,480, 5,330, 5,310, 3,830, 3,410, 3,830, 3,150, 5,810
1949–1958	2,030, 3,620, 4,920, 4,090, 5,570, 9,400, 32,900, 8,710, 3,850, 4,970
1959–1965	5,398, 4,780, 4,020, 5,790, 4,510, 5,520, 5,300

Table 15

The Buffalo max 24 hour snowfall for 63 years from 1950 to 2012 at Buffalo, NY

Year	Max 24 hour snowfall
1950–1959	6.1, 2.5, 2.3, 2.2, 2.5, 6.2, 6.2, 7.6, 8.3, 6.9
1960–1969	3.2, 3.3, 6.9, 7.8, 8.1, 3.3, 15, 1.9, 3.9, 7.8
1970–1979	5.8, 4.1, 3.5, 2, 4.9, 2.1, 5.3, 13.2, 8.8, 7.4
1980–1989	3.4, 3, 18.3, 3.1, 3.2, 15.6, 5.7, 5.2, 5, 1.7
1990–1999	3.6, 4.6, 4.5, 4.5, 7.1, 11.1, 10.1, 12.1, 5.5, 12.4
2000–2012	5.1, 3.6, 3.7, 9.1, 8.8, 6.7, 1.8, 5.4, 3, 7.1, 8.8, 3.6, 6.4

Table 16

The amount of snowfall for 102 winters from 1910/11 to 2011/12 at Buffalo, NY

Year	Amount of snowfall
1910–11 to 1919–20	126.4, 82.4, 78.1, 51.1, 90.9, 76.2, 104.5, 87.4, 110.5, 25.0
1921–22 to 1929–30	69.3, 53.5, 39.8, 63.6, 46.7, 72.9, 79.6, 83.6, 80.7, 60.3
1931–32 to 1939–40	79.0, 74.4, 49.6, 54.7, 71.8, 49.1, 103.9, 51.6, 82.4, 83.6
1941–42 to 1949–50	77.8, 79.3, 89.6, 85.5, 58.0, 120.7, 110.5, 65.4, 39.9, 40.1
1951–52 to 1959–60	88.7, 71.4, 83.0, 55.9, 89.9, 84.8, 105.2, 113.7, 124.7, 114.5
1961–62 to 1969–70	115.6, 102.4, 101.4, 89.8, 71.5, 70.9, 98.3, 55.5, 66.1, 78.4
1971–72 to 1979–80	120.5, 97.0, 110.0, 88.7, 95.6, 82.5, 199.4, 154.3, 97.3, 68.4
1980–81 to 1989–90	60.9, 112.4, 52.4, 132.5, 107.2, 114.7, 67.5, 56.4, 67.4, 93.7
1990–91 to 1999–00	57.5, 92.8, 93.2, 112.7, 74.6, 141.4, 97.6, 75.6, 100.5, 63.6
2000–01 to 2011–12	158.7, 132.4, 111.3, 100.9, 109.1, 78.2, 88.9, 103.8, 100.2, 74.1, 111.8, 36.7

Table 17

Quantile estimators for Feather River 50/1,000 year flood with 95% confidence intervals

Quantiles	50 year flood ($u = 0.98$)	1000 year flood ($u=0.999$)
$\hat{Q}_S$	203,000 (140,000,230,000)	230,000 (185,000,230,000)
$\hat{Q}_{TS}$	203,000 (140,000,267,665)	305,962 (185,000,645,053)
$\hat{Q}_{HD}$	213,732 (165,221,225,561)	229,572 (174,055,229,589)
$\hat{Q}_{THD}$	213,732 (165,221,283,774)	293,158 (188,928,586,929)
$\hat{Q}_K$	206,797 (156,342,214,245)	217,749 (165,989,216,204)
$\hat{Q}_{TK}$	206,797 (156,559,267,009)	285,008 (186,116,565,840)

Table 18

Quantile estimators for Blackstone River 50/1,000 year flood with 95% confidence intervals

Quantiles	50 year flood ($u = 0.98$)	1000 year flood ($u = 0.999$)
$\hat{Q}_S$	32,900 (9,400,32,900)	32,900 (9,400,32,900)
$\hat{Q}_{TS}$	37,785 (9,400,116,445)	37,785 (9,400,11,6445)
$\hat{Q}_{HD}$	28,128 (11,308,31,629)	28,128 (11,308,31,629)
$\hat{Q}_{THD}$	31,055 (11,571,91,165)	31,055 (11,576,91,165)
$\hat{Q}_K$	26,423 (9,692,30,633)	26,423 (9,692,30,633)
$\hat{Q}_{TK}$	28,782 (9,900,84,581)	28,782 (9,900,84,581)

Table 19

Quantile estimators for Buffalo 24 hour maximum snowfall amount with 95% confidence intervals

Quantiles	$u = 0.95$	$u = 0.975$	$u = 0.99$
$\hat{Q}_S$	13.20(9.10,15.60)	15.60(12.10,18.30)	18.30(13.20,18.30)
$\hat{Q}_{TS}$	13.20(9.10,15.60)	15.60(12.10,22.06)	19.50(13.20,32.86)
$\hat{Q}_{HD}$	14.12(10.99,16.76)	16.22(12.58,17.74)	17.65(13.37,18.11)
$\hat{Q}_{THD}$	14.12(11.02,18.32)	16.22(12.59,22.35)	18.34(13.54,28.73)
$\hat{Q}_K$	13.94(10.38,16.33)	15.86(12.15,16.96)	16.86(12.84,17.14)
$\hat{Q}_{TK}$	13.94(10.41,17.78)	15.86(12.17,21.38)	17.72(13.11,26.38)

Table 20

Quantile estimators for Buffalo annual snowfall amount with 95% confidence intervals

Quantiles	$u = 0.95$	$u = 0.975$	$u = 0.99$
$\hat{Q}_S$	132.40(114.70,154.30)	154.30(126.40,199.40)	158.70(132.40,199.40)
$\hat{Q}_{TS}$	132.40(114.70,154.30)	154.30(126.40,218.32)	158.70(132.40,256.03)
$\hat{Q}_{HD}$	135.23(121.16,163.34)	155.95(132.41,184.38)	182.53(141.16,194.95)
$\hat{Q}_{THD}$	135.23(121.22,167.85)	155.95(132.42,220.21)	182.53(141.16,293.63)
$\hat{Q}_K$	132.37(117.20,158.05)	152.98(128.42,175.42)	173.53(136.46,181.88)
$\hat{Q}_{TK}$	132.37(117.20,158.35)	152.98(128.65,212.31)	173.53(136.46,268.03)