Published in final edited form as:

J Subst Abuse Treat. 2016 April ; 63: 34–38. doi:10.1016/j.jsat.2015.12.005.

Gender Differences In The Life Concerns Of Persons Seeking Alcohol Detoxification

Michael D. Stein^{a,b}, Megan M. Risi^a, Jessica N. Flori^a, Micah T. Conti^a, Bradley J. Anderson^a, and Genie L. Bailey^{b,c}

^aGeneral Medicine Research Unit, Butler Hospital, Providence, RI, 02906

^bWarren Alpert Medical School of Brown University, Providence, RI, 02912

^cStanley Street Treatment and Resources, Inc., Fall River, Massachusetts 02720

Abstract

Background—This study explored the life concerns of persons seeking alcohol detoxification, a group with multiple life and psychosocial challenges. Gender may be an important contributor to the particular life concerns of persons with alcohol use disorders.

Methods—Using a 32-item, previously-validated life concerns survey that captures ten conceptual domains, we interviewed persons entering inpatient alcohol detoxification asking them to rate their level of concern about health and welfare items.

Results—Participants (n = 189) were 27% female, with a mean age of 43.5 years. Overall, concern about alcohol problems was perceived as the most serious, followed by mental health, cigarette smoking, financial, and relationship problems. Men were significantly more concerned than women about six of the ten domains including money, drug use, transmissible diseases, and physical illness.

Conclusions—Recognition of the daily worries of persons seeking inpatient alcohol detoxification persons could allow providers to better tailor their services to the context of their patients' lives. Focusing on pressing life concerns such as mental health, financial, relationship problems, and other drug use may influence detoxification services and aftercare treatment choices.

Keywords

alcohol; detoxification; self-reported health; life concerns

Corresponding Author: Michael D. Stein, M.D., Professor of Medicine, Health Services, Policy & Practice, Alpert School of Medicine at Brown University, Butler Hospital, 345 Blackstone Blvd., Providence, RI 02906, Telephone: (401) 455-6646, FAX: (401) 455-6685, Michael_Stein@brown.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1.0 INTRODUCTION

Alcohol Use Disorder (AUD) is pervasive and poses a significant public health concern (Whiteford, et al., 2013). Alcohol use results in nearly 88,000 deaths annually, making it the third leading cause of preventable death in the United States (Mokdad, Marks, Stroup, & Gerberding, 2004). The lives of persons with severe AUD are often challenged by economic burden (Navarro, Doran, & Shakeshaft, 2011), mental health disorders, traumatic events, unemployment, violence (O'Meara, Witherspoon, Hapangama, & Hyam, 2011), housing problems, stigma, relational disruptions, infectious disease risks (Claxton, DeLuca, & van Dulmen, 2015), and acute and chronic medical problems associated with heavy drinking (Chartier, Hesselbrock, & Hesselbrock, 2013; National Institute on Alcohol Abuse and Alcoholism, 2000a, 2000b; Pettinati, O'Brien, & Dundon, 2013; Rehm, 2011). The Substance Abuse and Mental Health Services Administration (SAMHSA) reports that in 2013, an estimated 17.3 million Americans met the criteria for AUD in the past year, but only 7.7% sought treatment (Substance Abuse and Mental Health Services Administration, 2014).

Among persons with AUD interested in long-term abstinence, approximately 30% are admitted as a first step to a medically supervised detoxification facility (Substance Abuse and Mental Health Services Administration, 2014). Inpatient detoxification provides patients with a medically supervised protocol to prevent or mitigate alcohol withdrawal symptoms, begin counseling, and access additional treatment modalities after discharge (Blondell, Smith, Canfield, & Servoss, 2006; Kosten & O'Connor, 2003; Merkx, et al., 2014). Aftercare planning is essential to reduce relapse risk, and detoxification program staff may be able to use knowledge of patient-specific life concerns in addition to abstinence to suggest aftercare treatment choices.

Previous work has described the life concerns of other urban populations with multiple life and psychosocial challenges (Carey, Braaten, Jaworski, Durant, & Forsyth, 1999; Carey & Senn, 2013), including opioid users (Stein, Anderson, Thurmond, & Bailey, 2015). Life concerns—subjective worries—are distinct from specific stressful or negative life events such as the death of a friend, divorce, or a hospitalization (Veenstra, et al., 2006). Negative life events have been examined in relation to alcohol use in cross-sectional and longitudinal studies of general population samples (Perreira & Sloan, 2001) and gender differences were notable. Among married couples, men reported more negative life events in the areas of work and finance whereas women noted more events related to their social networks (Conger, Lorenz, Elder, Simons, & Ge, 1993). Among past year drinkers, the impact of life events affected men and women differently with life events affecting the alcohol use of men more profoundly (Dawson, Grant, & Ruan, 2005; Frone, Cooper, & Russell, 1994).

Historically, men have higher rates of alcohol use disorders (AUDs) than women (Nolen-Hoeksema & Hilt, 2006), although the gap in the prevalence of alcohol use disorder has narrowed in the recent past, with the prevalence now two times more common in men (Keyes, Grant, & Hasin, 2008). The burden of morbidity is high across genders (Goldstein, Dawson, Chou, & Grant, 2012), but women are more susceptible than men to certain medical consequences such as liver disease, ulcers, and hypertension (Mann, et al., 2005;

Stewart, et al., 2009). While women may be less likely to enter alcohol treatment, gender is not a consistent predictor of outcome (Green, Polen, Lynch, Dickinson, & Bennett, 2004; Greenfield, et al., 2007).

Gender may also be an important contributor to the particular life concerns of persons with alcohol use disorders who seek inpatient detoxification, and such concerns could, in theory, drive aftercare choices. While the severity of alcohol use is likely to be the immediate and dominant focus of persons seeking detoxification, impairments in mental health and social functioning (Daepfen, Krieg, Burnand, & Yersin, 1998; Morgan, Landron, Leher, & New European Alcoholism Treatment Study, 2004; Polen, Green, Perrin, Anderson, & Weisner, 2010) may influence both treatment needs during detox and future service use. For example, women with AUD are more frequently severely depressed than men (Pettinati, Pierce, Wolf, Rukstalis, & O'Brien, 1997) and mental health concerns may guide treatment needs. Similarly, relationship concern may influence treatment. For women, marriage and marital stress are risk factors for alcohol relapse as alcoholic women are more likely to be married to heavy drinking partners than are alcoholic men; among men, marriage lowers relapse risk (Walitzer & Dearing, 2006); marital counseling may be an important part of aftercare. General quality of life in female alcohol misusers is lower than in males with comparable levels of dependency (Peters, Millward, & Foster, 2003), with psychiatric comorbidity and social environment being important contributing factors influencing quality of life (QoL) (Foster, Powell, Marshall, & Peters, 1999).

The spectrum of life concerns has never been comprehensively evaluated in persons entering an inpatient alcohol detoxification program. In this study we describe the relative importance of specific life concerns in this population and hypothesize that there will be gender differences, namely that women will be more concerned with relationship and mental health issues, and men will be more concerned with their use of drugs in addition to alcohol.

2.0 METHODS

Between September 2014 and March 2015, consecutive persons seeking alcohol detoxification were approached within the first 24 hours of admission by research staff to Stanley Street Treatment Addiction and Recovery, Inc. (SSTAR) in Fall River, Massachusetts to participate in a research survey. SSTAR's program, one of the largest in Southeastern New England, has 38 beds and is a 24-hour medically supervised treatment facility that provides evaluation and withdrawal management. The average length of stay for alcohol detoxification at SSTAR during the study period was 3.6 days.

Two hundred four patients who were 18 years or older and English-speaking were admitted to SSTAR seeking alcohol detoxification services during the recruitment period and therefore eligible to provide verbal informed consent as approved by the Butler Hospital Institutional Review Board. Fifteen refused participation. The remaining 189 persons completed a face-to-face, 15-minute, structured, standardized interview; no compensation was provided.

2.1 Measures

Sample descriptors included age, gender, race or ethnicity, employment (part or full-time vs. unemployed), homelessness (any nights on street or in a shelter in the prior 90 days), and years of education. Participants were asked if they had attended an outpatient primary care visit in the past year, if they had been hospitalized for mental health issues in the past year, and what treatment services, if any, they have received in the past year. Participants were classified as recent cocaine, benzodiazepine or marijuana users if they reported any use during the past 30 days. Participants also reported frequency and usual quantity of alcohol use during the past 30 days; binge drinking was defined as 4 drinks on a single occasion for females and 5 drinks on a single occasion for males (National Institute on Alcohol Abuse and Alcoholism, 2005).

We asked participants about life concern items previously validated with general urban population (Carey & Senn, 2013) and later used to characterize an opioid dependent population (Stein, et al., 2015). We asked each participant, “The following questions are related to concerns you may have about your health and welfare. Please rate how much the following are of concern to you.” Responses were scored on a 0, “no concern,” 1, “little concern,” 2, “some concern” to 3, “serious concern” four-point scale. All scales were constructed as the mean of included items and range from 0 to 3 with a mid-point of 1.5. Scale means are presented to describe the relative degree to which participants perceive these issues as serious concerns. We report Cronbach's alpha as a measure of internal consistency for multi-item scales. Using 32 survey items, we organized life concerns into seven conceptual multi-item domains; serious physical illness (5 items; $\alpha = .76$, e.g. high blood pressure), transmitted diseases (3 items; $\alpha = .87$, e.g. HIV/AIDS), mental health (8 items; $\alpha = .79$ e.g. feeling depressed), financial concerns (5 items; $\alpha = .82$, e.g. not having enough money for basic needs), health insurance (2 items; $\alpha = .92$, e.g., not having health insurance), concerns about community safety (4 items.; $\alpha = .82$, e.g. violence in my community), and relationship concerns (5 items; $\alpha = .85$, e.g. being cut off from my family) (Stein et al, 2015). Three additional single-item domains, concern about their alcohol problem, concern about smoking, and concern about drug use, were retained in subsequent analysis because of their combination of relevance to and high endorsement in this population.

2.2 Analytical methods

We report descriptive statistics to summarize the characteristics of the sample. T-tests for differences in means and χ^2 -tests for differences in counts were used to test for gender differences on demographics and background variables. For each life concern we report the mean, Standard Deviation, and the number and percentage of persons scoring 0. The distributions of life concern scales varied, many were distinctly non-normal and several clustered at 0. To test for gender differences in life concerns, we used the nonparametric Wilcoxon Rank-Sum test for equality of rank-ordered distributions.

3.0 RESULTS

Participants averaged 43.5 ($\pm$ 9.9) years of age and 163 (86.7%) were non-Latino Caucasian (Table 1). Mean years of education was 12.4 ($\pm$ 2.1), 46 (24.3%) were employed either part- or full-time, and 20 (10.6%) were homeless. One hundred twenty-nine (68.6%) said they had seen a primary care physician within the last year. Fifty-two (27.7%) reported no history of any prior alcohol detox, 82 (43.6%) had been in detoxification within the past year, and 54 (28.7%) had been in detoxification, but not within the past year. During the past month, participants reported consuming alcohol on 26.4 ($\pm$ 6.6) days on average; 127 (67.2%) reported daily alcohol use. Mean drinks / drinking day was 17.6 ($\pm$ 10.0; Median = 15.0), and the mean frequency of binge drinking was 20.5 ($\pm$ 11.1) days. One hundred fifty (79.4%) were current cigarette smokers, 68 (36.0%), 35 (18.5%), and 46 (24.3%) participants reported using marijuana, cocaine, and benzodiazepines, respectively, in the past month.

Compared to males, females had significantly higher mean educational attainment (Table 1) and there was a trend suggesting they were more likely (78.4% v 65.0%, $p = .077$) to say they had seen a primary care physician in the last year, but none of the other background characteristics described in Table 1 differed significantly by gender.

In Table 2, life domains are sorted from most to least concern based on means. In this cohort, alcohol problems were of most concern ($M = 2.86$), followed by mental health ($M = 1.65$), cigarette smoking ($M = 1.59$), money problems ($M = 1.7$), and relationship problems ($M = 1.48$). All other domains had mean concern ratings < 1.0 on a scale ranging from 0 to 3. Concern about community safety and transmitted disease were the domains of least concern in this cohort.

Males expressed significantly ($p = .003$) more concern about money problems than females (Table 2). Males also had significantly ($p = .005$) higher mean concern about drug problems, serious physical illness ($p = .007$), health insurance ($p = .006$), transmitted disease ($p = .05$), and community safety ($p = .03$). Males and females did not differ significantly with respect to mean concern about alcohol problems, mental health, cigarette smoking, or relationship problems (Table 2).

4.0 DISCUSSION

We have described ten life domains that concern alcohol dependent persons at the time they enter an inpatient detoxification program. Beyond the continued use of alcohol, study participants reported, being most concerned about mental health issues, cigarette use, and financial and relationship problems. In this cohort with alcohol use disorder, men reported greater concern than women in over half the domains assessed.

This is the first study classifying life concerns of persons with AUDs. Unlike alcohol severity measures which evaluate drinking consequences, or quality of life measures which focus on physical and mental health function and limitations, the current life concerns measure assesses an individual's perceptions of ongoing or impending life difficulties. Life concerns are a multidimensional construct that captures subjective worries meaningful to

this population and which may influence or interact with alcohol use behaviors. Life concerns could also affect completion of detoxification as well as decision-making about and engagement with aftercare plans. Notably, men and women had similar demographic characteristics and recent drinking profiles in this sample, suggesting that these factors did not contribute to gender differences in life concerns. Indeed, given the literature suggesting that similar quantities of alcohol tend to have more deleterious effects on women (certainly physical effects (National Institute on Alcohol Abuse and Alcoholism, 2000b)), it was surprising that men reported higher concerns across domains, including physical health. Men and women reported similarly high level of concern about their alcohol use, as expected.

AUD patients have often reported high levels of depressive symptoms at the onset of treatment (Foulds, et al., 2015; Pettinati, et al., 2013), and mental health is a featured concern of our study participants as well. Our findings are compatible with the National Comorbidity Survey, which reported that individuals with lifetime alcohol use disorder have a high probability of being diagnosed with another co-occurring psychiatric disorder (Kessler, et al., 1997). The National Epidemiological Survey on Alcohol and Related Conditions also reported the presence of mood and anxiety disorders, as well as other psychiatric diagnoses, in a national sample of persons with AUD (Glass, Williams, & Bucholz, 2014). Women with AUD report greater severity of childhood trauma, but similar lifetime exposure to traumatic events compared with men (Johnson, Heffner, Blom, & Anthenelli, 2010).

Nearly four in five of our participants were current smokers, and smoking was one of the highest rated life concerns reported. Approximately 6.2 million American adults have both an AUD and nicotine dependence (Falk, Yi, & Hiller-Sturmhofel, 2006). People with AUD are three times more likely to be smokers than those in the general population (Grant, Hasin, Chou, Stinson, & Dawson, 2004). With smoking rates far exceeding the general population, smokers with alcohol use disorder experience high rates of tobacco-related health consequences (Marrero, et al., 2005; Mukamal, 2006; Negri, La Vecchia, Franceschi, & Tavani, 1993; Pelucchi, Gallus, Garavello, Bosetti, & La Vecchia, 2006). Given that cigarette smoking was identified as one of the more concerning life domains, this patient population may be receptive to cessation interventions that begin during this brief supervised period. In a meta-analysis of smoking cessation interventions, Prochaska, et al., (Prochaska, Delucchi, & Hall, 2004) found that smoking cessation interventions during addiction treatments were correlated with an improved likelihood of long-term tobacco abstinence.

Economic concerns, such as not having a steady income and not having enough money for basic needs, would be expected given the high rates of unemployment reported by study participants: 76 % compared to a state average of 5.5–7.1% for the same time period (Labor and Workforce Development, 2014). Problematic substance use, including alcohol use disorder, is consistently more prevalent among the unemployed (Compton, Gfroerer, Conway, & Finger, 2014; Mullahy & Sindelar, 1996). Although it is unclear whether those with AUD are more likely to become unemployed, or whether those who are unemployed are more likely to develop alcohol use disorder, emerging evidence supports the latter (Mossakowski, 2008; Popovici & French, 2013). Alcohol is a common mechanism used for

- Carey MP, Braaten LS, Jaworski BC, Durant LE, Forsyth AD. HIV and AIDS relative to other health, social, and relationship concerns among low-income urban women: a brief report. *Journal of Women's Health and Gender-Based Medicine*. 1999; 8(5):657–661.
- Carey MP, Senn TE. Health and life concerns among patients attending a publicly funded sexually transmitted infection clinic. *Journal of Public Health Management and Practice*. 2013; 19(1):E30–34. doi: 10.1097/PHH.0b013e31822d4bd5.
- Chartier KG, Hesselbrock MN, Hesselbrock VM. Ethnicity and gender comparisons of health consequences in adults with alcohol dependence. *Substance Use & Misuse*. 2013; 48(3):200–210. doi: 10.3109/10826084.2013.747743. [PubMed: 23302062]
- Claxton SE, DeLuca HK, van Dulmen MH. The association between alcohol use and engagement in casual sexual relationships and experiences: a meta-analytic review of non-experimental studies. *Archives of Sexual Behavior*. 2015; 44(4):837–856. doi: 10.1007/s10508-014-0392-1.
- Compton WM, Gfroerer J, Conway KP, Finger MS. Unemployment and substance outcomes in the United States 2002–2010. *Drug and Alcohol Dependence*. 2014; 142:350–353. doi: 10.1016/j.drugalcdep.2014.06.012. [PubMed: 25042761]
- Conger RD, Lorenz FO, Elder GH Jr, Simons RL, Ge X. Husband and wife differences in response to undesirable life events. *Journal of Health and Social Behavior*. 1993; 34(1):71–88. [PubMed: 8463637]
- Daepfen JB, Krieg MA, Burnand B, ersin B. MOS-SF-36 in evaluating health-related quality of life in alcohol-dependent patients. *American Journal of Drug and Alcohol Abuse*. 1998; 24(4):685–694. [PubMed: 9849778]
- Dawson DA, Grant BF, Ruan WJ. The association between stress and drinking: modifying effects of gender and vulnerability. *Alcohol and Alcoholism*. 2005; 40(5):453–460. doi: 10.1093/alcal/agh176. [PubMed: 15972275]
- Falk DE, Yi HY, Hiller-Sturmhofel S. An epidemiologic analysis of co-occurring alcohol and tobacco use and disorders: findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *Alcohol Research & Health*. 2006; 29(3):162–171. [PubMed: 17373404]
- Foster JH, Powell JE, Marshall EJ, Peters TJ. Quality of life in alcohol-dependent subjects--a review. *Quality of Life Research*. 1999; 8(3):255–261. [PubMed: 10472156]
- Foulds JA, Douglas Sellman J, Adamson SJ, Boden JM, Mulder RT, Joyce PR. Depression outcome in alcohol dependent patients: an evaluation of the role of independent and substance-induced depression and other predictors. *Journal of Affective Disorders*. 2015; 174:503–510. doi: 10.1016/j.jad.2014.11.052. [PubMed: 25554995]
- Frone MR, Cooper LM, Russell M. Stressful life events, gender, and substance use: An application of tobit regression. *Psychology of Addictive Behaviors*. 1994; 8(2):59–69.
- Glass JE, Williams EC, Bucholz KK. Psychiatric comorbidity and perceived alcohol stigma in a nationally representative sample of individuals with DSM-5 alcohol use disorder. *Alcoholism, Clinical Experimental Research*. 2014; 38(6):1697–1705. doi: 10.1111/acer.12422.
- Goldstein RB, Dawson DA, Chou SP, Grant BF. Sex differences in prevalence and comorbidity of alcohol and drug use disorders: results from wave 2 of the National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Studies on Alcohol and Drugs*. 2012; 73(6):938–950. [PubMed: 23036212]
- Grant BF, Hasin DS, Chou SP, Stinson FS, Dawson DA. Nicotine dependence and psychiatric disorders in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Archives of General Psychiatry*. 2004; 61(11):1107–1115. doi: 10.1001/archpsyc.61.11.1107. [PubMed: 15520358]
- Green CA, Polen MR, Lynch FL, Dickinson DM, Bennett MD. Gender differences in outcomes in an HMO-based substance abuse treatment program. *Journal of Addictive Diseases*. 2004; 23(2):47–70. doi: 10.1300/J069v23n02_04. [PubMed: 15132342]
- Greenfield SF, Brooks AJ, Gordon SM, Green CA, Kropp F, McHugh RK, et al. Substance abuse treatment entry, retention, and outcome in women: a review of the literature. *Drug and Alcohol Dependence*. 2007; 86(1):1–21. doi: 10.1016/j.drugalcdep.2006.05.012. [PubMed: 16759822]

- Johnson CS, Heffner JL, Blom TJ, Anthenelli RM. Exposure to traumatic events among treatment-seeking, alcohol-dependent women and men without PTSD. *Journal of Traumatic Stress*. 2010; 23(5):649–652. doi: 10.1002/jts.20563. [PubMed: 20803578]
- Kessler RC, Crum RM, Warner LA, Nelson CB, Schulenberg J, Anthony JC. Lifetime co-occurrence of DSM-III-R alcohol abuse and dependence with other psychiatric disorders in the National Comorbidity Survey. *Archives of General Psychiatry*. 1997; 54(4):313–321. [PubMed: 9107147]
- Keyes KM, Grant BF, Hasin DS. Evidence for a closing gender gap in alcohol use, abuse, and dependence in the United States population. *Drug and Alcohol Dependence*. 2008; 93(1–2):21–29. doi: 10.1016/j.drugalcdep.2007.08.017. [PubMed: 17980512]
- Kosten TR, O'Connor PG. Management of drug and alcohol withdrawal. *New England Journal of Medicine*. 2003; 348(18):1786–1795. doi: 10.1056/NEJMra020617. [PubMed: 12724485]
- Labor and Workforce Development. Labor Force and Unemployment Data. 2014. http://lmi2.detma.org/lmi/lmi_lur_a.asp
- Levin JS, Taylor RJ, Chatters LM. Race and gender differences in religiosity among older adults: findings from four national surveys. *Journal of Gerontology*. 1994; 49(3):S137–145. [PubMed: 8169348]
- Liu XC, Keyes KM, Li G. Work stress and alcohol consumption among adolescents: moderation by family and peer influences. *BMC Public Health*. 2014; 14:1303. doi: 10.1186/1471-2458-14-1303. [PubMed: 25523951]
- Mann K, Ackermann K, Croissant B, Mundt G, Nakovics H, Diehl A. Neuroimaging of gender differences in alcohol dependence: are women more vulnerable? *Alcoholism, Clinical and Experimental Research*. 2005; 29(5):896–901.
- Marrero JA, Fontana RJ, Fu S, Conjeevaram HS, Su GL, Lok AS. Alcohol, tobacco and obesity are synergistic risk factors for hepatocellular carcinoma. *Journal Hepatology*. 2005; 42(2):218–224. doi: 10.1016/j.jhep.2004.10.005. [PubMed: 15962294]
- Martin RA, Ellingsen VJ, Tzilos GK, Rohsenow DJ. General and religious coping predict drinking outcomes for alcohol dependent adults in treatment. *The American Journal on Addictions*. 2015; 24(3):240–245. doi: 10.1111/ajad.12181. [PubMed: 25662479]
- Mauro PM, Canham SL, Martins SS, Spira AP. Substance-use coping and self-rated health among US middle-aged and older adults. *Addictive Behaviors*. 2015; 42:96–100. doi: 10.1016/j.addbeh.2014.10.031. [PubMed: 25437264]
- Merkx MJ, Schippers GM, Koeter MW, De Wildt WA, Vedel E, Goudriaan AE, et al. Treatment outcome of alcohol use disorder outpatients with or without medically assisted detoxification. *Journal of Studies on Alcohol and Drugs*. 2014; 75(6):993–998. [PubMed: 25343657]
- Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004; 291(10):1238–1245. doi: 10.1001/jama.291.10.1238. [PubMed: 15010446]
- Morgan MY, Landron F, Leher P, New European Alcoholism Treatment Study, G. Improvement in quality of life after treatment for alcohol dependence with acamprosate and psychosocial support. *Alcoholism, Clinical and Experimental Research*. 2004; 28(1):64–77. doi: 10.1097/01.ALC.0000108652.73143.4B.
- Mossakowski KN. Is the duration of poverty and unemployment a risk factor for heavy drinking? *Social Science & Medicine*. 2008; 67(6):947–955. doi: 10.1016/j.socscimed.2008.05.019. [PubMed: 18573582]
- Mukamal KJ. The effects of smoking and drinking on cardiovascular disease and risk factors. *Alcohol Research & Health*. 2006; 29(3):199–202. [PubMed: 17373409]
- Mulia N, Schmidt L, Bond J, Jacobs L, Korcha R. Stress, social support and problem drinking among women in poverty. *Addiction*. 2008; 103(8):1283–1293. doi: 10.1111/j.1360-0443.2008.02234.x. [PubMed: 18855817]
- Mullahy J, Sindelar J. Employment, unemployment, and problem drinking. *Journal of Health Economics*. 1996; 15(4):409–434. [PubMed: 10164037]
- National Institute on Alcohol Abuse and Alcoholism. Health risks and benefits of alcohol consumption. *Alcohol Research & Health*. 2000a; 24(1):5–11. [PubMed: 11199274]
- National Institute on Alcohol Abuse and Alcoholism. Medical consequences of alcohol abuse. *Alcohol Research & Health*. 2000b; 24(1):27–31. [PubMed: 11199271]

- National Institute on Alcohol Abuse and Alcoholism. Helping Patients Who Drink Too Much: A Clinician's Guide. NIH Publication No. 07-3769. National Institutes of Health; Rockville, MD: 2005.
- Navarro HJ, Doran CM, Shakeshaft AP. Measuring costs of alcohol harm to others: a review of the literature. *Drug and Alcohol Dependence*. 2011; 114(2-3):87-99. doi: 10.1016/j.drugalcdep.2010.11.009. [PubMed: 21193271]
- Negri E, La Vecchia C, Franceschi S, Tavani A. Attributable risk for oral cancer in northern Italy. *Cancer Epidemiology, Biomarkers & Prevention*. 1993; 2(3):189-193.
- Nolen-Hoeksema S, Hilt L. Possible contributors to the gender differences in alcohol use and problems. *The Journal of General Psychology*. 2006; 133(4):357-374. doi: 10.3200/GENP.133.4.357-374. [PubMed: 17128956]
- O'Meara C, Witherspoon R, Hapangama N, Hyam DM. Mandible fracture severity may be increased by alcohol and interpersonal violence. *Australian Dental Journal*. 2011; 56(2):166-170. doi: 10.1111/j.1834-7819.2011.01319.x. [PubMed: 21623808]
- Pelucchi C, Gallus S, Garavello W, Bosetti C, La Vecchia C. Cancer risk associated with alcohol and tobacco use: focus on upper aero-digestive tract and liver. *Alcohol Research & Health*. 2006; 29(3):193-198. [PubMed: 17373408]
- Perreira KM, Sloan FA. Life events and alcohol consumption among mature adults: a longitudinal analysis. *Journal of Studies on Alcohol*. 2001; 62(4):501-508. [PubMed: 11513228]
- Peters TJ, Millward LM, Foster J. Quality of life in alcohol misuse: comparison of men and women. *Archives of Women's Mental Health*. 2003; 6(4):239-243. doi: 10.1007/s00737-003-0012-x.
- Pettinati HM, O'Brien CP, Dundon WD. Current status of co-occurring mood and substance use disorders: a new therapeutic target. *The American Journal of Psychiatry*. 2013; 170(1):23-30. doi: 10.1176/appi.ajp.2012.12010112. [PubMed: 23223834]
- Pettinati HM, Pierce JD Jr, Wolf AL, Rukstalis MR, O'Brien CP. Gender differences in comorbidly depressed alcohol-dependent outpatients. *Alcoholism, Clinical and Experimental Research*. 1997; 21(9):1742-1746.
- Polen MR, Green CA, Perrin NA, Anderson BM, Weisner CM. Drinking Patterns, Gender and Health I: Attitudes and Health Practices. *Addiction Research & Theory*. 2010; 18(2):122-142. [PubMed: 23946720]
- Popovici I, French MT. Does Unemployment Lead to Greater Alcohol Consumption? *Industrial Relations*. 2013; 52(2):444-466. doi: 10.1111/irel.12019. [PubMed: 23543880]
- Prochaska JJ, Delucchi K, Hall SM. A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery. *Journal of Consulting and Clinical Psychology*. 2004; 72(6):1144-1156. doi: 10.1037/0022-006X.72.6.1144. [PubMed: 15612860]
- Rehm J. The risks associated with alcohol use and alcoholism. *Alcohol Research & Health*. 2011; 34(2):135-143. doi: Fea-AR&H-65 Fea-AR&H-65. [PubMed: 22330211]
- Reyes CD, Pagano ME, Ronis RJ. The Impact of Stressful Life Events on Alcohol Relapse: Findings from the Collaborative Longitudinal Personality Disorders Study. *Journal of Dual Diagnosis*. 2009; 5(2):226-232. doi: 10.1080/15504260902886505. [PubMed: 21170176]
- Stein MD, Anderson BJ, Thurmond P, Bailey GL. Comparing the life concerns of prescription opioid and heroin users. *Journal of Substance Abuse Treatment*. 2015; 48(1):43-48. doi: 10.1016/j.jsat.2014.07.001. [PubMed: 25171955]
- Stewart, SH.; Gavric, D.; Collins, P.; Brady, KT.; Back, SE.; Greenfield, SF. Women and Addiction: A Comprehensive Handbook. Guilford Press; New York, NY: 2009. Woman, girls, and alcohol; p. 341-359.
- Substance Abuse and Mental Health Services Administration. Results from the 2013 National Survey on Drug Use and Health: Summary of National Findings, NSDUH Series H-48, HHS Publication No. (SMA) 14-4863. Substance Abuse and Mental Health Services Administration; Rockville, MD: 2014.
- Veenstra MY, Lemmens PH, Friesema IH, Garretsen HF, Knottnerus JA, Zwietering PJ. A literature overview of the relationship between life-events and alcohol use in the general population. *Alcohol and Alcoholism*. 2006; 41(4):455-463. doi: 10.1093/alcalc/agl023. [PubMed: 16627624]

- Walitzer KS, Dearing RL. Gender differences in alcohol and substance use relapse. *Clinical Psychology Review*. 2006; 26(2):128–148. doi: 10.1016/j.cpr.2005.11.003. [PubMed: 16412541]
- Whiteford HA, Degenhardt L, Rehm J, Baxter AJ, Ferrari AJ, Erskine HE, et al. Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *Lancet*. 2013; 382(9904):1575–1586. doi: 10.1016/S0140-6736(13)61611-6. [PubMed: 23993280]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

HIGHLIGHTS

- Mental health, smoking, and finances are the most prevalent life concerns
- Gender affects the life concerns of persons with alcohol use disorders
- Men more concerned than women about money and drug use problems
- Men reported greater concern than women in most domains assessed

Table 1

Background Characteristics by Gender (n = 189).

	Cohort (n = 189)	Female (n =51)	Male (n = 138)	t or χ^2 (p =)
Age	43.5 (± 9.9)	43.2 (± 9.4)	43.6 (± 10.1)	−0.31 (.760)
Non-Latino Caucasian	163 (86.7%)	47 (92.2%)	116 (84.1%)	2.18 (.140)
Education	12.4 (± 2.1)	13.1 (± 2.0)	12.1 (± 2.1)	3.09 (.002)
Employed (Part- or Full-Time)	46 (24.3%)	14 (27.5%)	32 (23.2%)	0.37 (.544)
Homeless (Yes)	20 (10.6%)	5 (9.8%)	15 (10.9%)	0.04 (.833)
Seen PCP Last Year (Yes)	129 (68.6%)	40 (78.4%)	89 (65.0%)	3.13 (.077)
Detox History				
No Prior Detox	52 (27.7%)	17 (33.3%)	35 (25.6%)	
Recent (Last Year)	82 (43.6%)	22 (43.1%)	60 (43.8%)	1.48 (.478)
Less Recent (> 1 Year)	54 (28.7%)	12 (25.5%)	42 (28.7%)	
Days Used Alcohol (0–30)	26.4 (± 6.6)	25.7 (± 6.9)	26.6 (± 6.6)	−0.83 (.406)
Drinks / Drinking Day	17.6 (± 9.9)	17.1 (± 1.4)	17.8 (± 10.1)	−0.42 (.675)
Days Binge Alcohol Use (0–30)	20.5 (± 11.1)	21.0 (± 10.3)	20.4 (± 11.5)	0.35 (.725)
Current Cig. Smoker (Yes)	150 (79.4%)	40 (78.4%)	110 (79.7%)	0.04 (.847)
Recent Marijuana Use (Yes)	68 (36.0%)	18 (35.3%)	50 (36.2%)	0.01 (.905)
Recent Cocaine Use (Yes)	35 (18.5%)	10 (19.6%)	25 (18.1%)	0.05 (.815)
Recent Benzodiazepine Use (Yes)	46 (24.3%)	13 (25.5%)	33 (23.9%)	0.05 (.823)

Table 2Mean ($\pm$ SD) Life Concerns by Gender.

Life Concern	n % Reporting No Concern (n = 189)	Mean ($\pm$ SD)			z (p =) ^a
		Cohort (n = 189)	Female (n = 51)	Male (n = 138)	
Alcohol Problems (NA)	3 (1.6%)	2.86 ($\pm$ 0.49)	2.84 ($\pm$ 0.54)	2.86 ($\pm$ 0.47)	0.34 (.973)
Mental Health (.79) ^b	4 (2.1%)	1.65 ($\pm$ 0.72)	1.68 ($\pm$ 0.79)	1.64 ($\pm$ 0.70)	0.52 (.605)
Cigarette Smoking (NA)	61 (32.3%)	1.59 ($\pm$ 1.27)	1.35 ($\pm$ 1.35)	1.67 ($\pm$ 1.23)	-1.48 (.140)
Money Problems (.82) ^b	29 (15.3%)	1.57 ($\pm$ 1.03)	1.20 ($\pm$ 1.05)	1.71 ($\pm$ 0.99)	-3.01 (.003)
Rel. Problems (.85) ^b	25 (13.2%)	1.48 ($\pm$ 1.00)	1.41 ($\pm$ 0.98)	1.51 ($\pm$ 1.01)	-0.61 (.542)
Drug Problems (NA)	112 (59.3%)	0.97 ($\pm$ 1.29)	0.57 ($\pm$ 1.12)	1.12 ($\pm$ 1.32)	-2.76 (.005)
Serious Phys. Illness (.76) ^b	55 (29.1%)	0.70 ($\pm$ 0.76)	0.43 ($\pm$ 0.47)	0.80 ($\pm$ 0.82)	-2.72 (.007)
Health Insurance (.92) ^b	126 (66.7%)	0.68 ($\pm$ 1.10)	0.34 ($\pm$ 0.85)	0.81 ($\pm$ 1.16)	-2.77 (.006)
Transmitted Disease (.87) ^b	139 (73.5%)	0.39 ($\pm$ 0.81)	0.20 ($\pm$ 0.55)	0.46 ($\pm$ 0.87)	-1.98 (.050)
Community Safety (.82) ^b	130 (68.8%)	0.33 ($\pm$ 0.66)	0.17 ($\pm$ 0.42)	0.39 ($\pm$ 0.42)	-2.18 (.030)

^aNonparametric Wilcoxon Rank-Sum z-statistic testing the equality of rank-ordered distributions.^bChronbach's α .