Published in final edited form as:

Obes Rev. 2015 February ; 16(Suppl 1): 77–90. doi:10.1111/obr.12258.

Appetite and body weight regulation after bariatric surgery

Heike Münzberg, Amanda Laque, Sangho Yu, Kavon Rezai-Zadeh, and Hans-Rudolf Berthoud

Pennington Biomedical Research Center, Louisiana State University, Baton Rouge, LA 70808, USA

Abstract

Bariatric surgery continues to be remarkably efficient in treating obesity and T2DM and a debate has started whether it should remain the last resort only or also be used for the prevention of metabolic diseases. Intense research efforts in humans and rodent models are underway to identify the critical mechanisms underlying the beneficial effects with a view towards non-surgical treatment options. This non-systematic review summarizes and interprets some of this literature, with an emphasis on changes in the controls of appetite. Contrary to earlier views, surgery-induced reduction of energy intake and subsequent weight loss appear to be the main drivers for rapid improvements of glycemic control. The mechanisms responsible for suppression of appetite, particularly in the face of the large weight loss, are not well understood. Although a number of changes in food choice, taste functions, hedonic evaluation, motivation, and self-control have been documented in both humans and rodents after surgery, their importance and relative contribution to diminished appetite has not yet been demonstrated. Furthermore, none of the major candidate mechanisms postulated in mediating surgery induced changes from the gut and other organs to the brain, such as gut hormones and sensory neuronal pathways, have been confirmed yet. Future research efforts should focus on interventional rather than descriptive approaches in both humans and rodent models.

Keywords

Obesity; brain; gut hormones; gastric bypass; sleeve gastrectomy; food choice; food reward; executive control; taste responsiveness; aversive conditioning

Introduction

It is now generally acknowledged that bariatric surgery is the most effective intervention for the treatment of common obesity and its associated co-morbidities. Weight loss is larger and more sustained compared with intensive medical treatment, including drugs and lifestyle changes. Although still too early to tell, bariatric surgeries also seem to achieve these beneficial effects without serious side effects and increased mortality. To the contrary, the longest running study, the Swedish Obese Subjects Study (SOS) found decreased mortality and decreased risk for certain cancers in bariatric surgery patients. The fact that about half of

the 16,000 total articles on bariatric surgery (PubMed) were published in just the last 5 years clearly illustrates the rush to identify the biological mechanisms underlying this remarkably efficient obesity treatment. However, despite these intense research efforts in humans and rodent models, we are far from understanding the underlying mechanisms. Early hypotheses such as changes in gut hormone secretion have not yet been confirmed, and new candidate mechanisms are being added to the list. This non-systematic review highlights potential mechanisms contributing to the sustained change in energy balance regulation that allows bariatric surgery patients and rodents to remain at greatly reduced body weight levels.

The role of increased energy expenditure

It is important to distinguish at least two phases of body weight regulation after bariatric surgery, an initial phase of rapid weight loss and a subsequent phase of weight stability or weight regain. Few clinical and rodent studies have measured energy expenditure during the rapid weight loss phase. The weight loss nadir after Roux-en-Y Gastric Bypass (RYGB) is typically reached by 6-12 months in humans and 2-3 weeks in rodents. In most of the few human studies with energy expenditure (EE) measurements at 1-3 months after surgery, before reaching the weight loss nadir, resting energy expenditure (REE) was consistently decreased by 14-24% compared to before surgery¹⁻³ (Table 1). Only one study reported that REE was not significantly decreased ($\sim -2\%$) at 6 weeks after RYGB⁴. Unfortunately, there are no RYGB rodent studies which have measured REE during the 2-3 weeks of rapid weight loss. Thus, the limited clinical data available suggest that during the rapid weight loss phase, reduced EE is more or less commensurate with weight loss. Because weight loss induced by calorie restriction leads to an adaptive fall in EE⁵, it is however possible that this fall is blunted by gastric bypass surgery.

Many more studies measured EE during the phase of relative weight stability that follows the rapid weight loss phase after RYGB and other bariatric surgeries (Table 1). In a recent comprehensive review of this literature, Thivel et al. concluded that, at least in humans, total EE and REE are decreased after surgery compared with pre-surgical levels, commensurate with the decrease in both fat mass and fat free mass⁶. Furthermore, these authors concluded that: "it is not the nature of the bariatric surgery but rather factors such as energy balance status (weight loss, stability, or regain) or body composition that impact the postoperative change in REE" (p. 257). One of the most comprehensive and longitudinal studies measuring energy expenditure before and at several time points after RYGB surgery also concluded that "REE changes were predicted by loss of body tissue; thus, there was no significant long-term change in energy efficiency that would independently promote weight regain"⁷.

This conclusion is in stark contrast to at least some rodent studies that claimed significantly increased EE after RYGB but not sleeve gastrectomy in rats^{8,9} and mice^{10,11}. However, much of the difference may be explained by how EE is expressed and to which reference it is compared. Most human studies express EE uncorrected for body weight, while most rodent studies express it per either total body weight or a fractional power function of body weight such as kcal/kg^{0.75}. For example, if EE is 10% lower per animal after RYGB and they weigh 25% less compared with sham surgery, correcting for body weight would result

in 17% higher EE, and even if using the 0.75 power for body weight, would result in 8% higher EE. While most human studies compare EE after surgery to pre-surgical levels, post-surgical EE in rodent models is always compared to sham-operated or weight-matched animals, not to pre-surgical levels.

In some rodent studies, evidence for increased EE after RYGB is more convincing. First, pair-feeding sham-operated rats with the same amount of food eaten by surgical rats resulted in higher body weight of sham-operated animals. In fact, sham-operated rats had to eat less than surgical rats to maintain the same weight, implicating either increased EE and/or fecal energy loss after RYGB. Direct measurements revealed increased EE and only negligible malabsorption^{8,9}. Second, a more systematic assessment of energy balance by measuring energy intake, fecal energy loss, and the metabolic costs of fat and lean tissue in mice demonstrated higher EE in mice after RYGB¹¹. Together, these findings suggest that there might be important species differences in EE regulation after bariatric surgery. One caveat to consider is that pair-feeding or weight-matching experiments disrupt natural feeding patterns and that restricted animals typically eat their daily food ration within a short period of time often during the light cycle. This “un-natural” feeding pattern could have profound influences on energy fluxes and the thermic effect of food that is different from eating smaller amounts of food but throughout the day¹².

The most unbiased way of analyzing EE data is multiple linear regression, relating EE to age, fat free mass and fat mass, three variables most strongly related to EE^{13,14}. However, most EE data are collected by indirect calorimetry which calculates EE values using an equation established in healthy animals and based on the assumption that substrate interconversion is negligible¹³. There is no guarantee that this equation would work the same way for both morbidly obese, often diabetic patients and patients that have undergone bariatric surgery.

In summary, there is very limited evidence for increased EE as an important factor in weight loss and maintenance after bariatric surgeries in humans. There may be an important species difference, with rodents, particularly mice, possibly using increased EE as a strategy to maintain lower body weight after surgery. It will be important in future studies to report EE data always together with detailed body composition data, and to run pair-fed and/or weight matched control groups with normal diurnal intake patterns, because only such carefully controlled studies are able to isolate the net effects of surgery.

Mechanisms of increased energy expenditure

Several potential mechanisms for increasing EE after bariatric surgeries have been suggested, but there is no study that directly tested the role of any of them. Bile acids and fibroblast growth factors have been shown to stimulate EE in mice by acting on brown adipose tissue thermogenesis¹⁵, and circulating levels of bile acids and FGF19 and 21 are increased in both humans¹⁶⁻¹⁸ and rodents¹⁹ after RYGB or sleeve gastrectomy. However, in humans, there are contradictory reports regarding the association of bile acid levels and energy metabolism^{20,21}.

Intestinal hypertrophy after RYGB ^{8, 22-25} and the resulting increased glucose utilization ²⁶ have also been suggested as a possible mechanism for increased EE.

The role of reduced energy intake

Weight loss phase

There is no doubt that decreased energy intake is the main driver for the initial weight loss phase after all types of bariatric surgeries and in both humans and rodents, even though energy intake studies in humans are complicated by pre- and post-surgical behavioral counselling and by limits in caloric intake and macronutrient composition during the immediate post-surgical period. Thus, clean separation of behavioral modification from biological needs is difficult. Bariatric surgery patients receive a liquid low calorie diet for the first post-operative week and are instructed to eat low-fat food with low energy density. Also, food intake is usually measured by non-validated self-reporting which notoriously underestimates the actual food intake. Furthermore, sham surgeries are not feasible in humans. Rodent (and other) models do not have these limitations and are thus indispensable in the study of natural ingestive behavior after surgery.

Given these limitations in human studies, data from mainly RYGB patients demonstrate very large reductions in energy intake (compared to pre-surgical levels) of about – 55% (– 47 to – 66%) at 6 months and about – 40% (– 19 to – 49 %) at 2 - 3 years ^{4, 27-30}. Energy intake for the earlier post-surgical period may be even lower with around – 70% to – 80%, ^{4, 31} and reduced energy intake appears to last almost indefinitely, with a reported reduction of –21% at 10 years across all types of surgery ²⁷. Even though energy intake is not corrected for the substantial weight loss, it appears to be the major contributor to weight loss and prevention of weight regain.

Fecal energy loss due to malabsorption can additionally reduce metabolizable energy. Although a number of rodent studies report no significant fecal energy loss, a majority of human and a few rodent studies do find significant malabsorption particularly of fat. In one human study, the efficiency of fat absorption significantly decreased from 92% before surgery to 72% at 5 months and 68% at 14 months after RYGB, translating into losses of 124 and 172 kcal/day ³². Based on parallel measurements of energy intake, the study concluded that malabsorption accounted for about 6% at 5 months and 11% at 14 months, of the total reduction of metabolizable energy intake ³². Thus, while not negligible, the contribution of malabsorption to the total energy balance is relatively small compared to the reduction in energy intake.

Weight maintenance/regain phase

Existing long term data on energy intake after RYGB and other bariatric surgeries suggest fundamental differences between humans and rodents, particularly mice. While human studies show continued suppression of energy intake at 2 years and later after RYGB ^{4, 27-29, 33}, food intake suppression in rats and mice typically lasts only for about 2-4 weeks after surgery ^{9, 34-36}. The suppression of food intake in various rat models of RYGB is highly variable, with no changes or even slight increases in food intake even at early time points ³⁷, to lasting, although moderate, suppression for up to 10 months ³⁸. In the few

viable mouse models described, initial suppression of food intake is even shorter than in rats and during the weight maintenance phase it is typically not changed or slightly higher than in sham-operated controls ^{10, 11, 36, 39}.

In summary, energy intake and metabolizable energy are drastically reduced at least initially in both humans and rodents and are mainly responsible for the weight loss phase. It is not clear whether the more substantial and sustained reduction of food intake in humans is due to the same surgery-induced physiological factors responsible for the early suppression in rodents or the effects of the rigorous behavioral coaching before and after surgery. However, it is becoming clear that this drastic reduction in energy intake and its consequences on body weight are mainly responsible for the rapid improvement of glycemic control and resolution of diabetes. Therefore, the acute hypocaloric state during the initial weight loss phase is an important research target to understand its underlying mechanisms. Feeding the same low caloric diet ingested by RYGB patients to non-surgical control subjects results in rapid body weight loss. In one recent study, RYGB patients and non-surgical control obese subjects provided with 500 kcal/day of a liquid diet with a macronutrient content similar to that consumed by patients after RYGB for the first 21 days after surgery lost 8.1 and 7.2 %, respectively, of their pre-surgical body weight, with similar significant improvements of glycemic control ⁴⁰. In another study, RYGB surgery patients were subjected to the same very low calorie (~1700 kcal/7 days) dietary regimen for one week before surgery and after surgery with a washout period in between. Surprisingly, weight loss with the diet alone was significantly larger than with diet plus surgery (5.1 vs. 2.9 %) and again, there were similar significant improvements in glycemic control ⁴¹. Together with other short-term controlled studies ^{42, 43}, these findings strongly suggest that reduced energy intake after RYGB leads to a profound hypocaloric state followed by rapid weight loss that fully explains the rapid improvements in glycemic control.

If the initial reduction of food intake and weight loss is one key effect of bariatric surgery, maintenance and defense of this reduced body weight level is the other one. This is in stark contrast to caloric restriction-induced weight loss, which is followed by hyperphagia and prompt weight regain, even if pre-intervention weight was in the obese range. Successful bariatric surgery and particularly RYGB appears to neutralize the powerful counter-regulatory mechanisms that are engaged by weight loss. The major counter-regulatory response to weight loss, increased hunger, seems to be offset or defused after RYGB. Why do RYGB patients not return to pre-surgical levels of food intake to regain preoperative body weight? Why do rodents not become hyperphagic to regain pre-operative body weight?

Mechanisms of reduced food intake

Ingestive behavior is ultimately controlled by the brain, but this says little about the information used by the brain for making the decision to eat or not to eat. This information can be derived from internal as well as external signals, and signal processing can take place outside or inside awareness. For example, low leptin levels and previously rewarded food cues from the environment can both induce strong feelings of hunger and initiate ingestion. The strong hunger drive induced by fasting and starvation occurs mainly in the absence of awareness and at least partially bypasses human executive control, while much of the

regular daily food intake initiation and meal size is largely under executive control. The former system is also known as homeostatic regulator of energy balance with key components in the hypothalamus-brain stem axis. This is the system that defends an optimal level of body weight/adiposity for a given individual and environment by engaging hormonal and neural feedback mechanisms controlling the major effectors of energy intake and expenditure ⁴⁴. The latter is often referred to as non-homeostatic or hedonic system with key components in the limbic system and cortex. Importantly, both systems are highly interactive so that internal signals can modulate hedonic systems in a bottom-up fashion and hedonic as well as cognitive processing can override homeostatic functions in a top-down fashion ⁴⁵. It is within this framework that we discuss the potential mechanisms by which bariatric surgery suppresses food intake and reduces body weight.

Does bariatric surgery change the homeostatically defended body weight level?

Defense of a set point is indicated behaviorally when perturbed body weight (either downwards or upwards) promptly returns to its pre-perturbation level. There is now considerable evidence that rodents after bariatric surgery also defend their 'new' body weight this way. If rats with sleeve gastrectomy are exposed to additional exogenous food restriction to further lower their body weight, they rapidly return to their original body weight when unlimited food access resumes ⁴⁶. Similarly, elevated food intake and body weight in RYGB rats, achieved with blockade of central melanocortin-3/4 receptor signaling, promptly return to their original (low) body weight when the blockade is removed (Manuscript under review). Furthermore, female rats with sleeve gastrectomy increase body weight when pregnant and return to pre-pregnancy body weight after delivery ⁴⁷. In all these cases, rats are able to easily double food intake despite their surgical intervention, demonstrating their physical ability to eat more and highlighting that they chose to eat less and defend a lower body weight ⁴⁸. The neural mechanisms of set-point regulation are far from completely understood and there is no convenient measure to demonstrate the neurological correlate of regulation [for more in-depth discussions see ⁴⁹⁻⁵²]. The mechanisms by which RYGB offsets increased hunger is perhaps the most crucial question for translational research ultimately directed towards non-surgical therapies. A recent neuroimaging study, which examined neural responses to visual food stimuli throughout the brain in severely obese and normal weight women as well as in women 4 years after RYGB, may provide an important clue ⁵³. All the significant differences in neural activity, including the diminished response to high calorie food pictures in the hypothalamus between obese and normal weight subjects were normalized in RYGB patients. Importantly, because visual analog ratings of hunger were significantly lower and ratings of satiety were higher in RYGB patients compared to both other groups, these findings could be interpreted as evidence for a changed set-point, although they do not provide any clue as to the specific hypothalamic mechanism involved.

The expression level of the basomedial hypothalamic peptides AGRP/NPY and POMC/CART has often been used as a read-out for the homeostatic regulator, because of the strong anabolic and catabolic effects of manipulating these neurons. Increased AGRP/NPY-expression and/or decreased POMC-expression indicate an energy depleted or "hungry" state as seen after prolonged food deprivation ⁵⁴. Therefore, if this same pattern of gene

expression is observed after surgery-induced weight loss, it would indicate that the subject is metabolically “hungry”, and if the expression levels are unchanged, it would indicate that the subject is ‘satisfied’ with the metabolic state. As recently reviewed⁵⁵, the few studies addressing this question provide conflicting results with changes in both directions^{34, 56-58}. Future studies should measure gene expression at different time points after surgery and after a meal to obtain a more definitive answer.

Under normal healthy conditions, leptin is a master regulator of this hypothalamic ying and yang system. Most importantly, any decrease of circulating leptin or leptin-signaling in the basomedial hypothalamus strongly stimulates AgRP/NPY gene expression and neuronal activity and inhibits POMC/CART gene expression and neuronal activity, a pattern that leads to increased hunger and reduced energy expenditure and guarantees energy sufficiency. However, in obese humans and animals, circulating leptin is dramatically increased in proportion to the amount of body fat, but because of cellular leptin resistance, it is unable to generate a catabolic state that would lead to weight loss. Because calorie restriction-induced weight loss in obese subjects has been demonstrated to re-sensitize leptin action⁵⁹, it is conceivable that bariatric surgery-induced weight loss has a similar effect. However, in a recent direct test of this hypothesis, administration of leptin failed to reduce body weight further in RYGB patients⁶⁰, suggesting that the incomplete reversal of obesity from a BMI of ~48 to ~34 was not enough to restore leptin-sensitivity. A similar conclusion was reached in a study after vertical sleeve gastrectomy in rats³⁴.

Finally, the involvement of an energy balance regulator in the basomedial hypothalamus can also be measured in downstream signaling effects through melanocortin receptors. Similar to the inconsistent evidence for AGRP/NPY and POMC/CART expression levels discussed above, studies in melanocortin-4 receptor (MC4R) knockout mice provide inconsistent results. In RYGB mice, weight loss requires MC4R^{36, 61}, while vertical sleeve gastrectomy (VSG) rats show body weight loss independent of MC4R⁶². Human studies in subjects with heterozygous MC4R mutations showed that gastric banding and sleeve gastrectomy were equally effective in reducing body weight^{62, 63}, while another study showed that gastric banding was less effective⁶⁴. In a patient with complete functional loss of both alleles of the MC4R, adjustable gastric banding did not result in long-term weight loss⁶⁵. One reason for the conflicting results may lie in the redundancy of homeostatic feeding circuitry, which was at least demonstrated for AgRP neurons. AgRP neurons co-express the inhibitory acting neurotransmitter GABA and the orexigenic acting neuropeptide Y (NPY). Deletion of all these components in AgRP neurons showed robust effects on feeding behavior, while monogenic deletions had no effect on feeding behavior, because each of these components can compensate for each other⁶⁶.

In summary, although behavioral evidence shows that after RYGB and sleeve gastrectomy rodents appear to defend a new lower body weight set point, there is not much consistent evidence for an involvement of leptin-sensitive basomedial hypothalamic AGRP/NPY and POMC/CART neurons and their downstream signaling pathways in establishing this new set point. However, because of the crucial role of this circuitry in energy balance, its potential role should be further pursued. The ob/ob and db/db mouse models as well as assessments of

molecular determinants of leptin receptor signaling such as the induction of phospho-STAT3, PTB1B, and SOCS3 in relevant brain areas should be helpful in future studies.

Does bariatric surgery change food hedonics?

The ability to engage in survival promoting behaviors such as ingestive behavior is a fundamental evolutionary mechanism critically guided by the so called brain reward system. A considerable body of evidence has accumulated demonstrating differences in structure and function of the brain reward system between lean and obese humans and rodents [see ^{67, 68, 69} for comprehensive reviews]. An important question in defining the relationship between food reward and obesity is what comes first – are differences in reward functions driving development of obesity or is the obese state impairing reward functions. Bariatric surgery patients are a valuable resource for studying this question.

Bariatric surgery changes food choice—Many human and rodent studies report changes in food preferences and choice and most of the earlier findings were recently reviewed ⁷⁰⁻⁷² (Table 2). The most commonly observed effect in humans is a shift from overly sweet and fatty energy dense foods to less energy dense foods ⁷³. Several rodent studies find a shift from a very high preference for high-fat diets before surgery and in the obese state towards lower preference of such diets after both RYGB and sleeve gastrectomy in longer-term two- or three-choice paradigms ^{35, 38, 74-76}. However, the underlying mechanisms for these changes, both with regard to the specific neural components involved and the mediating signals from the gut or other organs, are not well understood.

Changes in taste sensitivity and preference—There is a growing number of studies investigating whether postoperative changes in taste perception following RYGB may influence eating behavior and by extension weight loss. Taste perception is classically broken into two components, taste acuity (sensitivity) and taste preference or palatability (hedonic value). Clinical and basic researchers alike have ventured to address if indeed RYGB augments one or both of these classic components of taste perception. The earliest work in this area was conducted by Scruggs and colleagues, who examined changes in taste acuity for sweet, salty, sour, and bitter stimuli in patients that were to undergo RYGB for clinically severe obesity ⁷⁷. They found a significant upregulation of taste detection and recognition of sour and bitter stimuli in postoperative RYGB patients. At the same time, patients exhibited a trend toward a reduction in the detection and recognition of sweet and salty stimuli. Three subsequent investigations into taste acuity also found that RYGB appeared to modify patient's detection and recognition of sweet stimuli, particularly sucrose. However, these studies did not find significant changes in patient's taste perception of other stimuli following RYGB and, in fact, did not reach a consensus of whether RYGB significantly increased ^{78, 79} or decreased ⁸⁰ sucrose taste sensitivity. Examinations of taste preference in these studies also produced discordant findings. Bueter and colleagues concluded that changes in taste sensitivity did not affect the hedonic value of sucrose ⁷⁹, while Pepino and colleagues found that RYGB patients shifted their responses to repeated sucrose exposure from pleasant to unpleasant during postoperative palatability trials ⁸⁰. Interestingly, Pepino and colleagues observed a threefold decrease in lingual fungiform papillae gene expression of α -gustducin in patients who underwent RYGB, which may

explain their reduced taste sensitivity to sucrose⁸⁰. Taken together, the clinical literature would suggest that RYGB alters patient's taste perception of sucrose.

Moving from the bedside to the bench top, multiple lines of basic research evidence also support a postoperative shift in the taste perception of sucrose and other sweet stimuli. Tichansky and colleagues examined a Sprague-Dawley rat model of RYGB for changes in sweet taste behavior and found a decreased sensitivity or preference for sucrose⁸¹. This data nicely complemented that of Bueter and colleagues, who observed a drop in postoperative sucrose intake in a Wistar rat model of RYGB with a two bottle choice challenge⁷⁹. These authors also noted decreased alimentary limb expression of T1R2 and T1R3 receptors along with increased plasma GLP-1. One tantalizing possibility, given the close relationship between T1R receptors, α -gustducin, and GLP-1, is that the elevated circulating GLP-1 that accompanies RYGB may influence T1R-related signaling pathways that are crucial for peripheral sweet taste perception⁷⁹. Another study, from Hajnal and colleagues, addressed whether RYGB altered the central nervous system circuits governing sucrose taste perception. In addition to confirming reduced sweet taste acuity or preference with a two bottle choice challenge, these authors also found postoperative decreases in pontine parabrachial nucleus (PBN) neural responses to sucrose in two rat models of RYGB⁸². Although the emerging picture is complex, it seems that both peripheral and central taste processing of sweet stimuli may be altered by RYGB. It would seem, that one component of taste perception is not necessarily dependent on the other and that both components may be independently affected by RYGB. Future studies are needed to determine exactly how altered taste processing of sweet stimuli may inform food selection, which may contribute to patients consuming a lower calorie diet and weight loss.

Changes in hedonic value of specific foods ('liking')—The food reward system is a complex neural system that can be divided both anatomically and operationally into several components. One plausible and influential operational differentiation has been to distinguish hedonic liking from wanting and learning as suggested by Berridge and Robinson⁸³. This distinction is based on the simple facts that a food item that is liked is not necessarily wanted at a given time and that learning is necessary to predict the reward value of a given food. Unfortunately, the three operational components do not simply segregate into three anatomically distinct neural circuits, but while 'liking' of food is mainly organized by relevant sensory pathways such as olfaction, taste, and vision, as well as their cortico-limbic representations, 'wanting' is mainly organized by the mesolimbic dopamine system consisting of midbrain, basal ganglia, cortex, and hypothalamus. For reward-related learning, the interaction of these above pathways is further enhanced by additional cortical and subcortical structures.

In rodents, implicit 'liking' of specific taste stimuli can be assessed by the taste reactivity test^{84, 85} and by the brief access lick test⁸⁶. Compared to high-fat diet-induced obese rats which 'like' high concentrations of sucrose and corn oil the most, rats after RYGB shift 'liking' from higher to lower concentrations of both sucrose and corn oil solutions, behaving similar to lean rats⁸⁷. However, because weight loss of similar magnitude induced by calorie-restriction led to the same shift, the mechanism appears to depend on weight loss rather than some other effect of the surgery⁸⁷. In humans, explicit liking can be assessed by

questionnaire and visual analog scale ⁸⁸. Similar to the findings in rats, RYGB patients preferentially reduced liking of high- vs. low-calorie foods as assessed before and after surgery ⁸⁹. However, because no calorie restriction-induced weight loss group was compared, the mechanism(s) for this change remain unclear.

Changes in motivation to eat specific foods ('wanting')—In humans, motivation to eat has been assessed either by questionnaire or by actual measurement of how much an individual will work to obtain a food reward ⁹⁰⁻⁹². When asked how much they want to eat foods that are directly in front of them or represented by pictures, RYGB patients show markedly decreased desire to eat (wanting) compared to before surgery ⁹³, particularly if the food items are high in calories ⁸⁹. Similar findings were obtained with assessment of food craving in RYGB patients compared to normal weight controls, except that in contrast to most other studies, craving for high fat foods was not different ⁹⁴.

The willingness to 'work' for a food reward by pressing a computer key on a progressive ratio scale was recently tested in obese gastric bypass patients both before and after surgery and in normal weight controls. The break point, a measure of wanting, was selectively decreased only in RYGB patients after surgery and only when they 'worked' for candy but not for vegetables ⁹⁵. In RYGB rats, the willingness to work for a food reward (fruit loop) was assessed in the running alley, in which completion time for running from start to goal box is a measure of implicit 'wanting' ⁹⁶. Surprisingly, and in stark contrast to the study in humans, 'wanting' was significantly lower in high-fat diet-induced obese vs. lean rats and this impairment was completely normalized in rats 5 months after RYGB ⁸⁷. One possible explanation for the discrepancy is the highly different "work load" used – key pad presses in human subjects vs. running in rats, and more research is needed to clarify the role of effort in motivated behaviors ⁹⁷.

Looking for potential neural mechanisms that underlie changes in 'wanting', particularly changes in activation of components of the mesolimbic dopamine system, there is a growing literature employing fMRI and PET neuroimaging as well as magnetoencephalography (MEG). The typical experimental paradigm consists in the presentation of food and non-food pictures, and in one study, direct oral stimulation, while lying in the magnet. In normal weight individuals these visual and gustatory appetitive stimuli elicit characteristic patterns of increased neural activity in areas related to sensory processing as well as key areas of the mesolimbic dopamine system such as the ventral tegmental area, ventral striatum, and various cortical areas. These responses are typically exaggerated in obese subjects and decreased after bariatric surgery ^{53, 98-102}, suggesting that external food cues lose their ability to drive eating after surgery. Specifically, the decreased striatal activity observed in fasted RYGB subjects correlated with reductions in "wanting" calorie-dense foods, but not "liking" for such foods ⁸⁹, reinforcing the idea of distinct neural systems mediating wanting and liking. Dopamine signaling through D1 and D2 receptors within target areas of the mesolimbic dopamine system is thought to be crucial for motivated behaviors such as food intake ¹⁰³ and a few studies have begun to examine components of dopamine receptor signaling after gastric bypass surgery in humans and rats ¹⁰⁴⁻¹⁰⁶.

In summary, the initial sketchy reports of changes in acceptance of, and preference for, specific foods in bariatric surgery patients have now been largely confirmed in easier to control rodent studies. Specifically, a number of studies in rodents show decreased preference for sweet and fatty foods after RYGB or sleeve gastrectomy. Studies designed to identify the neural component(s) responsible for these adaptive changes further suggest that 'liking' of, not only high-fat, but also high-sucrose taste stimuli, is decreased after RYGB. However, whether this shift towards low-calorie sweet and fatty stimuli is due to changes in taste perception or more central components of taste processing is not yet clear and needs further investigation. Similarly, although one recent study found RYGB patients to be less willing to work for high-calorie food, there is no clear consensus regarding changes in the motivation to obtain food rewards ('wanting') and its underlying mesolimbic dopamine system.

Importantly, the mechanisms responsible for any changes in these components of food hedonics after bariatric surgery are not known. Candidate mechanisms include changes in signaling by gut hormones and other gut factors such as GLP-1, PYY, ghrelin, bile acids, and microbiota-derived proteins, as well as changes in leptin signaling. None of these hypothesized mechanisms have been directly tested in interventional approaches, but the recent observations that both RYGB and sleeve gastrectomy are effective in reducing food intake and body weight in whole body knockout mice deficient in GLP-1 receptor³⁵ or ghrelin-signaling¹⁰⁷ have somewhat shortened this list of important candidate mechanisms. Just as recently shown for improvements of glycemic control⁴⁰, it is likely that surgery-induced calorie-restriction and weight loss are perhaps more important than gut-specific hormonal mechanisms.

Ideally, future studies should be longitudinal, allowing assessment of food hedonism in the same subjects before surgery (in the obese state) and at several time points after surgery. It should also include matched, calorie-restriction-induced weight loss groups for comparison, and food hedonics should be tested in both fasted and fed conditions to capture the full dynamic range.

The role of aversive learning and conditioned anorexia

The marked changes in ingestive behavior with smaller meals and shifts in preference observed in humans and rodents after RYGB^{28, 38, 71, 73, 76, 108, 109} strongly suggest the involvement of coping or learning mechanisms to avoid unpleasant gastrointestinal sensations such as fullness, nausea and pain when ingesting too much of certain foods. Indeed, bariatric surgery in humans often causes episodes of fullness, nausea, pain, and vomiting with different intensity and frequency for different bariatric procedures¹¹⁰⁻¹¹⁴. Furthermore, conditioned taste aversion to orally administered corn oil was demonstrated in rats after RYGB⁷⁶.

The physiological states and mechanisms of satiety and nausea are typically discussed as distinct, even though they are mediated by partially overlapping brain areas including the nucleus of the solitary tract, parabrachial nucleus, and amygdala¹¹⁵⁻¹²⁰ [and see¹²¹ for a recent review]. Earlier literature has identified the parabrachial nucleus (PBN) as a site of

integration of viscerosensory information, including gastric distension, and taste^{116, 117, 122}. The PBN, particularly its lateral subnuclei, is activated by a number of diverse stimuli including intraperitoneal administration of the satiety hormones CCK-8¹²³, GLP-1^{118, 124}, Exendin-4¹²⁵, PYY¹¹⁹, and amylin^{118, 126}, electrical stimulation of vagal afferents¹²⁷, the 5-HT reuptake inhibitor dexfenfluramine¹²³, systemic administration of LiCl^{120, 128}, lipopolysaccharide (LPS)¹²⁹ and other immune-activation signals¹³⁰, and the cancer chemotherapy drug cisplatin^{131, 132}. Importantly, a common effect of all these challenges is a reduction of food intake. Thus, there is a strong correlation between LPBN activity and anorexia.

Only recently has a more integrative picture emerged, with satiety, nausea, and anorexia, seen as a functional continuum that opposes hunger orchestrated by hypothalamic AGRP/NPY neurons. More specifically, a group of calcitonin-gene related peptide expressing neurons has been identified in the external lateral subnucleus of the PBN¹³³⁻¹³⁸. The profound anorexia and starvation of mice after AGRP neuron ablation in adult mice is surprisingly not only caused by food intake stimulatory actions of AGRP and NPY projections elsewhere¹³⁹⁻¹⁴¹, but also by withdrawal of GABA_A-ergic inhibitory input to these critical lateral parabrachial nucleus neurons. Activity within this specific group of neurons is positively correlated with the food intake suppressing effect of a variety of stimuli such as exogenous cholecystokinin and amylin, lithium chloride and lipopolysaccharide¹³⁷, and we recently found similarly exaggerated neural activity in these neurons in mice eating a high-fat meal after RYGB as compared to sham surgery (Berthoud, unpublished observations). These preliminary observations suggest that the LPBN anorexia pathway is strongly activated by eating a meal, particularly a high-fat meal, early after RYGB, potentially explaining the rapid behavioral change of eating smaller but more frequent meals which is evident soon after surgery³⁸. It is conceivable that this exaggerated activation leads to negative reinforcement of eating large meals and may be selective for specific food types, e.g. high fat diet^{74, 87, 95}. This could lead to permanent changes in food intake through plastic neural changes within the anorexia pathway and its interactions with the homeostatic regulator and the reward system.

In summary, there is clear evidence that ingestive behavior and food choice changes after bariatric surgeries and that eating “as usual” can cause discomfort and nausea. It is thus very plausible that animals and humans learn to avoid these negative consequences and thereby reduce food intake. The recent re-discovery of the lateral parabrachial nucleus as a hub of viscerosensory integration and the molecular identification of its major inputs and outputs offer the intriguing possibility that this anorexia pathway plays a crucial role in the food intake-suppressing effects of bariatric surgeries. This idea gains additional support from the demonstration that this anorexia pathway is part of the classic homeostatic feeding circuits in the basomedial hypothalamus and projects to important behavioral effector systems in the paraventricular nucleus of the hypothalamus and the mesolimbic dopamine pathways. It may thus be possible to leverage this system for the development of drug or behavioral therapies to suppress food intake without induction of nausea.

The role of executive control

When bariatric surgery patients are asked to describe their eating experience using interpretative phenomenological analysis, self-control was the central theme permeating all areas of the interviews ^{142, 143}. Most of these patients have been struggling all their life with control over eating and successful surgery appeared to make control easier. Ogden et al. concluded that successful surgery without weight regain brings the patient's mind 'in gear', while failed surgery is characterized by a continuing battle for control ¹⁴³. While the liking and wanting systems generate incentive salience and craving (as discussed above), the executive control system acts as a brake to align impulsive behavior with longer term goals. Inhibitory control is particularly important to resist temptation to eat as stimulated by ubiquitous food cues in the modern environment.

The neural system underlying executive control is not well-defined, but the dorsolateral prefrontal cortex (DLPFC) is thought to be an important component ¹⁴⁴⁻¹⁴⁷. Very little is known about how these executive control functions impact obesity or how it may change after RYGB. A recent study by Goldstone and colleagues may be among the first to address these questions ¹⁴⁸. These authors initially stratified a population of recent RYGB patients based solely on how successful they were at weight loss and then compared their neural responses to food cues during an executive control challenge. This challenge consisted of two parts, a crave phase in which patients were told to let themselves desire a given food after seeing its visual cue and a resist phase in which they were told to do the opposite. Functional MRI analysis of patient's brains revealed that the crave phase was associated with dorsomedial prefrontal cortex (DMPFC) activity and the resist phase was associated with DLPFC activity. While their entire RYGB patient population had similar levels of DMPFC activity during the crave phase, the most successful weight loss patients demonstrated significantly higher DLPFC activity during the resist phase. It would appear then that the success of RYGB surgery may be due, in part, to the postoperative ability of an individual to mobilize neural circuits involved in executive control ¹⁴⁹.

It is unclear, though, if executive functions are specifically impaired with obesity or even preoperatively in RYGB patients. The extent to which the hedonic value of a given food cue impacts executive control functions also remains to be determined. That is to say, RYGB postoperative changes in food hedonics may also facilitate executive control functions ⁸⁹. If this is the case, then by simply removing or reducing the hedonic value of notoriously high calorie foods through RYGB would make it easier to self-regulate their intake. Another intriguing possibility is that the ability to mobilize DLPFC activity is aided by signals from the parabrachial nucleus-amygdala anorexia pathway discussed above.

Summary and conclusions

During the past decade or so, there has been a surge in studies characterizing the effects of bariatric surgeries in humans and rodents. They have demonstrated numerous structural, functional, and molecular changes in the gut, the brain, and other organs as well as changes in energy metabolism, glucose homeostasis, and behavior. After going for the 'low-hanging fruit', it is now time to separate irrelevant changes from mechanistically relevant ones. The

marked and sustained body weight loss and concomitant correction of many obesity-related impairments in metabolism and behavior are well-documented, while other effects that do not depend on the hypo-caloric state and weight loss are variable and less clear. In general, while the sustained body weight loss in humans is mainly explained by reduced energy intake, not increased energy expenditure, the opposite is true for rodents; only temporary reduction in energy intake but increased energy expenditure, at least in the long term and particularly in mice. However, regardless of these differences, a key observation is that energy intake is not increased to re-gain lost body weight, even though food intake can be doubled if properly stimulated. This suggests active defense of a new lower body weight level after surgery. Thus, explaining the potential mechanisms for this sustained relative hypophagia is perhaps most crucial for future non-surgical treatments of obesity. While a number of candidate mechanisms have been proposed on the basis of changes in gut hormone secretion as well as changes in peripheral and central targets of such hormones, direct testing of individual signaling cascades was unable to confirm any of these hypotheses so far.

Acknowledgments

Supported by National Institutes of Health Grants DK 047348 (HRB), DK092587 (HM), F32DK097986 (KRZ), and 2P30 DK072476 (SY).

References

1. Carey DG, Pliego GJ, Raymond RL. Body composition and metabolic changes following bariatric surgery: effects on fat mass, lean mass and basal metabolic rate: six months to one-year follow-up. *Obes Surg*. 2006; 16:1602–8. [PubMed: 17217636]
2. van Gemert WG, Westerterp KR, van Acker BA, Wagenmakers AJ, Halliday D, Greve JM, et al. Energy, substrate and protein metabolism in morbid obesity before, during and after massive weight loss. *Int J Obes Relat Metab Disord*. 2000; 24:711–8. [PubMed: 10878677]
3. Bobbioni-Harsch E, Morel P, Huber O, Assimacopoulos-Jeannet F, Chassot G, Lehmann T, et al. Energy economy hampers body weight loss after gastric bypass. *J Clin Endocrinol Metab*. 2000; 85:4695–700. [PubMed: 11134130]
4. Flancbaum L, Choban PS, Bradley LR, Burge JC. Changes in measured resting energy expenditure after Roux-en-Y gastric bypass for clinically severe obesity. *Surgery*. 1997; 122:943–9. [PubMed: 9369895]
5. Maclean PS, Bergouignan A, Cornier MA, Jackman MR. Biology's response to dieting: the impetus for weight regain. *Am J Physiol Regul Integr Comp Physiol*. 2011; 301:R581–600. [PubMed: 21677272]
6. Thivel D, Brakonietki K, Duche P, Morio B, Boirie Y, Laferrere B. Surgical weight loss: impact on energy expenditure. *Obes Surg*. 2013; 23:255–66. [PubMed: 23224568]
7. Das SK, Roberts SB, McCrory MA, Hsu LK, Shikora SA, Kehayias JJ, et al. Long-term changes in energy expenditure and body composition after massive weight loss induced by gastric bypass surgery. *Am J Clin Nutr*. 2003; 78:22–30. [PubMed: 12816767]
8. Bueter M, Lowenstein C, Olbers T, Wang M, Cluny NL, Bloom SR, et al. Gastric bypass increases energy expenditure in rats. *Gastroenterology*. 2010; 138:1845–53. [PubMed: 19931268]
9. Stylopoulos N, Hoppin AG, Kaplan LM. Roux-en-Y gastric bypass enhances energy expenditure and extends lifespan in diet-induced obese rats. *Obesity (Silver Spring)*. 2009; 17:1839–47. [PubMed: 19556976]
10. Hao Z, Zhao Z, Berthoud HR, Ye J. Development and verification of a mouse model for roux-en-Y gastric bypass surgery with a small gastric pouch. *PLoS ONE*. 2013; 8:e52922. [PubMed: 23326365]

11. Nestoridi E, Kvas S, Kucharczyk J, Stylopoulos N. Resting Energy Expenditure and Energetic Cost of Feeding Are Augmented after Roux-en-Y Gastric Bypass in Obese Mice. *Endocrinology*. 2012
12. Rothwell NJ, Stock MJ. A paradox in the control of energy intake in the rat. *Nature*. 1978; 273:146–7. [PubMed: 643077]
13. Kaiyala KJ, Schwartz MW. Toward a more complete (and less controversial) understanding of energy expenditure and its role in obesity pathogenesis. *Diabetes*. 2011; 60:17–23. [PubMed: 21193735]
14. Rosenbaum M, Hirsch J, Gallagher DA, Leibel RL. Long-term persistence of adaptive thermogenesis in subjects who have maintained a reduced body weight. *Am J Clin Nutr*. 2008; 88:906–12. [PubMed: 18842775]
15. Watanabe M, Houten SM, Mataki C, Christoffolete MA, Kim BW, Sato H, et al. Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature*. 2006; 439:484–9. [PubMed: 16400329]
16. Jansen PL, van Werven J, Aarts E, Berends F, Janssen I, Stoker J, et al. Alterations of hormonally active fibroblast growth factors after Roux-en-Y gastric bypass surgery. *Dig Dis*. 2011; 29:48–51. [PubMed: 21691104]
17. Pournaras DJ, Glicksman C, Vincent RP, Kuganlipava S, Alaghband-Zadeh J, Mahon D, et al. The role of bile after Roux-en-Y gastric bypass in promoting weight loss and improving glycaemic control. *Endocrinology*. 2012; 153:3613–9. [PubMed: 22673227]
18. Simonen M, Dali-Youcef N, Kaminska D, Venesmaa S, Kakela P, Paakkonen M, et al. Conjugated Bile Acids Associate with Altered Rates of Glucose and Lipid Oxidation after Roux-en-Y Gastric Bypass. *Obes Surg*. 2012; 22:1473–80. [PubMed: 22638681]
19. Cummings BP, Bettaieb A, Graham JL, Stanhope KL, Kowala M, Haj FG, et al. Vertical sleeve gastrectomy improves glucose and lipid metabolism and delays diabetes onset in UCD-T2DM rats. *Endocrinology*. 2012; 153:3620–32. [PubMed: 22719048]
20. Brufau G, Bahr MJ, Staels B, Claudel T, Ockenga J, Boker KH, et al. Plasma bile acids are not associated with energy metabolism in humans. *Nutr Metab (Lond)*. 2010; 7:73. [PubMed: 20815878]
21. Ockenga J, Valentini L, Schuetz T, Wohlgemuth F, Glaeser S, Omar A, et al. Plasma bile acids are associated with energy expenditure and thyroid function in humans. *J Clin Endocrinol Metab*. 2012; 97:535–42. [PubMed: 22162464]
22. Stearns AT, Balakrishnan A, Tavakkolizadeh A. Impact of Roux-en-Y gastric bypass surgery on rat intestinal glucose transport. *Am J Physiol Gastrointest Liver Physiol*. 2009; 297:G950–7. [PubMed: 20501442]
23. le Roux CW, Borg C, Wallis K, Vincent RP, Bueter M, Goodlad R, et al. Gut hypertrophy after gastric bypass is associated with increased glucagon-like peptide 2 and intestinal crypt cell proliferation. *Ann Surg*. 2010; 252:50–6. [PubMed: 20562614]
24. Mumphrey MB, Patterson LM, Zheng H, Berthoud HR. Roux-en-Y gastric bypass surgery increases number but not density of CCK-, GLP-1-, 5-HT-, and neurotensin-expressing enteroendocrine cells in rats. *Neurogastroenterol Motil*. 2013; 25:e70–9. [PubMed: 23095091]
25. Hansen CF, Bueter M, Theis N, Lutz T, Paulsen S, Dalboge LS, et al. Hypertrophy Dependent Doubling of L-Cells in Roux-en-Y Gastric Bypass Operated Rats. *PLoS ONE*. 2013; 8:e65696. [PubMed: 23776529]
26. Saeidi N ML. Reprogramming of intestinal glucose metabolism and glycemic control in rats after gastric bypass. *Science (Washington, D C, 1883-)*. 2013
27. Sjostrom L, Lindroos AK, Peltonen M, Torgerson J, Bouchard C, Carlsson B, et al. Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *N Engl J Med*. 2004; 351:2683–93. [PubMed: 15616203]
28. Laurenus A, Larsson I, Bueter M, Melanson KJ, Bosaeus I, Forslund HB, et al. Changes in eating behaviour and meal pattern following Roux-en-Y gastric bypass. *Int J Obes (Lond)*. 2012; 36:348–55. [PubMed: 22124454]

29. Brolin RE, Robertson LB, Kenler HA, Cody RP. Weight loss and dietary intake after vertical banded gastroplasty and Roux-en-Y gastric bypass. *Ann Surg.* 1994; 220:782–90. [PubMed: 7986146]
30. Kumar R, Lieske JC, Collazo-Clavell ML, Sarr MG, Olson ER, Vrtiska TJ, et al. Fat malabsorption and increased intestinal oxalate absorption are common after Roux-en-Y gastric bypass surgery. *Surgery.* 2011; 149:654–61. [PubMed: 21295813]
31. Custodio Afonso Rocha V, Ramos de Arvelos L, Pereira Felix G, Nogueira Prado de Souza D, Bernardino Neto M, Santos Resende E, et al. Evolution of nutritional, hematologic and biochemical changes in obese women during 8 weeks after Roux-en-Y gastric bypass. *Nutr Hosp.* 2012; 27:1134–40. [PubMed: 23165553]
32. Odstrcil EA, Martinez JG, Santa Ana CA, Xue B, Schneider RE, Steffer KJ, et al. The contribution of malabsorption to the reduction in net energy absorption after long-limb Roux-en-Y gastric bypass. *Am J Clin Nutr.* 2010; 92:704–13. [PubMed: 20739420]
33. Ortega J, Ortega-Evangelio G, Cassinello N, Sebastia V. What are obese patients able to eat after Roux-en-Y gastric bypass? *Obes Facts.* 2012; 5:339–48. [PubMed: 22722236]
34. Stefater MA, Perez-Tilve D, Chambers AP, Wilson-Perez HE, Sandoval DA, Berger J, et al. Sleeve gastrectomy induces loss of weight and fat mass in obese rats, but does not affect leptin sensitivity. *Gastroenterology.* 2010; 138:2426–36. e1–3. 36. [PubMed: 20226189]
35. Ye J, Hao Z, Mumphy MB, Townsend RL, Patterson LM, Stylopoulos N, et al. GLP-1 receptor signaling is not required for reduced body weight after RYGB in rodents. *Am J Physiol Regul Integr Comp Physiol.* 2014
36. Zechner JF, Mirshahi UL, Satapati S, Berglund ED, Rossi J, Scott MM, et al. Weight-independent effects of roux-en-Y gastric bypass on glucose homeostasis via melanocortin-4 receptors in mice and humans. *Gastroenterology.* 2013; 144:580–90. e7. [PubMed: 23159449]
37. Furnes MW, Tommeras K, Arum CJ, Zhao CM, Chen D. Gastric bypass surgery causes body weight loss without reducing food intake in rats. *Obes Surg.* 2008; 18:415–22. [PubMed: 18247101]
38. Zheng H, Shin AC, Lenard NR, Townsend RL, Patterson LM, Sigalet DL, et al. Meal patterns, satiety, and food choice in a rat model of Roux-en-Y gastric bypass surgery. *Am J Physiol Regul Integr Comp Physiol.* 2009; 297:R1273–82. [PubMed: 19726714]
39. Seyfried F, Lannoo M, Gsell W, Tremoleda JL, Bueter M, Olbers T, et al. Roux-en-Y gastric bypass in mice--surgical technique and characterisation. *Obes Surg.* 2012; 22:1117–25. [PubMed: 22527601]
40. Jackness C, Karmally W, Febres G, Conwell IM, Ahmed L, Bessler M, et al. Very Low Calorie Diet Mimics the Early Beneficial Effect of Roux-en-Y Gastric Bypass on Insulin Sensitivity and Beta-Cell Function in Type 2 Diabetic Patients. *Diabetes.* 2013
41. Lingvay I, Guth E, Islam A, Livingston E. Rapid Improvement of Diabetes After Gastric Bypass Surgery: Is It the Diet or Surgery? *Diabetes Care.* 2013
42. Lips MA, de Groot GH, van Klinken JB, Aarts E, Berends FJ, Janssen IM, et al. Calorie Restriction is a Major Determinant of the Short-Term Metabolic Effects of Gastric Bypass Surgery in Obese Type 2 Diabetic Patients. *Clin Endocrinol (Oxf).* 2013
43. Isbell JM, Tamboli RA, Hansen EN, Saliba J, Dunn JP, Phillips SE, et al. The importance of caloric restriction in the early improvements in insulin sensitivity after Roux-en-Y gastric bypass surgery. *Diabetes Care.* 2010; 33:1438–42. [PubMed: 20368410]
44. Schwartz MW, Woods SC, Porte D Jr. Seeley RJ, Baskin DG. Central nervous system control of food intake. *Nature.* 2000; 404:661–71. [PubMed: 10766253]
45. Berthoud HR. Metabolic and hedonic drives in the neural control of appetite: who is the boss? *Curr Opin Neurobiol.* 2011; 21:888–96. [PubMed: 21981809]
46. Stefater MA, Sandoval DA, Chambers AP, Wilson-Perez HE, Hofmann SM, Jandacek R, et al. Sleeve gastrectomy in rats improves postprandial lipid clearance by reducing intestinal triglyceride secretion. *Gastroenterology.* 2011; 141:939–49. e1–4. [PubMed: 21699773]
47. Grayson BE, Schneider KM, Woods SC, Seeley RJ. Improved rodent maternal metabolism but reduced intrauterine growth after vertical sleeve gastrectomy. *Sci Transl Med.* 2013; 5:199ra12.

48. Stefater MA, Wilson-Perez HE, Chambers AP, Sandoval DA, Seeley RJ. All bariatric surgeries are not created equal: insights from mechanistic comparisons. *Endocr Rev.* 2012; 33:595–622. [PubMed: 22550271]
49. Shin AC, Zheng H, Berthoud HR. An expanded view of energy homeostasis: neural integration of metabolic, cognitive, and emotional drives to eat. *Physiol Behav.* 2009; 97:572–80. [PubMed: 19419661]
50. Keeseey RE, Hirvonen MD. Body weight set-points: determination and adjustment. *J Nutr.* 1997; 127:1875S–83S. [PubMed: 9278574]
51. Wirtshafter D, Davis JD. Set points, settling points, and the control of body weight. *Physiol Behav.* 1977; 19:75–8. [PubMed: 11803695]
52. Harris RB. Role of set-point theory in regulation of body weight. *FASEB J.* 1990; 4:3310–8. [PubMed: 2253845]
53. Frank S, Wilms B, Veit R, Ernst B, Thurnheer M, Kullmann S, et al. Altered brain activity in severely obese women may recover after Roux-en Y gastric bypass surgery. *Int J Obes (Lond).* 2013
54. Paulsen SJ, Larsen LK, Jelsing J, Janssen U, Gerstmayer B, Vrang N. Gene expression profiling of individual hypothalamic nuclei from single animals using laser capture microdissection and microarrays. *J Neurosci Methods.* 2009; 177:87–93. [PubMed: 18955080]
55. Shin AC, Berthoud HR. Obesity surgery: happy with less or eternally hungry? *Trends Endocrinol Metab.* 2013; 24:101–8. [PubMed: 23317811]
56. Nadreau E, Baraboi ED, Samson P, Blouin A, Hould FS, Marceau P, et al. Effects of the biliopancreatic diversion on energy balance in the rat. *Int J Obes (Lond).* 2006; 30:419–29. [PubMed: 16302016]
57. Warne JP, Padilla BE, Horneman HF, Ginsberg AB, Pecoraro NC, Akana SF, et al. Metabolic and neuroendocrine consequences of a duodenal-jejunal bypass in rats on a choice diet. *Ann Surg.* 2009; 249:269–76. [PubMed: 19212181]
58. Romanova IV, Ramos EJ, Xu Y, Quinn R, Chen C, George ZM, et al. Neurobiologic changes in the hypothalamus associated with weight loss after gastric bypass. *J Am Coll Surg.* 2004; 199:887–95. [PubMed: 15555972]
59. Rosenbaum M, Sy M, Pavlovich K, Leibel RL, Hirsch J. Leptin reverses weight loss-induced changes in regional neural activity responses to visual food stimuli. *J Clin Invest.* 2008; 118:2583–91. [PubMed: 18568078]
60. Korner J, Conroy R, Febres G, McMahon DJ, Conwell I, Karmally W, et al. Randomized double-blind placebo-controlled study of leptin administration after gastric bypass. *Obesity (Silver Spring).* 2013; 21:951–6. [PubMed: 23512892]
61. Hatoum IJ, Stylopoulos N, Vanhoose AM, Boyd KL, Yin DP, Ellacott KL, et al. Melanocortin-4 Receptor Signaling Is Required for Weight Loss after Gastric Bypass Surgery. *J Clin Endocrinol Metab.* 2012
62. Mul JD, Begg DP, Alsters SI, van Haaften G, Duran KJ, D'Alessio DA, et al. Effect of vertical sleeve gastrectomy in melanocortin receptor 4-deficient rats. *Am J Physiol Endocrinol Metab.* 2012; 303:E103–10. [PubMed: 22535749]
63. Aslan IR, Campos GM, Calton MA, Evans DS, Merriman RB, Vaisse C. Weight loss after Roux-en-Y gastric bypass in obese patients heterozygous for MC4R mutations. *Obes Surg.* 2011; 21:930–4. [PubMed: 20957447]
64. Potoczna N, Branson R, Kral JG, Piec G, Steffen R, Ricklin T, et al. Gene variants and binge eating as predictors of comorbidity and outcome of treatment in severe obesity. *J Gastrointest Surg.* 2004; 8:971–81. discussion 81–2. [PubMed: 15585384]
65. Aslan IR, Ranadive SA, Ersoy BA, Rogers SJ, Lustig RH, Vaisse C. Bariatric surgery in a patient with complete MC4R deficiency. *Int J Obes (Lond).* 2011; 35:457–61. [PubMed: 20733581]
66. Krashes MJ, Koda S, Ye C, Rogan SC, Adams AC, Cusher DS, et al. Rapid, reversible activation of AgRP neurons drives feeding behavior in mice. *J Clin Invest.* 2011
67. Berridge KC, Ho CY, Richard JM, DiFeliceantonio AG. The tempted brain eats: Pleasure and desire circuits in obesity and eating disorders. *Brain Res.* 2010; 1350:43–64. [PubMed: 20388498]

68. Berthoud HR, Lenard NR, Shin AC. Food reward, hyperphagia, and obesity. *Am J Physiol Regul Integr Comp Physiol.* 2011; 300:R1266–77. [PubMed: 21411768]
69. Stice E, Spoor S, Ng J, Zald DH. Relation of obesity to consummatory and anticipatory food reward. *Physiol Behav.* 2009; 97:551–60. [PubMed: 19328819]
70. Mathes CM, Spector AC. Food selection and taste changes in humans after Roux-en-Y gastric bypass surgery: a direct-measures approach. *Physiol Behav.* 2012; 107:476–83. [PubMed: 22366157]
71. Shin AC, Berthoud HR. Food reward functions as affected by obesity and bariatric surgery. *Int J Obes (Lond).* 2011; 35(Suppl 3):S40–4. [PubMed: 21912387]
72. Miras AD, le Roux CW. Bariatric surgery and taste: novel mechanisms of weight loss. *Curr Opin Gastroenterol.* 2010; 26:140–5. [PubMed: 19901832]
73. Laurenius A, Larsson I, Melanson KJ, Lindroos AK, Lonroth H, Bosaeus I, et al. Decreased energy density and changes in food selection following Roux-en-Y gastric bypass. *Eur J Clin Nutr.* 2013; 67:168–73. [PubMed: 23299713]
74. Wilson-Perez HE, Chambers AP, Sandoval DA, Stefater MA, Woods SC, Benoit SC, et al. The effect of vertical sleeve gastrectomy on food choice in rats. *Int J Obes (Lond).* 2013; 37:288–95. [PubMed: 22334194]
75. Saeidi N, Nestoridi E, Kucharczyk J, Uygun MK, Yarmush ML, Stylopoulos N. Sleeve gastrectomy and Roux-en-Y gastric bypass exhibit differential effects on food preferences, nutrient absorption and energy expenditure in obese rats. *Int J Obes (Lond).* 2012; 36:1396–402. [PubMed: 23044855]
76. le Roux CW, Bueter M, Theis N, Werling M, Ashrafi H, Lowenstein C, et al. Gastric bypass reduces fat intake and preference. *Am J Physiol Regul Integr Comp Physiol.* 2011; 301:R1057–66. [PubMed: 21734019]
77. Scruggs DM, Buffington C, Cowan GS Jr. Taste Acuity of the Morbidly Obese before and after Gastric Bypass Surgery. *Obes Surg.* 1994; 4:24–28. [PubMed: 10742759]
78. Burge JC, Schaumburg JZ, Choban PS, DiSilvestro RA, Flancbaum L. Changes in patients' taste acuity after Roux-en-Y gastric bypass for clinically severe obesity. *J Am Diet Assoc.* 1995; 95:666–70. [PubMed: 7759742]
79. Bueter M, Miras AD, Chichger H, Fenske W, Ghatei MA, Bloom SR, et al. Alterations of sucrose preference after Roux-en-Y gastric bypass. *Physiol Behav.* 2011; 104:709–21. [PubMed: 21827777]
80. Pepino MY, Bradley D, Eagon JC, Sullivan S, Abumrad NA, Klein S. Changes in taste perception and eating behavior after bariatric surgery-induced weight loss in women. *Obesity (Silver Spring).* 2013
81. Tichansky DS, Glatt AR, Madan AK, Harper J, Tokita K, Boughter JD. Decrease in sweet taste in rats after gastric bypass surgery. *Surg Endosc.* 2011; 25:1176–81. [PubMed: 20844896]
82. Hajnal A, Kovacs P, Ahmed TA, Meirelles K, Lynch CJ, Cooney RN. Gastric bypass surgery alters behavioral and neural taste functions for sweet taste in obese rats. *Am J Physiol Gastrointest Liver Physiol.* 2010; 299:G967–G79. [PubMed: 20634436]
83. Berridge KC, Robinson TE. Parsing reward. *Trends Neurosci.* 2003; 26:507–13. [PubMed: 12948663]
84. Grill HJ, Norgren R. The taste reactivity test. I. Mimetic responses to gustatory stimuli in neurologically normal rats. *Brain Res.* 1978; 143:263–79. [PubMed: 630409]
85. Berridge KC. Modulation of taste affect by hunger, caloric satiety, and sensory-specific satiety in the rat. *Appetite.* 1991; 16:103–20. [PubMed: 2064389]
86. Spector AC, Redman R, Garcea M. The consequences of gustatory nerve transection on taste-guided licking of sucrose and maltose in the rat. *Behav Neurosci.* 1996; 110:1096–109. [PubMed: 8919012]
87. Shin AC, Zheng H, Pistell PJ, Berthoud HR. Roux-en-Y gastric bypass surgery changes food reward in rats. *Int J Obes (Lond).* 2011; 35:642–51. [PubMed: 20805826]
88. Finlayson G, King N, Blundell JE. Liking vs. wanting food: importance for human appetite control and weight regulation. *Neurosci Biobehav Rev.* 2007; 31:987–1002. [PubMed: 17559933]

89. Ochner CN, Stice E, Hutchins E, Afifi L, Geliebter A, Hirsch J, et al. Relation between changes in neural responsivity and reductions in desire to eat high-calorie foods following gastric bypass surgery. *Neuroscience*. 2012; 209:128–35. [PubMed: 22406414]
90. Finlayson G, King N, Blundell J. The role of implicit wanting in relation to explicit liking and wanting for food: implications for appetite control. *Appetite*. 2008; 50:120–7. [PubMed: 17655972]
91. Lemmens SG, Schoffelen PF, Wouters L, Born JM, Martens MJ, Rutters F, et al. Eating what you like induces a stronger decrease of ‘wanting’ to eat. *Physiol Behav*. 2009; 98:318–25. [PubMed: 19545583]
92. Nasser JA, Evans SM, Geliebter A, Pi-Sunyer FX, Foltin RW. Use of an operant task to estimate food reinforcement in adult humans with and without BED. *Obesity (Silver Spring)*. 2008; 16:1816–20. [PubMed: 18535551]
93. Ullrich J, Ernst B, Wilms B, Thurnheer M, Schultes B. Roux-en Y gastric bypass surgery reduces hedonic hunger and improves dietary habits in severely obese subjects. *Obes Surg*. 2013; 23:50–5. [PubMed: 22941334]
94. Leahey TM, Bond DS, Raynor H, Roye D, Vithiananthan S, Ryder BA, et al. Effects of bariatric surgery on food cravings: do food cravings and the consumption of craved foods “normalize” after surgery? *Surg Obes Relat Dis*. 2012; 8:84–91. [PubMed: 21925967]
95. Miras AD, Jackson RN, Jackson SN, Goldstone AP, Olbers T, Hackenberg T, et al. Gastric bypass surgery for obesity decreases the reward value of a sweet-fat stimulus as assessed in a progressive ratio task. *Am J Clin Nutr*. 2012; 96:467–73. [PubMed: 22836034]
96. Pecina S, Cagniard B, Berridge KC, Aldridge JW, Zhuang X. Hyperdopaminergic mutant mice have higher “wanting” but not “liking” for sweet rewards. *J Neurosci*. 2003; 23:9395–402. [PubMed: 14561867]
97. Salamone JD, Correa M, Farrar AM, Nunes EJ, Pardo M. Dopamine, behavioral economics, and effort. *Front Behav Neurosci*. 2009; 3:13. [PubMed: 19826615]
98. Ochner CN, Kwok Y, Conceicao E, Pantazatos SP, Puma LM, Carnell S, et al. Selective reduction in neural responses to high calorie foods following gastric bypass surgery. *Ann Surg*. 2011; 253:502–7. [PubMed: 21169809]
99. Ochner CN, Laferrere B, Afifi L, Atalayer D, Geliebter A, Teixeira J. Neural responsivity to food cues in fasted and fed states pre and post gastric bypass surgery. *Neurosci Res*. 2012; 74:138–43. [PubMed: 22921709]
100. Scholtz S, Miras AD, Chhina N, Prechtel CG, Sleeth ML, Daud NM, et al. Obese patients after gastric bypass surgery have lower brain-hedonic responses to food than after gastric banding. *Gut*. 2013
101. Bruce AS, Bruce JM, Ness AR, Lepping RJ, Malley S, Hancock L, et al. A comparison of functional brain changes associated with surgical versus behavioral weight loss. *Obesity (Silver Spring)*. 2014; 22:337–43. [PubMed: 24115765]
102. Sweet LH, Hassenstab JJ, McCaffery JM, Raynor HA, Bond DS, Demos KE, et al. Brain response to food stimulation in obese, normal weight, and successful weight loss maintainers. *Obesity (Silver Spring)*. 2012; 20:2220–5. [PubMed: 22569002]
103. Stice E, Yokum S, Blum K, Bohon C. Weight gain is associated with reduced striatal response to palatable food. *J Neurosci*. 2010; 30:13105–9. [PubMed: 20881128]
104. Dunn JP, Cowan RL, Volkow ND, Feurer ID, Li R, Williams DB, et al. Decreased dopamine type 2 receptor availability after bariatric surgery: preliminary findings. *Brain Res*. 2010; 1350:123–30. [PubMed: 20362560]
105. Steele KE, Prokopowicz GP, Schweitzer MA, Magunson TH, Lidor AO, Kuwabawa H, et al. Alterations of central dopamine receptors before and after gastric bypass surgery. *Obes Surg*. 2010; 20:369–74. [PubMed: 19902317]
106. Davis JF, Tracy AL, Schurdak JD, Magrisso IJ, Grayson BE, Seeley RJ, et al. Roux en Y gastric bypass increases ethanol intake in the rat. *Obes Surg*. 2013; 23:920–30. [PubMed: 23440511]
107. Chambers AP, Kirchner H, Wilson-Perez HE, Willency JA, Hale JE, Gaylinn BD, et al. The effects of vertical sleeve gastrectomy in rodents are ghrelin independent. *Gastroenterology*. 2013; 144:50–52. e5. [PubMed: 22995675]

108. Shin AC, Zheng H, Pistell PJ, Berthoud HR. Roux-en-Y gastric bypass surgery changes food reward in rats. *Int J Obes (Lond)*. 2011; 35:642–51. [PubMed: 20805826]
109. Delin CR, Watts JM, Saebel JL, Anderson PG. Eating behavior and the experience of hunger following gastric bypass surgery for morbid obesity. *Obes Surg*. 1997; 7:405–13. [PubMed: 9730494]
110. Harbottle L. Audit of nutritional and dietary outcomes of bariatric surgery patients. *Obes Rev*. 2011; 12:198–204. [PubMed: 20406412]
111. Laurenus A, Olbers T, Naslund I, Karlsson J. Dumping syndrome following gastric bypass: validation of the dumping symptom rating scale. *Obes Surg*. 2013; 23:740–55. [PubMed: 23315151]
112. Banerjee A, Ding Y, Mikami DJ, Needleman BJ. The role of dumping syndrome in weight loss after gastric bypass surgery. *Surg Endosc*. 2013; 27:1573–8. [PubMed: 23233009]
113. Sirinek KR, O'Dorisio TM, Howe B, McFee AS. Neurotensin, vasoactive intestinal peptide, and Roux-en-Y gastrojejunostomy. Their role in the dumping syndrome. *Arch Surg*. 1985; 120:605–9. [PubMed: 3985800]
114. Suter M, Calmes JM, Paroz A, Giusti V. A new questionnaire for quick assessment of food tolerance after bariatric surgery. *Obes Surg*. 2007; 17:2–8. [PubMed: 17355761]
115. Seeley RJ, Blake K, Rushing PA, Benoit S, Eng J, Woods SC, et al. The role of CNS glucagon-like peptide-1 (7-36) amide receptors in mediating the visceral illness effects of lithium chloride. *J Neurosci*. 2000; 20:1616–21. [PubMed: 10662851]
116. Baird JP, Travers SP, Travers JB. Integration of gastric distension and gustatory responses in the parabrachial nucleus. *Am J Physiol Regul Integr Comp Physiol*. 2001; 281:R1581–93. [PubMed: 11641131]
117. Karimnamazi H, Travers SP, Travers JB. Oral and gastric input to the parabrachial nucleus of the rat. *Brain Res*. 2002; 957:193–206. [PubMed: 12445962]
118. Rowland NE, Crews EC, Gentry RM. Comparison of Fos induced in rat brain by GLP-1 and amylin. *Regul Pept*. 1997; 71:171–4. [PubMed: 9350975]
119. Baraboi ED, Michel C, Smith P, Thibaudeau K, Ferguson AV, Richard D. Effects of albumin-conjugated PYY on food intake: the respective roles of the circumventricular organs and vagus nerve. *Eur J Neurosci*. 2010; 32:826–39. [PubMed: 20646064]
120. Swank MW, Bernstein IL. c-Fos induction in response to a conditioned stimulus after single trial taste aversion learning. *Brain Res*. 1994; 636:202–8. [PubMed: 8012803]
121. Maniscalco JW, Kreisler AD, Rinaman L. Satiation and stress-induced hypophagia: examining the role of hindbrain neurons expressing prolactin-releasing Peptide or glucagon-like Peptide 1. *Front Neurosci*. 2012; 6:199. [PubMed: 23346044]
122. Baird JP, Travers JB, Travers SP. Parametric analysis of gastric distension responses in the parabrachial nucleus. *Am J Physiol Regul Integr Comp Physiol*. 2001; 281:R1568–80. [PubMed: 11641130]
123. Li BH, Rowland NE. Effects of vagotomy on cholecystokinin- and dexfenfluramine-induced Fos-like immunoreactivity in the rat brain. *Brain Res Bull*. 1995; 37:589–93. [PubMed: 7670882]
124. Degen L, Oesch S, Casanova M, Graf S, Ketterer S, Drewe J, et al. Effect of peptide YY3-36 on food intake in humans. *Gastroenterology*. 2005; 129:1430–6. [PubMed: 16285944]
125. Labouesse MA, Stadlbauer U, Weber E, Arnold M, Langhans W, Pacheco-Lopez G. Vagal afferents mediate early satiation and prevent flavour avoidance learning in response to intraperitoneally infused exendin-4. *J Neuroendocrinol*. 2012; 24:1505–16. [PubMed: 22827554]
126. Lutz TA. Pancreatic amylin as a centrally acting satiating hormone. *Curr Drug Targets*. 2005; 6:181–9. [PubMed: 15777188]
127. Gieroba ZJ, Blessing WW. Fos-containing neurons in medulla and pons after unilateral stimulation of the afferent abdominal vagus in conscious rabbits. *Neuroscience*. 1994; 59:851–8. [PubMed: 7914681]
128. Lamprecht R, Dudai Y. Differential modulation of brain immediate early genes by intraperitoneal LiCl. *Neuroreport*. 1995; 7:289–93. [PubMed: 8742472]
129. Gaykema RP, Daniels TE, Shapiro NJ, Thacker GC, Park SM, Goehler LE. Immune challenge and satiety-related activation of both distinct and overlapping neuronal populations in the

- brainstem indicate parallel pathways for viscerosensory signaling. *Brain Res.* 2009; 1294:61–79. [PubMed: 19646973]
130. Castex N, Fioramonti J, Ducos de Lahitte J, Luffau G, More J, Bueno L. Brain Fos expression and intestinal motor alterations during nematode-induced inflammation in the rat. *Am J Physiol.* 1998; 274:G210–6. [PubMed: 9458792]
 131. De Jonghe BC, Horn CC. Chemotherapy agent cisplatin induces 48-h Fos expression in the brain of a vomiting species, the house musk shrew (*Suncus murinus*). *Am J Physiol Regul Integr Comp Physiol.* 2009; 296:R902–11. [PubMed: 19225146]
 132. Horn CC, De Jonghe BC, Matyas K, Norgren R. Chemotherapy-induced kaolin intake is increased by lesion of the lateral parabrachial nucleus of the rat. *Am J Physiol Regul Integr Comp Physiol.* 2009; 297:R1375–82. [PubMed: 19710391]
 133. Wu Q, Zheng R, Srisai D, McNight GS, Palmiter RD. The NR2B subunit of the NMDA glutamate receptor in the parabrachial nucleus regulates appetite. *PNAS.* 2013 In Press.
 134. Wu Q, Boyle MP, Palmiter RD. Loss of GABAergic signaling by AgRP neurons to the parabrachial nucleus leads to starvation. *Cell.* 2009; 137:1225–34. [PubMed: 19563755]
 135. Wu Q, Clark MS, Palmiter RD. Deciphering a neuronal circuit that mediates appetite. *Nature.* 2012; 483:594–7. [PubMed: 22419158]
 136. Wu Q, Palmiter RD. GABAergic signaling by AgRP neurons prevents anorexia via a melanocortin-independent mechanism. *Eur J Pharmacol.* 2010
 137. Carter ME, Soden ME, Zweifel LS, Palmiter RD. Genetic identification of a neural circuit that suppresses appetite. *Nature (London).* 2013
 138. Palmiter RD. New game for hunger neurons. *Nat Neurosci.* 2012; 15:1060–1. [PubMed: 22837033]
 139. Aponte Y, Atasoy D, Sternson SM. AGRP neurons are sufficient to orchestrate feeding behavior rapidly and without training. *Nat Neurosci.* 2011; 14:351–5. [PubMed: 21209617]
 140. Atasoy D, Betley JN, Su HH, Sternson SM. Deconstruction of a neural circuit for hunger. *Nature.* 2012; 488:172–7. [PubMed: 22801496]
 141. Sternson SM. Hypothalamic survival circuits: blueprints for purposive behaviors. *Neuron.* 2013; 77:810–24. [PubMed: 23473313]
 142. Ogden J, Clementi C, Aylwin S. The impact of obesity surgery and the paradox of control: A qualitative study. *Psychol Health.* 2006; 21:273–93. [PubMed: 21985121]
 143. Ogden J, Avenell S, Ellis G. Negotiating control: patients' experiences of unsuccessful weight-loss surgery. *Psychol Health.* 2011; 26:949–64. [PubMed: 21827354]
 144. Volkow ND, Wang GJ, Fowler JS, Telang F. Overlapping neuronal circuits in addiction and obesity: evidence of systems pathology. *Philos Trans R Soc Lond B Biol Sci.* 2008; 363:3191–200. [PubMed: 18640912]
 145. DelParigi A, Chen K, Salbe AD, Hill JO, Wing RR, Reiman EM, et al. Successful dieters have increased neural activity in cortical areas involved in the control of behavior. *Int J Obes (Lond).* 2007; 31:440–8. [PubMed: 16819526]
 146. Hare TA, Camerer CF, Rangel A. Self-control in decision-making involves modulation of the vmPFC valuation system. *Science.* 2009; 324:646–8. [PubMed: 19407204]
 147. Uher R, Yoganathan D, Mogg A, Eranti SV, Treasure J, Campbell IC, et al. Effect of left prefrontal repetitive transcranial magnetic stimulation on food craving. *Biol Psychiatry.* 2005; 58:840–2. [PubMed: 16084855]
 148. Goldman RL, Canterberry M, Borckardt JJ, Madan A, Byrne TK, George MS, et al. Executive control circuitry differentiates degree of success in weight loss following gastric-bypass surgery. *Obesity (Silver Spring).* 2013; 21:2189–96. [PubMed: 24136926]
 149. Goldman VJ, Santos M, Shayegan M, Cunningham JE. Evidence for two-dimensional quantum Wigner crystal. *Phys Rev Lett.* 1990; 65:2189–92. [PubMed: 10042476]

Table 1

Differences in measurements of energy expenditure and intake at different time points after bariatric surgeries between human and rodent models

		Human	Rodent
Energy expenditure	Weight loss phase (1-6 months in humans and 1-3 weeks in rodents)	↓ ~16, 21, and 21% in BMR at 1, 3, and 6 months, respectively after RYGB ¹ ↓ ~24% in SMR at 3 months after VBG ² ↓ ~13 and 16% in REE at 3 and 6 months, respectively after RYGB ³ ↓ ~2, 8, and 7% in REE at 1.5, 3, and 6 months, respectively after RYGB ⁴	NA
	Weight maintenance/regain phase (after 6 months in humans and 3 weeks in rodents)	↓ ~20% at 12 months after RYGB ¹ ↓ ~28% in SMR at 12 months after VBG ² ↓ ~19% at 12 months after RYGB ³ ↓ ~8, 9 and 7% in REE at 1, 1.5 and 2 years, respectively, after RYGB ⁴ ↓ ~25% in REE at 14±2 months after RYGB ⁷	↑ ~5% in TEE at 6 weeks after RYGB in rats vs. sham-operated ad lib fed, ↑ ~13% in TEE at 11 weeks vs. sham-operated weight-matched ⁸ ↑ ~18 and 30% in REE at 12-15 weeks after RYGB in rats vs. sham-operated ad lib fed and vs. sham-operated weight-matched, respectively ⁹ ↑ ~26% in TEE at 8 weeks after RYGB in mice ¹⁰ ↑ ~22% in REE at 8 weeks after RYGB in mice ¹¹
Energy intake	Weight loss phase	↓ ~74, 70 and 63% at 1.5, 3 and 6 months, respectively, after RYGB ⁴ ↓ ~58% at 6 weeks after RYGB ²⁸ ↓ ~66% and 63% at 6 months after RYGB and VBG, respectively ²⁹ ↓ ~46% at 6 months after RYGB ³⁰ ↓ ~80 and 67% at 2 and 8 weeks ²⁹	↓ until ~4 weeks after VSG ³⁴ ↔ at 3 weeks after RYGB in rats ³⁷ ↑ ~9% between day 6 and 13 after RYGB in mice ³⁹
	Weight maintenance/regain phase	↓ ~58, 52, and 47% at 1, 1.5, and 2 years, respectively, after RYGB ⁴ ↓ ~29 and 21% at 2 and 10 years, respectively after RYGB, VSG, and GB ²⁷ ↓ ~43 and 34% at 1 and 2 year, respectively after RYGB ²⁸ ↓ ~57, 52, 49 and 47% at 1, 1.5, 2 and 3 years after RYGB ²⁹ ↓ ~33% at 1 year after RYGB ³⁰	↓ ~17% over 5 months after RYGB in rats ⁹ ↔ after 4 weeks after VSG in rats ³⁴ ↔ at 4 weeks after RYGB in mice ³⁶ ↔ at 14 weeks after RYGB in rats ³⁷ ↓ ~7% over 60 days after RYGB in mice ³⁹

Table 2

Similarities and differences in measurements of food hedonics after bariatric surgeries in humans and rodent models

		Human	Rodent
Food Hedonics	Intake of calorically dense vs. less dense foods	↓~27, 16, and 10% at 6 weeks, 1 year, and 2 years after RYGB, respectively ⁷³	↓~12% at 16 days after RYGB in rats ³⁵ ↓~24, 24, and 27% at days 22-40, 41-80, and 81-150 after RYGB in rats, respectively ³⁸ ↓ 5 weeks after VSG in rats ⁷⁴ No differences after VSG, ↓ after RYGB in rats at 8 weeks ⁷⁵ ↓10 days after RYGB in rats ⁷⁶
	Changes in taste sensitivity	↑ sour and bitter stimuli, ↓ sweet and salty after RYGB ⁷⁷ ↑ sweet stimuli after RYGB ^{78, 79} ↓ sweet stimuli after RYGB ⁸⁰	NA
	Changes in taste preference	No difference in sweet taste preference after RYGB ⁷⁹ ↓ in taste preference for sucrose after RYGB ⁸⁰	↓ in taste preference for sucrose after RYGB in rats ^{79, 81, 82}
	Changes in 'liking'	↓ liking of high vs. low-calorie foods after RYGB ⁹⁹	↓ liking of high vs. low-calorie foods (concentrations of sucrose or corn oil) after RYGB in rats ⁸⁷
	Changes in 'wanting'	↓ wanting of high vs. low-calorie foods after RYGB ^{93, 95, 99} No difference in wanting of high-fat foods after RYGB ⁹⁴	↑ wanting of high-fat foods in diet-induced obesity rats after RYGB ^{87, 87}