Published in final edited form as:

Fertil Steril. 2010 February ; 93(2): 382–390. doi:10.1016/j.fertnstert.2008.10.061.

Racial and ethnic disparities in assisted reproductive technology outcomes in the United States

Victor Y. Fujimoto, M.D.^a, Barbara Luke, Sc.D., M.P.H.^b, Morton B. Brown, Ph.D.^c, Tarun Jain, M.D.^d, Alicia Armstrong, M.D.^e, David A. Grainger, M.D., M.P.H.^f, and Mark D. Hornstein, M.D.^g a Society for Assisted Reproductive Technology Writing Group

^aDepartment of Obstetrics, Gynecology, and Reproductive Sciences, University of California at San Francisco, San Francisco, California ^bDepartment of Obstetrics, Gynecology, and Reproductive Biology and Department of Epidemiology, Michigan State University, East Lansing, Michigan ^cDepartment of Biostatistics, School of Public Health, University of Michigan, Ann Arbor, Michigan ^dDivision of Reproductive Endocrinology and Infertility, Department of Obstetrics and Gynecology, University of Illinois at Chicago, Chicago, Illinois ^eProgram in Reproductive and Adult Endocrinology, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland ^fCenter for Reproductive Medicine and Department of Obstetrics and Gynecology, University of Kansas School of Medicine, Wichita, Kansas ^gDepartment of Obstetrics, Gynecology and Reproductive Biology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts

Abstract

Objective—To evaluate ethnic differences in assisted reproductive technology (ART) outcomes in the United States.

Design—Historical cohort study.

Setting—Clinic-based data.

Patient(s)—A total of 139,027 ART cycles from the Society for Assisted Reproductive Technology Clinic Outcome Reporting System online database for 2004–2006, limited to white, Asian, black, and Hispanic women.

Intervention(s)—None.

Main Outcome Measure(s)—Logistic regression was used to model the odds of pregnancy and live birth; among singletons and twins, the odds of preterm birth and fetal growth restriction. Results are presented as adjusted odds ratios, with white women as the reference group.

Reprint requests: Victor Y. Fujimoto, M.D., UCSF Center for Reproductive Health, 2356 Sutter Street, 7th Floor, San Francisco, CA 94115-0916 (FAX: 415-353-3065; fujimotov@obgyn.ucsf.edu).

M.D.H. is on the medical advisory committee for WIN Fertility. V.Y.F. is on the medical advisory board for EMD Serono and has received support from IBSA. B.L. has nothing to disclose. M.B.B. has nothing to disclose. T.J. has nothing to disclose. A.A. has nothing to disclose. D.A.G. has nothing to disclose.

Presented at the 64th Annual Meeting of the American Society for Reproductive Medicine, November 11–13, 2008, San Francisco, California.

Result(s)—The odds of pregnancy were reduced for Asians (0.86), and the odds of live birth were reduced for all groups: Asian (0.90), black (0.62), and Hispanic (0.87) women. Among singletons, moderate and severe growth restriction were increased for all infants in all three minority groups (Asians [1.78, 2.05]; blacks [1.81, 2.17]; Hispanics [1.36, 1.64]), and preterm birth was increased among black (1.79) and Hispanic women (1.22). Among twins, the odds for moderate growth restriction were increased for infants of Asian (1.30) and black women (1.97), and severe growth restriction was increased among black women (3.21). The odds of preterm birth were increased for blacks (1.64) and decreased for Asians (0.70).

Conclusion(s)—There are significant disparities in ART outcomes according to ethnicity.

Keywords

Ethnic disparities; racial disparities; Asian; black; Hispanic; ART outcomes; IVF outcomes; IVF pregnancy rates; IVF live birth rates; prematurity; growth restriction

Patient demographics are known to influence IVF pregnancy success but have largely focused on maternal age associations (1). There is an increasing body of evidence with the acknowledgment that ethnic disparities exist in reproductive outcomes spanning the obstetrical and gynecologic populations. With respect to IVF outcomes, several studies have found reduced pregnancy rates in infertile populations consisting of Asian and black women (2–6). The lower live birth rates in both Asian and black populations exist despite similar numbers of embryos transferred (2–5). However, other studies comparing IVF outcomes between black and white women have not shown such a disparity (7–9). Several fecundity-based studies have also demonstrated differences in ethnic groups, although these studies are confounded by potential region- and culture-specific differences in populations, independent of ethnicity (10, 11).

Recent studies support increased rates of small-for-gestational-age infants, preterm delivery, and perinatal morbidity associated with the IVF process compared with spontaneously conceived pregnancies (12–16). However, no study to date has addressed potential ethnic differences in perinatal outcomes within infertility cohorts of white, Asian, black, and Hispanic women after IVF treatment.

In an obstetrical population, Indian and Pakistani women had higher risks of low birth weight at term compared with other subpopulations of Asian and Pacific Islander women; however, no comparisons of Asian and Pacific Islander perinatal outcomes with a white population have been published (17). Other studies have also demonstrated low birth weight in Asian populations (18–21). Further studies describing the relationship between ethnicity and birth weight differences reveal reduced birth weights in black and Hispanic populations (22–29).

Hence, we hypothesized that differences in live birth rates and perinatal outcomes would exist within three minority ethnic groups in the United States as defined by the U.S. Census Bureau (30). Using the Society for Assisted Reproductive Technology Clinic Outcome Reporting System (SART-CORS) national database, we conducted this data analysis to identify such differences.

MATERIALS AND METHODS

The data source for this study was the SART-CORS database, which contains comprehensive data from more than 90% of all reporting clinics performing more than 90% of the ART cycles in the United States and included 391 clinics in 2004, 394 in 2005, and 392 in 2006. This database contains data collected and verified by SART, which are reported to the Centers for Disease Control and Prevention in compliance with the Fertility Clinic Success Rate and Certification Act of 1992 (Public Law 102-493, October 24, 1992). The study was reviewed by the Committee for the Protection of Human Subjects at Brigham and Women's Hospital in Boston, Massachusetts, and allowed exemption from approval. This article was written in conjunction with the SART Research Committee.

The study population was limited to all ART cycles using non-donor oocytes and partner semen among women of the four major racial and ethnic groups as reported by participating clinics: white, Asian, black, and Hispanic. Because no specific instructions for SART reporting of ethnicity are given, there may be reporting biases of specific race or ethnicity that can vary from clinic to clinic and patient to patient. According to U.S. Census Bureau categorization, white, Asian, and black are considered racial groups, whereas Hispanic is considered an ethnic category of white. Dependent variables included the odds of pregnancy (presence of gestational sac on early ultrasound) as the treatment outcome, live birth as the pregnancy outcome, and, among singleton and twin live births, the odds of prematurity (<29 weeks, <32 weeks, and <37 weeks) and growth restriction (birth-weight-for-gestation z scores as moderate [<-1] or severe [<-2]), adjusting for other significant factors. White women were the reference group, and each ethnic group was compared separately. A single-ethnicity birth weight reference (Canadian live births, 1994–1996 [31]) was used to calculate birth weight z scores. Plurality-specific birth weight z scores were calculated to evaluate adequacy of weight for age (31, 32), as recommended by Land (33), and modeled as continuous and categorical variables (<-1.0 and <-2.0).

Birth weights at each gestational age are normally distributed, with a reference mean of zero and an SD of 1. A z score (or SD score) is the deviation of the value for an individual from the mean value of the reference population divided by the SD for the reference population. Z scores have a direct relationship with percentiles, with z scores from -1 to $+1$ and from -2 to $+2$ representing 68% and 95%, respectively, of the population distribution. The z score is useful to describe how far the observed birth weight for gestational age is from its expected value. For this study, we used birth weight z scores of -1 and $+2$, corresponding to the 16th and 2.5th percentiles, respectively.

Maternal demographic factors, reproductive history, ART cycle-specific parameters, and ART treatment and pregnancy outcomes were compared across the four ethnic groups using the χ^2 test and analysis of variance. Logistic regression analyses were used for dichotomized outcomes. Models were adjusted for maternal age, number of embryos transferred, and the infertility diagnoses of male factor, endometriosis, polycystic ovarian syndrome, diminished ovarian reserve, tubal factors, and other factors. Models of birth weight z scores and prematurity outcomes were limited to pregnancies of ≥ 154 days (22 weeks) and >300 g.

Data were analyzed using the Statistical Package for the Social Sciences, version 16.0 (SPSS, Chicago, IL).

RESULTS

The four ethnic groups differed significantly in their age distribution, with Asian and black women being older and Hispanic women younger than their white counterparts (Table 1). Asian women were more likely and black women less likely to be nulligravidas. The four groups also differed in their infertility diagnoses. Black women were less likely to be diagnosed with endometriosis, polycystic ovary syndrome, and unexplained factors and more likely to have tubal or uterine factors. Hispanic women were more likely to have tubal factors and less likely to have unexplained factors. Other demographic differences, including prior term birth, prior preterm birth, and prior spontaneous abortion, are also described in Table 1. Although data regarding sperm parameters were not available in the dataset, the relative percentages of those patients undergoing intracytoplasmic sperm injection or conventional insemination (IVF) are also listed for each racial or ethnic group in Table 1. Asian women tended to have fewer embryos transferred, whereas black and Hispanic women had more embryos transferred compared with white women (Table 1).

Treatment outcomes also varied significantly across the four groups. Black women were more likely to develop ovarian hyperstimulation syndrome (any and severe) and were less likely to achieve a clinical pregnancy; however, the absolute risk of ovarian hyperstimulation syndrome remains low. Among those who did become pregnant, black women were least likely to have a live birth (Table 2). We also examined multiple pregnancy rates by ethnicity (Table 2). Asian and black women were also less likely than white or Hispanic women to become pregnant with twins, but among those who did achieve a pregnancy, their live birth rates were comparable to those for white women, without an increase in pregnancy loss. Length of gestation and birth weight by plurality also differed significantly across the four groups, as shown in Table 3. Infants of black women averaged the shortest gestations and lowest birth weights, whereas infants of Asian women averaged the longest gestations and highest birth weights within each plurality. Infants of Asian women averaged the lowest singleton birth weight z scores, and black infants the lowest twin birth weight z scores.

The results of the logistic regression models are shown in Table 4. Asian women were 14% less likely to achieve a pregnancy as the treatment outcome. All three minority groups were less likely to have a live birth as the pregnancy outcome compared with white women: 10% less for Asian, 13% less for Hispanic, and 38% less for black women. Among women with singleton live births, black women were approximately four-, three-, and twofold as likely to deliver very early (<29 weeks), early (<32 weeks), or preterm (<37 weeks), respectively, compared with white women; Hispanic women were approximately 22% more likely to deliver preterm. Moderate and severe growth restriction were significantly more likely among the singleton infants of all three minority groups. Among women with twin births, black women were more likely and Asian women less likely to deliver preterm. Moderate growth restriction was more likely among Asian and black twin infants, and severe growth restriction among black twins.

DISCUSSION

The major findings of our study of the U.S. ART patient population are [1] the reduced clinical pregnancy rate in Asian women compared with white women, [2] the reduced live birth rates in all three racial and ethnic minority groups compared with white women, [3] the increased preterm delivery rate in black and Hispanic women compared with white women, and [4] the increase in fetal growth restriction seen in all three minority groups of women compared with white women. Our findings confirm that Asian and black women have reduced live birth rates after ART treatment when compared with white women and demonstrate for the first time that Hispanic women also have reduced live birth rates after ART.

The study of racial and ethnic disparities is complicated by a myriad of factors: social, cultural, nutritional, anthropometric, environmental, physical, metabolic, and genetic. No single study can address all of these factors, and any findings are limited to factors that can be measured or assessed. Although clinical pregnancy rates were similar between black, Hispanic, and white women, Asian women had a lower clinical pregnancy rate compared with white women in our study. Black and Hispanic women had significantly higher rates of fetal loss that subsequently reduced their live birth rates significantly compared with white women. Finally, all three minority ethnic groups had higher rates of moderate and severe growth restriction at delivery compared with white women, independent of gestational age. Collectively, these data describe significant disparities in the reproductive health outcomes of minority women undergoing ART treatment in the United States.

Our confirmation of reduced clinical pregnancy rates in Asian women compared with white women confirms disparities reported in prior published work (3, 4). Our finding of a reduction in pregnancy success in Asian women is lower than previously reported by Purcell et al. (4), representing SART National data from 1999–2000, for which the odds ratio for clinical pregnancy rate in Asian women was 0.71. It is important to note that only first-cycle IVF data were analyzed by Purcell et al., whereas our dataset includes all IVF cycles generated over the 3-year reporting period (4). The higher odds ratio seen in our study may be explained by several possible etiologies: [1] reporting bias influenced by demographic differences, and [2] repeated-measure bias represented by a multiple-cycle dataset. In 2005 SART mandated the reporting of female race and ethnicity data for all IVF cycles generated by reporting clinics. It is possible that this change in reporting could have altered the ethno-demographic distribution of reporting Asian populations in the United States and thus influenced the statistical differences in pregnancy outcomes. Another possible explanation for the difference may be that Asian and white women have comparable outcomes after multiple IVF treatment cycles. The possible influence of diagnosis on Asian reproductive outcomes is demonstrated by Palep-Singh et al. (3), who reported that Asian women with polycystic ovarian syndrome undergoing IVF had reduced clinical pregnancy and live birth outcomes compared with white women, whereas Asian women with tubal factor infertility had similar clinical pregnancy and live birth outcomes as white women (3). Various potential mechanisms could be influencing the ability of Asian women to conceive with ART. It is known that sociocultural factors reduce health care utilization in Asian populations, which has been demonstrated in an infertility population as well (34). Although

other studies have not shown differences in ovarian stimulation characteristics (4), the concept that Asian women may suffer from an acceleration in ovarian aging has been proposed to explain the disparity seen in Asian women undergoing IVF (35). Genetic differences have been demonstrated in Asian populations, which may be relevant in IVF success rates (36). Finally, environmental exposures may also partially contribute to the disparities seen in clinical pregnancy and live birth rates seen in Asian women (37).

Contrary to the findings of prior studies (2, 5, 6), our dataset did not demonstrate a lower clinical pregnancy rate for black women after adjusting for maternal age, number of embryos transferred, and infertility diagnoses. However, our observation of reduced live birth rates associated with black women is consistent with prior studies (2, 5, 6). Feinberg et al. (2) attributed the reduced pregnancy rate in part to the increased incidence of uterine fibroids present in the black population (2). After controlling for this factor, that study found that black women experienced a higher spontaneous miscarriage rate than white women (2). The increased fetal loss rate was confirmed in our study, with black women experiencing a 21.8% miscarriage rate, significantly higher than the rates seen in the other three racial or ethnic groups. To what extent uterine fibroids are contributing to this observation in our black population is unknown because this SART dataset did not include the presence or absence of fibroids and other contributing factors, such as the risk of aneuploidy (2). Black women also have more tubal disease, and it is well accepted that hydrosalpinges contribute to a lower IVF success rate (38, 39). The increased fetal loss rate resulted in the lowest live birth rate in the black population in our study. Although several studies have not confirmed this disparity in live birth outcomes within the black population, these studies were limited by sample size (7–9). Additionally, Asian and Hispanic live birth rates were also significantly lower than white live birth rates in our study after IVF treatment. The observation of a lower Hispanic live birth rate and higher fetal loss rate has not been previously demonstrated with IVF treatment.

There are several limitations to this study. It is important to clarify that these populations do not represent the overall reproductive-aged populations in the United States for each ethnic group, because a disproportionately higher socioeconomic status exists for each ethnicity in our patient population (40, 41). Unfortunately, specific socioeconomic data were not available for analyses within this dataset. Another limitation of this dataset is the lack of body mass indices, which have been consistently associated with pregnancy outcome disparities (42–47). Finally, we acknowledge that a significant limitation of this study is the potential variability in the reporting and classification of race and ethnicity, which may vary from clinic to clinic. Although SART requires ethnicity reporting with each registered cycle, there are no specific guidelines for the documentation of clinic-specific race and ethnicity. That said, race and ethnicity may be a proxy for some other factors that remain unmeasured in our database. The racial and ethnic minority disparities that exist within this dataset are hypothesis-generating and raise questions regarding the role of environmental and genetic susceptibility factors that may be influencing live birth rates in ART.

Our observations of increased fetal growth restriction in all three racial and ethnic minority groups, independent of gestational number, are interesting. There is precedence for the increased growth restriction seen in Asian, black, and Hispanic infants (23). Nationally,

black women have higher rates of preterm delivery and low-birth-weight births compared with white and Hispanic women. Whereas nationally, Hispanic women did not have low-birth-weight births, a study based in California found that both foreign-born and U.S.-born Hispanic women had lower-birth-weight infants compared with white women (24). In that study, it was also demonstrated that Asian women had lower-birth-weight infants compared with white women (24). The etiology of lower birth weight in minority populations after adjustment for maternal age, gestational age, and limited to singleton births is unknown. Although this study is based on an IVF population, the disparities cannot be attributed to the contributing effect of the IVF process on reduced singleton birth weight (12–16). There is increasing evidence that environmental exposures (e.g., organic solvents in the workplace) may negatively influence the birth weight of infants born to exposed women (48). It is possible that various environmental exposures may differ within race-based populations that may ultimately influence birth outcomes.

Intrauterine growth and birth weight vary with well-established maternal characteristics, including prepregnancy or first-trimester body weight, height, parity, ethnicity, and gestational weight gain, as well as by the gender of the baby (49–53). Prepregnancy weight, height, and gestational weight gain were factors not available in the SART-CORS database; also lacking were data on pregnancy complications (preeclampsia and gestational diabetes), which could also have influenced fetal growth and length of gestation (54). In addition, there are known racial and ethnic differences in visceral fat distribution (55, 56), as well as carbohydrate and insulin metabolism (57–63). Research has indicated an even greater level of insulin resistance within certain ethnic groups and specific infertility diagnoses (64).

Another potential limitation in our results is that we used a single-ethnicity birth weight reference (Canadian live births, 1994–1996 [31]) to calculate birth weight z scores, representing predominately white populations. The use of ethnic-specific birth weight references may have altered our findings, particularly moderate growth restriction (birth weight z scores <-1) but probably not severe growth restriction (birth weight z scores <-2). The few birth weight references that are ethnic-specific (black and white [29], Mexican American [27], Hispanic [22], and Asian [18–21]) fail to account for other important maternal characteristics, such as parity, height, and prepregnancy weight.

In summary, our findings confirm the racial and ethnic disparities seen in prior studies with respect to pregnancy rates and live birth rates resulting from ART treatment in the United States. In addition to the current concern regarding the use of ART and reduced singleton birth weight, the differences in birth weight seen in this study between the four major racial and ethnic groups further illustrates the diversity of influences, both genetic and environmental, that may impact children born from ART treatment. This study further demonstrates the need to document racial and ethnic origins within the framework of the infertile patient evaluation for patient education purposes. More accurate and detailed recording of race and ethnicity would also provide further opportunity to understand better the ART outcome disparities of an increasingly diverse population here in the United States. Further studies are clearly needed to explore the etiologies of these disparities within all minority racial and ethnic groups described in this study.

Acknowledgments

SART thanks all of its members for providing clinical information to the SART-CORS database for use by patients and researchers. Without the efforts of our members, this research would not have been possible. The authors thank the SART Research Review Committee for its review of the manuscript.

Supported by the Society for Assisted Reproductive Technology.

References

1. Lalwani S, Timmreck L, Friedman R, Penzias A, Alper M, Reindollar RH. Variations in individual physician success rates within an in vitro fertilization program might be due to patient demographics. *Fertil Steril*. 2004; 81:944–6. [PubMed: 15066443]
2. Feinberg EC, Larsen FW, Catherino WH, Zhang J, Armstrong AY. Comparison of assisted reproductive technology utilization and outcomes between Caucasian and African American patients in an equal-access-to-care setting. *Fertil Steril*. 2006; 85:888–94. [PubMed: 16580370]
3. Palep-Singh M, Picton HM, Vrotsou K, Maruthini D, Balen AH. South Asian women with polycystic ovary syndrome exhibit greater sensitivity to gonadotropin stimulation with reduced fertilization and ongoing pregnancy rates than their Caucasian counterparts. *Eur J Obstet Gynecol Reprod Biol*. 2007; 134:202–7. [PubMed: 17367914]
4. Purcell K, Schembri M, Frazier LM, Rall MJ, Shen S, Croughan M, et al. Asian ethnicity is associated with reduced pregnancy outcomes after assisted reproductive technology. *Fertil Steril*. 2007; 87:297–302. [PubMed: 17081529]
5. Seifer DB, Frazier LM, Grainger DA. Disparity in assisted reproductive technologies outcomes in black women compared with white women. *Fertil Steril*. 2008; 90:1701–10. [PubMed: 17980873]
6. Sharara FI, McClamrock HD. Differences in in vitro fertilization (IVF) outcome between white and black women in an inner-city, university-based IVF program. *Fertil Steril*. 2000; 73:1170–3. [PubMed: 10856477]
7. Bendikson K, Cramer DW, Vitonis A, Hornstein MD. Ethnic background and in vitro fertilization outcomes. *Int J Gynaecol Obstet*. 2005; 88:342–6. [PubMed: 15733901]
8. Dayal MB, Gindoff P, Dubey A, Spitzer TLB, Bergin A, Peak D, et al. Does ethnicity influence in vitro fertilization (IVF) birth outcomes? *Fertil Steril*. In press.
9. Nichols JE Jr, Higdon HL 3rd, Crane MM 4th, Boone WR. Comparison of implantation and pregnancy rates in African American and white women in an assisted reproductive technology practice. *Fertil Steril*. 2001; 76:80–4. [PubMed: 11438323]
10. Jensen TK, Slama R, Ducot B, Suominen J, Cawood EH, Andersen AG, et al. Regional differences in waiting time to pregnancy among fertile couples from four European cities. *Hum Reprod*. 2001; 16:2697–704. [PubMed: 11726598]
11. Tuntiseranee P, Olsen J, Chongsuvivatwong V, Limbutara S. Fecundity in Thai and European regions: results based on waiting time to pregnancy. *Hum Reprod*. 1998; 13:471–7. [PubMed: 9557860]
12. Helmerhorst FM, Perquin DAM, Donker D, Keirse MJNC. Perinatal outcome of singletons and twins after assisted conception: a systematic review of controlled studies. *BMJ*. 2004; 328:261–5. [PubMed: 14742347]
13. Jackson RA, Gibson KA, Wu YW, Croughan MS. Perinatal outcomes in singletons following in vitro fertilization: a meta-analysis. *Obstet Gynecol*. 2004; 103:551–63. [PubMed: 14990421]
14. Reddy UM, Wapner RJ, Rebar RW, Tasca RJ. Infertility, assisted reproductive technology, and adverse pregnancy outcomes: executive summary of a National Institute of Child Health and Human Development Workshop. *Obstet Gynecol*. 2007; 109:967–77. [PubMed: 17400861]
15. Schieve LA, Meikle SF, Ferre C, Peterson HB, Jeng G, Wilcox LS. Low and very low birth weight in infants conceived with use of assisted reproductive technology. *N Engl J Med*. 2002; 346:731–7. [PubMed: 11882728]
16. Shih W, Rushford DD, Bourne H, Garrett C, McBain JC, Healy DL, et al. Factors affecting low birthweight after assisted reproduction technology: difference between transfer of fresh and

cryopreserved embryos suggests an adverse effect of oocyte collection. *Hum Reprod.* 2008; 23:1644–53. [PubMed: 18442997]

17. Rao AK, Cheng YW, Caughey AB. Perinatal complications among different Asian-American subgroups. *Am J Obstet Gynecol.* 2006; 194:e39–41. [PubMed: 16579923]
18. Wang X, Guyer B, Paige DM. Differences in gestational age-specific birthweight among Chinese, Japanese and white Americans. *Int J Epidemiol.* 1994; 23:119–28. [PubMed: 8194906]
19. Wen SW, Kramer MS, Usher RH. Comparison of birth weight distributions between Chinese and Caucasian infants. *Am J Epidemiol.* 1995; 141:1177–87. [PubMed: 7771456]
20. Williams RL. Intrauterine growth curves: intra- and international comparisons with different ethnic groups in California. *Prev Med.* 1975; 4:163–72. [PubMed: 1153395]
21. Yip R, Li Z, Chong WH. Race and birth weight: the Chinese example. *Pediatrics.* 1991; 87:688–93. [PubMed: 2020515]
22. Aguilar T, Teberg AJ, Chan L, Hodgman J. Intrauterine growth curves of weight, length, and head circumference for a predominantly Hispanic infant population. *Public Health Rep.* 1995; 110:327–32. [PubMed: 7610225]
23. Alexander GR, Kogan M, Bader D, Carlo W, Allen M, Mor J. US birth weight/gestational age-specific neonatal mortality: 1995–1997 rates for whites, Hispanics, and blacks. *Pediatrics.* 2003; 111:e61–6. [PubMed: 12509596]
24. Fuentes-Afflick E, Hessol NA, Perez-Stable EJ. Maternal birthplace, ethnicity, and low birth weight in California. *Arch Pediatr Adolesc Med.* 1998; 152:1105–12. [PubMed: 9811289]
25. Gould JB, Madan A, Qin C, Chavez G. Perinatal outcomes in two dissimilar immigrant populations in the United States: a dual epidemiologic paradox. *Pediatrics.* 2003; 111:e676–82. [PubMed: 12777585]
26. Madan A, Palaniappan L, Urizar G, Wang Y, Fortmann SP, Gould JB. Sociocultural factors that affect pregnancy outcomes in two dissimilar immigrant groups in the United States. *J Pediatrics.* 2006; 148:341–6.
27. Overpeck MD, Hediger ML, Zhang JUN, Trumble AC, Klebanoff MA. Birth weight for gestational age of Mexican American infants born in the United States. *Obstet Gynecol.* 1999; 93:943–7. [PubMed: 10362159]
28. Singh GK, Yu SM. Adverse pregnancy outcomes: differences between US- and foreign-born women in major US racial and ethnic groups. *Am J Public Health.* 1996; 86:837–43. [PubMed: 8659659]
29. Zhang J, Bowes WA Jr. Birth-weight-for-gestational-age patterns by race, sex, and parity in the United States population. *Obstet Gynecol.* 1995; 86:200–8. [PubMed: 7617350]
30. Wendler D, Kington R, Madans J, Wye GV, Christ-Schmidt H, Pratt LA, et al. Are racial and ethnic minorities less willing to participate in health research? *PLoS Med.* 2006; 3:e19. [PubMed: 16318411]
31. Kramer MS, Platt RW, Wen SW, Joseph KS, Allen A, Abrahamowicz M, et al. A new and improved population-based Canadian reference for birth weight for gestational age. *Pediatrics.* 2001; 108:E35. [PubMed: 11483845]
32. Min SJ, Luke B, Gillespie B, Min L, Newman RB, Mauldin JG, et al. Birth weight references for twins. *Am J Obstet Gynecol.* 2000; 182:1250–7. [PubMed: 10819867]
33. Land JA. How should we report on perinatal outcome? *Hum Reprod.* 2006; 21:2638–9. [PubMed: 16829595]
34. Lamb JD, Purcell KJ, Modan AE, Vittinghoff E, Fujimoto VY. Asian ethnicity is associated with longer duration of infertility and decreased pregnancy rates following intrauterine insemination (IUI) [abstract]. *Fertil Steril.* 2007; 88:S260.
35. Gleicher N, Weghofer A, Li J, Barad D. Differences in ovarian function parameters between Chinese and Caucasian oocyte donors: do they offer an explanation for lower IVF pregnancy rates in Chinese women? *Hum Reprod.* 2007; 22:2879–82. [PubMed: 17855412]
36. Sudo S, Kudo M, Wada S, Sato O, Hsueh A, Fujimoto S. Genetic and functional analyses of polymorphisms in the human FSH receptor gene. *Mol Hum Reprod.* 2002; 8:893–9. [PubMed: 12356937]

37. Hightower JM, O'Hare A, Hernandez GT. Blood mercury reporting in NHANES: identifying Asian, Pacific Islander, Native American, and multiracial groups. *Environ Health Perspect.* 2006; 114:173–5. [PubMed: 16451850]
38. Green JA, Robins JC, Scheiber M, Awadalla S, Thomas MA. Racial and economic demographics of couples seeking infertility treatment. *Am J Obstet Gynecol.* 2001; 184:1080–2. [PubMed: 11349163]
39. Strandell A, Lindhard A, Waldenstrom U, Thorburn J. Hydrosalpinx and IVF outcome: cumulative results after salpingectomy in a randomized controlled trial. *Hum Reprod.* 2001; 16:2403–10. [PubMed: 11679529]
40. Jain T. Socioeconomic and racial disparities among infertility patients seeking care. *Fertil Steril.* 2006; 85:876–81. [PubMed: 16580368]
41. Jain T, Hornstein MD. Disparities in access to infertility services in a state with mandated insurance coverage. *Fertil Steril.* 2005; 84:221–3. [PubMed: 16009188]
42. Bellver J, Rossal LP, Bosch E, Zúñiga A, Corona JT, Meléndez F, et al. Obesity and the risk of spontaneous abortion after oocyte donation. *Fertil Steril.* 2003; 79:1136–40. [PubMed: 12738508]
43. Maheshwari A, Stofberg L, Bhattacharya S. Effect of overweight and obesity on assisted reproductive technology: a systematic review. *Hum Reprod Update.* 2007; 13:433–44. [PubMed: 17584821]
44. Sneed ML, Uhler ML, Grotjan HE, Rapisarda JJ, Lederer KJ, Beltsos AN. Body mass index: impact on IVF success appears age-related. *Hum Reprod.* 2008; 23:1835–9. [PubMed: 18503054]
45. Thum MY, El-Sheikhah A, Faris R, Parikh J, Wren M, Ogunyemi T, et al. The influence of body mass index to in-vitro fertilisation treatment outcome, risk of miscarriage and pregnancy outcome. *J Obstet Gynaecol.* 2007; 27:699–702. [PubMed: 17999296]
46. Veleva Z, Tiitinen A, Vilska S, Hyden-Granskog C, Tomas C, Martikainen H, et al. High and low BMI increase the risk of miscarriage after IVF/ICSI and FET. *Hum Reprod.* 2008; 23:878–84. [PubMed: 18281684]
47. Wei H-J, Young R, Kuo IL, Liaw C-M, Chiang H-S, Yeh C-Y. Abnormal preconception oral glucose tolerance test predicts an unfavorable pregnancy outcome after an in vitro fertilization cycle. *Fertil Steril.* 2008; 90:613–8. [PubMed: 17980878]
48. Ahmed P, Jaakkola JJK. Exposure to organic solvents and adverse pregnancy outcomes. *Hum Reprod.* 2007; 22:2751–7. [PubMed: 17725989]
49. Cogswell ME, Yip R. The influence of fetal and maternal factors on the distribution of birthweight. *Semin Perinatol.* 1995; 19:222–40. [PubMed: 7570074]
50. Gardosi J, Chang A, Kalyan B, Sahota D, Symonds EM. Customised antenatal growth charts. *Lancet.* 1992; 339:283–7. [PubMed: 1346292]
51. Gardosi J, Mongelli M, Wilcox M, Chang A. An adjustable fetal weight standard. *Ultrasound Obstet Gynecol.* 1995; 6:168–74. [PubMed: 8521065]
52. Rosenberg TJ, Garbers S, Chavkin W, Chiasson MA. Prepregnancy weight and adverse perinatal outcomes in an ethnically diverse population. *Obstet Gynecol.* 2003; 102:1022–7. [PubMed: 14672480]
53. World Health Organization. Maternal anthropometry and pregnancy outcomes. A WHO collaborative study. *Bull World Health Organ.* 1995; 73(Suppl):1–98.
54. Silva JK, Kaholokula JK, Ratner R, Mau M. Ethnic differences in perinatal outcome of gestational diabetes mellitus. *Diabetes Care.* 2006; 29:2058–63. [PubMed: 16936153]
55. Conway JM, Yanovski SZ, Avila NA, Hubbard VS. Visceral adipose tissue differences in black and white women. *Am J Clin Nutr.* 1995; 61:765–71. [PubMed: 7702017]
56. Kanaley JA, Giannopoulou I, Tillapaugh-Fay G, Nappi JS, Ploutz-Snyder LL. Racial differences in subcutaneous and visceral fat distribution in postmenopausal black and white women. *Metabolism.* 2003; 52:186–91. [PubMed: 12601630]
57. Green JR, Schumacher LB, Pawson IG, Partridge JC, Kretchmer N. Influence of maternal body habitus and glucose tolerance on birth weight. *Obstet Gynecol.* 1991; 78:235–40. [PubMed: 2067768]

58. Jiang X, Srinivasan SR, Radhakrishnamurthy B, Dalferes ER, Berenson GS. Racial (black-white) differences in insulin secretion and clearance in adolescents: the Bogalusa heart study. *Pediatrics*. 1996; 97:357–60. [PubMed: 8604270]
59. Kieffer E, Nolan G, Carman W, Sanborn C, Guzman R, Ventura A. Glucose tolerance during pregnancy and birth weight in a Hispanic population. *Obstet Gynecol*. 1999; 94:741–6. [PubMed: 10546721]
60. Manolio TA, Savage PJ, Burke GL, Liu KA, Wagenknecht LE, Sidney S, et al. Association of fasting insulin with blood pressure and lipids in young adults. The CARDIA study. *Arteriosclerosis*. 1990; 10:430–6. [PubMed: 2188641]
61. Nahum GG, Huffaker BJ. Racial differences in oral glucose screening test results: establishing race-specific criteria for abnormality in pregnancy. *Obstet Gynecol*. 1993; 81:517–22. [PubMed: 8459959]
62. Scholl TO, Chen X, Gaughan C, Smith WK. Influence of maternal glucose level on ethnic differences in birth weight and pregnancy outcome. *Am J Epidemiol*. 2002; 156:498–506. [PubMed: 12225997]
63. Velasquez-Mieyer PA, Cowan PA, Umpierrez GE, Lustig RH, Cashion AK, Burghen GA. Racial differences in glucagon-like peptide-1 (GLP-1) concentrations and insulin dynamics during oral glucose tolerance test in obese subjects. *Int J Obes Relat Metab Disord*. 2003; 27:1359–64. [PubMed: 14574347]
64. Kauffman RP, Baker VM, DiMarino P, Gimpel T, Castracane VD. Polycystic ovarian syndrome and insulin resistance in white and Mexican American women: a comparison of two distinct populations. *Am J Obstet Gynecol*. 2002; 187:1362–9. [PubMed: 12439532]

TABLE 1

Characteristics and therapy of women by race and ethnicity.

Characteristic	All Groups	Racial and ethnic groups				P value
		White	Asian	Black	Hispanic	
n	139,027	107,484	13,671	8,903	8,969	
% racial or ethnic	100	77.2	9.8	6.5	6.5	
Maternal age (y), mean (SD)	35.3 (4.6)	35.3 (4.6)	35.8 (4.6)	35.9 (4.7)	35.0 (4.8)	<.0001
<30	11.2	11.5	8.6	9.9	13.6	<.0001
30–34	31.3	31.7	30.1	26.5	31.9	
35–39	37.2	37.1	38.1	38.8	35.4	
40–44	19.0	18.4	21.4	23.2	17.7	
45	1.3	1.2	1.8	1.6	1.3	
Nulligravida	46.8	46.9	53.2	35.4	46.5	<.0001
No prior full-term birth	55.5	54.3	64.6	58.7	52.0	<.0001
No prior preterm birth	96.5	96.6	97.5	94.9	96.2	<.0001
No prior spontaneous abortion	48.8	48.6	50.3	46.2	51.5	<.0001
Infertility diagnosis						
Male factor	38.2	38.7	36.8	34.2	38.3	<.0001
Endometriosis	14.4	15.3	12.4	7.8	12.5	<.0001
Polycystic ovary syndrome	15.2	15.7	14.2	10.6	15.0	<.0001
Diminished ovarian reserve	16.8	16.7	19.0	14.9	15.8	<.0001
All tubal factors	21.1	18.5	17.8	46.7	32.0	<.0001
Uterine factors	5.2	4.5	5.8	13.1	5.3	<.0001
Other factors	14.2	14.4	15.4	12.4	12.3	<.0001
Unexplained factors	11.7	12.2	13.5	6.3	8.5	<.0001
No. of fresh embryos transferred						
1	8.9	8.7	11.2	9.0	8.0	<.0001
2	43.7	44.4	40.2	41.9	41.7	
3	30.3	30.4	28.7	30.9	30.6	
4	11.7	11.4	13.0	12.0	13.6	
5	5.4	5.1	7.0	6.2	6.1	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Characteristic	Racial and ethnic groups				P value
	All Groups	White	Asian	Black	Hispanic
ICSI (all or some)	70.0	70.0	70.0	67.6	71.9
Assisted hatching (all or some)	46.8	46.1	53.1	46.6	45.7

Note: Values are percentages unless otherwise noted. ICSI = intracytoplasmic sperm injection.
Fujimoto. Racial and ethnic disparities in ART outcomes. Fertil Steril 2010.

TABLE 2

Treatment outcomes by racial and ethnic group.

Characteristic	All groups	Racial and ethnic groups				P value
		White	Asian	Black	Hispanic	
n	139,027	107,484	13,671	8,903	8,969	
% racial or ethnic	100	77.2	9.8	6.5	6.5	
Complications						
Any hyperstimulation	1.6	1.6	1.3	2.7	1.6	<.0001
Severe hyperstimulation	0.4	0.4	0.3	0.9	0.4	<.0001
Treatment outcome						
Not pregnant	53.8	51.9	61.8	62.2	55.9	<.0001
Biochemical	6.8	7.1	6.4	5.0	5.9	
Ectopic	0.8	0.8	0.9	0.8	0.9	
Clinical intrauterine gestation	38.5	40.1	30.9	32.0	37.3	
Heterotopic	0.1	0.0	0.1	0.0	0.0	
Fetal hearts on ultrasound						
0	5.8	5.7	6.6	6.8	5.7	<.0001
1	61.1	60.9	63.6	61.9	59.8	
2	28.7	29.0	25.7	27.6	29.6	
3	4.0	4.0	3.9	3.1	4.2	
4	0.4	0.4	0.3	0.6	0.7	
Pregnancy Outcome (%)						
Livebirth	83.0	83.7	81.6	75.0	82.2	<.0001
Stillbirth or fetal loss	17.0	16.3	18.4	25.0	17.8	
Plurality of liveborn						
Pregnancies (n)	44,508	36,178	3,445	2,155	2,730	<.0001
Singleton	68.1	67.7	71.6	70.9	66.8	
Twins	29.5	29.9	26.8	27.0	30.0	
Triplets	2.3	2.3	1.6	2.1	2.9	
Quadruplets	0.1	0.1	0.0	0.0	0.3	

Note: Values are percentages unless otherwise noted.

Fujimoto. Racial and ethnic disparities in ART outcomes. *Fertil Steril* 2010.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

TABLE 3

Length of gestation and birth weight by plurality and racial and ethnic groups.

Characteristic	All groups	Racial and ethnic groups				P value
		White	Asian	Black	Hispanic	
n	44,508	36,178	3,445	2,155	2,730	
% racial or ethnic	100	81.3	7.7	4.8	6.1	
Length of gestation						
Singletons (d), mean (SD) ^a	259 (21)	260 (21)	262 (19)	254 (26)	257 (22)	<.0001
22–28 wk (%)	1.3	1.1	0.9	4.6	1.4	<.0001
29–32 wk (%)	1.9	1.8	2.0	3.5	2.0	
33–36 wk ^v	15.2	14.8	13.9	21.0	17.7	
37 wk (%)	81.6	82.3	83.1	70.9	78.9	
Twins (d), mean (SD) ^a	243 (21)	243 (21)	247 (19)	236 (26)	241 (22)	<.0001
22–28 wk (%)	5.5	5.2	3.3	12.0	7.4	<.0001
29–32 wk (%)	12.6	12.5	11.7	15.6	12.3	
33–36 wk (%)	59.0	59.4	54.6	56.3	59.7	
37 wk (%)	23.0	22.9	30.4	16.1	20.6	
Triplets (d), mean (SD) ^a	224 (21)	224 (20)	230 (21)	217 (21)	222 (20)	.02
22–28 wk (%)	14.3	14.4	5.6	28.6	11.7	.008
29–32 wk (%)	40.0	39.1	35.2	42.9	50.6	
33–36 wk (%)	44.5	45.5	55.6	28.6	35.1	
37 wk (%)	1.2	1.0	3.7	0.0	2.6	
Birth weight (g), mean (SD) ^b						
Singletons	2,900 (745)	2,966 (743)	2,906 (678)	2,735 (816)	2,861 (766)	<.0001
Twins	2,271 (603)	2,287 (599)	2,281 (545)	2,047 (685)	2,189 (618)	<.0001
Triplets	1,678 (501)	1,688 (509)	1,783 (472)	1,569 (456)	1,567 (435)	.03
Singleton birth weight z score, mean (SD) ^b	0.258 (1.15)	0.299 (1.11)	−0.003 (1.07)	0.106 (1.65)	0.186 (1.10)	<.0001
Singleton <−1	6.9	6.2	11.0	10.9	8.0	<.0001
Singleton <−2	0.8	0.7	1.4	1.6	1.1	<.0001
Twin birth weight z score, mean (SD) ^b	0.685 (1.34)	0.728 (1.31)	0.448 (1.30)	0.408 (1.71)	0.568 (1.40)	<.0001

Characteristic	Racial and ethnic groups				P value
	All groups	White	Asian	Black	Hispanic
Twin <-1	7.6	7.1	9.2	13.0	8.6
Twin <-2	1.6	1.4	1.8	4.3	2.2

^aLimited to pregnancies of 154 days (22 weeks).

^bLimited to birth weights 300 g.

Fujimoto. Racial and ethnic disparities in ART outcomes. Fertil Steril 2010.

TABLE 4
Regression Models on the Impact of Race and Ethnicity on the Response and Outcome to ART treatment*

Dependent	Independent	AOR	95% CI	P Value	Dependent	Independent	AOR	95% CI	P Value
Treatment Outcome: Pregnancy	White	1.00	Reference		Pregnancy Outcome: Live birth	White	1.00	Reference	
	Asian	0.86	0.80–0.93	<0.0001		Asian	0.90	0.82–0.97	0.01
	Black	1.09	0.99–1.20	0.09		Black	0.62	0.56–0.68	<0.0001
	Hispanic	1.06	0.96–1.16	0.25		Hispanic	0.87	0.79–0.96	0.005
Singleton Pregnancies**					Twin Pregnancies**				
Very Early Preterm Birth (<29 weeks)	White	1.00	Reference		Very Early Preterm Birth (<29 weeks)	White	1.00	Reference	
	Asian	0.77	0.48–1.25	0.30		Asian	0.62	0.42–0.92	0.02
	Black	4.25	3.14–5.76	<0.0001		Black	2.46	1.84–3.29	<0.0001
	Hispanic	1.38	0.91–2.09	0.13		Hispanic	1.36	1.01–1.82	0.04
Early Preterm Birth (<32 weeks)	White	1.00	Reference		Early Preterm Birth (<32 weeks)	White	1.00	Reference	
	Asian	1.01	0.78–1.31	0.93		Asian	0.83	0.69–1.01	0.07
	Black	2.72	2.19–3.38	<0.0001		Black	1.76	1.44–2.15	<0.0001
	Hispanic	1.19	0.91–1.56	0.21		Hispanic	1.10	0.92–1.33	0.30
Preterm Birth (<37 weeks)	White	1.00	Reference		Preterm Birth (<37 weeks)	White	1.00	Reference	
	Asian	0.95	0.85–1.06	0.35		Asian	0.70	0.60–0.82	<0.0001
	Black	1.79	1.59–2.03	<0.0001		Black	1.64	1.29–2.08	<0.0001
	Hispanic	1.22	1.08–1.37	0.001		Hispanic	1.14	0.95–1.37	0.15
Term Birth (≥ 37 weeks)	White	1.00	Reference		Term Birth (≥ 37 weeks)	White	1.00	Reference	
	Asian	1.06	0.94–1.18	0.35		Asian	1.42	1.22–1.65	<0.0001
	Black	0.56	0.49–0.63	<0.0001		Black	0.61	0.48–0.78	<0.0001
	Hispanic	0.82	0.73–0.93	0.001		Hispanic	0.88	0.73–1.05	0.15
Birth Weight Z-Score <–1	White	1.00	Reference		Birth Weight Z-Score <–1	White	1.00	Reference	
	Asian	1.78	1.58–2.01	<0.0001		Asian	1.30	1.03–1.66	0.03
	Black	1.81	1.56–2.11	<0.0001		Black	1.97	1.50–2.57	<0.0001
	Hispanic	1.36	1.17–1.58	<0.0001		Hispanic	1.26	0.97–1.64	0.08
Birth Weight Z-Score <–2	White	1.00	Reference		Birth Weight Z-Score <–2	White	1.00	Reference	
	Asian	2.05	1.50–2.80	<0.0001		Asian	1.29	0.76–2.17	0.35

Dependent	Independent	AOR	95% CI	P Value	Dependent	Independent	AOR	95% CI	P Value
	Black	2.17	1.47–3.19	<0.0001		Black	3.21	2.02–5.11	<0.0001
	Hispanic	1.64	1.11–2.42	0.01		Hispanic	1.57	0.94–2.62	0.08

* Models adjusted for maternal age; number of embryos transferred; and diagnoses of male factor, endometriosis, polycystic ovarian syndrome, diminished ovarian reserve, tubal factors, uterine factors, and other factors.

** Pregnancy outcomes limited to gestations 154 days (22 weeks) and birth weights >300 grams.
Fujimoto. Racial and ethnic disparities in ART outcomes. Fertil Steril 2010.