Genetics of Lipid-Storage Management in *Caenorhabditis elegans* Embryos

Verena Schmökel,* Nadin Memar,^{†,1} Anne Wiekenberg,[†] Martin Troitzmüller,^{*,§}
Ralf Schnabel,[†] and Frank Döring^{*,2}

*Department of Molecular Prevention, Institute of Human Nutrition and Food Science, Christian-Albrechts University of Kiel, 24118 Germany, [†]Institute of Genetics, Technical University (TU) Braunschweig, 38106 Germany, [‡]Core Facility for Mass Spectrometry, Lipidomics and Metabolomics, Centre for Medical Research (ZMF), University of Graz, A-8010 Austria, and [§]Omic Center Graz, A-8010 Austria

ABSTRACT Lipids play a pivotal role in embryogenesis as structural components of cellular membranes, as a source of energy, and as signaling molecules. On the basis of a collection of temperature-sensitive embryonic lethal mutants, a systematic database search, and a subsequent microscopic analysis of >300 interference RNA (RNAi)-treated/mutant worms, we identified a couple of evolutionary conserved genes associated with lipid storage in *Caenorhabditis elegans* embryos. The genes include *cpl-1* (cathepsin L-like cysteine protease), *ccz-1* (guanine nucleotide exchange factor subunit), and *asm-3* (acid sphingomyelinase), which is closely related to the human Niemann-Pick disease-causing gene *SMPD1*. The respective mutant embryos accumulate enlarged droplets of neutral lipids (*cpl-1*) and yolk-containing lipid droplets (*ccz-1*) or have larger genuine lipid droplets (*asm-3*). The *asm-3* mutant embryos additionally showed an enhanced resistance against C band ultraviolet (UV-C) light. Herein we propose that *cpl-1*, *ccz-1*, and *asm-3* are genes required for the processing of lipid-containing droplets in *C. elegans* embryos. Owing to the high levels of conservation, the identified genes are also useful in studies of embryonic lipid storage in other organisms.

KEYWORDS *C. elegans*; embryonic lipid droplets; embryo development; essential genes; yolk

AFTER fertilization, an embryo undergoes cell cleavage, cellular differentiation, gastrulation, and morphogenesis. These fundamental processes require the proper metabolism of carbohydrates and lipids (Biggers *et al.* 1967; Harvey and Orbidans 2011; Leese 2012; Minevich *et al.* 2012). For example, the maturation of oocytes and developing embryos depends on the metabolism of glucose and pyruvate (Gardner *et al.* 2000; Sutton-McDowall *et al.* 2010). However, the exact functions of lipids in embryogenesis are not clear (Dunning *et al.* 2010, 2014; McKeegan and Sturmey 2011). Generally, lipids function as rich sources of energy, signaling molecules, and structural components of plasma and organelle membranes. Given their diverse and important functions,

the formation, accumulation, and storage of lipids during oogenesis and embryogenesis have to be precisely regulated. Females of nearly all oviparous species synthesize vitellogenin, a yolk precursor lipoprotein, which is incorporated in oocytes (Byrne *et al.* 1989; Chen *et al.* 1997; Sappington and Raikhel 1998; Wallace 1985). In insects, vitellogenins are synthesized primarily in fat cells (Pan *et al.* 1969; Tufail and Takeda 2009); in vertebrates, the lipoprotein originates from the liver (Deeley *et al.* 1975; Wangh and Knowland 1975; Wahli *et al.* 1981; Mosconi *et al.* 2002). Yolk complexes undergo catabolism in order to provide raw materials, such as amino acids, carbohydrates, and lipids, for the developing embryo (Reimer and Crawford 1995; Jorgensen *et al.* 2009). In mammals, lipids are also directly provided by cumulus cells surrounding the maturing oocyte (Gilchrist *et al.* 2008; Prates *et al.* 2014). Inside the oocyte and embryo, yolk-storing platelets and other lipid-containing droplets have been observed, but their mechanistic roles and significance have yet to be elucidated (Reimer and Crawford 1995; Grant and Hirsh 1999; McEvoy *et al.* 2000; Ferguson and Leese 2006; Aardema *et al.* 2011).

Copyright © 2016 by the Genetics Society of America
doi: 10.1534/genetics.115.179127

Manuscript received July 2, 2015; accepted for publication January 4, 2016; published Early Online January 12, 2016.

Supporting information is available online at www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.179127/-/DC1.

¹Present address: Biozentrum der Ludwig-Maximilians Universität (LMU) München, Großhaderner Str. 2, 82152 Planegg-Martinsried, Germany.

²Corresponding author: Department of Molecular Prevention, Institute of Human Nutrition and Food Science, Christian-Albrechts University of Kiel, Heinrich-Hecht-Platz 10, 24118 Kiel, Germany. E-mail: sek@molprev.uni-kiel.de

The free-living nematode *Caenorhabditis elegans* serves as an excellent genetic model organism for the study of diverse mechanisms, including embryonic development and lipid metabolism (Byerly *et al.* 1976; Hodgkin and Barnes 1991; Muschiol *et al.* 2009; Harvey and Orbidans 2011). Most studies have examined lipid metabolism in adult worms (Ashrafi *et al.* 2003; Mullaney and Ashrafi 2009; Watts 2009; Klapper *et al.* 2011;). In the *C. elegans* embryo, lipid metabolism has been studied primarily with respect to the processing of yolk platelets. *C. elegans* vitellogenins are synthesized in the intestine of adult hermaphrodites, secreted into the pseudocoelomic space, and transported to the oocytes in the proximal gonadal arm (Kimble and Sharrock 1983; Hall *et al.* 1999; Kuo *et al.* 2013). The growing oocyte takes up plenty of yolk through receptor-mediated endocytosis and stores the yolk in yolk platelets (Sharrock *et al.* 1990; Grant and Hirsh 1999). In *C. elegans*, yolk seems to be only partially degraded during embryogenesis, as evidenced by the presence of yolk particles in larval intestines (Bossinger and Schierenberg 1992, 1996; Grant and Hirsh 1999). Further, lipid-containing droplets have been partially characterized in the *C. elegans* embryo (Levitte *et al.* 2010; Ehmke *et al.* 2014). In this study, we used *C. elegans* to identify genes that function in the processing of lipid-containing droplets, especially in the embryo. Our study identified candidate genes, including *cpl-1* (cathepsin L-like cysteine protease), *ccz-1* (guanine nucleotide exchange factor subunit), and *asm-3* (acid sphingomyelinase 3). These genes were further characterized.

Materials and Methods

C. elegans strains and maintenance

Bristol N2 was used as a wild-type reference strain. The mutant strains *rrf-3(pk1426)*, *asm-3(ok1744)*, *hjls67[atgl-1p::atgl-1::GFP]*, *hjls9[pges-1::glo-1::GFP]*, *wels15[pie-1p::GFP::eea-1(FYVEx2)+unc-119(+)]*, and *pwls23[vit-2::GFP]* were provided by the *C. elegans* Genetic Center (CGC), which is funded by the National Institutes of Health Office of Research Infrastructure Programs (P40 OD010440). The strains *asm-3(tm2384)*, *asm-1(tm5267)*, *asm-2(tm3764)*, and *hosl-1(tm2369)* were received from the National BioResource Project (NBRP). Twenty-eight temperature-sensitive mutants (Supporting Information, Table S1) were generated in the laboratory of R. Schnabel (TU Braunschweig) (R. Schnabel *et al.*, unpublished material). All mutant strains were backcrossed to wild-type N2. *ccz-1(t3403)*; *atgl-1p::atgl-1::GFP*, *ccz-1(t3213)*; *pwls23[vit-2::GFP]*, *ccz-1(t3403)*; *pwls23[vit-2::GFP]*, *ccz-1(t3403)*; *hjls9[pges-1::glo-1::GFP]*, *cpl-1(t3423)*; *atgl-1p::atgl-1::GFP*, *cpl-1(t3423)*; *pwls23[vit-2::GFP]*, *asm-3(ok1744)*; *atgl-1p::atgl-1::GFP*, *asm-3(ok1744)*; *pwls23[vit-2::GFP]*, and *asm-3(ok1744)*; *wels15[pie-1p::GFP::eea-1(FYVEx2)+unc-119(+)]* double mutants were generated by standard genetic techniques, and the presence of the homozygous mutant alleles was confirmed by PCR, embryonic phenotype, and/or GFP expression. Nematodes were maintained at 15° (ts mutants) or 20° on NGM agar plates with *Escherichia*

coli OP50 lawns as food sources using standard methods (Brenner 1974). Experiments with the ts mutants were performed at 15° (permissive temperature) and 25° (restrictive temperature). For all experiments, nematodes were synchronized via the hypochlorite treatment of gravid adults.

Temperature-sensitive mutant screen

We screened a collection of 1669 ts mutants obtained from the laboratory of R. Schnabel (TU Braunschweig) [partially described in Ehmke *et al.* (2014)]. We selected 28 of these mutants showing either an enlargement or a depletion of droplets as observed in four-dimensional (4D) microscopic analyses for further characterization. The embryos of self-fertilizing hermaphrodites of the selected mutants were scored for enlarged cytoplasmic droplets, droplet depletion, embryonic lethality, and brood size. Eight promising candidates were chosen from this group, and the mutations responsible for the droplet phenotype were identified. For this step, we combined SNP mapping (Davis *et al.* 2005) with whole-genome sequencing (GATC NextGen Sequencing, GATC Biotech, Konstanz, Germany). The resulting data sets were analyzed on the Galaxy Web platform (<http://usegalaxy.org/cloudmap>) established by the Hobert Laboratory (Minevich *et al.* 2012). We identified the phenotype-causing mutations for four alleles, two alleles each of *cpl-1* and *ccz-1*. For the remaining four ts alleles, we found relevant mutations in more than one candidate gene (Table S1).

Database search

The WormMart (BioMart v 0.7, <http://caprica.caltech.edu:9002/biomart/martview/>), PhenomicDB (v 3.9.7, <http://phenomicdb.info/>), RNAi Database (release 5 <http://rna.org/>), WormBase (v WS244, <http://www.wormbase.org/>), Tissue-specific expression predictions for *C. elegans* (v 1.0 <http://worm-tissue.princeton.edu/search>), KEGG Pathway Database (<http://www.genome.jp/kegg/pathway.html>), Expression Patterns for *C. elegans* promoter::GFP fusions Database (Hunt-Newbury *et al.* 2007), DAVID Gene Functional Annotation Tool (resource 6.7, <https://david.ncicrf.gov/>), and National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov/>) GenBank databases were used to identify candidate genes that met one or more of the following criteria: embryonic lethal with unknown or predicted function, embryonic lethal or embryonic expression and known function in lipid metabolism, known phenotypes that are associated with lipid storage (e.g., lipid depleted, lipid content reduced or increased), and orthologs of human genes implicated in lipid diseases.

RNAi screen by feeding

For the interference RNA (RNAi) screen, clones from the Ahringer *C. elegans* RNAi library, obtained from Gene Services (Source BioScience, Berlin, Germany), were used (Kamath and Ahringer 2003). Bacteria containing the 316 different RNAi clones were cultured in Luria broth medium with 50 µg/ml ampicillin and 5 µg/ml tetracycline at 37° for

16–18 hr. Afterwards, 50 μ l of the culture was seeded on NGM plates containing 2.5 mM IPTG and 50 μ g/ml ampicillin. The RNAi plates were incubated at 37° overnight. The RNAi-hypersensitive *rrf-3* strain was used for the screen. Ten synchronized L₁ larvae were placed on each RNAi plate and incubated at 20°. After 48 hr, Nile Red was added on the top of each RNAi plate to a final concentration of 0.05 μ g/ml. Embryos of the F₁ generation were scored for their overall droplet phenotype and Nile Red incorporation into the droplets after 48 and 72 hr, respectively, using an Axio Zoom.V16 Stereo Microscope (Zeiss, Jena, Germany). Additionally, the phenotypes of the embryos in the F₂ generation were determined. In each experiment, an L4440 empty vector and OP50 bacteria were used as negative controls, and the RNAi clones *T03E6.7* (coding for *cpl-1*) and *C49F5.1* (coding for *sams-1*) were used as positive controls, showing the droplet phenotype and embryonic lethality. RNAi clones resulting in a droplet phenotype were rescreened twice.

Microscopy and staining methods

Synchronized adult worms were harvested and washed in M9 buffer. Embryos were obtained by dissecting adult worms at the vulva and subsequently releasing the embryos from the uterus. The embryos were transferred on top of a small 5% agar pad on an objective glass slide. For vital staining, the embryos were directly incubated with BODIPY 493/503 (1:300, 15 min), LipidTOX Red (1:100, 40 min), or Lyso-Tracker (1:50, 20 min) at room temperature. For fixative BODIPY 493/503 and LipidTOX Red staining, nematodes were washed in M9, incubated in 4% paraformaldehyde (PFA) solution for 15 min, followed by three freeze-thaw cycles in liquid nitrogen. After the PFA was removed by three washing steps with M9, the worms were incubated in 500 μ l of 1 μ g/ml dye in M9 for 1 hr at room temperature. Subsequently, the nematodes were washed three times with M9 buffer. Finally, the worms were mounted for microscopic analysis. Images were taken either by laser-scanning confocal fluorescence microscopy (Zeiss Axio Imager Z2 Upright Microscope equipped with a Zeiss LSM 700 scanning confocal imaging system) or by an Axio Imager Z1 Microscope (Zeiss) equipped with ApoTome.2, Colibri.2, and an AxioCam MRm camera. For fluorescent images, an 81 HE DAPI/FITC/Rhodamine/Cy5 filter (Zeiss) was used. Objects were magnified using a 63 $\times$ /1.4 oil objective. All experiments were performed three to five times.

Brood size and lethality assays

Brood size and lethality assays were performed on NGM plates at 15 and 25°, respectively. Three synchronized L₄ larvae were placed on the NGM plates, grown to adults, and transferred to new NGM plates twice a day during the egg-laying period. The number of eggs laid was counted, and the eggs were allowed to develop for 24 hr. Residual eggs were counted. The overall brood size per worm and the lethality rate were determined. All experiments were performed three to five times.

Temperature-shift assay

Adult worms were synchronized by hypochlorite treatment. The embryos obtained were either shifted directly to the restrictive temperature (25°) or maintained at the permissive temperature (15°) to the L₄ stage. Afterwards, larvae were shifted to 25°. The overall larval development into fertile adults and the embryonic lethality rate of subsequent generations were determined.

UV survival assay

The UV survival assay was performed on NGM plates at 20°. Embryos were collected from gravid adults by hypochlorite treatment, and ~50–60 embryos were transferred to plates seeded with OP50 bacteria. Animals were irradiated directly after hypochlorite treatment for the indicated periods of time (0–180 min). For irradiation, a Philips G30T8 lamp (30 W) emitting a constant C band ultraviolet (UV-C) dose was used. The exact number of irradiated eggs was determined. After 24 hr of development, the remaining eggs were counted, and the lethality rate was calculated. The experiment was performed three to four times.

Analysis of embryonic lipid composition

For analyzing the lipid composition of the wild-type N2 and *asm-3(ok1744)* embryos, ~100,000 eggs were obtained by hypochlorite treatment of gravid adults. M9 buffer was removed, and the samples were stored at –80°. Five biological samples were collected per strain, and lipidomic analyses were performed on a high-resolution LC-MS/MS platform according to Fauland *et al.* (2011). Data processing was accomplished by a Lipid Data Analyzer (Hartler *et al.* 2011). The following lipid classes were quantified: triacylglycerol (TG), sphingomyelin (SM), phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylserine (PS), phosphoinositol (PI), and plasmalogen (aPE). The results were normalized to class amount, protein content, and sample weight. The distribution of lipid species was analyzed using GraphPad Prism (v 4.0) and Microsoft Excel (v 2013).

Protein alignment

We used the SIAS tool (imed.med.ucm.es) with the BLSOSUM62 matrix to calculate the overall protein identity and similarity. The alignment of catalytic domains was performed using the GENTle tool (v 1.9.4). The catalytic regions of human SMPD1 (aa 203–498) and *C. elegans* ASM-3 (aa 143–439) were aligned using the Clustal-W algorithm. Identical amino acids are indicated with asterisks.

Statistical analyses

Statistical analyses were performed using GraphPad Prism (v 4.0). The data presented represent means $\pm$ SEM of results from three or more independent experiments. Significances were calculated using Student's unpaired *t*-test or one-way ANOVA. Differences were considered to be significant when **P* < 0.05, ***P* < 0.005, and ****P* < 0.001.

Figure 1 Identification of genes associated with lipid metabolism and/or embryonic lethality. (A) Schematic representation of the approaches performed. The screening of ts alleles resulted in identification of eight possible candidates. The database research identified 1085 candidate genes, of which 316 were further analyzed by functional RNAi testing and/or mutant analysis. (B) Three different approaches were used to identify genes associated with lipid droplets in embryos. First, 28 ts alleles were screened for embryonic lethality and an enlargement or reduction of lipid droplets in embryos (I). Second, 1085 candidate genes of the database screen (II) were grouped in three partly overlapping categories: embryonic lethal, known function, and unknown/predicted function. Third, candidates from the database screen were functionally tested by RNAi (III). See *Materials and Methods* and *Results* for more information.

Results

Screening of mutant alleles altering the phenotype of putative lipid-containing droplets present in the *C. elegans* embryo

We searched for *C. elegans* mutant embryos with altered droplet phenotypes. We assumed that the screening and subsequent analyses of the mutants would lead to identification of genes functioning in the processing of lipid-containing droplets. Our first approach was based on a collection of 1669 ts embryonic lethal mutants and 4D DIC recordings of embryos (R. Schnabel *et al.*, unpublished material). We have already shown that one of these mutants, *sams-1(t3210)*, displayed an expansion of genuine lipid droplets (Ehmke *et al.* 2014). In this study, the embryos of self-fertilizing hermaphrodites of 28 selected mutants were scored for both embryonic lethality and the appearance and size of droplet structures (Figure 1 and Table S1). By combining SNP mapping with whole-genome sequencing, we identified two new alleles of *cpl-1*, *t3423* and *t3438*, and of *ccz-1*, *t3403* and *t3213* (Figure S1). The pronounced and enlarged droplets in the *cpl-1* and *ccz-1* mutant embryos (Figure 2) prompted us to further characterize the phenotypes.

To identify the additional genes influencing the droplet phenotype of the *C. elegans* embryo, we collected and combined phenotypic information from several databases (see *Materials and Methods*). This resulted in a list of 1085 candidate genes, of which 725 were genes with known functions and 360 had unknown or predicted functions (Figure 1 and Table S2). The latter group was of particular interest. From

316 selected genes (Table S3), we identified RNAi-treated and/or mutant embryos showing altered droplet structures (e.g., size, distribution) by staining with dyes to trace the lipids. Of 22 candidate genes (Table S4), the most robust droplet phenotype was observed in embryos lacking *asm-3*, *hosl-1*, or *E01A2.1* (Figure 2). We focused on the acid sphingomyelinase-encoding gene *asm-3* because this gene has not been associated with lipid storage in embryos previously.

Mutant embryos of *cpl-1* exhibit large droplets containing neutral lipids

The *cpl-1* gene encodes a member of the cathepsin L-like cysteine protease family that seems to be involved in the proteolytic processing of yolk proteins (Hashmi *et al.* 2002; Britton and Murray 2004). The high embryonic lethality of the identified alleles, *cpl-1(t3423)* and *cpl-1(t3438)*, at the restrictive temperature (Figure 3A and Table S1) correlated with the reported lethality of the *cpl-1(ok360)* null mutant (Britton and Murray 2004). In agreement with the strictly maternal function of *cpl-1* (Hashmi *et al.* 2002; Britton and Murray 2004), a temperature shift during gametogenesis drastically enhanced the embryonic lethality of the F₁ generation of *cpl-1(t3423)* (Figure 3B). Both *cpl-1(t3423)* and *cpl-1(t3438)* showed a minor increase in brood size compared with wild type at the restrictive temperature (+26.3%, *P* = 0.2343, and +20.6%, *P* = 0.361, respectively) (Figure 3C). Based on these results, the *cpl-1* mutants exhibited normal or even enhanced fertility, whereas their embryonic development was disturbed.

Figure 2 Identified candidate genes associated with a droplet phenotype in the embryo. (A–I) DIC images of homozygous embryos. (A) Wild-type N2. Homozygous *cpl-1(t3423)* (B) and *cpl-1(t3438)* (C) embryos at the restrictive temperature. Homozygous *ccz-1(t3403)* (D) and *ccz-1(t3213)* (E) embryos at the restrictive temperature. (F) *asm-3(ok1744)* embryo. (G) *asm-3(tm2384)* mutant. (H) *hosl-1(tm2369)* embryo. (I) Downregulation of *E01A2.1* transcript by RNAi in *rrf-3(pk1426)* embryo. Scale bar, 10 μ m.

Regarding the function of *cpl-1* during gametogenesis and embryogenesis, the *cpl-1(t3423)* and *cpl-1(t3438)* mutant alleles showed normal droplet size in the intestines of adults (Figure 3D) but enlarged droplet sizes in oocytes (Figure 3E) and embryos (Figure 3F). To ascertain the mechanism responsible for the formation of the large droplets, we performed vital and fixative staining experiments with dyes (BODIPY493/503 and LipidTOX Red) tracing neutral lipids. Similar to other vital staining methods (O'Rourke *et al.* 2009; Wahlby *et al.* 2014), our short-term (20-min) vital staining approach (Klapper *et al.* 2011) was capable of visualizing lipid droplets. The fat-staining signals of the dyes did not co-localize with the autofluorescent signals of lysosome-related organelles (LROs) in the intestine (Figure 3D) or with the signals of the yolk marker VIT-2::GFP in oocytes (Figure 3E). Based on our vital fat-staining method, we showed that the expanded droplets present in the oocytes (Figure 3E) and embryos (Figure 3F) of the *cpl-1(ts)* mutants indeed contained neutral lipids. The fixative fat-staining method confirmed the lipid-containing character of the enlarged droplets of the *cpl-1(ts)* mutant embryos (Figure 3G). Signals from the ATGL-1::GFP fusion protein, a marker of canonical lipid

droplets, were not detectable on the expanded droplets of the *cpl-1(t3423)* mutant embryos (Figure 3H). However, small embryonic droplets were visualized by ATGL-1::GFP (Figure 3H). According to a previous report on a *cpl-1* null mutant (Britton and Murray 2004), the VIT-2::GFP signal was not visible on the large droplets of the *cpl-1(ts)* mutant embryos (Figure 3I). In conclusion, we found that the *cpl-1* mutant embryos accumulated enlarged lipid-containing droplets that did not co-localize with marker proteins for yolk or lipid droplets.

Mutant embryos of *ccz-1* accumulate enlarged yolk-containing lipid droplets

The novel alleles of *ccz-1*, *t3213* and *t3403*, produced mostly dead embryos at the restrictive temperature (Figure 4A and Table S1). At the permissive temperature, the two mutants showed a lethality of approximately 50%. We further observed a high level of embryonic lethality (>95%) independent of whether we shifted the *ccz-1(ts)* mutants to the restrictive temperature early in development or after reaching the L₄ stage (Figure 4B). The two *ccz-1(ts)* mutants produced considerably fewer embryos at the restrictive

Figure 3 *cpl-1* mutant embryos exhibit enlarged lipid-laden droplets. (A) Embryonic lethality at permissive temperature (15°, white bar) and restrictive temperature (25°, gray bar). (B) Temperature-shift experiment. *cpl-1(t3423)* mutants are shifted to restrictive temperature (25°) at the indicated stages. The level of embryonic lethality (emb) in the F₁ generation is stated. (C) Brood size at permissive temperature (15°) and restrictive temperature (25°). (***P* < 0.005; ****P* < 0.001). (D–I) DIC images of *cpl-1(t3423)* mutant embryos and adults at the restrictive temperature. (D–D'') Intestinal region of adult homozygous *cpl-1(t3423)* mutant stained with LipidTOX (red, autofluorescence in blue). (E–E'') Gonadal tissue of adult *cpl-1* mutant expressing *vit-2::GFP* and stained with the lipid marker LipidTOX. (F–F'') *cpl-1* mutant embryo stained with BODIPY to visualize neutral lipids. (G–G'') Fixative LipidTOX staining of *cpl-1* mutant embryo. (H–H'') *cpl-1* mutant embryo expressing the lipid droplet marker *atgl-1::GFP*. (I–I'') Embryo expressing *vit-2::GFP* in *cpl-1* mutant background. Higher-magnification images represent the area outlined in the dotted box. Arrows point to enlarged droplets. Scale bar, 10 µm; scale bar in magnification images, 2 µm.

temperature than at the permissive temperature (Figure 4C). The high embryonic lethality and impaired fertility of the *ccz-1(ts)* mutants confirmed the essential role of CCZ-1 in gametogenesis and embryogenesis (Nieto *et al.* 2010). It has been reported that *ccz-1* is involved in the digestion of apoptotic corpses, phagosome maturation, and processing of LROs in the intestine (Poteryaev *et al.* 2007; Nieto *et al.* 2010; Delahaye *et al.* 2014). We found large droplets in the intestine, oocytes, and embryos of the *ccz-1(ts)* mutants (Figure 4, D–F, and Table S1). Both vital and fixative fat-staining methods revealed that the expanded droplets contained neutral lipids (Figure 4, F and G). The signals of the lipid droplet marker protein ATGL-1::GFP and the yolk marker VIT-2::GFP are clearly visible on the large droplets of the *ccz-1(ts)* mutant (Figure 4, H and I). In contrast, the LRO marker GLO-1::GFP

is not present on the large intestinal droplets of the *ccz-1(ts)* mutant (Figure 4J). Taken together, the *ccz-1* mutant embryos accumulated enlarged lipid-containing droplets, which are characterized by the presence of marker proteins for both yolk and lipid droplets.

Mutant embryos of *asm-3* are more resistant to UV-C exposure, accumulate enlarged genuine lipid droplets, and show distinct changes in their lipid profiles

The *asm-3* gene is closely related to the *C. elegans* acid sphingomyelinase genes *asm-1* and *asm-2* and to the human Niemann-Pick disease gene *SMPD1* (Figure S2A). The large-droplet phenotype of the *asm-3* mutant embryo (Figure 2, F and G) was not observable in the *C. elegans* mutants *asm-1* and *asm-2* (Figure S2, B and C). The *asm-3(ok1744)* mutant

Figure 4 *ccz-1* mutant embryos possess enlarged yolk-enriched lipid droplets. (A) Overall embryonic lethality and (B) lethality in temperature-shift assay at permissive temperature (15°, white bar) and restrictive temperature (25°, gray bar). *ccz-1(t3403)* mutants are shifted to the restrictive temperature at the indicated stages. The level of embryonic lethality (emb) in the F₁ generation is stated. (C) Number of progeny per worm (**P* < 0.05; ***P* < 0.005). (D–J) DIC images of *ccz-1* mutant embryos or adults at the restrictive temperature. (D–D'') Intestinal region of adult homozygous *ccz-1(t3403)* mutant stained with LipidTOX (red, autofluorescence in blue). (E–E'') Gonadal tissue of adult *ccz-1(tm3403)* mutant expressing *vit-2::GFP* and stained with the lipid marker LipidTOX. (F–F'') *ccz-1(t3403)* mutant embryo stained with BODIPY to visualize neutral lipids. (G–G'') Fixative BODIPY staining of *ccz-1* mutant embryo. (H–H'') Embryo expressing *atgl-1::GFP* in *ccz-1(t3403)* mutant background. (I–I'') *ccz-1(t3213)* mutant embryo expressing *vit-2::GFP* as yolk marker. (J–J'') Intestinal region of adult *ccz-1(t3403)* mutant expressing *glo-1::GFP* and stained with the lipid marker LipidTOX (autofluorescence in blue). Higher-magnification images represent the area outlined in the dotted box. Arrows point to enlarged droplets. Scale bar, 10 μm; scale bar in magnification images, 2 μm.

displayed a 56% reduced brood size and a slightly increased embryonic lethality compared with wild-type embryos (Figure 5, A and B). A previous study showed that adult *asm-3* mutant worms were characterized by an increased stress resistance (Kim and Sun 2012). We found that the embryos of the *asm-3(ok1744)* mutant survived for longer times than

wild-type embryos under UV-C irradiation (Figure 5C). Thus, the *asm-3* mutant embryos were more resistant to UV-C stress.

The *asm-3(ok1744)* mutant was characterized by the presence of enlarged droplets in all embryonic stages and in maturing oocytes (Figure S3), indicating a maternal

Figure 5 Lipid droplets are enlarged in *asm-3* mutant embryos. The brood size (A) and embryonic lethality (B) of wild-type N2 and *asm-3(ok1744)* mutants are shown. (C) Survival rate of wild-type N2 and *asm-3(ok1744)* embryos after UV-C irradiation (*** $P < 0.001$). (D–I) DIC images of homozygous *asm-3(ok1744)* mutant embryos and adults. (D–D'') *asm-3* mutant embryo stained with BODIPY. (E–E'') Intestinal region of adult *asm-3* mutant stained with BODIPY. (F–F'') Embryo expressing the lipid droplet marker *atgl-1::GFP* in *asm-3* mutant background. (G–G'') *asm-3* mutant embryo expressing *vit-2::GFP*. (H–H'') *asm-3* mutant embryo expressing the early endosome marker *pie-1p::GFP::eea-1*. (I–I'') *asm-3* mutant embryo stained with the lysosomal marker LysoTracker. Higher-magnification images represent the areas in the dotted box. Arrows point to enlarged droplets. Scale bar, 10 μm ; scale bar in magnification images, 2 μm .

contribution to *asm-3* expression. To validate this observation, we crossed homozygous *asm-3* mutant hermaphrodites with wild-type N2 males and examined the offspring of the F₁ and F₂ generations for the incidence of large droplets (Figure S4). The results demonstrated a maternal contribution to *asm-3* gene expression. To analyze whether the large droplets of the *asm-3(ok1744)* mutant embryo contained neutral lipids, we used the vital BODIPY 493/503 staining method. This approach showed that the large droplets of the *asm-3(ok1744)* mutant embryo were BODIPY 493/503 positive (Figure 5D). In the *asm-3(ok1744)* adult worms, we also found enlarged BODIPY 493/503-positive droplets (Figure 5E). Moreover, the lipid droplet marker protein ATGL-1::GFP was clearly visible on the surfaces of the enlarged droplets in the *asm-3(ok1744)* mutant embryos (Figure 5F). In contrast,

the expanded droplets were not marked by reporters for yolk (VIT-2::GFP), a subset of early endosomes (*pie-1p::GFP::EEA-1*), or lysosome (LysoTracker) (Figure 5, G–I). We concluded that the loss of *asm-3* resulted in the accumulation of genuine lipid droplets in the embryo.

To identify the changes in the lipid profile of *asm-3* mutant embryos, we performed a lipidomics analysis and identified 265 relevant individual lipid species. The total lipid amount (N2: 4999 $\mu\text{mol/g}$ protein; *asm-3*: 4566 $\mu\text{mol/g}$ protein) and the relative distribution of the seven lipid classes were comparable between the *asm-3(ok1744)* and wild-type embryos (Figure 6, A and B). In the sphingomyelin group, the most prominent effects were observed for SM 15:0 [–2.9-fold lower in *asm-3(ok1744)* embryos compared with wild type], SM 17:0 (–5.8-fold), and SM 20:0 (–2.2-fold) (Figure

Figure 6 Lipidomic analysis of sphingomyelin and triglyceride in *N2* and *asm-3* mutant embryos. (A–D) Results of lipidomic analysis of wild-type *N2* and *asm-3(ok1744)* mutant embryos. (A) Quantification of lipid classes. (B) Distribution of lipids in different classes. (C) Comparison of sphingomyelin fatty acid species. (D) Heat map of significantly regulated lipid species with fold changes. Indicated numbers refer to fatty acids (* $P < 0.05$; ** $P < 0.005$).

6C). In the triglyceride species, we observed an overall shift toward long-chain and polyunsaturated fatty acids in the *asm-3(ok1744)* mutants (Figure S5). When considering the

other lipid classes, we found robust differences for PC 36:0 (–19-fold), PC 31:0 (–6.6-fold), PE 40:8 (+5.9-fold), and PS 38:6 (+5.9-fold) (Figure 6D). Overall, 84 lipid species

(31.7%) showed significantly higher or lower levels (44 and 40 compounds, respectively) in the *asm-3(ok1744)* embryos compared with wild-type embryos (Figure 6D). The results of the lipidomics analyses showed that the *asm-3* mutant embryo was characterized by certain changes in the lipid profile.

Discussion

Yolk complexes and lipid droplets are distinct lipid storage organelles in C. elegans embryos

Yolk complexes are lipoprotein complexes that serve as energy reserves for the nourishment of developing embryos (Wallace 1985; Byrne *et al.* 1989; Tufail and Takeda 2009). The precursors of the yolk proteins, vitellogenins, are typically synthesized in organs distant from the gonad and are then endocytosed by the developing oocytes (Mosconi *et al.* 2002; Tufail and Takeda 2009). Here they are stored in yolk platelets until they are degraded (Reimer and Crawford 1995; Jorgensen *et al.* 2009). In *C. elegans*, the cathepsin L-like cysteine protease CPL-1 is involved in the degradation of yolk proteins during embryogenesis (Hashmi *et al.* 2002; Britton and Murray 2004). In agreement with previous studies (Hashmi *et al.* 2002; Britton and Murray 2004), we observed enlarged droplets in both oocytes and embryos in the newly identified *cpl-1(ts)* mutants. Although the yolk marker protein VIT-2::GFP was not detectable on the large droplets, Britton and Murray (2004) showed that the droplets still contained yolk. Similarly, we found that the enlarged droplets of the *cpl-1(ts)* mutant embryos also contained neutral lipids not marked by the lipid droplet reporter protein ATGL-1::GFP. These findings suggested that the loss of function of *cpl-1* causes an enlargement of lipid-containing yolk complexes distinct from genuine lipid droplets.

It has been shown that CCZ-1 acts in complex with SAND-1 as a guanine nucleotide exchange factor for RAB-7 and is involved in the digestion of apoptotic corpses, phagosome maturation, and the processing of lysosome-related organelles (Poteryaev *et al.* 2007; Nieto *et al.* 2010; Delahaye *et al.* 2014). As found in the isolated *ccz-1(ts)* mutant embryos, *sand-1* mutants also accumulate yolk-enriched large droplets in oocytes and embryos (Poteryaev and Spang 2005; Poteryaev *et al.* 2007). Our further examination of the *ccz-1(ts)* mutant embryos revealed that the enlarged droplets contained neutral lipids and were marked with reporter proteins for genuine lipid droplets and yolk complexes. Therefore, we proposed that ATGL-positive lipid droplets undergo fusion with yolk complexes to transiently form a mixture of genuine lipid droplets and yolk. These structures might be involved in separating the lipid moiety from vitellogenins and in the subsequent storage of lipids in genuine lipid droplets.

Involvement of the asm-3 gene in altering the size of lipid droplets and in the stress response

Loss of the human sphingomyelinase-encoding gene *SMPD1* causes Niemann-Pick disease types A and B, which are lyso-

somal storage diseases (Schuchman *et al.* 1992; Jones *et al.* 2008; Jenkins *et al.* 2009). In Niemann-Pick patients, lipid-laden foam cells can be found in bone marrow, spleen, and lymph nodes (Viana *et al.* 1990; Putterman *et al.* 1992; Thurberg *et al.* 2012). At a cellular level, loss of *SMPD1* results in the accumulation of SM in large droplets within endolysosomal compartments (Schuchman *et al.* 1992; Jones *et al.* 2008; Jenkins *et al.* 2009). *C. elegans* encodes three acidic sphingomyelinases: *asm-1*, *asm-2*, and *asm-3* (Lin *et al.* 1998; Kim and Sun 2012). In this study, we showed that *asm-3* mutant oocytes and embryos exclusively accumulated enlarged droplets containing neutral lipids marked by the lipid droplet marker ATGL-1::GFP. Because the droplets are not marked by reporter proteins for lysosomes, yolk, or a subset of early endosomes, we concluded that loss of *asm-3* function causes an enlargement of genuine lipid droplets.

The reduction of PC levels in the *asm-3(ok1744)* mutant embryos was remarkable because a decrease of this lipid class increases the size of the lipid droplets in the intestine of adult worms (Krahmer *et al.* 2011; Xu *et al.* 2012; Ehmke *et al.* 2014). This also could be the reason for lipid droplet enlargement in the *asm-3* mutant embryos. Acid sphingomyelinases are crucial in establishing a well-balanced level between SM and ceramide production in the plasma membrane (Goni and Alonso 2002; Jenkins *et al.* 2009; Young *et al.* 2013). However, SM has been discovered in lipid droplet fractions in a variety of different cell types (Hood and Patton 1973; Ishii *et al.* 1995; Tauchi-Sato *et al.* 2002; McIntosh *et al.* 2010; Storey *et al.* 2011). Moreover, SMs, PCs, and PIs are enriched in functionally active lipid droplets (Storey *et al.* 2011). The conversion of SMs to ceramide causes structural changes in membranes, facilitating the formation of vesicles (Wheeler *et al.* 2009; Mitsutake *et al.* 2011). Indeed, the addition of bacterial sphingomyelinase to giant unilamellar vesicles results in the formation of intraluminal vesicles (Trajkovic *et al.* 2008; Ibarguren *et al.* 2013), indicating an involvement of sphingomyelinases in vesicle formation. Thus, sphingomyelinase-induced SM hydrolysis also could be involved in determining the size of lipid droplets. We further observed a shift toward long-chain and polyunsaturated fatty acids (PUFAs) in triglyceride species, indicating an upregulation of the desaturase genes (*i.e.*, *fat-1*, *fat-3*, and *fat-4*). This likely increases the fluidity of membranes (Dietrich *et al.* 2001; Roux *et al.* 2005) and facilitates the enlargement of lipid droplets.

Acid sphingomyelinases have been implicated in mediating apoptotic and stress responses to Fas ligands and ionizing and UV irradiation (Lin *et al.* 2000; Garcia-Barros *et al.* 2003; Zeidan *et al.* 2008; Appelqvist *et al.* 2013). The *C. elegans* sphingomyelinase functions as a positive regulator of the DAF-2/insulin signaling pathway, which is involved in the regulation of lifespan, metabolism, reproduction, and development (Kimura *et al.* 1997; Kim and Sun 2012). Accordingly, we observed a drastically reduced brood size and slightly increased embryonic lethality in *asm-3* mutants.

Reduced fertility also has been documented in mouse models for Niemann-Pick disease and in other lysosomal storage diseases (Trasler *et al.* 1998; Adamali *et al.* 1999; Butler *et al.* 2007). Kim and Sun (2012) showed that *asm-3(ok1744)* mutant worms have a prolonged lifespan, their dauer arrest is promoted, and they are more resistant to oxidative or heat stresses. We could show that UV-induced embryonic lethality was lower in the *asm-3* null allele *ok1744*, and thus, stress resistance was already established in embryos. By searching for lipid species that might be important for this phenotype, we found that GPEtn (40:8) was increased in the *asm-3* mutant embryo. Assuming that this lipid species is composed of two arachidonic acids (20:4), this fatty acid could contribute to the enhanced stress resistance (O'Rourke *et al.* 2013) of the *asm-3* mutant embryos.

Conclusion

We identified a few evolutionarily conserved and essential genes that seem to be involved in the processing of lipid-containing droplets in *C. elegans* embryos. Using marker proteins and vital and fixative fat-staining methods, we found that the mutant embryos of the genes *cpl-1* (cathepsin L-like cysteine protease), *ccz-1* (guanine nucleotide exchange factor subunit), and *asm-3* (acid sphingomyelinase 3) showed accumulations of three distinct types of enlarged neutral lipid-containing droplets. These mutant phenotypes indicated that *asm-3*, *ccz-1*, and *cpl-1* are critical for the distinct steps in the processing of lipid-containing droplets in the *C. elegans* embryo. The identified genes may be useful for studying embryonic lipid storage in other organisms as well.

Literature Cited

Aardema, H., P. L. Vos, F. Lolicato, B. A. Roelen, H. M. Knijn *et al.*, 2011 Oleic acid prevents detrimental effects of saturated fatty acids on bovine oocyte developmental competence. *Biol. Reprod.* 85: 62–69.

Adamali, H. I., I. H. Somani, J. Q. Huang, D. Mahuran, R. A. Gravel *et al.*, 1999 I. Abnormalities in cells of the testis, efferent ducts, and epididymis in juvenile and adult mice with beta-hexosaminidase A and B deficiency. *J. Androl.* 20: 779–802.

Appelqvist, H., P. Waster, I. Eriksson, I. Rosdahl, and K. Ollinger, 2013 Lysosomal exocytosis and caspase-8-mediated apoptosis in UVA-irradiated keratinocytes. *J. Cell Sci.* 126: 5578–5584.

Ashrafi, K., F. Y. Chang, J. L. Watts, A. G. Fraser, R. S. Kamath *et al.*, 2003 Genome-wide RNAi analysis of *Caenorhabditis elegans* fat regulatory genes. *Nature* 421: 268–272.

Biggers, J. D., D. G. Whittingham, and R. P. Donahue, 1967 The pattern of energy metabolism in the mouse oocyte and zygote. *Proc. Natl. Acad. Sci. USA* 58: 560–567.

Bossinger, O., and E. Schierenberg, 1992 Cell-cell communication in the embryo of *Caenorhabditis elegans*. *Dev. Biol.* 151: 401–409.

Bossinger, O., and E. Schierenberg, 1996 The use of fluorescent marker dyes for studying intercellular communication in nematode embryos. *Int. J. Dev. Biol.* 40: 431–439.

Brenner, S., 1974 The genetics of *Caenorhabditis elegans*. *Genetics* 77: 71–94.

Britton, C., and L. Murray, 2004 Cathepsin L protease (CPL-1) is essential for yolk processing during embryogenesis in *Caenorhabditis elegans*. *J. Cell Sci.* 117: 5133–5143.

Butler, A., R. E. Gordon, S. Gatt, and E. H. Schuchman, 2007 Sperm abnormalities in heterozygous acid sphingomyelinase knockout mice reveal a novel approach for the prevention of genetic diseases. *Am. J. Pathol.* 170: 2077–2088.

Byerly, L., R. C. Cassada, and R. L. Russell, 1976 The life cycle of the nematode *Caenorhabditis elegans*. I. Wild-type growth and reproduction. *Dev. Biol.* 51: 23–33.

Byrne, B. M., M. Gruber, and G. Ab, 1989 The evolution of egg yolk proteins. *Prog. Biophys. Mol. Biol.* 53: 33–69.

Chen, J. S., T. W. Sappington, and A. S. Raikhel, 1997 Extensive sequence conservation among insect, nematode, and vertebrate vitellogenins reveals ancient common ancestry. *J. Mol. Evol.* 44: 440–451.

Davis, M. W., M. Hammarlund, T. Harrach, P. Hullett, S. Olsen *et al.*, 2005 Rapid single nucleotide polymorphism mapping in *C. elegans*. *BMC Genomics* 6: 118.

Deeley, R. G., D. P. Mullinix, W. Wetekam, H. M. Kronenberg, M. Meyers *et al.*, 1975 Vitellogenin synthesis in the avian liver: vitellogenin is the precursor of the egg yolk phosphoproteins. *J. Biol. Chem.* 250: 9060–9066.

Delahaye, J. L., O. K. Foster, A. Vine, D. S. Saxton, T. P. Curtin *et al.*, 2014 *Caenorhabditis elegans* HOPS and CCZ-1 mediate trafficking to lysosome-related organelles independently of RAB-7 and SAND-1. *Mol. Biol. Cell* 25: 1073–1096.

Dietrich, C., L. A. Bagatolli, Z. N. Volovyk, N. L. Thompson, M. Levi *et al.*, 2001 Lipid rafts reconstituted in model membranes. *Biophys. J.* 80: 1417–1428.

Dunning, K. R., K. Cashman, D. L. Russell, J. G. Thompson, R. J. Norman *et al.*, 2010 Beta-oxidation is essential for mouse oocyte developmental competence and early embryo development. *Biol. Reprod.* 83: 909–918.

Dunning, K. R., D. L. Russell, and R. L. Robker, 2014 Lipids and oocyte developmental competence: the role of fatty acids and beta-oxidation. *Reproduction* 148: R15–R27.

Ehmke, M., K. Luthe, R. Schnabel, and F. Doring, 2014 S-Adenosyl methionine synthetase 1 limits fat storage in *Caenorhabditis elegans*. *Genes Nutr.* 9: 386.

Fauland, A., H. Kofeler, M. Trotschmuller, A. Knopf, J. Hartler *et al.*, 2011 A comprehensive method for lipid profiling by liquid chromatography-ion cyclotron resonance mass spectrometry. *J. Lipid Res.* 52: 2314–2322.

Ferguson, E. M., and H. J. Leese, 2006 A potential role for triglyceride as an energy source during bovine oocyte maturation and early embryo development. *Mol. Reprod. Dev.* 73: 1195–1201.

Garcia-Barros, M., F. Paris, C. Cordon-Cardo, D. Lyden, S. Rafii *et al.*, 2003 Tumor response to radiotherapy regulated by endothelial cell apoptosis. *Science* 300: 1155–1159.

Gardner, D. K., T. B. Pool, and M. Lane, 2000 Embryo nutrition and energy metabolism and its relationship to embryo growth, differentiation, and viability. *Semin. Reprod. Med.* 18: 205–218.

Gilchrist, R. B., M. Lane, and J. G. Thompson, 2008 Oocyte-secreted factors: regulators of cumulus cell function and oocyte quality. *Hum. Reprod. Update* 14: 159–177.

Goni, F. M., and A. Alonso, 2002 Sphingomyelinases: enzymology and membrane activity. *FEBS Lett.* 531: 38–46.

Grant, B., and D. Hirsh, 1999 Receptor-mediated endocytosis in the *Caenorhabditis elegans* oocyte. *Mol. Biol. Cell* 10: 4311–4326.

Hall, D. H., V. P. Winfrey, G. Blaeuer, L. H. Hoffman, T. Furuta *et al.*, 1999 Ultrastructural features of the adult hermaphrodite gonad of *Caenorhabditis elegans*: relations between the germ line and soma. *Dev. Biol.* 212: 101–123.

Hartler, J., M. Trotschmuller, C. Chittraju, F. Spener, H. C. Kofeler *et al.*, 2011 Lipid Data Analyzer: unattended identification

- and quantitation of lipids in LC-MS data. *Bioinformatics* 27: 572–577.
- Harvey, S. C., and H. E. Orbidans, 2011 All eggs are not equal: the maternal environment affects progeny reproduction and developmental fate in *Caenorhabditis elegans*. *PLoS One* 6: e25840.
- Hashmi, S., C. Britton, J. Liu, D. B. Guiliano, Y. Oksov *et al.*, 2002 Cathepsin L is essential for embryogenesis and development of *Caenorhabditis elegans*. *J. Biol. Chem.* 277: 3477–3486.
- Hodgkin, J., and T. M. Barnes, 1991 More is not better: brood size and population growth in a self-fertilizing nematode. *Proc. Biol. Sci.* 246: 19–24.
- Hood, L. F., and S. Patton, 1973 Isolation and characterization of intracellular lipid droplets from bovine mammary tissue. *J. Dairy Sci.* 56: 858–863.
- Hunt-Newbury, R., R. Viveiros, R. Johnsen, A. Mah, D. Anastas *et al.*, 2007 High-throughput in vivo analysis of gene expression in *Caenorhabditis elegans*. *PLoS Biol.* 5: e237.
- Ibarguren, M., J. Sot, L. R. Montes, A. I. Vasil, M. L. Vasil *et al.*, 2013 Recruitment of a phospholipase C/sphingomyelinase into non-lamellar lipid droplets during hydrolysis of lipid bilayers. *Chem. Phys. Lipids* 166: 12–17.
- Ishii, I., R. Onozaki, E. Takahashi, S. Takahashi, N. Fujio *et al.*, 1995 Regulation of neutral cholesterol esterase activity by phospholipids containing negative charges in substrate liposome. *J. Lipid Res.* 36: 2303–2310.
- Jenkins, R. W., D. Canals, and Y. A. Hannun, 2009 Roles and regulation of secretory and lysosomal acid sphingomyelinase. *Cell. Signal.* 21: 836–846.
- Jones, I., X. He, F. Katouzian, P. I. Darroch, and E. H. Schuchman, 2008 Characterization of common SMPD1 mutations causing types A and B Niemann-Pick disease and generation of mutation-specific mouse models. *Mol. Genet. Metab.* 95: 152–162.
- Jorgensen, P., J. A. Steen, H. Steen, and M. W. Kirschner, 2009 The mechanism and pattern of yolk consumption provide insight into embryonic nutrition in *Xenopus*. *Development* 136: 1539–1548.
- Kamath, R. S., and J. Ahringer, 2003 Genome-wide RNAi screening in *Caenorhabditis elegans*. *Methods* 30: 313–321.
- Kim, Y., and H. Sun, 2012 ASM-3 acid sphingomyelinase functions as a positive regulator of the DAF-2/AGE-1 signaling pathway and serves as a novel anti-aging target. *PLoS One* 7: e45890.
- Kimble, J., and W. J. Sharrock, 1983 Tissue-specific synthesis of yolk proteins in *Caenorhabditis elegans*. *Dev. Biol.* 96: 189–196.
- Kimura, K. D., H. A. Tissenbaum, Y. Liu, and G. Ruvkun, 1997 *daf-2*, an insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science* 277: 942–946.
- Klapper, M., M. Ehmke, D. Palgunow, M. Bohme, C. Matthaus *et al.*, 2011 Fluorescence-based fixative and vital staining of lipid droplets in *Caenorhabditis elegans* reveal fat stores using microscopy and flow cytometry approaches. *J. Lipid Res.* 52: 1281–1293.
- Krahmer, N., Y. Guo, F. Wilfling, M. Hilger, S. Lingrell *et al.*, 2011 Phosphatidylcholine synthesis for lipid droplet expansion is mediated by localized activation of CTP:phosphocholine cytidyltransferase. *Cell Metab.* 14: 504–515.
- Kuo, Y., T. Y. Hsu, Y. C. Wu, and H. C. Chang, 2013 Fluorescent nanodiamond as a probe for the intercellular transport of proteins in vivo. *Biomaterials* 34: 8352–8360.
- Leese, H. J., 2012 Metabolism of the preimplantation embryo: 40 years on. *Reproduction* 143: 417–427.
- Levitte, S., R. Salesky, B. King, S. Coe Smith, M. Depper *et al.*, 2010 A *Caenorhabditis elegans* model of orotic aciduria reveals enlarged lysosome-related organelles in embryos lacking *umps-1* function. *FEBS J.* 277: 1420–1439.
- Lin, T., L. Genestier, M. J. Pinkoski, A. Castro, S. Nicholas *et al.*, 2000 Role of acidic sphingomyelinase in Fas/CD95-mediated cell death. *J. Biol. Chem.* 275: 8657–8663.
- Lin, X., M. O. Hengartner, and R. Kolesnick, 1998 *Caenorhabditis elegans* contains two distinct acid sphingomyelinases. *J. Biol. Chem.* 273: 14374–14379.
- McEvoy, T. G., G. D. Coull, P. J. Broadbent, J. S. Hutchinson, and B. K. Speake, 2000 Fatty acid composition of lipids in immature cattle, pig and sheep oocytes with intact zona pellucida. *J. Reprod. Fertil.* 118: 163–170.
- McIntosh, A. L., S. M. Storey, and B. P. Atshaves, 2010 Intracellular lipid droplets contain dynamic pools of sphingomyelin: ADRP binds phospholipids with high affinity. *Lipids* 45: 465–477.
- McKeegan, P. J., and R. G. Sturmey, 2011 The role of fatty acids in oocyte and early embryo development. *Reprod. Fertil. Dev.* 24: 59–67.
- Minevich, G., D. S. Park, D. Blankenberg, R. J. Poole, and O. Hobert, 2012 CloudMap: a cloud-based pipeline for analysis of mutant genome sequences. *Genetics* 192: 1249–1269.
- Mitsutake, S., K. Zama, H. Yokota, T. Yoshida, M. Tanaka *et al.*, 2011 Dynamic modification of sphingomyelin in lipid microdomains controls development of obesity, fatty liver, and type 2 diabetes. *J. Biol. Chem.* 286: 28544–28555.
- Mosconi, G., O. Carnevali, H. R. Habibi, R. Sanyal, and A. M. Polzonetti-Magni, 2002 Hormonal mechanisms regulating hepatic vitellogenin synthesis in the gilthead sea bream, *Sparus aurata*. *Am. J. Physiol. Cell Physiol.* 283: C673–C678.
- Mullaney, B. C., and K. Ashrafi, 2009 *C. elegans* fat storage and metabolic regulation. *Biochim. Biophys. Acta* 1791: 474–478.
- Muschiol, D., F. Schroeder, and W. Trautspurger, 2009 Life cycle and population growth rate of *Caenorhabditis elegans* studied by a new method. *BMC Ecol.* 9: 14.
- Nieto, C., J. Almendinger, S. Gysi, E. Gomez-Orte, A. Kaech *et al.*, 2010 *ccz-1* mediates the digestion of apoptotic corpses in *C. elegans*. *J. Cell Sci.* 123: 2001–2007.
- O'Rourke, E. J., A. A. Soukas, C. E. Carr, and G. Ruvkun, 2009 *C. elegans* major fats are stored in vesicles distinct from lysosome-related organelles. *Cell Metab.* 10: 430–435.
- O'Rourke, E. J., P. Kuballa, R. Xavier, and G. Ruvkun, 2013 ω -6 Polyunsaturated fatty acids extend life span through the activation of autophagy. *Genes Dev.* 27: 429–440.
- Pan, M. L., W. J. Bell, and W. H. Telfer, 1969 Vitellogenic blood protein synthesis by insect fat body. *Science* 165: 393–394.
- Poteryaev, D., and A. Spang, 2005 A role of SAND-family proteins in endocytosis. *Biochem. Soc. Trans.* 33: 606–608.
- Poteryaev, D., H. Fares, B. Bowerman, and A. Spang, 2007 *Caenorhabditis elegans* SAND-1 is essential for RAB-7 function in endosomal traffic. *EMBO J.* 26: 301–312.
- Prates, E. G., J. T. Nunes, and R. M. Pereira, 2014 A role of lipid metabolism during cumulus-oocyte complex maturation: impact of lipid modulators to improve embryo production. *Mediators Inflamm.* 2014: 692067.
- Putterman, C., J. Zelingher, and D. Shouval, 1992 Liver failure and the sea-blue histiocyte/adult Niemann-Pick disease: case report and review of the literature. *J. Clin. Gastroenterol.* 15: 146–149.
- Reimer, C. L., and B. J. Crawford, 1995 Identification and partial characterization of yolk and cortical granule proteins in eggs and embryos of the starfish, *Pisaster ochraceus*. *Dev. Biol.* 167: 439–457.
- Roux, A., D. Cuvelier, P. Nassoy, J. Prost, P. Bassereau *et al.*, 2005 Role of curvature and phase transition in lipid sorting and fission of membrane tubules. *EMBO J.* 24: 1537–1545.
- Sappington, T. W., and A. S. Raikhel, 1998 Molecular characteristics of insect vitellogenins and vitellogenin receptors. *Insect Biochem. Mol. Biol.* 28: 277–300.
- Schuchman, E. H., O. Levrán, L. V. Pereira, and R. J. Desnick, 1992 Structural organization and complete nucleotide sequence

- of the gene encoding human acid sphingomyelinase (SMPD1). *Genomics* 12: 197–205.
- Sharrock, W. J., M. E. Sutherlin, K. Leske, T. K. Cheng, and T. Y. Kim, 1990 Two distinct yolk lipoprotein complexes from *Caenorhabditis elegans*. *J. Biol. Chem.* 265: 14422–14431.
- Storey, S. M., A. L. McIntosh, S. Senthivinayagam, K. C. Moon, and B. P. Atshaves, 2011 The phospholipid monolayer associated with perilipin-enriched lipid droplets is a highly organized rigid membrane structure. *Am. J. Physiol. Endocrinol. Metab.* 301: E991–E1003.
- Sutton-McDowall, M. L., R. B. Gilchrist, and J. G. Thompson, 2010 The pivotal role of glucose metabolism in determining oocyte developmental competence. *Reproduction* 139: 685–695.
- Tauchi-Sato, K., S. Ozeki, T. Houjou, R. Taguchi, and T. Fujimoto, 2002 The surface of lipid droplets is a phospholipid monolayer with a unique fatty acid composition. *J. Biol. Chem.* 277: 44507–44512.
- Thurberg, B. L., M. P. Wasserstein, T. Schiano, F. O'Brien, S. Richards *et al.*, 2012 Liver and skin histopathology in adults with acid sphingomyelinase deficiency (Niemann-Pick disease type B). *Am. J. Surg. Pathol.* 36: 1234–1246.
- Trajkovic, K., C. Hsu, S. Chiantia, L. Rajendran, D. Wenzel *et al.*, 2008 Ceramide triggers budding of exosome vesicles into multivesicular endosomes. *Science* 319: 1244–1247.
- Trasler, J., F. Saberi, I. H. Somani, H. I. Adamali, J. Q. Huang *et al.*, 1998 Characterization of the testis and epididymis in mouse models of human Tay Sachs and Sandhoff diseases and partial determination of accumulated gangliosides. *Endocrinology* 139: 3280–3288.
- Tufail, M., and M. Takeda, 2009 Insect vitellogenin/lipophorin receptors: molecular structures, role in oogenesis, and regulatory mechanisms. *J. Insect Physiol.* 55: 87–103.
- Viana, M. B., R. Giugliani, V. H. Leite, M. L. Barth, C. Lekhwani *et al.*, 1990 Very low levels of high density lipoprotein cholesterol in four sibs of a family with non-neuropathic Niemann-Pick disease and sea-blue histiocytosis. *J. Med. Genet.* 27: 499–504.
- Wahlby, C., A. L. Conery, M. A. Bray, L. Kamentsky, J. Larkins-Ford *et al.*, 2014 High- and low-throughput scoring of fat mass and body fat distribution in *C. elegans*. *Methods* 68: 492–499.
- Wahli, W., I. B. Dawid, G. U. Ryffel, and R. Weber, 1981 Vitellogenesis and the vitellogenin gene family. *Science* 212: 298–304.
- Wallace, R. A., 1985 Vitellogenesis and oocyte growth in non-mammalian vertebrates. *Dev. Biol.* 1: 127–177.
- Wang, L. J., and J. Knowland, 1975 Synthesis of vitellogenin in cultures of male and female frog liver regulated by estradiol treatment in vitro. *Proc. Natl. Acad. Sci. USA* 72: 3172–3175.
- Watts, J. L., 2009 Fat synthesis and adiposity regulation in *Caenorhabditis elegans*. *Trends Endocrinol. Metab.* 20: 58–65.
- Wheeler, D., E. Knapp, V. V. Bandaru, Y. Wang, D. Knorr *et al.*, 2009 Tumor necrosis factor- α -induced neutral sphingomyelinase-2 modulates synaptic plasticity by controlling the membrane insertion of NMDA receptors. *J. Neurochem.* 109: 1237–1249.
- Xu, N., S. O. Zhang, R. A. Cole, S. A. McKinney, F. Guo *et al.*, 2012 The FATP1-DGAT2 complex facilitates lipid droplet expansion at the ER-lipid droplet interface. *J. Cell Biol.* 198: 895–911.
- Young, M. M., M. Kester, and H. G. Wang, 2013 Sphingolipids: regulators of crosstalk between apoptosis and autophagy. *J. Lipid Res.* 54: 5–19.
- Zeidan, Y. H., B. X. Wu, R. W. Jenkins, L. M. Obeid, and Y. A. Hannun, 2008 A novel role for protein kinase C δ -mediated phosphorylation of acid sphingomyelinase in UV light-induced mitochondrial injury. *FASEB J.* 22: 183–193.

Communicating editor: M. P. Colaiácovo

GENETICS

Supporting Information

www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.179127/-/DC1

Genetics of Lipid-Storage Management in *Caenorhabditis elegans* Embryos

Verena Schmökel, Nadin Memar, Anne Wiekenberg, Martin Troitzmüller,
Ralf Schnabel, and Frank Döring

A					
Gene ID	Gene name	ts-allele	Mutation	Possible function	
T03E6.7	<i>cpl-1</i>	t3423	Arg ³²⁰ → Cys	cathepsin L-like cysteine protease	
		t3438	Ala ¹⁴⁸ → Val		
F58G11.6	<i>ccz-1</i>	t3403	Gln ³³² → Stop	guanine nucleotide exchange factor subunit for RAB-7	
		t3213	splice site donor G → A		

B		C	
001	MNRFILLALV AAVVAVNSAK LSRQIESAIE KWDYYKEDFD	001	MESTIANPLYF FSNLSLNPPS CSMVVREIDV LEFFVVAHPL SGRKEGEEHN RVMYFHPKGE
041	KKEYSEERQT YMEAGVMSMT HIEHNRDHR LGRATFEMGL	061	QLERQTEITG FAEAVVNFTF NPLSTSRREI SERISENEDG FDFRTVTTQR TERVYIRTED
081	NHIADLPFSQ YKELNGYHRL FGDRIENSS SFLAFFNVQV	121	DQFILGVSSIS KQLSNVSDYP LFQPAIRSEL SDAYKMFHMF FGTFFSFIEN VFDDIEFKKE
121	PDEVDRDITH LVTGVNQQH CGSCHAFSAT GALEGQHARK	181	RIDFFFSKYI PLLKVNOMPL LQHLGGVEFL RMGGPLYLNV VSLLESLRES FQVIEKIMFL
161	LQQLVSLSEQ NLVDCSTNYG NHGNGGLMD GAFETIRDNH	241	YQDKLLYYQL SKRDLPSLFH YLTHNLLFTT LAPELEHSGR NASKGRYLRG PTDLTTDAPL
201	GVDTESYPY KGRDHKCHF KRTVGADNKG YVDTFGDES	301	LGDSESLVNH LHSDDGTEFQ EELIGYQNIY YACLNATVCM FVRNCPANGG LEIRDSESS
241	QLKIAVATQG PISIAIDAGH RSPQLYKGV YYDEACSSSS	361	QSRATPSQN GVVSRLLRN IDQFLETELS QVASKIGDEI GDEKYPEAPD PHYIYFNPS
281	LHNGVLLVGY QTDPEKQDYH IVNHRGAGW GENGYTRIAE	421	LSMTSSLSTS PELPVAASGS NPTLKVPLPF VDVNRINCDT MSNFVSETEE FGECFVSSSS
321	KRNHROGVAT KASYPLV	481	DAWIVIKKVN SRLLVLIIPP SNYTSIADLV QSKTATIVRS HFEAIFFS

Figure S1: Associated mutations of genes identified in the temperature-sensitive screen.

(A) Whole-genome sequencing of candidates of the ts-screen resulted in the identification of two new alleles for both *cpl-1* and *ccz-1*. The corresponding mutations are listed. (B) The protein sequence of *C. elegans* CPL-1 isoform a comprises of 377 amino acids. The signal peptide is lined; boxed residues correspond to the ERFNIN motif. Residues of the catalytic triad (cysteine, histidine, and asparagine) are marked with an asterisk. The alleles identified in this study are listed together with the corresponding amino acid exchange (triangle). (C) The *C. elegans* CCZ-1 protein sequence comprises of 528 amino acids. The alleles identified in this study are listed together with the corresponding mutations (triangle).

A

ASM-3	ALHLTDLHVD	MFYTVGLEAD	CGTPQCCRPO	DMNVEIVENG	DVKQPAGPWG
SMPD1	ILFLTDLHWD	HDYLEGTDPD	CADPLCCR-R	GSGLPFPAS--	--RPGAGYWG
Identity	* * * * *	* * * * *	* * * * *		* * * *
ASM-3	SVGSCDTPYW	LLTNMLQNI	STAGKLDYIM	VSGDLVSHTV	WAYTPETHSF
SMPD1	EYSKCDLPLR	TLESLLSGLG	-PAGPFDMVY	WTGDIPAHDV	WHQTRQDQLR
Identity	** *	* *	** *	** *	* * *
ASM-3	MVKNLSDTIR	SYFPKTPVYF	AVGNHEGVFV	DNIAPHFTPK	KYHMDWLYKA
SMPD1	ALTTVTALVR	KFLGPVPVYP	AVGNHESTPV	NSFPPPFIEG	NHSSRWLYEA
Identity		*	***	*****	**
ASM-3	MSNAWQGWIP	ADQEKSLIYN	GCYMKKIYDG	LRMISLNNVY	GDRINFWLYI
SMPD1	MAKAWEPWLP	AEALRTLRI	GFYALSPYPG	LRLISLNMNF	CSRENFWLLI
Identity	* ** *	*	* *	* * *	***
ASM-3	NQTDPDGTQ	WLINQLQDAE	NVGDKVHIVA	HIPGSDGEAL	EGYALNYYKI
SMPD1	NSTDPAGQLQ	WLVGELQAAE	DRGDKVHIIG	HIP--PGHCL	KSWSWNYRI
Identity	* *** *	**	*****	*** *	*** *
ASM-3	INRYANTVVG	QFFGHTHSEK	FYMMYANPDD	YKSTPTNVVY	SAPSVTPYSY
SMPD1	VARYENTLAA	QFFGHTHVDE	FEVFYD--EE	TLSPRLAVAF	LAPSATTYIG
Identity	** *	*****	* *	* * *	*** *
ASM-3	YFPAYR				
SMPD1	LNPGRYR				
Identity	* **

Figure S2: Only *asm-3* mutant embryos possess large droplets.

(A) Clustal-W protein alignment of the catalytic region of *C. elegans* ASM-3 (amino acids 143-439) and the human homolog SMPD1 (amino acids 203-498). Identical amino acids are marked with an asterisk. (B and C) DIC images of homozygous mutant embryos of 30-50 cell-stage. (B) *asm-1*(*tm5267*). (C) *asm-2*(*tm3746*). Scale bar: 10 μ m.

Figure S3: The droplet phenotype of *asm-3* is present in oocytes and in all embryonic stages.
(A-F) DIC images of homozygous *asm-3(ok1744)* mutants at various developmental stages. Maturing oocytes (A), zygote (B), 2-cell stage (C), 50-cell stage (D), 200-cell stage (E), and 2-fold stage embryos (F) are shown.

Figure S4: The *asm-3* gene has a maternal and a zygotic component.

(A) Crossing scheme. (B, D, F) DIC images of embryos in early stages. (C, E, G) DIC images of adult gonadal tissue. Homozygous *asm-3(ok1744)* mutant 2-cell embryo (B) and adult oocytes (C). (D) One- to four-cell stage heterozygous *asm-3/+* embryos. (E) The gonadal tissue of heterozygous *asm-3/+* adults. One- and four-cell stage *asm-3* embryos of the F2 generation (F) and adult gonadal tissue (G).

Figure S5: Lipidomic analysis of triglycerides in N2 and *asm-3* mutant embryos.

(A and B) Results of lipidomic analysis of wild-type N2 and *asm-3(ok1744)* mutant embryos. Number of C-atoms (A) and double bonds (B) per triglyceride molecule. Indicated numbers refer to fatty acids. (* $p < 0.05$, ** $p < 0.005$, *** $p < 0.001$).

Table S1: List of candidate ts-alleles from ts-screen

Pt, phenotype
LD, lipid droplets

Gene ID	Gene name	Allele	Droplet Pt at 15°C (diameter)	Droplet Pt at 25°C (diameter)	Droplet Pt in oocytes at 25°C	Droplet Pt in adult intestine at 25°C	Brood size at 15°C	Brood size at 25°C	Lethality rate at 15°C	Lethality rate at 25°C	Embryo size (length)	Other
-	<i>h2</i> (wild-type)	-	< 1.5 µm	< 2.5 µm	< 1.0 µm	mostly < 1.5 µm	274	186	1.6 %	2.7%	50-54 µm	
T03E6.7	<i>cpl-1</i>	<i>t3423</i>	< 1.5 µm	< 3.0 µm (mostly 2.2-2.4 µm) Enlarged LD size	< 1.0 µm	Slightly enlarged LDs	299	235	1.2 %	81.7 %	~ 55 µm	
		<i>t3438</i>	< 1.5 µm	< 2.7 µm (mostly 1.7-2.2 µm) Enlarged LD size	< 1.5 µm Normal LD size	Some enlarged, some normal LD size	244	224	13.3 %	98.2 %	49-53 µm	variable Pt in progeny, incomplete penetrance
F58G11.6	<i>ccz-1</i>	<i>t3403</i>	< 4.0 µm (mostly 2.4-2.8 µm) Enlarged LD size	< 3.3 µm (mostly 2.4-2.8 µm) Enlarged LD size	Enlarged LD size, sometimes oocytes mispositioned	Normal LD size	214	92	59.6 %	98.6 %	~51-54 µm	very slow growth, some adults dumpy, some have protruding vulva
		<i>t3213</i>	< 3.6 µm (mostly 2.6-3.0 µm) Enlarged LD size	< 3.6 µm (mostly 2.4-2.8 µm) Enlarged LD size	Enlarged LD size	Normal LD size	202	65	46.3 %	99.3 %	~ 53 µm (some ~ 45 µm)	very slow growth, sometimes larval arrest, some adults dumpy, some have protruding vulva
K08D10.8 F20C5.1 K07H8.5	<i>scrm-5</i> <i>pme-3</i> <i>K07H8.5</i>	<i>t3425</i>	< 1.0 µm Reduced LD size or number	< 1.0 µm Reduced LD size or number	< 1.0 µm	Normal LD size	114	45	5.4 %	84.3 %	47-50 µm	
K12D12.2 Y17G7B.13	<i>npp-3</i> <i>Y17G7B.13</i>	<i>t3430</i>	< 1.0 µm Reduced LD size or number	< 1.0 µm Reduced LD size or number	< 1.0 µm	n.d.	124	11	6.2 %	97.1%	~ 50 µm	
ZK180.5 F35F11.2	<i>ZK180.5</i> <i>F35F11.2</i>	<i>t3434</i>	< 1.0 µm Reduced LD size or number	< 1.0 µm Reduced LD size or number	< 1.0 µm	n.d.	66	30	11.7 %	22.0 %	~ 55 µm	
M88.3 C15H7.3	<i>M88.3</i> <i>C15H7.3</i>	<i>t3218</i>	< 1.0 µm	< 1.0 µm	< 1.0 µm	n.d.	50	13	6.7 %	28.0 %	~ 52 µm	slightly more granules
n.d.	n.d.	<i>t3426</i>	< 1.0 µm	< 1.0 µm	< 1.0 µm	n.d.	n.d.	n.d.	n.d.	43.4 %	~50 µm	sick' adults at 25°C, larval lethal at 25°C
n.d.	n.d.	<i>t3427</i>	< 1.0 µm	< 1.5 µm	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	
n.d.	n.d.	<i>t3241</i>	< 1.0 µm	< 2.0 µm	n.d.	n.d.	386	412	2.4 %	100.0 %	n.d.	
n.d.	n.d.	<i>t3428</i>	< 1.5 µm Normal LD size	< 2.4 µm (mostly < 2.0 µm)	< 1.5 µm	Normal LD size	80	71	16.7 %	90.4 %	50-54 µm	
n.d.	n.d.	<i>t3270</i>	< 1.5 µm	< 2.0 µm	n.d.	n.d.	n.d.	n.d.	4.5 %	100.0 %	n.d.	
n.d.	n.d.	<i>t3298</i>	< 1.5 µm	< 2.4 µm	n.d.	n.d.	40	14	66.6 %	100.0 %	n.d.	
n.d.	n.d.	<i>t3429</i>	< 1.5 µm	< 2.0 µm	n.d.	n.d.	n.d.	n.d.	17.5 %	30.0 %	n.d.	
n.d.	n.d.	<i>t3431</i>	< 1.5 µm	< 1.5 µm	n.d.	n.d.	n.d.	n.d.	50.0 %	72.2%	n.d.	
n.d.	n.d.	<i>t3432</i>	< 1.0 µm	< 1.5 µm	n.d.	n.d.	n.d.	n.d.	n.d.	50.0 %	n.d.	
n.d.	n.d.	<i>t3322</i>	< 1.5 µm	< 1.5 µm	n.d.	n.d.	n.d.	n.d.	n.d.	50.0 %	n.d.	
n.d.	n.d.	<i>t3433</i>	< 1.0 µm	< 1.5 µm	n.d.	n.d.	159	72	17.8 %	100.0 %	n.d.	
n.d.	n.d.	<i>t3331</i>	< 2.8 µm	< 3.0 µm	n.d.	n.d.	n.d.	n.d.	52.5 %	63.3 %	n.d.	
n.d.	n.d.	<i>t3333</i>	< 1.0 µm	< 1.5 µm	n.d.	n.d.	n.d.	n.d.	n.d.	20.0%	n.d.	
n.d.	n.d.	<i>t3343</i>	< 2.0 µm (mostly < 1.0 µm)	< 2.5 µm (mostly < 1.5 µm)	n.d.	n.d.	n.d.	n.d.	0.0 %	n.d.	n.d.	
n.d.	n.d.	<i>t3435</i>	< 1.5 µm	<2.5 µm	n.d.	n.d.	n.d.	n.d.	n.d.	50.0 %	n.d.	
n.d.	n.d.	<i>t3436</i>	< 1.5 µm	< 2.0 µm	n.d.	n.d.	n.d.	n.d.	n.d.	100.0 %	n.d.	
n.d.	n.d.	<i>t3437</i>	< 1.5 µm	<2.5 µm	n.d.	n.d.	n.d.	n.d.	n.d.	60.0 %	n.d.	
n.d.	n.d.	<i>t3351</i>	< 2.5 µm (mostly < 1.0 µm)	< 2.8 µm (mostly < 1.5 µm)	n.d.	n.d.	235	187	10.0 %	97.8 %	n.d.	
n.d.	n.d.	<i>t3439</i>	< 1.5 µm	< 2.0 µm	n.d.	n.d.	73	84	73.1 %	97.6 %	n.d.	
n.d.	n.d.	<i>t3440</i>	< 1.5 µm	< 2.6 µm (mostly < 1.5 µm)	n.d.	n.d.	162	98	0.0 %	87.5 %	n.d.	

Table S2: List of candidate genes from database screen (.xlsx, 179 KB)

Available for download as a .xlsx file at

<http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.179127/-/DC1/TableS2.xlsx>

Table S3: List of candidate genes in RNAi screen

Emb pt, embryonic lethal

Droplet pt, occurrence of large droplets

F1, F1 generation

F2, F2 generation

Gene ID	Gene name	RNAi clone (Geneservices)	Emb pt	Droplet pt	
				F1	F2
E01A2.1	<i>E01A2.1</i>	I-1F04		x	x
Y23H5A.3	<i>Y23H5A.3</i>	I-1G14	x		
C32E8.8	<i>ptr-2</i>	I-1J07	x	x	x
ZK973.11	<i>ZK973.11</i>	I-1L22			
W01B11.5	<i>pqn-72</i>	I-1O20			
C46H11.2	<i>C46H11.2</i>	I-2A16			
C44E4.5	<i>C44E4.5</i>	I-2C23			
B0207.9	<i>B0207.9</i>	I-2D12			
Y76G2A.2	<i>Y76G2A.2</i>	I-2D22			
C06A5.10	<i>C06A5.10</i>	I-2F18			
B0261.1	<i>B0261.1</i>	I-2G18			
T19B4.5	<i>T19B4.5</i>	I-2J19			
T09B4.8	<i>T09B4.8</i>	I-2N04			
T09B4.9	<i>T09B4.9</i>	I-2N06	x		
C30H7.2	<i>C30H7.2</i>	I-2O04	x		
F27C1.4	<i>F27C1.4</i>	I-2O12			
F27C1.6	<i>F27C1.6</i>	I-2O14			
H37N21.1	<i>hpo-11</i>	I-3D03			
ZC328.3	<i>ZC328.3</i>	I-3E23			
T10E9.6	<i>T10E9.6</i>	I-3M03			
C30F12.1	<i>C30F12.1</i>	I-3M04			
H05L14.2	<i>H05L14.2</i>	I-3N22			
K07A1.13	<i>K07A1.13</i>	I-4D16			
C03D6.1	<i>C03D6.1</i>	I-4F18	x		
C03D6.3	<i>cel-1</i>	I-4F20			
DY3.2	<i>lmn-1</i>	I-4G24			
F48F5.5	<i>fce-1</i>	I-4H24	x		
F36A2.7	<i>F36A2.7</i>	I-4I22	x		
F26H9.6	<i>Rab-5</i>	I-4J01	x		
F10D11.6	<i>F10D11.6</i>	I-4K23	x		
F26E4.6	<i>F26E4.6</i>	I-4L02			
C17E4.3	<i>C17E4.3</i>	I-4L09			
Y67A6A.1	<i>Y67A6A.1</i>	I-4L24	x		
K02B12.3	<i>sec-12</i>	I-4M19	x		
F59C6.5	<i>F59C6.5</i>	I-5C22			
ZK39.6	<i>clec-97</i>	I-5H11			
ZK39.7	<i>clec-98</i>	I-5H13			
F25D7.4	<i>F25D7.4</i>	I-5O15			
K11D2.2	<i>asah-1</i>	I-6G18			
H28O16.1	<i>H28O16.1</i>	I-6M02	x		
T26E3.7	<i>T26E3.7</i>	I-6M18			
W04A4.5	<i>W04A4.5</i>	I-6P02			
W09G3.1	<i>W09G3.1</i>	I-7A09			
W09G3.6	<i>W09G3.6</i>	I-7A09			
W09G3.5	<i>fbxa-93</i>	I-7A15			
ZK909.3	<i>ZK909.3</i>	I-7K24			
Y47G6A.23	<i>lpd-3</i>	I-7P03			
K07A1.2	<i>dut-1</i>	I-9C01			

B0205.6	<i>B0205.6</i>	I-9F20			
D1007.5	<i>D1007.5</i>	I-9I-06			
B0252.2	<i>asm-1</i>	II-10A04			
T10D4.1	<i>T10D4.1</i>	II-10B21			
C41C4.8	<i>cdc-48.2</i>	II-10C11			
C17C3.1	<i>C17C3.1</i>	II-10M03			
Y38E10A.2	<i>Y38E10A.2</i>	II-10O20			
C17G10.7	<i>C17G10.7</i>	II-11A06			
F12E12.2	<i>F12E12.2</i>	II-11A08			
F22E5.1	<i>F22E5.1</i>	II-11B24			
Y48C3A.1	<i>Y48C3A.1</i>	II-11D14			
ZK1290.1	<i>ZK1290.1</i>	II-11E10			
W07E6.4	<i>prp-21</i>	II-1A22			
F23F1.5	<i>F23F1.5</i>	II-1C03	x		
F23F1.6	<i>F23F1.6</i>	II-1C05			
C23H3.2	<i>C23H3.2</i>	II-1C13			
C23H3.5	<i>C23H3.5</i>	II-1C19	x	x	x
C50D2.1	<i>C50D2.1</i>	II-1E03	x		
T02H6.4	<i>T02H6.4</i>	II-1I16			
T02H6.7	<i>T02H6.7</i>	II-1I22			
C08G5.2	<i>fbxc-48</i>	II-1M12			
F46F5.10	<i>F46F5.10</i>	II-1O22			
K02E7.11	<i>K02E7.11</i>	II-1P23			
F52C6.12	<i>F52C6.12</i>	II-2A10	x		
F08D12.4	<i>F08D12.4</i>	II-2D12			
C16C4.2	<i>math-12</i>	II-2K21			
C41H7.4	<i>C41H7.4</i>	II-2L10	x		
F52C6.2	<i>F52C6.2</i>	II-2O13	x		
T10D4.14	<i>T10D4.14</i>	II-2P14	x		
Y27F2A.6	<i>Y27F2A.6</i>	II-3A15		x	x
C01F1.1	<i>C01F1.1</i>	II-3B24	x		
B0286.4	<i>ntl-2</i>	II-3F06			
C46E10.9	<i>C46E10.9</i>	II-3O02			
R03H10.4	<i>R03H10.4</i>	II-3P07			
B0034.2	<i>B0034.2</i>	II-4B15			
F21H12.3	<i>F21H12.3</i>	II-4F19			
C29H12.6	<i>C29H12.6</i>	II-4H11			
ZK546.3	<i>ZK546.3</i>	II-4I17			
T27F7.1	<i>T27F7.1</i>	II-4K19			
K05F1.6	<i>K05F1.6</i>	II-4L08			
F13H8.9	<i>F13H8.9</i>	II-4L19	x		
F33G12.2	<i>F33G12.2</i>	II-4M15	x		
F33G12.4	<i>lrr-1</i>	II-4M19			
F12A10.8	<i>F12A10.8</i>	II-4O04			
C32D5.11	<i>C32D5.11</i>	II-4P07			x
F55C12.7	<i>tag-234</i>	II-4P12			
E04F6.6	<i>E04F6.6</i>	II-5B13			
C56E6.4	<i>C56E6.4</i>	II-5E15			
F18C5.2	<i>wrn-1</i>	II-5E23			
C56C10.3	<i>vps-32.1</i>	II-5G17	x		
C56C10.10	<i>C56C10.10</i>	II-5I07			
F54H5.3	<i>F54H5.3</i>	II-5I15			
T19D12.1	<i>T19D12.1</i>	II-5I21			
T19D12.3	<i>T19D12.3</i>	II-5K01			
T02G5.4	<i>T02G5.4</i>	II-5K10			
C18H9.5	<i>C18H9.5</i>	II-5M03			
T14B4.8	<i>T14B4.8</i>	II-5O01			
T07F8.1	<i>T07F8.1</i>	II-5O04			
C27H5.4	<i>C27H5.4</i>	II-5O18			

T05H10.6	<i>pdha-1</i>	II-5P10		x	x
F54C9.9	<i>F54C9.9</i>	II-6A22			
C18E9.5	<i>C18E9.5</i>	II-6B09			
T05C12.1	<i>T05C12.1</i>	II-6C03			
F37B12.3	<i>F37B12.3</i>	II-6D13			
E02H1.1	<i>E02H1.1</i>	II-6H24			
F14E5.1	<i>F14E5.1</i>	II-6K13			
C01G6.5	<i>C01G6.5</i>	II-6L05	x		
C01G6.9	<i>C01G6.9</i>	II-6L13		x	x
F46C5.6	<i>F46C5.6</i>	II-6M06			
Y53C12B.2	<i>Y53C12B.2</i>	II-6P04			
ZK1321.1	<i>ZK1321.1</i>	II-6P14			x
E04D5.1	<i>E04D5.1</i>	II-7E08	x		
C06A1.1	<i>cdc-48.1</i>	II-7K02			
C06A1.3	<i>C06A1.3</i>	II-7K06			
C14A4.12	<i>C14A4.12</i>	II-7M12			
T06D8.5	<i>T06D8.5</i>	II-7P09			
K12D12.4	<i>K12D12.4</i>	II-8E11			
Y38F1A.9	<i>oig-2</i>	II-8F17			
F08G2.7	<i>F08G2.7</i>	II-8L18			
F49C5.6	<i>str-223</i>	II-8M02			
Y57A10A.27	<i>Y57A10A.27</i>	II-8M03			
Y38E10A.3	<i>Y38E10A.3</i>	II-8M12			
Y57A10A.14	<i>Y57A10A.14</i>	II-8O09	x		
Y54G11A.7	<i>Y54G11A.7</i>	II-9I03	x		
Y51H7BR.8	<i>Y51H7BR.8</i>	II-9N21			
M01F1.3	<i>M01F1.3</i>	III-1D22	x		
E02H9.3	<i>E02H9.3</i>	III-1H09			
C36A4.5	<i>C36A4.5</i>	III-1N08			
F56F3.4	<i>F56F3.4</i>	III-2A24			
C03C10.3	<i>rnr-2</i>	III-2C17	x		
W03A5.4	<i>W03A5.4</i>	III-2H16			
T23F11.4	<i>T23F11.4</i>	III-2I04	x		
F09F7.4	<i>F09F7.4</i>	III-2N02			
R07E5.1	<i>R07E5.1</i>	III-2O01	x		
T12A2.1	<i>T12A2.1</i>	III-3C20	x		
F20H11.2	<i>let-765</i>	III-3D01	x		
B0280.6	<i>B0280.6</i>	III-3D06	x		
T12A2.5	<i>T12A2.5</i>	III-3E02			
R05H11.1	<i>R05H11.1</i>	III-3H07	x		
T04C9.2	<i>T04C9.2</i>	III-3K19			
F57B9.3	<i>F57B9.3</i>	III-3L15			
C18H2.1	<i>C18H2.1</i>	III-4E13	x		
F02A9.6	<i>glp-1</i>	III-4L02	x		
ZC84.5	<i>ZC84.5</i>	III-4N24	x		
M01A8.1	<i>M01A8.1</i>	III-4P16	x		
K01G5.8	<i>K01G5.8</i>	III-5B14			
C07A9.2	<i>C07A9.2</i>	III-5M11	x	x	x
Y43F4B.5	<i>Y43F4B.5</i>	III-6H21	x		
Y37D8A.8	<i>Y37D8A.8</i>	III-6M22	x	x	x
Y71H2B.8	<i>Y71H2B.8</i>	III-7K12			
Y53G8AL.2	<i>Y53G8AL.2</i>	III-7O21			
T17H7.1	<i>T17H7.1</i>	III-8K14			
F40F12.7	<i>F40F12.7</i>	III-9A01			
F42A6.3	<i>F42A6.3</i>	IV-1B20	x		
F56B3.2	<i>F56B3.2</i>	IV-1E04	x		
K02D7.4	<i>dsc-4</i>	IV-1G17			
C09G12.5	<i>C09G12.5</i>	IV-1H22	x		x
F52C12.2	<i>F52C12.2</i>	IV-1J01			

C09G12.9	<i>tsg-101</i>	IV-1J06			
M70.1	<i>M70.1</i>	IV-1L17	x	x	x
M70.3	<i>M70.3</i>	IV-1L21			
W03G1.7	<i>asm-3</i>	IV-1M17	x		
T23E1.1	<i>T23E1.1</i>	IV-1N21	x		
T26C12.1	<i>T26C12.1</i>	IV-1P01	x		
F36H12.15	<i>F36H12.15</i>	IV-2B24			
Y24D9A.6	<i>Y24D9A.6</i>	IV-2C02			
Y17G9A.5	<i>str-172</i>	IV-2D17	x		
R105.1	<i>R105.1</i>	IV-2I18			
F29B9.11	<i>F29B9.11</i>	IV-2K16	x		
F38A5.1	<i>odr-8</i>	IV-3A12			
F45E4.6	<i>F45E4.6</i>	IV-3H24			
C33H5.17	<i>C33H5.17</i>	IV-3P16	x		
F13B6.1	<i>F13B6.1</i>	IV-3P23			
C06A6.2	<i>C06A6.2</i>	IV-3P24			
Y11D7A.8	<i>Y11D7A.8</i>	IV-4B14			
Y11D7A.9	<i>Y11D7A.9</i>	IV-4B16	x	x	x
F20D12.2	<i>F20D12.2</i>	IV-4E03	x		
F49C12.11	<i>F49C12.11</i>	IV-4F06	x		
F49C12.15	<i>F49C12.15</i>	IV-4F14			
C53B4.5	<i>col-119</i>	IV-4H13	x		
F42G8.5	<i>islo-1</i>	IV-4I03	x	x	x
F56D5.9	<i>F56D5.9</i>	IV-4J12	x		
B0218.5	<i>B0218.5</i>	IV-4K01			
F21D5.5	<i>F21D5.5</i>	IV-4O02	x	x	x
F56H11.6	<i>F56H11.6</i>	IV-4P02			
C04G2.2	<i>C04G2.2</i>	IV-5C20			
C04G2.5	<i>C04G2.5</i>	IV-5C24			
R10H10.3	<i>R10H10.3</i>	IV-5D01	x		
ZK593.3	<i>ZK593.3</i>	IV-5D20	x		
T13F2.6	<i>T13F2.6</i>	IV-5E23	x		
K07F5.13	<i>npp-1</i>	IV-5K01			
K07F5.14	<i>K07F5.14</i>	IV-5K03	x		
C24F3.2	<i>C24F3.2</i>	IV-5K18			
F25H8.1	<i>F25H8.1</i>	IV-5M03			
F25H8.2	<i>F25H8.2</i>	IV-5M05			
F01G4.6	<i>F01G4.6</i>	IV-5N16			
C42C1.6	<i>C42C1.6</i>	IV-6D19			
F55B11.4	<i>F55B11.4</i>	IV-7D11			
LLC1.2	<i>LLC1.2</i>	IV-7D15			
H25K10.1	<i>H25K10.1</i>	IV-7H22	x		
B0513.7	<i>B0513.7</i>	IV-7I04	x		
F52B11.2	<i>F52B11.2</i>	IV-7M04			
Y41E3.11	<i>Y41E3.11</i>	IV-7N09			
Y40H7A.1	<i>str-78</i>	IV-7P01			
Y51H4A.15	<i>Y51H4A.15</i>	IV-7P20	x		
T28F3.5	<i>T28F3.5</i>	IV-8A18	x		
Y41D4A.6	<i>Y41D4A.6</i>	IV-8D07			
Y67D8C.7	<i>Y67D8C.7</i>	IV-8F10			
ZK550.6	<i>ZK550.6</i>	IV-8O09			
Y9C9A.9	<i>str-158</i>	IV-8P16			
T11G6.8	<i>T11G6.8</i>	IV-9C07			
F58F9.1	<i>F58F9.1</i>	IV-9K16			
F11G11.5	<i>F11G11.5</i>	none			
F58E10.3	<i>F58E10.3</i>	none			
ZK455.4	<i>asm-2</i>	none			
C33E10.7	<i>C33E10.7</i>	none			
F53H1.2	<i>F53H1.2</i>	none			

F58H1.6	<i>F58H1.6</i>	none			
R07B7.9	<i>R07B7.9</i>	none			
Y38H8A.7	<i>Y38H8A.7</i>	none			
C04G2.3	<i>C04G2.3</i>	none			
T28C6.4	<i>col-117</i>	none			
C29F4.1	<i>col-125</i>	none			
C04F5.8	<i>C04F5.8</i>	none			
F10A3.11	<i>F10A3.11</i>	V-10D14			
F10A3.13	<i>str-108</i>	V-10D18			
K06B4.3	<i>K06B4.3</i>	V-10E22			
F11A5.3	<i>F11A5.3</i>	V-10F14			
F11A5.4	<i>F11A5.4</i>	V-10F14			
T26E4.10	<i>T26E4.10</i>	V-10M14			
C14A6.6	<i>C14A6.6</i>	V-11P04			
F58G4.2	<i>str-90</i>	V-14P21			
B0554.6	<i>dod-20</i>	V-15K16			
F47B8.3	<i>F47B8.3</i>	V-16A17			
C14C6.8	<i>C14C6.8</i>	V-16K15			
F07C4.1	<i>str-47</i>	V-16K19			
B0554.7	<i>B0554.7</i>	V-1K21		x	x
T08B1.1	<i>T08B1.1</i>	V-1P06	x		
F56A4.3	<i>F56A4.3</i>	V-2C24			
Y73C8C.4	<i>Y73C8C.4</i>	V-2F06			
K03B4.1	<i>K03B4.1</i>	V-3D18	x		
C37H5.3a	<i>C37H5.3</i>	V-3J12			
C37H5.8	<i>hsp-6</i>	V-3J20			
T22F3.3	<i>T22F3.3</i>	V-3O11			
R01B10.6	<i>R01B10.6</i>	V-4F19			
CD4.4	<i>vps-37</i>	V-4G04			
K09H11.15	<i>K09H11.15</i>	V-4K12			
F31F7.1	<i>F31F7.1</i>	V-4L07			
K12B6.8	<i>K12B6.8</i>	V-4N03			
F26D11.10	<i>che-7 / inx-4</i>	V-5B14			
ZC513.1	<i>ZC513.1</i>	V-5D02			
ZK682.2	<i>ZK682.2</i>	V-6D21			
F59E11.5	<i>F59E11.5</i>	V-6G06			
F25G6.9	<i>F25G6.9</i>	V-6I23			
F11A3.2	<i>F11A3.2</i>	V-6L21			
T27E4.7	<i>T27E4.7</i>	V-6M08			
T06E8.1	<i>acl-2</i>	V-7A03			
Y32F6A.4	<i>Y32F6A.4</i>	V-7A04			x
T11F9.10	<i>T11F9.10</i>	V-7F18			
T21C9.5	<i>lpd-9</i>	V-7G06			
T21C9.11	<i>T21C9.11</i>	V-7G18			
H27A22.1	<i>H27A22.1</i>	V-7G24	x		
H09F14.1	<i>H09F14.1</i>	V-7I02			
C32C4.1	<i>C32C4.1</i>	V-7I04			
W05E10.2	<i>W05E10.2</i>	V-7N08		x	x
F18E2.1	<i>F18E2.1</i>	V-8D01			
T28H10.3	<i>T28H10.3</i>	V-8I24			
C14C10.4	<i>C14C10.4</i>	V-8M04	x		
F15B9.10	<i>F15B9.10</i>	V-8N05		x	x
C56A3.7	<i>cav-1, cav-2</i>	V-8N20			
F53F4.11	<i>F53F4.11</i>	V-9A05	x		
H37A05.1	<i>lpin-1</i>	V-9A13	x		
F08H9.5	<i>clec-227</i>	V-9D09			
R11H6.5	<i>R11H6.5</i>	V-9H21			
R11H6.6	<i>R11H6.6</i>	V-9H23			
F47B8.4	<i>F47B8.4</i>	V-9M12			

R10D12.10	<i>R10D12.10</i>	V-9O13	x	x	x
T19D7.7	<i>T19D7.7</i>	X-1I11			
K02E10.4	<i>K02E10.4</i>	X-1L08			
K02E10.5	<i>K02E10.5</i>	X-1L10			
F19G12.2	<i>F19G12.2</i>	X-1M16			
T02C5.3	<i>igcm-3</i>	X-1P12			
Y71H10B.1	<i>Y71H10B.1</i>	X-2A13			
T19D2.2	<i>prl-1</i>	X-2B03			
T19D2.3	<i>T19D2.3</i>	X-2B05			
C15C7.5	<i>C15C7.5</i>	X-2I23			
R160.4	<i>R160.4</i>	X-2L13			
K09C4.5	<i>K09C4.5</i>	X-2M13			
F15A8.6	<i>F15A8.6</i>	X-2N07			
C31H2.3	<i>C31H2.3</i>	X-2P04			
C05E11.2	<i>C05E11.2</i>	X-2P23			
H05L03.3	<i>H05L03.3</i>	X-3A08			
F22A3.5	<i>ceh-60</i>	X-3B19			
T14E8.4	<i>T14E8.4</i>	X-3D03			
C56E10.3	<i>C56E10.3</i>	X-3F03			
C54G7.1	<i>C54G7.1</i>	X-3G23			
F41C6.6	<i>F41C6.6</i>	X-3L19			
F46G11.3	<i>tag-257</i>	X-3M23			
ZK154.4	<i>ZK154.4</i>	X-4C17			
T14B1.1	<i>T14B1.1</i>	X-4F08			
F53A9.4	<i>jbts-14</i>	X-4M12			
T06H11.2	<i>T06H11.2</i>	X-5A01			
F59F5.1	<i>slcf-1</i>	X-5I23			
C34F6.2	<i>col-178</i>	X-5K06			
C34F6.3	<i>col-179</i>	X-5K08			
C34F6.11	<i>C34F6.11</i>	X-5K08			
F54F7.2	<i>F54F7.2</i>	X-5L17			
F13E6.1	<i>F13E6.1</i>	X-5M13			
T01C1.2	<i>mbr-1</i>	X-5O03			
R07A4.2	<i>R07A4.2</i>	X-5O09			
F28H6.2	<i>ceh-89</i>	X-6J03			
R03E1.2	<i>R03E1.2</i>	X-6J17			
F31F6.2	<i>F31F6.2</i>	X-6L16			
K02D3.2	<i>K02D3.2</i>	X-6O04			
F20D1.9	<i>F20D1.9</i>	X-6P14			
T27A8.4	<i>T27A8.4</i>	X-7E18			
F39D8.2	<i>psa-3</i>	X-7G21			
C30G4.5	<i>C30G4.5</i>	X-7L17			
C08A9.2	<i>C08A9.2</i>	X-7L23			
F38E9.4	<i>F38E9.4</i>	X-7M06			

T03E6.7	<i>cpl-1</i>	V-11I07	positive control
C49F5.1	<i>sams-1</i>	X-5P21	positive control
-	-	L4440	negative control

Table S4: 22 genes of the database screen are associated with a droplet phenotype in the embryo.

The genes listed below show a droplet phenotype in embryos. Droplet pt, occurrence of large droplets in embryos of the F1 or F2 generation; Emb pt, embryonic lethal phenotype

Gene ID	Gene name	Allele/RNAi	Droplet pt	Emb pt	Possible function
B0554.7	<i>B0554.7</i>	RNAi	F1	-	Ion channel regulatory protein (UNC-93 like)
C01G6.9	<i>C01G6.9</i>	RNAi	F1	-	unknown
C07A9.2	<i>C07A9.2</i>	RNAi	F1	+	transcription factor, BUD 31 family
C09G12.5	<i>C09G12.5</i>	RNAi	F2	+	unknown
C23H3.5	<i>C23H3.5</i>	RNAi	F1	+	unknown
C32D5.11	<i>C32D5.11</i>	RNAi	F2	-	peroxisomal protein
C32E8.8	<i>ptr-2</i>	RNAi	F1	+	ortholog of Drosophila and human PTCHD3 (sterol sensing domain protein)
E01A2.1	<i>E01A2.1</i>	RNAi	F2	-	Glutamate-cysteine ligase modifier subunit
F15B9.10	<i>F15B9.10</i>	RNAi	F1	-	component of cytoskeleton
F21D5.5	<i>F21D5.5</i>	RNAi	F1	+	DNA-repair
F42G8.5	<i>islo-1</i>	RNAi	F1	+	Interactor with SLO-1
M70.1	<i>M70.1</i>	RNAi	F1	+	unknown
R10D12.10	<i>R10D12.10</i>	RNAi	F1	+	tau-tubulin kinase
T05H10.6	<i>pdha-1</i>	RNAi	F1	-	Pyruvate dehydrogenase (lipoamide) ALPHA 1
W05E10.2	<i>W05E10.2</i>	RNAi	F1	-	unknown
Y11D7A.9	<i>Y11D7A.9</i>	RNAi	F1	+	unknown
Y27F2A.6	<i>Y27F2A.6</i>	RNAi	F1	-	unknown
Y32F6A.4	<i>Y32F6A.4</i>	RNAi	F2	-	unknown
Y37D8A.8	<i>Y37D8A.8</i>	RNAi	F1	+	lysosomal membrane protein
ZK1321.1	<i>ZK1321.1</i>	RNAi	F2	-	unknown
C46C11.1	<i>hosl-1</i>	<i>tm2369</i>	+	-	ortholog of hormon-sensitive lipase (LIPE)
W03G1.7	<i>asm-3</i>	<i>ok1744</i>	+	-	ortholog of sphingomyelin phosphodiesterase 1 (SMPD1)