Unique and Overlapping Functions of Formins Frl and DAAM During Ommatidial Rotation and Neuronal Development in *Drosophila*

Gretchen Dollar,^{*,†} Rita Gombos,[‡] Austen A. Barnett,^{*,†} David Sanchez Hernandez,^{*,†}

Saw M. T. Maung,^{*,†} Jozsef Mihály,^{‡,§} and Andreas Jenny^{*,†,1}

^{*}Department of Developmental and Molecular Biology and [†]Department of Genetics, Albert Einstein College of Medicine, Bronx, New York 10461, [‡]Office for Subsidized Research Units, Hungarian Academy of Sciences (HAS), H-1051 Budapest, Hungary, and

[§]Institute of Genetics, Biological Research Centre, HAS, Szeged H-6726, Hungary

ORCID ID: 0000-0001-5989-8212 (A.J.)

ABSTRACT The noncanonical Frizzled/planar cell polarity (PCP) pathway regulates establishment of polarity within the plane of an epithelium to generate diversity of cell fates, asymmetric, but highly aligned structures, or to orchestrate the directional migration of cells during convergent extension during vertebrate gastrulation. In *Drosophila*, PCP signaling is essential to orient actin wing hairs and to align ommatidia in the eye, in part by coordinating the movement of groups of photoreceptor cells during ommatidial rotation. Importantly, the coordination of PCP signaling with changes in the cytoskeleton is essential for proper epithelial polarity. Formins polymerize linear actin filaments and are key regulators of the actin cytoskeleton. Here, we show that the diaphanous-related formin, Frl, the single fly member of the FMNL (formin related in leukocytes/formin-like) formin subfamily affects ommatidial rotation in the *Drosophila* eye and is controlled by the Rho family GTPase Cdc42. Interestingly, we also found that *frl* mutants exhibit an axon growth phenotype in the mushroom body, a center for olfactory learning in the *Drosophila* brain, which is also affected in a subset of PCP genes. Significantly, Frl cooperates with Cdc42 and another formin, DAAM, during mushroom body formation. This study thus suggests that different formins can cooperate or act independently in distinct tissues, likely integrating various signaling inputs with the regulation of the cytoskeleton. It furthermore highlights the importance and complexity of formin-dependent cytoskeletal regulation in multiple organs and developmental contexts.

KEYWORDS planar cell polarity; formin; cytoskeleton; neural development; noncanonical Wnt signaling

In multicellular organisms, epithelia are polarized along the apical–basal axis as well as within the epithelial plane, imparting cells with a sense of direction. The latter polarity, planar cell polarity (PCP), can be propagated over hundreds of cell diameters and is evident in many vertebrate and non-vertebrate organs. In its simplest form, PCP is seen in cuticular structures of insects such as *Drosophila melanogaster*, where sensory bristles on the thorax or actin hairs produced by each wing cell all point in the same direction, posteriorly on the thorax and distally in the wing (Goodrich and Strutt

2011; Maung and Jenny 2011). Analogously in vertebrates, hair follicles, cilia in the lung and brain ventricles, and stereocilia in the sensory cells of the inner ear are arranged in a stereotypic manner that is crucial for proper organ function ((Montcouquiol *et al.* 2003; Devenport and Fuchs 2008; Tissir *et al.* 2010; Vladar *et al.* 2012; reviewed in Goodrich and Strutt 2011; Devenport 2014). More complex examples of PCP signaling include coordination of cell fate specification and ommatidial orientation in the *Drosophila* compound eye, the orientation of cell division, and cell migration during vertebrate gastrulation (reviewed in Jenny 2010; Goodrich and Strutt 2011; Devenport 2014). One of the main signaling pathways involved in the establishment of PCP is the non-canonical Wnt/Frizzled (Fz)-PCP pathway that was originally identified in *Drosophila* and is highly conserved in vertebrates. Mutations in PCP genes cause misaligned ommatidia or wing hairs in flies and severe developmental abnormalities such as

Copyright © 2016 by the Genetics Society of America

doi: 10.1534/genetics.115.181438

Manuscript received July 30, 2015; accepted for publication January 18, 2016; published Early Online January 20, 2016.

Supporting information is available online at www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.181438/-/DC1.

¹Corresponding author: Albert Einstein College of Medicine, Chanin 503, 1300 Morris Park Ave., Bronx, NY 10461. E-mail: andreas.jenny@einstein.yu.edu

Figure 1 Loss of *fri* causes a PCP-like defect in the eye. Tangential sections of adult eyes are shown with corresponding schematic representation of ommatidial orientations underneath. Anterior is to the left and dorsal is up in all sections. Black and red arrows represent dorsal and ventral chiral forms of ommatidia, and circles mark lost R-cells in all figures. (A) Control eyes expressing *Dcr2* under control of the *sevenless* (*sev*) promoter. The inset shows an enlarged single ommatidium with numbered rhabdomeres, the light-sensitive organelles of R-cells (note that only seven rhabdomeres are visible at once in a section, as R7 lies on top of R8). (B) Compared to control (A), knock down of *fri* causes ommatidial rotation defects. (C–F) *fri*^{Ex83} (C) and *fri*^{Ex88} (E) clones marked by the absence of pigment show PCP-like defects (due to the *Minute* heterozygous background, most of the sections are homozygous mutant). (D and F) Lethality as well as orientation defects of *fri*^{Ex83} (D) and *fri*^{Ex88} (F) are rescued by a genomic copy of *fri* (*Fos*^{Frl}). Note that throughout this work, all alleles of *fri* correspond to the excision alleles indicated in the background of a *FRT80* chromosome with a fosmid adding back all additional genes removed by the excisions (*Fos*^{AB}).

neural-tube and left-right defects, and ciliopathies in mammals (Murdoch *et al.* 2001; Ross *et al.* 2005; Borovina *et al.* 2010; Song *et al.* 2010).

Core PCP factors including the transmembrane proteins Frizzled (Fz), Flamingo [Fmi; also known as (aka) Stan], Van Gogh (Vang, aka Stbm), the adaptor proteins Dishevelled (Dsh), and Prickle are required for PCP establishment in all tissues (reviewed in Strutt and Strutt 2009; Vladar *et al.* 2009; Maung and Jenny 2011). Fz and Dsh transduce the PCP sig-

nal to Rho family GTPases, including RhoA and Rac, leading to changes in the cytoskeleton (Strutt *et al.* 1997; Boutros *et al.* 1998; Fanto *et al.* 2000; Maung and Jenny 2011). Additionally, at least in the *Drosophila* eye, Rac activates a JNK-type MAP kinase cascade leading to a transcriptional response (Fanto *et al.* 2000; Jenny 2010). Overall, however, the activation of Rho GTPases and their mechanism of action downstream of Dsh during PCP signaling are only poorly understood.

Figure 2 Frl likely is autoinhibited. (A) Schematic of domain structure of Frl with amino acid number of predicted domains indicated (relative to version Frl-PC). GBD, GTPase binding domain; DID, diaphanous inhibitory domain; DD, dimerization domain; FH1 and FH2, formin homology 1/2 domains; DAD, diaphanous autoregulatory domain. Underneath: schematic and coordinates of constructs used in this study; note that Flag Δ FH2 is 11 aa shorter than the Gst- Δ FH2. (B–D) Images of eyes of flies overexpressing full-length Frl (B), constitutively active Frl (Frl Δ GBD-DID; C), or dominant negative Frl (Frl Δ FH2; D) under the control of *sev-Gal4*. Note that in contrast to full-length Frl that does not affect eye formation, Frl Δ GBD-DID and Frl Δ FH2 cause severely rough eyes. (E) Gst pull-down experiment of *in vitro* translated proteins indicated on the left with Gst-fusion proteins indicated on top. While only a small amount of full-length Frl is binding to the FH2-CT domain, much stronger binding is found between the GBD-DID-DD-CC region with Gst-FH2-CT and Gst-CT, consistent with an intramolecular interaction of the N and C termini of Frl. (F) Coomassie staining of a typical pull-down gel of the assays in E with corresponding Gst proteins indicated on top.

In the *Drosophila* eye, PCP is manifest in the mirror-symmetric arrangement of its ~800 ommatidia or facets dorsal and ventral to the midline (equator) (Wolff and Ready 1993; Singh *et al.* 2005; reviewed in Jenny 2010). This arrangement is established in third instar larvae, after ommatidial preclusters emerge from the morphogenetic furrow (MF), a transient indentation that sweeps from posterior to anterior across the eye imaginal disc (Treisman and Heberlein 1998; Roignant and Treisman 2009; Jenny 2010). Fz-PCP signaling determines the R3 vs. R4 photoreceptor (R-cell) fates in the precluster consisting of the R8, R3/4, and R2/5 cells, with the precursor

closer to the equator becoming R3. This in turn triggers rotation of the clusters by 90° clockwise on the dorsal and counterclockwise on the ventral side of the equator, a process requiring changes in the cytoskeleton mediated by Rho kinase (Winter *et al.* 2001). During pupal stages, R-cell clusters remodel to form adult ommatidia with six outer photoreceptors (R1–R6) arranged in a trapezoid around the two inner photoreceptors (R7/8; see insert in Figure 1A) that are aligned in a stereotypic parallel and antiparallel arrangement. In core PCP mutants, R3/4 fates and ommatidial rotation are randomized.

Members of the seven formin families are important regulatory proteins of the cytoskeleton characterized by the presence of two formin homology (FH) domains. The FH1 domain binds profilin in complex with G-actin, while the FH2 domain nucleates and polymerizes unbranched actin filaments (reviewed in Higgs 2005; Faix and Grosse 2006; Liu *et al.* 2010). Although all formins affect actin filaments, little is known about their cooperation in the regulation of the cytoskeleton. In addition to the FH1 and FH2 domains, the subfamily of diaphanous-related formins (DRFs) such as Diaphanous, Daam (Dishevelled-associated activator of morphogenesis), and Frl (formin related in leukocytes; FMNLs in humans) also contain an N-terminal Rho family GTPase binding domain (GBD), a diaphanous inhibitory domain (DID), and a dimerization domain (DD) followed by a coiled coil domain (CC; see also schematic in Figure 2A). C terminal to the FH2 domain, there is the diaphanous autoregulatory domain (DAD) that binds to the GBD-DID domain, in most cases keeping the formin in an inactive conformation (Higgs 2005; Vaillant *et al.* 2008; Liu *et al.* 2010). DRFs are usually activated by a Rho family GTPase binding to the GBD domain, thereby alleviating autoinhibition. In addition, factors binding to the C-terminal DAD domain can also contribute to DRF activation. For example, Dsh is able to activate Daam1 by binding to its C-term/DAD domain in tissue culture and during convergence and extension in *Xenopus* (Habas *et al.* 2001; Liu *et al.* 2008). Surprisingly, *Drosophila* DAAM mutants show no PCP defects in the eye or wing. Instead, DAAM is required to pattern the tracheal cuticle via the modification of the actin cytoskeleton, for sarcomeric thin filament formation, and for axon growth, the latter function under control of the Rac GTPase (Matusek *et al.* 2006, 2008; Molnar *et al.* 2014; Gombos *et al.* 2015). Thus, the role of formins and the extent of their cooperation in PCP signaling in fruit flies remained elusive.

Here, we find that *Drosophila* Frl (CG32138), the sole representative of the Frl formin subfamily in flies, likely contributes to ommatidial rotation during eye development, as RNAi-mediated knockdown of *frl* or eyes mutant for *frl* show mild rotation defects. Furthermore, binding assays and *in vivo* overexpression studies suggest that full-length Frl is autoinhibited. Coimmunoprecipitation and pull-down assays showed that Frl strongly binds to GTP-bound forms of Cdc42 and, more weakly, to Rac, but genetic interaction assays suggest that, *in vivo*, Frl is activated by Cdc42. Although Frl can bind Dsh *in vitro*, we found no *in vivo* evidence for such an

interaction during PCP establishment in the eye. Interestingly, while we found no obvious redundancy between *frl* and *DAAM* during PCP signaling in the eye, they cooperate during R-cell morphogenesis or maintenance and during axon growth in the mushroom body, a process also requiring a subset of the PCP genes (Shimizu *et al.* 2011; Ng 2012). Overall, our data suggest that Frl may contribute to ommatidial rotation in the eye and, importantly, has context-dependent, redundant, and nonredundant functions with *DAAM* during neural development in *Drosophila*.

Materials and Methods

Fly strains

*dsh*¹, *sep-Gal4*, *sev-Gal4*, *FRTG13 shg^{p34-1}*, *FRTG13 shg¹⁶²⁹* (Mirkovic and Mlodzik 2006), *svp-lacZ⁰⁷⁴⁸²*, *sev > Fz* and *sev-Dsh* (Boutros *et al.* 1998), and *sev-RacV12* (Luo *et al.* 1994) were kind gifts from M. Mlodzik (Mount Sinai Hospital) and L. Luo (Stanford University). *nmo^{DB}* (amorphic), *nmo^{P1}*, and *CadNΔ14* (removing N-cadherin 1 and 2) (Prakash *et al.* 2005) were kind gifts from E. Verheyen (Simon Fraser University) and T. Clandinin (Stanford University). *VDRC^{GD34412}*, *frl^{HMS00445}* (BL no. 32447), *UAS-Cdc42* (BL no. 28873), *Df(2L)BSC148* (uncovering several genes including the N-cadherins), *cdc42¹*, *cdc42²*, *cdc42⁵*, *cno²*, *aos^{Δ7}*, *phyl¹²²⁴⁵*, and *sty^{Δ5}* were obtained from the Vienna *Drosophila* RNAi Center (VDRC) or Bloomington Stock Centers. *rac1^{J11}*, *rac2^Δ*, *rac1^{J11} rac2^Δ FRT2A*, *rac1^{J11} rac2^Δ FRT2A mtl^Δ* (all from the Bloomington Stock Center), *UAS-Dcr2*, and *DAAM^{Ex68} FRT19A* (a null allele) and *DAAM^{Ex1}* and *DAAM^{Ex4}* (both hypomorphic alleles giving rise to viable hemizygous males) were described previously (Ng *et al.* 2002; Matusek *et al.* 2006; Dietzl *et al.* 2007; Ng 2012; Gombos *et al.* 2015).

Fly genetics

frl^{Ex83} and *frl^{Ex88}* mutations were generated by standard excision using Δ2–3 to remobilize the isogenized *P*-element insertions P[d06832] (after recombining away a second site lethal; see Supporting Information, Figure S1B). *frl^{ExK62}* was generated by imprecise excision of isogenized P[HP36775]. *frl^{Ex83}* and *frl^{Ex88}* retain various amounts of scrambled *P*-element sequences (including some UAS sequences for *frl^{Ex83}*) at the original 5' *P* end in intron 1 of *frl*, but delete the start codon (schematic in Figure S1B). The upstream breakpoints of *frl^{Ex83}* and *frl^{Ex88}* are in *hsc70Cb* and *neur14* and delete 21.4 kb and 16 kb, respectively. *frl^{ExK62}* deletes 11.4 kb starting in *CG6833* and ending in intron 2 of *frl*, thus removing the first two exons of *frl*, including the start codon (Figure S1B). While *frl^{Ex83}* and *frl^{Ex88}* are of different genetic background than *frl^{ExK62}*, all excisions obtained delete additional upstream genes. To prevent any phenotypes that may be due to these genes, FlyFOS 027901 (*Fos^{AB}*, add back; Figure S1B) (Ejsmont *et al.* 2009) encoding all missing upstream genes but ending in intron 1 of *frl* was integrated into landing site VK00027 (Venken and Bellen 2005) and recombined onto a *FRT80* chromosome with each excision. *frl^{Ex} FRT80 Fos^{AB}* were used throughout this work as

frl alleles. FlyFOS 023275 encoding *frl* (and other genes) integrated in VK00002 on 2L rescues lethality and eye phenotype of all *frl* alleles.

To generate mutant clones in the mushroom body (MB), we used the mosaic analysis with a repressible cell marker (MARCM) technique (Lee *et al.* 1999). For *frl* mutant clones, *elav-Gal4*, *UAS-mCD8::GFP*, *hsFLP/+*; *FRT80B tubP-Gal80/frl^{Ex83} FRT80B Fos^{AB}*, *elav-Gal4*, *UAS-mCD8::GFP*, *hsFLP/+*; *FRT80B*, *tubP-Gal80/frl^{Ex88} FRT80B Fos^{AB}*, and *elav-Gal4*, *UAS-mCD8::GFP*, *hsFLP/+*; *FRT80B*, *tubP-Gal80/frl^{ExK62} FRT80B Fos^{AB}*, and for wild-type control clones *elav-Gal4*, *UAS-mCD8::GFP*, *hsFLP/+*; *FRT80B tubP-Gal80/FRT80B* pupae were heat shocked at 37° for 30 min at 50 hr after puparium formation (APF). The morphology of the MB clones was analyzed in flies 3–7 days old.

Immunohistochemistry

Third instar eye discs were fixed and stained as described (Freeman *et al.* 1992; Weber *et al.* 2000). For analysis of mushroom bodies, adult brains were dissected in cold PBS, fixed in 4% paraformaldehyde (diluted in PBS) at room temperature (RT) for 20 min. Primary antibodies were applied overnight at 4°. After the secondary antibodies and standard washing steps, samples were mounted in PBS:glycerol (1:4). Confocal images were captured on an Olympus FV1000 LSM microscope, images were edited with Adobe Photoshop 7.0CE and Olympus FW10-ASW. Antibodies used were rabbit anti-Sal (1:50; a kind gift from C. Desplan (New York University)), mouse α-β-Gal (1:200; Promega), rat α-Elav (1:200; DSHB 7E8A10), rabbit α-GFP (1:1000; Life Technologies), and mouse α-FasII (1:50; DSHB 1D4) as primary antibodies, and the corresponding Alexa-488 or Alexa-546 coupled secondary antibodies (Life Technologies).

Eye analysis

Mutant eye clones were induced with *eyFLP* (Newsome *et al.* 2000) or *hsFLP* (1 hr 15 min at 38° at 0–72 hr after egg laying) using the Minute technique (Morata and Ripoll 1975) as indicated. Adult heads were embedded and sectioned as described (Jenny 2011). Ommatidia in three to six eyes were scored for quantifications with an ommatidium considered misrotated if misaligned by >20°. For *DAAM frl* double mutant analysis, an *eyFLP* insertion on the third chromosome was recombined onto a *W⁺[70c] FRT80* chromosome (note that the location of the *eyFLP* transgene on chromosome 3 is unknown), which was then combined with a *FRT19A* (*W⁺*). *FRT19A*; *W⁺[70c] FRT80 eyFLP* males were crossed with *DAAM^{Ex68} FRT19A/ FM6*; *frl^{Ex} FRT80 Fos^{AB}/ TM6B* virgins.

Plasmids

All PCR amplified fragments were sequence verified. Frl constructs are based on cDNA clone AT04875 (encoding protein variant PC). pSca-Frl was assembled in multiple steps from a N-terminal PCR product starting with CG32138_CT_for_Mfe to an internal BmgBI site, a BmgBI/BsrGI fragment of AT04875,

and a C-terminal PCR product starting at the *Bsr*GI site and ending with CG32138_rev2_Xba. pSca-FrlΔGBD-DID and pSca-FrlΔFH2 correspond to PCR products amplified with primers CG32138_DeltaGBD_for and CG32138_rev2_Xba, and CG32138_for_MfeNco and CG32138_DeltaFH2_rev_Xba, respectively. The constitutively active FrlΔGBD-DID thus starts with amino acid 346 ([M]KGGH. . .) and the dominant negative FrlΔFH2 ends with amino acid 677 (GAMT; see also schematic in Figure 2A).

Gst fusion constructs: Gst-Dsh fragments are as in Jenny *et al.* (2005). pGexKG-GBD-DID and pGexKG-GBD-DID-DD-CC were cloned as PCR fragments using primers FrlGBD_Xba_f and FrlGBD_Hind3_r and FrlGBDplusFH3_Hind3_r, respectively into the *Xba*I/*Hind*III sites of pGexKG (see schematic in Figure 2A). The FH2 domain with and without Cterm was cloned as PCR product with primers FH2plusC Bam_F and CterXhoR_Frl and FH2minusC_Xho_R into the *Bam*HI/*Xho*I sites of pGex4T1. pGex4T1-Cterm was cloned as PCR product amplified with primers CterBamF_Frl and CterXhoIR_Frl into the *Bam*HI/*Xho*I sites of pGex4T1. pGex-3X-Rac encoding *Drosophila* Rac1 was a kind gift from S. Parkhurst (Fred Hutchinson Cancer Research Center). *Drosophila* Cdc42 was amplified with Cdc42_Gex_for_Bam and Cdc42_Gex_rev_Sal and cloned into the corresponding sites of pGex4T1.

In vitro translation constructs: Full-length Frl was *in vitro* translated from cDNA clone AT04875 using SP6 polymerase. The GBD-DID, GBD-DID-DD-CC, DD-CC, FH2, and FH2-CT domains were cloned as PCR products (see schematic in Figure 2A) using primers FrlGBD_Nsi_f/FrlGBD_Nco_r, FrlGBD_Nsi_f/FrlGBDplusFH3_Nco_r, Frl_FH3_for_Nsi/FrlGBDplusFH3_Nco_r, FrlFH2plusC_Nsi_F/FrlFH2minusC_Nco_R, FrlFH2plusC_Nsi_F/FrlCterm_Nco_R, respectively, into the *Nsi*I/*Nco*I sites of pBTH (Jenny *et al.* 2003).

Cell culture plasmids: pRK5_Myc_RhoA_Q63L, pRK5_Myc_Rac1_Q61L, pCR_Myc_Rac1_N17, pDNA3_Myc_Cdc42_Q61L, and pCR_Myc_Cdc42_N17 encoding Myc tagged constitutively active and dominant negative human Rho GTPase versions were kind gifts from D. Cox (Developmental and Molecular Biology, Albert Einstein College of Medicine) (Cox *et al.* 1997; Abou-Kheir *et al.* 2008; Cammer *et al.* 2009). pCS2(105)-Dsh-Flag was described previously (Yanfeng *et al.* 2011). pCaspTubPA-Frl was generated by inserting the *Kpn*I/*Xba*I fragment of pSca-Frl into the corresponding site of pCaspTubPA. pAcFlag3 was made by inserting oligos triple flag upper and lower into the *Kpn*I/*Eco*RI sites of a version of pAc5.1 (Invitrogen) in which the internal *Bam*HI had previously been destroyed. pAcFlag3-Frl was generated by cloning a *Mfe*I/*Xba*I fragment of pCaspTubPA-Frl into the *Eco*RI/*Xba*I sites of pAcFlag3. pCS2(105)-Flag3-FrlΔFH2 was cloned by inserting a *Kpn*I(blunt)/*Nae*I fragment of pAcFlag3-Frl into the blunted *Eco*RI site of pCS2(105). Note that this construct is 11-aa shorter than the GST version and ends at amino acid 666. pCsMT-Frl and pCsMT-FrlΔFH2 were

made by inserting *Nco*I/*Xba*I fragments of pSca-Frl and pSca-FrlΔFH2 into the corresponding sites of pCsMT (six N-terminal Myc tags). pCsMT-FrlΔGBD-DID was cloned as a *Hind*III(blunt)/*Xba*I fragment of pSca-FrlΔGBD-DID into the *Nco*I(blunt)/*Xba*I sites of pCsMT.

Transgenics: pUAST versions of Frl, FrlΔGBD-DID, and FrlΔFH2 were made by cloning the *Mfe*I/*Xba*I fragments of the corresponding pSca constructs into the *Eco*RI/*Xba*I sites of pUAST (Brand and Perrimon 1993). Transgenes were injected by Genetic Services (Sudbury, MA) and Rainbow Transgenics (Camarillo, CA).

Coimmunoprecipitation and pulldown assays

Coimmunoprecipitations were done with a modified version of Jenny *et al.* (2005). Briefly, HEK293 cells were transfected with 3 μg of each plasmid using polyethylenimine (PEI). The cells were collected after 48 hr and washed with ice cold 1× PBS and lysed in 1 ml lysis buffer (50 mM Tris pH 8.0, 150 mM NaCl, 1% Triton X-100, 1 mM EDTA supplemented with 1:1000 dilution of 1 mM benzamidine, and 1:100 dilutions of 10 μM leupeptin, and 1 μM pepstatin). Cleared supernatants were precipitated with 1 μg of mouse anti-FlagM2 antibody (Sigma) that was bound to 20 μl of 50% Protein G beads overnight at 4°. Immunoprecipitates were washed three times with cold wash buffer (lysis buffer with 0.1% Triton) and resuspended in 15 μl of 2× Laemmli buffer. Samples were separated on a 12% SDS gel and Western blot analysis was performed using standard protocols.

GST fusion protein expression and purification and GST pull-down assays for Frl and Dsh interactions were performed as described (Jenny *et al.* 2003) using 1 μg of GST fusion protein and 3 μl of a standard 50 μl TNT coupled transcription/translation reaction (Promega). Gst-Cdc42 and Gst-Rac pull-downs were modified from Magie *et al.* (1999). Per reaction, 1 μg of respective Gst protein was bound to 15 μl of Gst Sepharose beads (Amersham) in PD buffer (Jenny *et al.* 2003). After three washes, beads were washed twice more with exchange buffer (EB) (50 mM Hepes pH 7.08, 50 mM NaCl, 5 mM EDTA, 0.1 mM EGTA, 0.1 mM DTT) and resuspended in 100 μl EB supplemented with either 100 μM GMP-PNP or GDP and 25 mM MgCl₂ for nucleotide loading and incubated for 30 min at 30° while rocking. Excess EB was then removed. A total of 3 μl *in vitro* translated Frl-GBD-DID-DD-CC were precleared with 10 μl GST-Sepharose beads in 100 μl HEPES PD buffer (20 mM Hepes 7.5, 250 mM NaCl, 0.5% Triton, 10 mM MgCl₂, 1 mM EDTA, 1 mM DTT, 1 μM pepstatin, 10 μM leupeptin, 1 mM benzamidine) for 30 min at 4°. The precleared Frl-GBD-DID-DD-CC was then combined with the nucleotide-loaded Gst protein beads with an additional 400 μl HEPES PD buffer and rotated for 45 min at 4° followed by three washes in Hepes PD buffer. Proteins were eluted by addition of 15 μl 2× SDS loading buffer and resolved on 12% SDS PAGE gels. After Coomassie staining, the gels were dried and exposed and scanned on a phosphorimager (Fuji 9000). Signals were quantified using MultiGauge V3.0.

Reagent availability

All strains and plasmids constructed and used are available from the indicated *Drosophila* stock centers or upon request.

Oligonucleotides

CG32138_for_MfeNco, TATCAATTGGCCACCATGGGCGCCG TCAAATCG; CG32138_rev2_Xba, ATATCTAGACTAACAATC CATTTCCTCCAGG; CG32138_ΔFH2_rev_Xba, ATATTCTAG ATTAGGTCATGGCGCCATCGGGA; CG32138_ΔGBD_for, AT CAATTGGGATCCGCCACCATGAAGGGAGGACACGAAATC; FrlGBD_Xba_f, GATCTCTAGAGAGCCGGAGCCAGCAACCCATG; FrlGBD_Hind3_r, GATCAAGCTTCTACTTTACCAGACAGATGGC TGC; FrlGBDplusFH3_Hind3_r, GATCAAGCTTCTAAGAACC ATCGCCGCTGGAAGC; FH2plusC Bam_F, GATCGGATCCGC CATGACCATCAAACGCAAG; FH2minusC_Xho_R, GATCCTC GAGCTACAGCTTTTTCTCGGCACCGA; CterXhoR_Frl, GATC CTCGAGCTAACAATCCATTTCTCCAGGGT; CterBamF_Frl, GA TCGGATCCGCGCACTCGGTGCGCGAGAAAA; FrlGBD_Nsi_f, GA TCATGCATAGCCGGAGCCAGCAACCCATG; FrlGBD_Nco_r, GAT CCCATGGCTACTTTACCAGACAGATGGCTGC; FrlGBDplusFH3_Nco_r, GATCCCATGGCTAAGAACCATCGCCGCTGGAAGC; Frl_FH3_for_Nsi, GATCATGCATGGAGGACACGAAATCATTT; FrlFH2plusC_Nsi_F, GATCATGCATGCCATGACCATCAAACG CAAG; FrlFH2minusC_Nco_R, GATCCCATGGCTACAGCTTTT TCTCGGCACCGA; FrlCterm_Nco_R, GATCCCATGGCTAACAA TCCATTTCTCCAGGGT; Cdc42_Gex_for_Bam, TATAGGATC CATGCAAACCATCAAGTGCG; Cdc42_Gex_rev_Sal, ATATGTC GACTTATAAGAATTTGCACTTCC; triple flag lower, AATTCGGA TCCCTTGTCTGTCGTCTCTTGTAGTCGATGTCGTGGTCTTGT TAGTCGCCGTCGTGGTCTTGTAGTCCATGGTGGTAC; triple flag upper, CACCATGGACTACAAGGACCACGACGGCGACTA CAAGGACCACGACATCGACTACAAGGACGACGACGACGACAA GGGATCCG.

Results

Loss of *frl* affects ommatidial rotation

The *Drosophila* genome encodes one gene each for six of the seven families of formins, including CG32138, the representative of the Frl/Fmnl subfamily (Higgs 2005). To test whether Frl may be involved in PCP signaling in the *Drosophila* eye, we knocked down *frl* in the eye using RNAi under the control of *sevenless*-Gal4 (*sev*) (Basler *et al.* 1991), which at the time of photoreceptor recruitment is expressed mainly in R3/R4. Compared to *sev* > *Dcr2* that was coexpressed to enhance knockdown (Figure 1A) (Dietzl *et al.* 2007), RNAi-mediated knockdown of *frl* showed mild, but clear PCP-like defects (Figure 1B and Figure S1A). We then generated *frl* mutants by imprecise excision of *P*-elements, HP36775 and d06832, located upstream of *frl* or within the first intron of *frl*, respectively (see *Materials and Methods* and Figure S1B). To prevent potential defects caused by additional deletion of upstream genes annotated in FlyBase (FlyBase Consortium 2003), we integrated a genomic rescue DNA fragment encoding all upstream genes, but terminating after exon 1 of *frl* (FlyFOS 027901;

Fos^{AB} for add back; Figure S1B) (Ejsmont *et al.* 2009) on the right arm of chromosome 3. The *Fos*^{AB} transgene was then recombined onto each *frl* excision that had previously been recombined with *FRT80*. All three *frl*^{Ex} *FRT80* *Fos*^{AB} chromosomes are thus mutant only for *frl* and are homozygous lethal (and used as *frl* mutants throughout this work). Loss of *frl* in a *Minute* heterozygous background to more easily obtain large mutant areas (Morata and Ripoll 1975) caused weak rotation defects in all alleles with *frl*^{Ex83}, *frl*^{Ex88}, and *frl*^{ExK62} showing 1.8% (*n* = 620 mutant ommatidia), 1.9% (*n* = 371), and 1.7% (*n* = 946) ommatidia with PCP-like defects, respectively [Figure 1, C and E and not shown; note that wild-type *FRT80* *M* clones show no defects (*n* = 293)]. Importantly, viability and the eye phenotype were rescued by FlyFOS 023275, which encodes the complete *frl* gene (*Fos*^{Frl}; Figure 1, D and F; Figure S1B and not shown). Frl may thus contribute to ommatidial rotation in the *Drosophila* eye.

Dominant negative Frl causes ommatidial rotation defects

Constitutively active (CA) and dominant negative (DN) DRF forms can be constructed by deleting the N-terminal regulatory GBD and DID domains (ΔGBD-DID) or FH2 domain and C terminus (ΔFH2), respectively (Habas *et al.* 2001; Liu *et al.* 2008; Vaillant *et al.* 2008). To test whether Frl may be regulated by autoinhibition, we made transgenic flies expressing full-length Frl, CA-Frl (FrlΔGBD-DID), and DN-Frl (FrlΔFH2) under the control of *UAS-Gal4* (Figure 2A). While the eyes of flies overexpressing full-length Frl under control of *sev-Gal4* (*sev* > *Frl*) looked normal (Figure 2B; several independent transgenic lines showed similar results), eyes of flies expressing the putative CA-Frl (Figure 2C) or DN-Frl (Figure 2D) were severely rough. These results are consistent with Frl, like other DRFs being autoinhibited. Indeed, Gst pull-down experiments showed that an *in vitro* translated GBD-DID-DD-CC fragment strongly bound to the Gst-FH2-CT (including the DAD domain), but more weakly to the C terminus only fused to Gst (Figure 2E; schematic in Figure 2A). Consistent with an intramolecular interaction, full-length Frl only very weakly bound to Gst-FH2-CT. Neither the GBD-DID nor the DD-CC domains alone were sufficient to bind to Gst-FH2-CT nor were the *in vitro* translated FH2 or FH2-CT fragments able to bind to Gst-GBD-DID (Figure 2, E and F shows a Coomassie stained gel of a typical pull-down; note that we were unable to purify soluble Gst-DD-CC). Thus, the absence of a GOF phenotype of full-length Frl and the ability of N- and C-terminal fragments of Frl to interact indicate that *Drosophila* Frl is likely to be autoinhibited.

To examine the eye phenotypes in more detail, we analyzed sectioned eyes of animals expressing CA- and DN-Frl under the control of *sep-Gal4*, which is weaker than *sev-Gal4*, as it uses the enhancer and promoter of the *sep* gene instead of a heat-shock minimal promoter (Basler and Hafen 1990; Basler *et al.* 1991; Strutt and Strutt 2002). Overexpression of *sep* > CA-Frl caused loss of R-cells (Figure 3A). In contrast, overexpression of *sep* > DN-Frl consistently also caused rotation

defects in adult eyes (Figure 3B; note that stronger lines also lead to gaps that may be due to the loss of R-cell clusters; Figure 5G). Coexpression of CA-Frl (using several independent transgenic lines) with DN-Frl did not alter the phenotype of DN-Frl (not shown), most likely due to inadequate expression levels. The phenotype of DN-Frl is stronger than of *frl* mutant clones, which could be due to expression of DN-Frl being more acute than a loss of *frl* in mutants, which may be able to partially compensate for its loss. Alternatively, DN-Frl may also inhibit other formins (see below). Importantly, however, the defects arose at the time of PCP signaling in third instar larvae, as, compared to control (Figure 3C), misrotating ommatidial clusters and clusters lacking R-cells were seen in eye discs stained for *seven-up* (*svp*)-*lacZ*, which is most strongly expressed in the R3/4 pair and more weakly in R1/6 during the early rotation period (Figure 3D) (Mlodzik *et al.* 1990; Fanto *et al.* 1998; Fanto *et al.* 2000; Gaengel and Mlodzik 2003; Jenny *et al.* 2003). Similar results were obtained upon staining of third instar eye discs with antibodies against Salm (Spalt-major) marking R3/4 early in rotation (Figure 3, E and F) (Domingos *et al.* 2004). Thus, CA-Frl affects R-cell morphogenesis (or maintenance), and DN-Frl, similar to *frl* mutants, implicates Frl in ommatidial rotation in the eye.

Frl is regulated by Cdc42 in vivo

DRFs can be activated by GTP-bound Rho family GTPase binding to the N-terminal GBD-DID domain (Habas *et al.* 2001; Vaillant *et al.* 2008; Liu *et al.* 2010). To determine which of the Rho family GTPases could bind Frl, we cotransfected Myc-tagged constitutively active human Rho, Rac, and Cdc42 and dominant negative Rac and Cdc42 with Flag-tagged FrlΔFH2 (schematic in Figure 2A) into HEK293 cells. Immunoprecipitation of Flag-FrlΔFH2 revealed a strong interaction with CA Cdc42 Q61L, but not with its DN version, Cdc42 N17 (Figure 4A). Upon prolonged exposure, we reproducibly also found a weaker coprecipitation of CA Rac Q61L (Figure 4A). No interaction was identified with RhoA, nor between Dsh (as a negative control) and any of the GTPases. To confirm Frl interactions, we used *Drosophila* Cdc42 and Rac1 fused to Gst loaded with the nonhydrolyzable GTP analog GMP-PNP or GDP to pull down *in vitro* translated ³⁵S-methionine-labeled FrlGBD-DID-DD-CC. FrlGBD-DID-DD-CC bound to Cdc42 with about an eightfold increase of the active, GTP-bound form of Cdc42 over GDP-Cdc42 (Figure 4B; quantified in Figure 4D). Again, weaker binding was found for GTP-Rac1 with about a fourfold enrichment over GDP-Rac1 (Figure 4C; quantified in Figure 4E). These findings are consistent with human FMNL1/2 or zebrafish Fmnl3 activation by Cdc42 (Block *et al.* 2012; Kuhn *et al.* 2015; Richards *et al.* 2015; Wakayama *et al.* 2015) and mouse Frl1/Fmnl1 showing a weak preference for Rac-GTP over Rac-GDP, although the latter failed to bind Cdc42 (Yayoshi-Yamamoto *et al.* 2000).

To test whether the interactions between Frl and Cdc42 and Rac were physiologically relevant, we tested for genetic interactions between Frl and Rho GTPases. Similar to *sep-Gal4*

driven overexpression of CA-Frl, overexpression of Cdc42 caused R-cell loss (compare Figure 5, A and E with Figure 3A; quantified in Figure 5D). Importantly, this R-cell loss was partially, but significantly suppressed by removal of one gene dose of any of the three *frl* excision alleles (Figures 5, B and C and not shown; quantified in Figure 5D) and completely suppressed by coexpression of DN FrlΔFH2 (Figure 5F; quantified in Figure 5I), suggesting that *cdc42* can act as an upstream activator of *frl*. In addition, coexpression of Cdc42 with *sep > DN-Frl* also fully suppressed the rotation phenotypes of *sep > DN-Frl* (compare Figure 5F with 5G; quantified in Figure 5I), while removing one gene dose of *cdc42* significantly enhanced *sep > DN-Frl* (Figure 5H; quantified in Figure 5I). We did not detect a genetic interaction between *cdc42* and *frl* in *frl Minute* clones in the background of hemizygous males of the hypomorphic *cdc42⁵* allele (not shown; however, see below for a genetic interaction between *cdc42* and *frl* in the mushroom body). These data are consistent with potential Frl activation by Cdc42 *in vivo*, whereby DN-Frl competes with endogenous Frl for Cdc42. Thus overexpressed Cdc42 would not only activate endogenous Frl, but would also titrate DN-Frl. Similarly, removal of one gene dose of *cdc42* would shift the balance toward DN-Frl and thus would enhance the phenotype of dominant negative Frl. Our *in vitro* binding and *in vivo* genetic interaction data suggest that Cdc42 can activate Frl in the eye.

In contrast, we did not observe genetic interactions between *frl* and *rac* *in vivo*. Activated Rac (*sev-Rac^{V12}*) was shown to cause PCP phenotypes in the eye that can be modified by removal of one gene dose of PCP pathway components (Fanto *et al.* 2000). However, the *frl* excision alleles did not consistently suppress or enhance *sev-Rac^{V12}* (Figure S2), nor was overexpression of *Rac^{V12}* and DN-Frl synergistic, suggesting that Rac binding to Frl may not be relevant in the eye.

Dsh binds Frl in vitro

The core PCP protein Dsh activates the formin xDaam1 during PCP signaling in *Xenopus* gastrula by associating with the DAD domain (Habas *et al.* 2001; Liu *et al.* 2008). Similar to xDaam1 (Habas *et al.* 2001), we found that *in vitro* translated full-length Frl, but not the Frl-FH2 domain bound to Gst-tagged full-length Dsh and its basic PDZ region (Figure 6A; the core PCP protein Pk served as positive control) (Jenny *et al.* 2003). *In vitro* translated Frl-FH2-CT also bound to Gst-Dsh-bPDZ over background (Figure 6A). We confirmed this interaction by cotransfecting HEK293 cells with Dsh-GFP and Myc-tagged Frl, FrlΔGBD-DID, and FrlΔFH2, respectively. Consistent with the above results, FrlΔGBD-DID lacking the N-terminal region of Frl, but not FrlΔFH2 lacking the C-terminal part was able to coprecipitate Dsh-GFP (Figure 6B). The inability of full-length Frl to coprecipitate Dsh may indicate that in tissue culture cells (or a lysate thereof), Dsh on its own may not be able to overcome the autoinhibitory interaction of Frl.

We then tested for genetic interactions between *frl* and *dsh*. First, we tested if *frl* alleles could dominantly modify the PCP

Figure 3 Eye phenotypes of constitutively active and dominant negative Frl. (A) Expression of activated *sep* > CA-Frl (Δ GBD-DID) induces a loss of R-cells. (B) *sep* > DN-Frl (Δ FH2) causes PCP-like defects reflected mainly by misrotated ommatidia and leads to loss of some R-cells. (C–F) PCP defects caused by *sep* > *Frl* Δ FH2 arise in third instar eye discs. Confocal images of *w*¹¹¹⁸ (C and E) and *sep* > *Frl* Δ FH2 (D and F) third instar eye discs stained for nuclear β -Gal expressed from *svp-lacZ*²⁰⁷⁴⁸² enhancer trap line (green in C and D) or stained for Salm marking R3/4 at early stages of rotation (E and F). R-cells are counterstained with ELAV (red). Dorsal parts of the disc just posterior to the MF are shown (top). Schematic of the cluster orientation (bottom).

phenotype of males hemizygous for the viable, PCP-specific allele *dsh*¹ (Fahmy and Fahmy 1959; Boutros *et al.* 1998; Axelrod 2001). Neither removal of one gene dose of *frl* by *frl*^{Ex83} nor *frl*^{ExK62} was able to modify the PCP phenotype of *dsh*¹ males (Figure 6, C–E; quantified in Figure 6F; note that *dsh*¹ can be suppressed by the core PCP allele *pk*¹) (Jenny *et al.* 2005). Similarly, the phenotype of homozygous mutant *frl*^{Ex83} or *frl*^{ExK62} clones in *dsh*¹ hemizygotes looked similar to *dsh*¹/Y alone (Figure 6F) and the phenotype of *frl*^{Ex83} or *frl*^{ExK62} clones was not enhanced in females lacking one gene dose of *dsh* (not shown). We then tested whether the GOF PCP phenotypes of *sev* > *Fz* or *sev*-*Dsh* (Boutros *et al.* 1998) could be modified by removal of a gene dose of *frl* or a deficiency uncovering the *frl* locus. However, similar to *DAAM* (Matusek *et al.* 2006), *frl* showed no such dominant interaction. Thus, despite Frl binding to Dsh *in vitro*, we were not able to find *in vivo* evidence to support a functional requirement for such an interaction during PCP signaling in the eye and Frl may thus act downstream of Cdc42 rather than the core PCP factors during eye polarization.

We then tested whether DN-Frl genetically interacts with other components known to be required for ommatidial rotation (reviewed in Jenny 2010; Goodrich and Strutt 2011; Devenport 2014). We did not observe a dominant genetic interaction of DN-Frl with EGF pathway-related genes [*argos* (*aos*), *sprouty* (*spy*), *phyllopod* (*phyl*), or *canoe*; not shown] or the rotation-specific kinase *nemo* (*nmo*; Figure S3) or E-cadherin (*shg*; Figure S3). In contrast, simultaneous removal of one gene dose of both *Drosophila* N-cadherin genes (*CadN1/2*) known to control the speed and extent of ommatidial rotation (Mirkovic and Mlodzik 2006) enhanced DN-Frl, again suggesting that DN-Frl affects ommatidial rotation

(Figure 7; quantified in Figure S3; note that *CadN1/2* may also show a trend to enhance R-cell loss).

Frl cooperates with DAAM during axon growth in the mushroom body

Xenopus Daam1 is required for PCP signaling during convergent extension (Habas *et al.* 2001), but *Drosophila* *DAAM* mutants show no PCP-related eye or wing phenotypes (Matusek *et al.* 2006), whereas *frl* mutants exhibit a mild PCP phenotype in the eye and thus could act redundantly with *DAAM* during PCP establishment in the eye. We thus generated *DAAM*; *frl* double mutant eye clones in which homozygous double mutant R-cells are the only cells that lack the rhabdome-associated pigment granules (e.g., magenta arrowheads in Figure 8, A–C; note that heterozygous or homozygous single mutant cells cannot be distinguished). The recovered double mutant clones were small, but mosaic regions showed no evidence for redundancy between *DAAM* and *frl* in PCP signaling in the eye (not shown). However, compared to *DAAM* (Figure 8A) or *frl* (Figure 1, C and E) single mutants, *DAAM*; *frl* double homozygous R-cells showed a partially penetrant, cell autonomous small rhabdome phenotype (Figure 8, B and C). Our data thus suggest that Frl and *DAAM* have redundant roles in R-cell morphogenesis or maintenance.

We then investigated whether *frl* and *DAAM* cooperate in other neural contexts. A subset of PCP signaling genes have been implicated in axon growth and/or guidance control in the fly MB, an integration and memory center for odorant input (Heisenberg *et al.* 1985; Shimizu *et al.* 2011; Ng 2012; reviewed in Zou 2012), and we recently have shown that *DAAM* functions as a molecular effector of the PCP system in this context (Gombos *et al.* 2015). The MB is composed of three types of neurons, the γ , α' / β' and α / β neurons (Kenyon cells),

Figure 4 Frl preferentially binds to Cdc42 in its GTP bound form. (A) Coimmunoprecipitation of the indicated, Myc-tagged constitutively active (Q) or dominant negative (N) human Rho family GTPases by Flag-Frl Δ FH2 from HEK293 cell lysates shows that Frl strongly binds Cdc42 Q61L and, as visible on the darker exposure, more weakly to Rac Q61L, but not to their dominant negative versions. Flag-Dsh was used as negative control. (Top) Immunoprecipitations. (Bottom) Cell lysates. (B) *In vitro* translated [35 S]-labeled Frl-GBD-DID-DD-CC binds to purified *Drosophila* Cdc42 loaded with the nonhydrolyzable GTP analog GMP-PNP, but not to Gst-Cdc42 loaded with GDP or Gst alone. (C) Frl-GBD-DID-DD-CC more weakly binds to purified *Drosophila* GMP-PNP-Rac1, but only barely to GDP-Rac1 or Gst alone. (D and E) Quantification of typical Cdc42 (D) and Rac1 (E) Gst pull downs performed in triplicates of assays analogous to the ones shown in B and C. *t*-test; ****P* < 0.001; **P* < 0.05.

that develop in a stereotypic way (Lee *et al.* 1999). Each Kenyon cell extends a neurite that splits into a dendritic and an axonal part. The axons extend anteroventrally forming a prominent fiber tract called the peduncle (Figure 8D). In adults, after reaching the anterior side, axons of the γ neurons run medially and form the γ lobe, whereas axons of the α'/β' and α/β neurons bifurcate and their dorsal branches form the α and α' lobes while their medially growing branches form the β and β' lobes (see schematic in Figure 8D). After initial neurite outgrowth, axonal extension toward the appropriate target areas requires the core PCP genes *vang*, *fz*, *dsh*, *fmi*, and *pk* (Shimizu *et al.* 2011; Ng 2012). In addition, we recently showed that *DAAM* functions as cytoskeletal effector of the PCP guidance system in the MBs. In ~70% of the *DAAM* mutant lobes, MB axons either fail to form, terminate their growth before reaching the tip of the neuropile lobes, or fail to project into the proper lobes, indicating that while *DAAM* is required for axon formation in the MB, additional formins might cooperate with *DAAM* for proper MB development (Gombos *et al.* 2015).

We therefore tested whether *frl* also affects MB axons and, if so, whether it cooperates with *DAAM*. We induced positively marked single cell clones that lack *frl* in α/β neurons using the

MARCM technique (see *Materials and Methods* for details) and assessed axon defects in adult brains. Compared to wild-type axons that bifurcate and project to the ends of the α/β lobes (counterstained with FasII in Figure 8E), 14% of *frl*^{Ex83}, 16.7% of *frl*^{Ex88}, and 21.6% of *frl*^{Ex62} mutant neurons showed premature axon growth arrest (but no bifurcation defects; Figure 8, F and G, and not shown; quantified in Figure 8H), suggesting that Frl is required for correct growth of MB axons, but may not be required for or act redundantly with other genes for their correct guidance. We then tested for genetic interaction between *frl* and *DAAM*. Hemizygous males of the hypomorphic alleles *DAAM*^{Ex1} and *DAAM*^{Ex4} are viable and show moderate MB axon growth and guidance defects (Gombos *et al.* 2015). Importantly, the growth phenotypes of *DAAM*^{Ex1} and *DAAM*^{Ex4} were dominantly and strongly enhanced by the removal of one gene dose of *frl* by all three alleles (quantified in Figure 9, A and B for the α and β lobes, respectively). Furthermore, these phenotypes in *DAAM*^{Ex1} and *DAAM*^{Ex4} hemizygotes are also enhanced by RNAi-mediated knockdown of *frl*^{TRIPHMS00445} expressed under control of *OK107-Gal4* in the β lobe (Figure 9D) and for *DAAM*^{Ex4} also in the α lobe (Figure 9C). Distinct from *DAAM* (Gombos *et al.* 2015), but consistent with the lack of genetic interaction between

Figure 5 Frl can be activated by Cdc42 *in vivo*. (A–C) The R-cell loss of *sep > Cdc42* (A) is dominantly suppressed by removal of one gene dose of each *frl* allele (B: *frl^{Ex83}*; C: *frl^{ExK62}* and not shown; quantification in D). (D) Quantification (percentage ommatidia with wild-type R-cell complement) of *sep > Cdc42* lacking one gene dose of the alleles indicated underneath. (E–H) *sep > Cdc42* induced R-cell loss (E shows the phenotype of a different chromosomal insertion than used in A) is fully suppressed by coexpression of dominant negative *sep > DN-Frl (ΔFH2)* (F; quantified in I). Conversely, the phenotype of *sep > DN-Frl* (G; note that this stronger line also causes gaps in between ommatidia) is suppressed by coexpression of *sep > Cdc42* (F), but dominantly enhanced by *cdc42¹*, a null allele (H; quantified in I). (I) Quantification of PCP and R-cell loss phenotypes of indicated genotypes (gaps: whole ommatidia missing). *t*-test; ****P* < 0.001; ***P* < 0.02.

frl and *dsh* during PCP signaling in the eye, removal of one gene dose of *frl* did not alter the axon growth or guidance phenotype of the PCP-specific *dsh¹* allele (quantified in Figure 9, E and F). Collectively, our data suggest that *frl* and *DAAM* are both required for proper axonal development of the MB neurons. Their strong

dominant genetic interaction indicates a cooperative mode of action between these two formins in this tissue, although they are likely controlled by different signals.

We also tested whether *frl* interacted with *cdc42* in the MB. We found that hemizygous males of the hypomorphic allele

Figure 6 Frl interacts with Dsh *in vitro*, but not *in vivo*. (A) Gst pull-down assay of *in vitro* translated Frl fragments indicated on the left by purified Gst fusion proteins indicated on top. Full-length Frl and Frl-FH2-CT, but not the FH2 domain alone, bind to Gst-Dsh and its basic PDZ region, but not to Gst alone or the Dsh DEP-Cterm domain. The core PCP gene Pk is used as positive control for Dsh binding (Jenny *et al.* 2005). (Top) Autoradiographs. (Bottom-most) Coomassie stained gel of a typical pull-down showing the Gst proteins. (B) Coimmunoprecipitations of Dsh-GFP by the indicated Myc-tagged Frl fragments from HEK293 cells. Only Frl Δ GBD-DID is able to interact with Dsh-GFP. A fragment of the Wnk kinase was used as negative control (Wnk-NT2). (Top) Immunoprecipitations. (Bottom) Cell lysates. (C–E) The PCP phenotype of *dsh*¹ hemizygous males (C) is not altered by the removal of one gene dose of *frl* (D: *frl*^{Ex83}; E: *frl*^{ExK62}; quantified in F). Tangential eye sections are shown above schematics. (F) Quantification of PCP phenotypes of indicated genotypes.

*cdc42*² show growth and guidance defects in the α and β lobes (quantified in Figure 9, G and H). Importantly, all three *frl* alleles dominantly enhance the growth phenotype of *cdc42*² hemizygotes (Figure 9, G and H), strongly suggesting that Cdc42 controls Frl also in this context.

Discussion

In this report, we have analyzed the role of the diaphanous-related formin Frl during ommatidial rotation in the *Drosophila* eye and axon outgrowth during mushroom body development. RNAi-mediated knockdown or mutations in *frl* cause mild

ommatidial rotation defects. Biochemical and genetic interaction experiments are consistent with Frl being autoinhibited and activated by Cdc42 *in vivo*. Significantly, we showed that Frl and DAAM, members of two distinct formin subfamilies, cooperate in R-cell morphogenesis or maintenance and have distinct and overlapping functions in axon growth in the mushroom body, for which a subset of PCP genes is also known to be required (Shimizu *et al.* 2011; Ng 2012).

Activation of Frl

Overexpression of full-length Frl does not cause a phenotype in the eye, indicating that Frl, similar to other DRFs, likely

Figure 7 The phenotype of DN-Frl (A) is dominantly enhanced by concurrent removal of one gene dose of both N-cadherins (*CadNΔ14*; B). Above: tangential eye sections; below: schematics. See Figure S3 for quantification.

needs to be activated. Many DRFs such as mDia2, Daam, or mFrl2 are autoinhibited by binding of the C-terminal DAD domain to the GBD-DID domains (Higgs 2005; Otomo *et al.* 2005, 2010; Vaillant *et al.* 2008; Liu *et al.* 2010). Consistent with a need for activation, the GBD-DID containing N-terminal region of Frl can bind the FH2-CT and the CT domain much more efficiently than full-length Frl (Figure 2E), although for unknown reasons, Frl-GBD-DID is not sufficient for this interaction *in vitro*. In addition, constitutively active Frl and dominant negative Frl cause severe rough eye phenotypes and, when expressed in a more restricted way using *sep-Gal4*, loss of R-cells and PCP-like phenotypes, respectively (note that although a “weaker” transgenic line mostly shows rotation defects (Figure 3B), we cannot exclude that additional morphological defects also contribute to the ommatidial rotation defects caused by DN-Frl). These data are thus consistent with Frl being autoinhibited like most DRFs (Campellone and Welch 2010; Bogdan *et al.* 2013).

Drosophila Frl strongly binds to Cdc42 with a significant preference for the active vs. the inactive form (Figure 4), suggesting that Frl is activated by Cdc42. Consistently, we observed that overexpression of Cdc42 and the CA-Frl cause a similar R-cell loss and that the R-cell loss upon Cdc42 expression is dominantly suppressed by removal of one gene dose of all *frl* alleles (and by DN-Frl; Figure 5). Conversely,

we find that ectopic activation of Cdc42 suppressed the PCP defects of DN-Frl, likely by competing with DN-Frl and thus allowing a higher activation of endogenous Frl by Cdc42. Overall, our data strongly suggest that Frl is activated by Cdc42 *in vivo*, which is also supported by our finding that the phenotype of a hypomorphic *cdc42* mutation in the mushroom body is enhanced by all three *frl* alleles (Figure 9, G and H). Our observations are consistent with recent data demonstrating that Cdc42 activates human FMNL2 (Frl3) during cell migration in culture and that Cdc42 activates Fmnl3 (Frl2) to promote filopodia formation during angiogenic sprouting (Block *et al.* 2012; Kuhn *et al.* 2015; Richards *et al.* 2015; Wakayama *et al.* 2015).

Originally, mouse Frl1 was shown to bind to Rac1 but not to RhoA or Cdc42 with a slight preference for the active, GTP-bound form over the GDP-Rac1 in lysates of a P388D1 macrophage cell line (Yayoshi-Yamamoto *et al.* 2000), although no functional data supporting this interaction were presented. Despite the weak, but reproducible physical interaction between Rac1 and Frl in coimmunoprecipitation and Gst pull-down experiments, we were unable to identify genetic interactions between *rac* and *frl* in the eye. Thus, the physiological relevance of Rac binding to Frl remains an open question.

Generally, the involvement of Rho-family GTPases in PCP signaling is controversial and less well understood (reviewed in Schlessinger *et al.* 2009). Loss of *cdc42* in the mouse inner ear has recently been shown to cause misalignment of sensory stereocilia similar to loss of core PCP genes such as Vangl2 (Montcouquiol *et al.* 2003; Kirjavainen *et al.* 2015). Additionally, while Daam1 activates Rho and Rac, but not Cdc42 in *Xenopus* tissue undergoing gastrulation (Habas *et al.* 2001), DN-Cdc42 can suppress the convergence and extension phenotypes caused by overexpressed Wnt11 or Fz7, but not Dsh (Djiane *et al.* 2000; Choi and Han 2002), indicating that Cdc42 may act in a parallel pathway to classical PCP signaling downstream of Dsh. In *Drosophila*, the involvement of Rho GTPases had originally been identified in *rhoA* mutant clones (Strutt *et al.* 1997) but had mainly been characterized using genetic interaction assays with PCP GOF scenarios and overexpression of constitutively active or dominant negative GTPase variants (Strutt *et al.* 1997; Boutros *et al.* 1998; Fanto *et al.* 2000). Unfortunately, eye clones of *cdc42* null mutant cells cannot be recovered (Genova *et al.* 2000; Muñoz-Descalzo *et al.* 2007), but a hypomorphic *cdc42* allele shows mild rotation defects in the eye similar in strength to loss of *frl* ($2.9 \pm 3.8\%$), and the mild PCP phenotype of *rac1* and its paralogs *rac2* and *mtl* in triple mutant clones are enhanced in a *cdc42* hypomorphic background, suggesting that these GTPases contribute to PCP signaling in a partially redundant manner (Muñoz-Descalzo *et al.* 2007). Cdc42 also acts together with Mtl (Muñoz-Descalzo *et al.* 2007; Muñoz-Soriano *et al.* 2011), which is regulated by EGFR signaling that is known to control ommatidial rotation (Brown and Freeman 2003; Gaengel and Mlodzik 2003; Strutt and Strutt 2003). While we present evidence that Cdc42 regulates Frl during PCP signaling in the eye, we failed to detect

Figure 8 Frl cooperates with DAAM during neural development. (A–C) Compared to *frl* (Figure 1) and *DAAM^{Ex68}* single (null) mutant R-cells (A), *DAAM^{Ex68}; frl^{Ex83}* (B) and *DAAM^{Ex68}; frl^{ExK62}* (C) double mutant R-cells show a cell-autonomous small rhabdomere phenotype, suggesting that DAAM and Frl have a redundant function in R-cell morphogenesis or maintenance. Double mutant cells lack rhabdomere-associated pigment granules (examples marked with magenta arrowheads), while R-cells of all other genotypes contain pigment (examples marked with blue arrowheads). (D) Schematic of a right part of a mushroom body (MB) of an adult brain with the neuropile lobes growing from the Kenyon cells. Axons of α/β and α'/β' neurons bifurcate with one axon projecting into a dorsal (α/α') and one growing into a medial (β/β') lobe. Magenta frame depicts area shown in confocal images in E–G. (E–G) Positively marked MARCM clones labeled by GFP of wild-type (E), *frl^{Ex88}* mutant (F), and *frl^{ExK62}* mutant (G) neurons in the α/β lobes counterstained with Fasciclin II (FasII, red). Wild-type neurons grow axons that terminate (yellow arrowheads in E) at the tip of the α/β lobes (white arrowheads), while axons of *frl* mutants often arrest prematurely (yellow arrowheads in F and G). (H) Quantification of axons with growth defects (in percentage) of the indicated *frl* mutant neurons. White numbers indicate number of counted clones. Dorsal is up and medial is to the right in E–G.

genetic interactions between *sep* > *DN-Frl* and EGFR signaling pathway components such as *aos*, *cno*, *phyl*, or *sprouty*. In support of Cdc42 affecting PCP signaling independent of Dsh, we found that similar to *cdc42* (Boutros *et al.* 1998), *frl* alleles are unable to suppress *sev* > *Fz* or *sev-Dsh* (not shown). Interestingly, however, we found that *sep* > *DN-Frl* is enhanced by removal of a gene dose of both N-cadherins (Figure 7) that are known to control speed and extent of ommatidial rotation downstream of the Fz-PCP and EGFR pathways (Mirkovic and Mlodzik 2006), suggesting a complicated relationship between genes involved in rotation.

Frl and DAAM cooperate during neural development

Daam1 was originally identified in a two-hybrid screen with a fragment of Dsh containing its central PDZ domain and later

was shown to be activated by Dsh interacting with the C terminus of Daam1, thereby regulating PCP signaling during convergence and extension in *Xenopus* (Habas *et al.* 2001; Liu *et al.* 2008). An interaction between Dvl3 and Daam2 in chicken was also identified and is considered crucial for the cofactor function of Daam1 during D/V patterning of the chick neural tube by canonical Wnt signaling (Lee and Deneen 2012). Because *Drosophila* DAAM has no unique function during PCP signaling in *Drosophila*, we initially hypothesized that Frl may cooperate with DAAM in that process. Indeed, we found that, similar to xDaam1, Frl can interact with the PDZ region of Dsh *in vitro* in a manner that is dependent on the FH2-CT region (Figure 6, A and B). However, we were unable to find *in vivo* evidence supporting such an interaction. Neither did we detect an obvious

Figure 9 *frl* and *DAAM* cooperate in MB axon growth. In all panels, quantifications of defects are shown in percentage for the indicated genotypes with growth defects (shortened, thinner, or absent lobes) in blue and guidance defects (thickened or misprojecting lobes) depicted in red. Numbers indicate MBs scored per genotype (note that α and β lobes were scored in the same samples). (A and B) All three *frl* alleles dominantly enhance the axon growth defects in α lobes (A) and β lobes (B) of *DAAM^{Ex1}* and *DAAM^{Ex4}* hemizygous males. (C and D) Similarly, RNAi-mediated knock down of *frl^{TRIP}* under control of *OK107-Gal4* enhances the axon growth defects of *DAAM^{Ex4}* in α lobes (C) and *DAAM^{Ex1}* and *DAAM^{Ex4}* in β lobes (D). (E and F) The MB phenotype of *dsh¹* hemizygous males is not altered by removal of a gene dose of *frl* (E: α lobes; F: β lobes). (G and H) All three *frl* alleles dominantly enhance the axon growth defects in α lobes (G) and β lobes (H) of *cdc42²* hemizygous males. Fisher's exact test; * $P < 0.05$; ** $P < 0.005$; *** $P < 0.0001$.

redundant function of *frl* and *DAAM* in PCP signaling in the eye in double mutants (although the double mutant clones are small only). Intriguingly, however, *DAAM frl* double mutant R-cells show abnormal, small rhabdomeres in adult eye sections, a phenotype that is not seen in either single mutant, suggesting that *frl* and *DAAM* indeed can exhibit overlapping functions in neurons (Figure 8, B and C).

It has recently been shown that *DAAM* is required for axon growth and pathfinding in the mushroom body, where it genetically interacts with the core PCP genes including *fz*, *vang*, and *dsh*, as well as *Rac1* (Gombos *et al.* 2015). Similar to *DAAM* mutants, *frl* single mutants display an axon growth phenotype (Figure 9), and heterozygosity for *frl* enhances the axon growth phenotype of hypomorphic *cdc42²* mutant

males, consistent with Cdc42-activating Frl. However, distinct from *DAAM*, *frl* mutants do not affect axon guidance. Nevertheless, we again found evidence for cooperation between *DAAM* and Frl, as *frl* strongly dominantly enhanced the growth phenotype of *DAAM* hypomorphic mutants. Again, however, no genetic interaction was found between *dsh* and *frl*, indicating that, in contrast to *DAAM*, Frl is likely regulated in a core PCP-independent manner. Thus, both formins cooperate during and are required for precise formation of the MB axons. However, they probably are under distinct regulatory control and thus may integrate various signaling inputs with the regulation of the cytoskeleton. Overall, our data suggest that Frl and *DAAM*, members of distinct formin subfamilies, can act together in specific cellular processes, likely reflecting the need for precise control of the complex and context-dependent regulation of the actin cytoskeleton by various upstream signaling systems.

Acknowledgments

We are very grateful to M. Mlodzik (Mount Sinai Hospital), L. Luo (Stanford University), T. Clandinin (Stanford University), E. Verheyen (Simon Fraser University), D. Cox (Albert Einstein College of Medicine), S. Parkhurst (Fred Hutchinson Cancer Research Center), and the Bloomington and Vienna Stock Centers for fly stocks, plasmids, and advice, and Ke Li for help with the generation of the *frl^{ExK62}* allele. We also thank the Berkeley Drosophila Genome Project and Drosophila Genomics Resource Center for plasmids, C. Desplan (New York University) and the Developmental Studies Hybridoma Bank for antibodies, and P. Tomancak (Max Planck Institute of Molecular Cell Biology and Genetics) for the FlyFOS clones. We are grateful to F. Marlow, A. Santos Ledo, and H. Bülow for discussions and comments on the manuscript. This work was supported by National Institutes of Health grant (GM088202 to A.J.), the Hungarian Brain Research Program (KTIA_NAP_13-2-2014-0007 to J.M.), and the Hungarian Scientific Research Foundation (OTKA grants K82039 and 109330 to J.M.).

Literature Cited

- Abou-Kheir, W., B. Isaac, H. Yamaguchi, and D. Cox, 2008 Membrane targeting of WAVE2 is not sufficient for WAVE2-dependent actin polymerization: a role for IRSp53 in mediating the interaction between Rac and WAVE2. *J. Cell Sci.* 121: 379–390.
- Axelrod, J. D., 2001 Unipolar membrane association of Dishevelled mediates Frizzled planar cell polarity signaling. *Genes Dev.* 15: 1182–1187.
- Basler, K., and E. Hafen, 1990 Receptor tyrosine kinases mediate cell-cell interactions during *Drosophila* development. *Prog. Growth Factor Res.* 2: 15–27.
- Basler, K., B. Christen, and E. Hafen, 1991 Ligand-independent activation of the sevenless receptor tyrosine kinase changes the fate of cells in the developing *Drosophila* eye. *Cell* 64: 1069–1081.
- Block, J., D. Breitsprecher, S. Kuhn, M. Winterhoff, F. Kage *et al.*, 2012 FMNL2 drives actin-based protrusion and migration downstream of Cdc42. *Curr. Biol.* 22: 1005–1012.
- Bogdan, S., J. Schultz, and J. Grosshans, 2013 Formin' cellular structures: physiological roles of Diaphanous (Dia) in actin dynamics. *Commun. Integr. Biol.* 6: e27634.
- Borovina, A., S. Superina, D. Voskas, and B. Ciruna, 2010 Vangl2 directs the posterior tilting and asymmetric localization of motile primary cilia. *Nat. Cell Biol.* 12: 407–412.
- Boutros, M., N. Paricio, D. I. Strutt, and M. Mlodzik, 1998 Dishevelled activates JNK and discriminates between JNK pathways in planar polarity and *wingless* signaling. *Cell* 94: 109–118.
- Brand, A. H., and N. Perrimon, 1993 Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* 118: 401–415.
- Brown, K. E., and M. Freeman, 2003 Egfr signalling defines a protective function for ommatidial orientation in the *Drosophila* eye. *Development* 130: 5401–5412.
- Cammer, M., J. C. Gevrey, M. Lorenz, A. Dovas, J. Condeelis *et al.*, 2009 The mechanism of CSF-1-induced Wiskott-Aldrich syndrome protein activation in vivo: a role for phosphatidylinositol 3-kinase and Cdc42. *J. Biol. Chem.* 284: 23302–23311.
- Campellone, K. G., and M. D. Welch, 2010 A nucleator arms race: cellular control of actin assembly. *Nat. Rev. Mol. Cell Biol.* 11: 237–251.
- Choi, S. C., and J. K. Han, 2002 *Xenopus* Cdc42 regulates convergent extension movements during gastrulation through Wnt/Ca2+ signaling pathway. *Dev. Biol.* 244: 342–357.
- Cox, D., P. Chang, Q. Zhang, P. G. Reddy, G. M. Bokoch *et al.*, 1997 Requirements for both Rac1 and Cdc42 in membrane ruffling and phagocytosis in leukocytes. *J. Exp. Med.* 186: 1487–1494.
- Devenport, D., 2014 The cell biology of planar cell polarity. *J. Cell Biol.* 207: 171–179.
- Devenport, D., and E. Fuchs, 2008 Planar polarization in embryonic epidermis orchestrates global asymmetric morphogenesis of hair follicles. *Nat. Cell Biol.* 10: 1257–1268.
- Dietzl, G., D. Chen, F. Schnorrrer, K. C. Su, Y. Barinova *et al.*, 2007 A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature* 448: 151–156.
- Djiane, A., J. Riou, M. Umbhauer, J. Boucalt, and D. Shi, 2000 Role of frizzled 7 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. *Development* 127: 3091–3100.
- Domingos, P. M., S. Brown, R. Barrio, K. Ratnakumar, B. J. Frankfort *et al.*, 2004 Regulation of R7 and R8 differentiation by the spalt genes. *Dev. Biol.* 273: 121–133.
- Ejsmont, R. K., M. Sarov, S. Winkler, K. A. Lipinski, and P. Tomancak, 2009 A toolkit for high-throughput, cross-species gene engineering in *Drosophila*. *Nat. Methods* 6: 435–437.
- Fahmy, O. G., and M. Fahmy, 1959 [New mutants report.] *D. I. S.* 33: 82–94.
- Faix, J., and R. Grosse, 2006 Staying in shape with formins. *Dev. Cell* 10: 693–706.
- Fanto, M., C. A. Mayes, and M. Mlodzik, 1998 Linking cell-fate specification to planar polarity: determination of the R3/R4 photoreceptors is a prerequisite for the interpretation of the Frizzled mediated polarity signal. *Mech. Dev.* 74: 51–58.
- Fanto, M., U. Weber, D. I. Strutt, and M. Mlodzik, 2000 Nuclear signaling by Rac and Rho GTPases is required in the establishment of epithelial planar polarity in the *Drosophila* eye. *Curr. Biol.* 10: 979–988.
- FlyBase Consortium, 2003 The FlyBase database of the *Drosophila* genome projects and community literature. *Nucleic Acids Res.* 31: 172–175.
- Freeman, M., C. Klämbt, C. S. Goodman, and G. M. Rubin, 1992 The *argos* gene encodes a diffusible factor that regulates cell fate decisions in the *Drosophila* eye. *Cell* 69: 963–975.
- Gaengel, K., and M. Mlodzik, 2003 Egfr signaling regulates ommatidial rotation and cell motility in the *Drosophila* eye via

- MAPK/Pnt signaling and the Ras effector Canoe/AF6. *Development* 130: 5413–5423.
- Genova, J. L., S. Jong, J. T. Camp, and R. G. Fehon, 2000 Functional analysis of Cdc42 in actin filament assembly, epithelial morphogenesis, and cell signaling during *Drosophila* development. *Dev. Biol.* 221: 181–194.
- Gombos, R., E. Migh, O. Antal, A. Mukherjee, A. Jenny *et al.*, 2015 The Formin DAAM functions as molecular effector of the planar cell polarity pathway during axonal development in *Drosophila*. *J. Neurosci.* 35: 10154–10167.
- Goodrich, L. V., and D. Strutt, 2011 Principles of planar polarity in animal development. *Development* 138: 1877–1892.
- Habas, R., Y. Kato, and X. He, 2001 Wnt/Frizzled activation of rho regulates vertebrate gastrulation and requires a novel formin homology protein Daam1. *Cell* 107: 843–854.
- Heisenberg, M., A. Borst, S. Wagner, and D. Byers, 1985 *Drosophila* mushroom body mutants are deficient in olfactory learning. *J. Neurogenet.* 2: 1–30.
- Higgs, H. N., 2005 Formin proteins: a domain-based approach. *Trends Biochem. Sci.* 30: 342–353.
- Jenny, A., 2010 Planar cell polarity signaling in the *Drosophila* eye. *Curr. Top. Dev. Biol.* 93: 189–227.
- Jenny, A., 2011 Preparation of adult *Drosophila* eyes for thin sectioning and microscopic analysis. *J. Vis. Exp.* 54: pii: 2959. DOI: 10.3791/2959.
- Jenny, A., R. S. Darken, P. A. Wilson, and M. Mlodzik, 2003 Prickle and Strabismus form a functional complex to generate a correct axis during planar cell polarity signaling. *EMBO J.* 22: 4409–4420.
- Jenny, A., J. Reynolds-Kenneally, G. Das, M. Burnett, and M. Mlodzik, 2005 Diego and Prickle regulate Frizzled planar cell polarity signalling by competing for Dishevelled binding. *Nat. Cell Biol.* 7: 691–697.
- Kirjavainen, A., M. Laos, T. Anttonen, and U. Pirvola, 2015 The Rho GTPase Cdc42 regulates hair cell planar polarity and cellular patterning in the developing cochlea. *Biol. Open.* 4: 516–526.
- Kuhn, S., C. Erdmann, F. Kage, J. Block, L. Schwenkmezger *et al.*, 2015 The structure of FMNL2-Cdc42 yields insights into the mechanism of lamellipodia and filopodia formation. *Nat. Commun.* 6: 7088.
- Lee, H. K., and B. Deneen, 2012 Daam2 is required for dorsal patterning via modulation of canonical Wnt signaling in the developing spinal cord. *Dev. Cell* 22: 183–196.
- Lee, T., A. Lee, and L. Luo, 1999 Development of the *Drosophila* mushroom bodies: sequential generation of three distinct types of neurons from a neuroblast. *Development* 126: 4065–4076.
- Liu, R., E. V. Linardopoulou, G. E. Osborn, and S. M. Parkhurst, 2010 Formins in development: orchestrating body plan origami. *Biochim. Biophys. Acta* 1803: 207–225.
- Liu, W., A. Sato, D. Khadka, R. Bharti, H. Diaz *et al.*, 2008 Mechanism of activation of the Formin protein Daam1. *Proc. Natl. Acad. Sci. USA* 105: 210–215.
- Luo, L., Y. J. Liao, L. Y. Jan, and Y. N. Jan, 1994 Distinct morphogenetic functions of similar small GTPases: *Drosophila* Drac1 is involved in axonal outgrowth and myoblast fusion. *Genes Dev.* 8: 1787–1802.
- Magie, C. R., M. R. Meyer, M. S. Gorsuch, and S. M. Parkhurst, 1999 Mutations in the Rho1 small GTPase disrupt morphogenesis and segmentation during early *Drosophila* development. *Development* 126: 5353–5364.
- Matusek, T., A. Djiane, F. Jankovics, D. Brunner, M. Mlodzik *et al.*, 2006 The *Drosophila* formin DAAM regulates the tracheal cuticle pattern through organizing the actin cytoskeleton. *Development* 133: 957–966.
- Matusek, T., R. Gombos, A. Szecsenyi, N. Sanchez-Soriano, A. Czibula *et al.*, 2008 Formin proteins of the DAAM subfamily play a role during axon growth. *J. Neurosci.* 28: 13310–13319.
- Maung, S. M., and A. Jenny, 2011 Planar cell polarity in *Drosophila*. *Organogenesis* 7: 165–179.
- Mirkovic, I., and M. Mlodzik, 2006 Cooperative activities of *Drosophila* DE-cadherin and DN-cadherin regulate the cell motility process of ommatidial rotation. *Development* 133: 3283–3293.
- Mlodzik, M., Y. Hiromi, U. Weber, C. S. Goodman, and G. M. Rubin, 1990 The *Drosophila* seven-up gene, a member of the steroid receptor gene superfamily, controls photoreceptor cell fates. *Cell* 60: 211–224.
- Molnar, I., E. Migh, S. Szikora, T. Kalmar, A. G. Vegh *et al.*, 2014 DAAM is required for thin filament formation and Sarcomerogenesis during muscle development in *Drosophila*. *PLoS Genet.* 10: e1004166.
- Montcouquiol, M., R. A. Rachel, P. J. Lanford, N. G. Copeland, N. A. Jenkins *et al.*, 2003 Identification of Vangl2 and Scrb1 as planar polarity genes in mammals. *Nature* 423: 173–177.
- Morata, G., and P. Ripoll, 1975 *Minutes*: mutants of *Drosophila* autonomously affecting cell division rate. *Dev. Biol.* 42: 211–221.
- Muñoz-Descalzo, S., A. Gómez-Cabrero, M. Mlodzik, and N. Paricio, 2007 Analysis of the role of the Rac/Cdc42 GTPases during planar cell polarity generation in *Drosophila*. *Int. J. Dev. Biol.* 51: 379–387.
- Muñoz-Soriano, V., Y. Belacortu, F. C. Durupt, S. Muñoz-Descalzo, and N. Paricio, 2011 Mtl interacts with members of Egfr signaling and cell adhesion genes in the *Drosophila* eye. *Fly (Austin)* 5: 88–101.
- Murdoch, J. N., K. Doudney, C. Paternotte, A. J. Copp, and P. Stanier, 2001 Severe neural tube defects in the loop-tail mouse result from mutation of Lpp1, a novel gene involved in floor plate specification. *Hum. Mol. Genet.* 10: 2593–2601.
- Newsome, T. P., B. Asling, and B. J. Dickson, 2000 Analysis of *Drosophila* photoreceptor axon guidance in eye-specific mosaics. *Development* 127: 851–860.
- Ng, J., 2012 Wnt/PCP proteins regulate stereotyped axon branch extension in *Drosophila*. *Development* 139: 165–177.
- Ng, J., T. Nardine, M. Harms, J. Tzu, A. Goldstein *et al.*, 2002 Rac GTPases control axon growth, guidance and branching. *Nature* 416: 442–447.
- Otomo, T., C. Otomo, D. R. Tomchick, M. Machius, and M. K. Rosen, 2005 Structural basis of Rho GTPase-mediated activation of the formin mDia1. *Mol. Cell* 18: 273–281.
- Otomo, T., D. R. Tomchick, C. Otomo, M. Machius, and M. K. Rosen, 2010 Crystal structure of the Formin mDia1 in auto-inhibited conformation. *PLoS One* 5: e12896.
- Prakash, S., J. C. Caldwell, D. F. Eberl, and T. R. Clandinin, 2005 *Drosophila* N-cadherin mediates an attractive interaction between photoreceptor axons and their targets. *Nat. Neurosci.* 8: 443–450.
- Richards, M., C. Hetheridge, and H. Mellor, 2015 The formin FMNL3 controls early apical specification in endothelial cells by regulating the polarized trafficking of podocalyxin. *Curr. Biol.* 25: 2325–2331.
- Roignant, J. Y., and J. E. Treisman, 2009 Pattern formation in the *Drosophila* eye disc. *Int. J. Dev. Biol.* 53: 795–804.
- Ross, A. J., H. May-Simera, E. R. Eichers, M. Kai, J. Hill *et al.*, 2005 Disruption of Bardet-Biedl syndrome ciliary proteins perturbs planar cell polarity in vertebrates. *Nat. Genet.* 37: 1135–1140.
- Schlessinger, K., A. Hall, and N. Tolwinski, 2009 Wnt signaling pathways meet Rho GTPases. *Genes Dev.* 23: 265–277.
- Shimizu, K., M. Sato, and T. Tabata, 2011 The Wnt5/planar cell polarity pathway regulates axonal development of the *Drosophila* mushroom body neuron. *J. Neurosci.* 31: 4944–4954.
- Singh, A., J. Lim, and K. Choi, 2005 Dorsal-ventral boundary for organizing growth and planar polarity in the *Drosophila* eye, pp. 59–90 in *Planar Cell Polarization During Development*, edited by M. Mlodzik. Elsevier, San Diego.

- Song, H., J. Hu, W. Chen, G. Elliott, P. Andre *et al.*, 2010 Planar cell polarity breaks bilateral symmetry by controlling ciliary positioning. *Nature* 466: 378–382.
- Strutt, H., and D. Strutt, 2002 Nonautonomous planar polarity patterning in *Drosophila*: dishevelled-independent functions of frizzled. *Dev. Cell* 3: 851–863.
- Strutt, H., and D. Strutt, 2003 EGF signaling and ommatidial rotation in the *Drosophila* eye. *Curr. Biol.* 13: 1451–1457.
- Strutt, H., and D. Strutt, 2009 Asymmetric localisation of planar polarity proteins: mechanisms and consequences. *Semin. Cell Dev. Biol.* 20: 957–963.
- Strutt, D. I., U. Weber, and M. Mlodzik, 1997 The role of RhoA in tissue polarity and Frizzled signalling. *Nature* 387: 292–295.
- Tissir, F., Y. Qu, M. Montcouquiol, L. Zhou, K. Komatsu *et al.*, 2010 Lack of cadherins Celsr2 and Celsr3 impairs ependymal ciliogenesis, leading to fatal hydrocephalus. *Nat. Neurosci.* 13: 700–707.
- Treisman, J. E., and U. Heberlein, 1998 Eye development in *Drosophila*: formation of the eye field and control of differentiation. *Curr. Top. Dev. Biol.* 39: 119–158.
- Vaillant, D. C., S. J. Copeland, C. Davis, S. F. Thurston, N. Abdennur *et al.*, 2008 Interaction of the N- and C-terminal autoregulatory domains of FRL2 does not inhibit FRL2 activity. *J. Biol. Chem.* 283: 33750–33762.
- Venken, K. J., and H. J. Bellen, 2005 Emerging technologies for gene manipulation in *Drosophila melanogaster*. *Nat. Rev. Genet.* 6: 167–178.
- Vladar, E. K., D. Antic, and J. D. Axelrod, 2009 Planar cell polarity signaling: the developing cell's compass. *Cold Spring Harb. Perspect. Biol.* 1: a002964.
- Vladar, E. K., R. D. Bayly, A. M. Sangoram, M. P. Scott, and J. D. Axelrod, 2012 Microtubules enable the planar cell polarity of airway cilia. *Curr. Biol.* 22: 2203–2212.
- Wakayama, Y., S. Fukuhara, K. Ando, M. Matsuda, and N. Mochizuki, 2015 Cdc42 mediates Bmp-induced sprouting angiogenesis through Fmnl3-driven assembly of endothelial filopodia in zebrafish. *Dev. Cell* 32: 109–122.
- Weber, U., N. Paricio, and M. Mlodzik, 2000 mediates Frizzled induced R3/R4 cell fate distinction and planar polarity determination in the *Drosophila* eye. *Development* 127: 3619–3629.
- Winter, C. G., B. Wang, A. Ballew, A. Royou, R. Karess *et al.*, 2001 *Drosophila* Rho-associated kinase (Drok) links Frizzled-mediated planar cell polarity signaling to the actin cytoskeleton. *Cell* 105: 81–91.
- Wolff, T., and D. F. Ready, 1993 Pattern formation in the *Drosophila* retina, pp. 1277–1326 in *The Development of Drosophila melanogaster*, edited by M. B. A. Martinez-Arias. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Yanfeng, W. A., H. Berhane, M. Mola, J. Singh, A. Jenny *et al.*, 2011 Functional dissection of phosphorylation of Dishevelled in *Drosophila*. *Dev. Biol.* 360: 132–142.
- Yayoshi-Yamamoto, S., I. Taniuchi, and T. Watanabe, 2000 FRL, a novel formin-related protein, binds to Rac and regulates cell motility and survival of macrophages. *Mol. Cell. Biol.* 20: 6872–6881.
- Zou, Y., 2012 Does planar cell polarity signaling steer growth cones? *Curr. Top. Dev. Biol.* 101: 141–160.

Communicating editor: H. J. Bellen

GENETICS

Supporting Information

www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.181438/-/DC1

Unique and Overlapping Functions of Formins Frl and DAAM During Ommatidial Rotation and Neuronal Development in *Drosophila*

Gretchen Dollar, Rita Gombos, Austen A. Barnett, David Sanchez Hernandez,
Saw M. T. Maung, Jozsef Mihály, and Andreas Jenny

sev>,ey>Dcr2, frl^{HMS000445}

B

Fosmid 027901 (add back)

Rescue Fosmid 023275

Figure S2. *frl* does not genetically interact with *sev-Rac^{V12}*. Quantification of PCP phenotypes of *sev-Rac^{V12}* lacking one gene dose of or overexpressing the indicated gene. The stronger defects due to co-expression of *sevRac^{V12}* with dominant negative *sep>Fr Δ FH2* is likely an additive effect. T-Test; ***p<0.001; **p<0.02; *p<0.05.

Figure S3. Dominant negative *frl* is dominantly enhanced by simultaneous removal of a gene dose of both N-Cadherins. Quantification of PCP phenotypes of *sep>DN-Frl* ($\Delta FH2$) lacking one gene dose of the indicated alleles. Note that (A) and (B) were separate experiments and thus each have their own baseline and that *CadN^{Δ14}* may also show a trend to enhance R-cell loss (although this is statistically not significant). The deficiency *BSC148* uncovers a larger genomic region including both N-Cadherin genes. T-Test; * $p < 0.05$.