The Double-Strand Break Landscape of Meiotic Chromosomes Is Shaped by the Paf1 Transcription Elongation Complex in *Saccharomyces cerevisiae*

Santosh K. Gothwal,* Neem J. Patel,[†] Meaghan M. Colletti,[†] Hiroyuki Sasanuma,*¹ Miki Shinohara,*

Andreas Hochwagen,[†] and Akira Shinohara*²

*Institute for Protein Research, Graduate School of Science, Osaka University, Suita, Osaka 565-0871, Japan, and [†]Department of Biology, New York University, New York, New York 10003-6688

ABSTRACT Histone modification is a critical determinant of the frequency and location of meiotic double-strand breaks (DSBs), and thus recombination. *Set1*-dependent histone H3K4 methylation and *Dot1*-dependent H3K79 methylation play important roles in this process in budding yeast. Given that the RNA polymerase II associated factor 1 complex, Paf1C, promotes both types of methylation, we addressed the role of the Paf1C component, *Rtf1*, in the regulation of meiotic DSB formation. Similar to a *set1* mutation, disruption of *RTF1* decreased the occurrence of DSBs in the genome. However, the *rtf1 set1* double mutant exhibited a larger reduction in the levels of DSBs than either of the single mutants, indicating independent contributions of *Rtf1* and *Set1* to DSB formation. Importantly, the distribution of DSBs along chromosomes in the *rtf1* mutant changed in a manner that was different from the distributions observed in both *set1* and *set1 dot1* mutants, including enhanced DSB formation at some DSB-cold regions that are occupied by nucleosomes in wild-type cells. These observations suggest that *Rtf1*, and by extension the Paf1C, modulate the genomic DSB landscape independently of H3K4 methylation.

KEYWORDS meiosis; recombination; H3K4 methylation; PAF; Rtf1; double-strand breaks

DNA double-strand breaks (DSBs) are a kind of DNA damage that is deleterious to cells unless it is repaired. Unrepaired DSBs lead to genome instability by forming broken chromosomes. DSBs are repaired by either homologous recombination or nonhomologous end joining (Krogh and Symington 2004; Symington and Gautier 2011). During vegetative growth, cells have to repair DSBs, which are accidentally created by exogenous impacts, such as ionizing radiation, as well as by internal sources, *e.g.*, due to oxidation. Meiotic cells are equipped with a program to introduce DSBs along the genome for the induction of homologous recombination. Recombination generates a crossover, a reciprocal exchange of parental DNAs, which is essential for the

correct segregation of homologous chromosomes during meiosis I (Petronczki *et al.* 2003; Lichten and de Massy 2011). Diversity in the gamete genome is also produced by meiotic recombination.

Programmed formation of DSBs on meiotic chromosomes is catalyzed by the topoisomerase II-like protein *Spo11* and its binding partner *Ski8* (Keeney *et al.* 1997; Lam and Keeney 2015). The activity of *Spo11* is also regulated by essential accessory proteins or protein complexes: the *Mre11–Rad50–Xrs2* (MRX) complex, the *Rec114–Mer2–Mei4* (RMM) complex, *Rec102*, and *Rec104* (de Massy 2013). In addition, DSB formation during meiosis is controlled by the chromosomal structure. Three meiosis-specific chromosomal proteins, *Hop1*, *Red1*, and *Mek1/Mre4*, are necessary for efficient DSB formation (Xu *et al.* 1997). Furthermore, a meiosis-specific cohesin complex containing *Rec8* kleisin regulates the distribution of DSBs in the genome (Klein *et al.* 1999; Kugou *et al.* 2009; Sun *et al.* 2015). Interestingly, the majority of proteins required for DSB formation, including components of the RMM complex, are localized on chromosome axes rather than chromatin loops. In meiotic nuclei, chromosome axes with multiple chromatin loops are a basic

Copyright © 2016 by the Genetics Society of America

doi: 10.1534/genetics.115.177287

Manuscript received April 12, 2015; accepted for publication November 19, 2015; published Early Online December 2, 2015.

Supporting information is available online at www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.177287/-DC1.

¹Present address: Department of Radiation Genetics, School of Medicine, Kyoto University, Yoshida Konoe, Sakyo ku, Kyoto, 606-8501, Japan.

²Corresponding author: Institute for Protein Research, Osaka University, 3-2

Yamadaoka, Suita, Osaka 565-0871, Japan. E-mail: ashino@protein.osaka-u.ac.jp

structural unit of the meiotic chromosome and are involved in various meiotic chromosomal events, including the regulation of DSB formation (Blat *et al.* 2002; Panizza *et al.* 2011).

Meiotic DSBs are nonuniformly distributed along the genome (Borde and de Massy 2013; de Massy 2013; Lam and Keeney 2015). In some chromosomal regions, known as recombination hotspots, DSBs are more frequent than in other parts of the genome. Moreover, the genome contains loci, recombination cold spots, where DSB frequency is lower than in other regions. In the budding yeast, most of the DSBs are introduced in nucleosome-free sites associated with 5'-regulatory regions, which contain the transcription start site of genes (Pan *et al.* 2011). This observation suggests that an open chromatin structure is essential for DSB formation. In the context of chromatin structure, histone modifications, which modulate the structure and dynamics of chromatin, also play a critical role in Spo11-dependent DSB formation during meiosis. Among these modifications, histone H3K4 trimethylation (H3K4me3) is a key determinant of DSB formation in budding yeast (Sollier *et al.* 2004; Borde *et al.* 2009). H3K4me3 is introduced by a protein complex referred to as complex associated with *Set1* (COMPASS) (Shilatifard 2012). Among the eight subunits of COMPASS, *Set1* is the catalytic subunit responsible for mono-, di-, and trimethylation of H3K4 (Jaehning 2010; Shilatifard 2012). Deletion of the *SET1* gene, as well as point mutations in H3K4, reduce the frequency of DSBs and change their distribution in a region-specific manner (Sollier *et al.* 2004; Borde *et al.* 2009).

The mechanism of H3K4me3 involvement in the chromatin-based regulatory circuitry for the generation of DSBs is described as the “axis-tethering” model, which was originally proposed by Kleckner and colleagues (Blat *et al.* 2002; Panizza *et al.* 2011; Acquaviva *et al.* 2013; Sommermeyer *et al.* 2013). The H3K4 trimethyl mark is recognized by the PHD finger domain of *Spp1*, a component of COMPASS. Importantly, *Spp1* is localized to chromosome axes while DSB regions are located on chromatin loops (Acquaviva *et al.* 2013; Sommermeyer *et al.* 2013). Thus, a hotspot region is recruited to chromosome axes through the interaction of H3K4me3 on the loop with *Spp1* on the axis, which, in turn, is a region enriched with the RMM complex. Indeed, the RMM component *Mer2* binds to *Spp1* (Acquaviva *et al.* 2013; Sommermeyer *et al.* 2013). In this model, COMPASS plays a dual role in the regulation of DSB formation: methylation of H3K4 by *Set1* and recruitment of DSB sites to chromatin axes by *Spp1*.

Spo11 is an evolutionarily conserved protein in most organisms with sexual reproduction (Keeney 2001). The regulatory pathway for recombination hotspots is also highly conserved. H3K4 methylation is associated with recombination hotspot activities in several higher eukaryotes, including human and mouse (Smagulova *et al.* 2011; Pratto *et al.* 2014). In these two species, H3K4 methylation at hotspots is catalyzed by the meiosis-specific histone methyltransferase PR domain-containing protein 9 (PRDM9) (Baudat *et al.* 2010; Myers *et al.* 2010; Parvanov *et al.* 2010), which specifically

recognizes a particular DNA sequence through its multiple zinc finger arrays. The types and numbers of zinc fingers of PRDM9 determine the diversity of recombination hotspots in these organisms.

In budding yeast, *Set1*-catalyzed H3K4 methylation is promoted by another histone modification, monoubiquitination of histone H2BK123, through *trans*-histone crosstalk (Shilatifard 2012). H2BK123 ubiquitination is mediated by the coordinated action of the E3 ligase *Bre1* and the E2-conjugating enzyme *Rad6*, as well as by the RNA polymerase II associated factor 1 complex (Paf1C). Paf1C, containing the subunits *Rtf1*, *Cdc73*, *Paf1*, *Ctr9*, and *Leo1*, is known to regulate transcription elongation by RNA polymerase II through interaction with *Set1*/COMPASS as well as with proteins involved in the termination and processing of messenger RNAs (mRNAs) (Jaehning 2010). Deletions of Paf1C components differentially affect H3K4 methylation levels. The *rtf1* deletion abolishes H3K4me and severely decreases H3K79me, while the *leo1* mutation does not affect either H3K4me or H3K79me (Krogan *et al.* 2003; Ng *et al.* 2003). Previously, we showed that *Rad6* and *Bre1* are critically important for efficient DSB formation and entry of budding yeast into meiosis I (Yamashita *et al.* 2004). However, the role of Paf1C during meiosis remains largely unknown. Interestingly, human Paf1C contains *SKI8*, whose yeast ortholog is essential for meiotic DSB formation, in addition to the five conserved components of Paf1C: RTF1, CDC73, PAF1, CTR9, and LEO1 (Kim *et al.* 2010).

In this study, we found that the Paf1C component *Rtf1*, which is essential for methylation of both H3K4 and H3K79, is also required for efficient DSB formation during meiosis. However, the distribution of meiotic DSBs along chromosomes in the *rtf1* mutant was different from that in the *set1* and *set1 dot1* mutants. In the *rtf1 set1* double mutant, the pattern of DSBs was similar to the one in the *rtf1* mutant and DSB frequency was less than that in either *set1* or *rtf1* single mutants. Location analysis of lost and gained DSB hotspots indicates a role of *Rtf1* in promoting DSBs in divergent promoter regions and preventing DSBs in regions that are occupied by nucleosomes in wild-type cells. These observations suggest that *Rtf1* and, probably, the Paf1C, modulate the generation of DSBs even in the absence of *Set1* and concomitant H3K4 methylation.

Materials and Methods

Yeast strains and strain construction

All strains used in this study were derived from the *Saccharomyces cerevisiae* SK1 diploid strains NKY1551 (*MAT α /MATa*, *ho::LYS2*⁺, *lys2*⁻, *ura3*⁻, *leu2::hisG*⁻, *his4X-LEU2-URA3/his4B-LEU2*, *arg4-nsp/arg4-bgl*) or MSY832/833 (*MAT α /MATa*, *ho::LYS2*⁺, *lys2*⁻, *ura3*⁻, *leu2::hisG*⁻, *trp1::hisG*⁻). Genotypes of each strain used in this study are listed in Supporting Information, Table S1. Deletion alleles of *CDC73*, *CTR9*, *LEO1*, *PAF1*, and *RTF1* were constructed using PCR-mediated gene deletion (Wach *et al.* 1994). The *set1* and *dot1*

deletion mutants were described previously (Bani Ismail *et al.* 2014). The primers used for strain construction are listed in Table S2.

Cytological analysis and antibodies

Chromosome spreads were prepared as described (Shinohara *et al.* 2000). Immunostaining analysis of Rad51 or Mre11 on meiotic chromosome spreads was conducted as described previously (Shinohara *et al.* 2008). Briefly, after staining, chromosome spreads were observed using an epifluorescence microscope (BL51; Olympus) with a $\times 100$ objective (N.A. 1.4). Images were captured by a CCD camera (Cool Snap; Photometrics) at room temperature and then processed using iVision (Silicon Software). Pseudo-coloring was performed using iVision and Photoshop (Adobe) software. At each time point, ~ 100 spreads were image captured and analyzed for counting foci. Primary antibodies directed against Rad51 (guinea pig, 1:500 dilution) and Mre11 (rabbit, 1:500 dilution) were used. Secondary antibodies (Alexa Fluor-488 or -594, GE Healthcare) directed against primary antibodies from the different species were used at a 1:2000 dilution.

Western blotting

For Western blotting, cell precipitates were washed twice with 20% (w/v) trichloroacetic acid (TCA) and then disrupted with glass beads (0.5 mm in diameter) using a bead beater (Yasui Kikai). Precipitated proteins were recovered by centrifugation and then suspended in the sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer. After adjusting pH to 8.8, samples were heated to 95° for 2 min. Protein samples were subjected to SDS-PAGE and transferred to PVDF membranes (Immobilon P, Millipore). Antibodies against Cdc5 (sc-33625, SantaCruz; 1:1000), Clb1 (sc-50440, SantaCruz; 1:1000), Zip1, Red1, and the α -subunit of yeast tubulin (YOL1/34, Serotec; 1:1000) were used. Anti-Red1 and anti-Zip1 antisera were described previously (Zhu *et al.* 2010). Antibodies against histone H3K4-me3 (ab8580; 1:1000), -me2 (07-030; 1:1000), -me1 (ab8895; 1:1000), and H3K79-me3 (ab2621; 1:1000) were from Abcam or Upstate Biotechnology.

Pulsed field gel electrophoresis

For pulsed field gel electrophoresis (PFGE), chromosomal DNA was prepared in agarose plugs as described (Farmer *et al.* 2011; Bani Ismail *et al.* 2014) and run at 14° in a clamped homogeneous electric field (CHEF) DR-III apparatus (Bio-Rad) using the field 6V/cm at a 120° angle. Switching times followed a ramp from 15.1 to 25.1 sec. Durations of electrophoresis were 41 hr for chromosome III and 44 hr for chromosomes V and XII.

Southern blotting

Southern blotting was performed as described previously (Storlazzi *et al.* 1995; Shinohara *et al.* 2003). DNA molecules were transferred to a nylon membrane (Hybond N, GE Healthcare). Restriction enzymes digesting DNA used for

Southern blotting were as follows: the *HIS4-LEU2* locus, PstI; *YCR048W* locus, BglII; *GAT1* locus, PstI; *PES4* locus, StuI; and *ELO2* locus, PstI. “Probe 291” was used for Southern blotting at the *HIS4-LEU2* locus (Storlazzi *et al.* 1995). Locations of probes for detecting DSBs at other loci are as follows: *YCR048W* (215,426–216,686; yeast genome coordinate of chromosome III), *GAT1* (97,171–97,980 on chromosome VI), *PES4* (194,848–196,286 on chromosome VI) and *ELO2* (186,502–187,165 on chromosome III). For DSBs located along chromosomes III, V, and XII, *CHA1*, *RMD6*, and *AYT1* were used as probes, respectively. Image Gauge software (Fujifilm) was used to quantify bands. The background of obtained images was subtracted using the software, and the lane profiles were exported to and analyzed by Excel (Microsoft).

Single-stranded DNA mapping

The genome-wide analysis of DSB positions using single-stranded DNA (ssDNA) enrichments in *dmc1* mutants was performed as described previously (Blitzblau *et al.* 2007; Blitzblau and Hochwagen 2011). Total genomic DNA for comparative hybridization was collected from a synchronous meiotic culture at 0 and 5 hr after induction of meiosis. ssDNA was enriched on benzoylated naphthoylated DEAE (BND) cellulose. Approximately 1.5 μ g of ssDNA from the two time points were differentially labeled with Cy3-dUTP and Cy5-dUTP (GE Healthcare) and cohybridized to a customized 44K tiled microarray (Agilent). Each experiment was performed in duplicate and with a dye swap between experiments.

Microarray data analysis

Probes were mapped to the annotated S288C_R64-1 reference genome (*Saccharomyces* Genome Database). Cy3 and Cy5 levels were calculated using Agilent Feature Extractor CGH software. Background normalization, log2 ratios, and scale normalizations across each set of duplicated experiments were calculated with the sma (Statistics for Microarray Analysis) package in R. Hotspot signals were defined as more than three features within 1 kb of each other that all had *P*-values < 0.15 (using the pnorm function in R) for both sets of experiments. Hotspots were then defined as the merged region from start to end of all significant data points. Datasets were intersected to obtain hotspots shared and unique between *dmc1* and *rtf1 dmc1*. For shared hotspots, the midpoint between the respective summits was calculated and used for further analysis. To assay hotspot enrichment over background relative to gene bodies, the expected number of peaks was calculated assuming a random distribution of hotspots and compared to the observed peak numbers. To determine nucleosome occupancy of the three hotspot classes (*i.e.*, peaks maintained, lost, and gained in *rtf1 dmc1* mutants), published high-resolution *Spo11* oligonucleotide data of wild-type cells (Pan *et al.* 2011; GEO accession nos. GSE26449 and GSE26452) were used to identify the expected DSB summit underlying each ssDNA peak. These data were then used to calculate average nucleosome occupancy as a function of distance from the hotspot

Figure 1 Paf1C components are necessary for meiosis. (A) Detection of several histone methylation marks during meiosis. Levels of histone H3K4-me1, -me2, and -me3, H3K79-me3, and α -tubulin as a loading control were verified by Western blotting. At each time point, cells were fixed with TCA and cell lysates were subjected to experimental analysis. Representative images are shown. (B) Spore viability of various strains was measured after dissecting tetrads. Spores were incubated at 30° for 3 days. Each bar indicates the percentage of spore viability and total number of dissected tetrads (in parentheses). Wild type, NKY1303/1543; *rtf1* mutant, SGY3/4; *leo1* mutant, SGY7/8; *cdc73* mutant, SGY5/6; *set1* mutant, SGY45/46; *set1 rtf1* mutant, SGY47/48; *set1 cdc73* mutant, SGY51/52 were used. (C) First meiotic cell division was analyzed by DAPI staining in wild type (blue circles; NKY1303/1543), *rtf1* (red circles; SGY3/4), *set1* (green circles; SGY45/46), and *rtf1 set1* (purple circles; SGY47/48) cells. At least 100 cells were counted for each time point. Each plotted value represents the mean values $\pm$ SD from three independent time courses. (D) Expression of various meiotic proteins was verified by Western blotting. Representative images are shown. Phosphorylated species of Zip1, Red1, Cdc5, and Clb1 are indicated by arrows. Wild type, NKY1303/1543; *rtf1*, SGY3/4; *set1* mutant, SGY45/46; *set1 rtf1* mutant, SGY47/48 were used.

summits. Relative replication timing of hotspot classes was determined using published replication timing data of wild-type cells (Blitzblau *et al.* 2012; GEO accession no. GSE35667).

Data availability

The datasets are available at the Gene Expression Omnibus (GEO) under accession no. GSE72827.

Results

Yeast Paf1C components are necessary for meiosis

To examine a possible role of Paf1C during meiosis, we constructed deletion mutants for five genes, *PAF1*, *CDC73*, *RTF1*, *LEO1*, and *CTR9* encoding various components of the complex in the background of the SK1 strain, which enters meiosis in an efficient and synchronous manner. As has been

previously demonstrated (Krogan *et al.* 2002; Mueller and Jaehning 2002; Squazzo *et al.* 2002), all five deletion mutants were viable, but showed different mitotic growth rates (data not shown). Among them, the *paf1* and *ctr9* deletion mutants grew poorly in the vegetative phase and entered meiosis inefficiently and asynchronously (H. Sasanuma, unpublished results), consistent with previous reports (Foreman and Davis 1996; Shi *et al.* 1996; Koch *et al.* 1999). Therefore, we focused our further studies on the three remaining deletion mutants, *rtf1*, *cdc73*, and *leo1*. First, we checked levels of the two types of histone H3 methylation promoted by Paf1C, H3K4me (me1, me2, and me3) and H3K79me (me3), in cell lysates by Western blotting (Figure 1A). As a control, we also included a *set1* deletion mutant, which is defective in H3K4 methylation. In wild-type cell lysates, levels of both H3K4me3 and H3K79me3 were almost constant during meiosis with similar levels observed during vegetative growth, which corresponds to 0-hr time point of the meiotic time course. On the other hand, levels of H3K4me1 and me2 increased slightly 2 hr after induction of meiosis in wild-type cells. The *set1* deletion completely abolished H3K4 methylation, but did not affect levels of H3K79me3. As reported previously (Jaehning 2010), the *rtf1* mutation abolished detectable levels of the H3K4me3/me2/me1 and strongly reduced H3K79me3 compared to the levels detected in wild-type cells. Defective H3K4 methylation in the *rtf1* mutant might be due to reduced Set1/COMPASS binding to the promoter (Krogan *et al.* 2003; Ng *et al.* 2003). The *cdc73* mutation decreased H3K4me3 to about two-thirds of the wild-type level, while intensities of H3K4me1, H3K4me2, and H3K79me3 were reduced by half (Figure S1). The deletion of the *LEO1* gene did not affect either H3K4 or H3K79 methylation (Figure S1).

We checked spore viability of the above three mutants (Figure 1B). The *rtf1* and *cdc73* mutants had significantly reduced spore viability levels of 83.4 and 87.9%, respectively, relative to the wild-type viability level of 97.8% (chi-square test, P -value = 7.6×10^{-13} and 2.6×10^{-8}). At the same time, the *leo1* mutant had spore viability of 95.0%, which was nearly equal to that in the wild type. The distribution of viable spores per tetrad in the *rtf1* and *cdc73* mutants did not show any bias (Figure 1B). Since the dependence of DSB formation on the activity of Set1 is critical for meiosis (Sollier *et al.* 2004), we expected to observe decreased spore viability (89.5%) in the *set1* mutant in accordance with an earlier report (Bani Ismail *et al.* 2014). To ask whether the *paf1C* mutations are epistatic to the *set1* mutation with respect to their effects on spore viability, we constructed several double mutants that combined the *set1* mutation with mutations in the genes encoding Paf1C components. The *set1 cdc73* double mutant exhibited 90.2% spore viability, which was similar to levels seen in the *cdc73* single and the *set1* single mutant. However, the *rtf1 set1* double mutant had a reduced spore viability of 77.5%, which was less than the levels detected in either *set1* or *rtf1* single mutants (chi-square test, P -values = 4.5×10^{-5} for *set1* and 0.031 for *rtf1*). This observation

suggests that Set1 and/or Rtf1 also affect meiosis independently of H3K4 methylation.

To avoid complicated interpretations due to differential contributions of each Paf1C component to histone modification, we focused our further analysis on the *rtf1* mutation, which abolished detectable levels of H3K4me and strongly reduced H3K79me relative to the levels seen in the wild-type yeast. The entry into meiosis I (MI) division was analyzed by 4',6-diamidino-2-phenylindole (DAPI) staining. Compared to the time of MI onset in wild-type cells, *rtf1* mutant cells entered MI with an ~2.5-hr delay. The delay in the *rtf1* mutant was ~2 hr shorter than the delay in the *set1* mutant (Figure 1C) (Sollier *et al.* 2004; Bani Ismail *et al.* 2014). Importantly, the *rtf1 set1* double mutant showed similar kinetics of entry into MI as the *rtf1* single mutant, suggesting that Rtf1 works upstream of Set1. To examine the progression of meiotic prophase in a different way, we analyzed the timing of expression of meiosis-specific chromosome proteins Red1 and Zip1 as early prophase markers and Cdc5 Polo-like kinase and Clb1 cyclin as markers of pachytene exit (Figure 1D). In the *rtf1* mutant, the emergence of Red1 and Zip1 proteins was similar to that in wild-type cells. However, the timing of their disappearance was clearly decelerated compared to that observed in wild-type cells. An even slower timing of Red1 and Zip1 disappearance was noted in the *set1* mutant. As for the expression of Clb1 and Cdc5 proteins (Chu and Herskowitz 1998), their appearance in the *rtf1* mutant peaked at ~8 hr, which is roughly a 2-hr delay in comparison to the time of their emergence in wild-type cells. The *set1* mutant showed a similar delay in the initial expression of the two pachytene-exit markers, but it also exhibited a retarded disappearance of these proteins in relation to their dynamics in the *rtf1* mutant. Total expression levels of the four marker proteins in the *rtf1 set1* double mutant were similar to those in the *rtf1* mutant. These results showed that, like Set1, Rtf1 is necessary for timely expression of proteins at pachytene exit and support the hypothesis that Rtf1 functions upstream of Set1 during meiosis.

The Paf1C component Rtf1 plays a role in DSB formation

Given the established role of Paf1C-dependent histone H3K4 and H3K79 methylation marks in the formation of meiotic DSBs (Sollier *et al.* 2004; Borde *et al.* 2009; Bani Ismail *et al.* 2014), we evaluated the effect of the *rtf1* mutation on the formation of meiotic DSBs. First, we studied DSB repair at the recombination hotspot *HIS4-LEU2* (Figure 2A). In the wild type, DSBs appear at 3 hr, peak at 4 hr, and then gradually disappear (Figure 2, B and C). The *rtf1* mutant showed a delay of ~1 hr in DSB appearance and a peak at 5 hr with reduced steady-state levels of DSBs to 18% of the levels seen in the wild type (at 4 hr). This finding shows that Rtf1 regulates the efficiency of DSB formation.

We next analyzed Rad51 foci on meiotic chromosomes (Figure 2D). Immunostaining of the Rad51 protein, which binds ssDNA of DSB ends for strand exchange in meiotic recombination (Shinohara *et al.* 1992), shows punctate

Figure 2 Rtf1 promotes DSB formation. (A) Schematic representation of the *HIS4-LEU2* locus. Sizes of fragments for DSB and recombinant analysis are shown with lines below. (B) DSB formation and repair at the *HIS4-LEU2* locus in the wild type and the *rtf1* mutant strains were analyzed by Southern blotting. The experiments were independently performed in triplicate and representative blots are shown. Genomic DNA was digested with *PstI*. (C) The bands of DSB I were quantified. The symbols represent the wild type (blue circles; NKY1303/1543) and *rtf1* mutant (red circles; SGY3/4). Each plotted value represents the mean values $\pm$ SD from three independent time courses. (D) Immunostaining analysis of Rad51 (green) for the wild type (NKY1303/1543) and *rtf1* (SGY3/4) mutant strains was carried out. Representative images are shown: wild type, 4 hr; *rtf1*, 4 and 8 hr. Bar, 2 μ m. (E) Kinetics of Rad51-focus positive cells in various strains. A focus-positive cell was defined as a cell with more than five foci. At least 100 nuclei were counted at each time point. The symbols represent the wild type (blue circles; NKY1303/1543) and *rtf1* mutant (red circles; SGY3/4) strains were used. (F) Kinetics of the number of Rad51 foci was analyzed in different strains. The symbols represent the wild type (blue circles; NKY1303/1543) and *rtf1* mutant (red circles; SGY3/4) strains. The number of Rad51 foci in Rad51-positive nuclei (with more than five foci) were counted at each time point. The average number of foci per positive nucleus ($\pm$ SEM from three independent experiments) is shown on the top.

staining as foci on chromosome spreads (Figure 2D) (Bishop 1994). The number of Rad51 foci is a good estimate of DSB numbers in a single nucleus. The *rtf1* mutant exhibited a delayed assembly and disassembly of Rad51 foci. The number of Rad51-focus positive cells peaked at 6 hr, i.e., with an $\sim$ 2-hr delay compared to the peak time in the wild type (Figure 2E). The disappearance of Rad51-focus positive cells was also delayed by $\sim$ 2 hr in the mutant. At 5 hr, the average number of Rad51 foci per nucleus in the *rtf1* mutant was 21.6 ± 4.1 (mean $\pm$ SE, $n = 3$; for each repeat

>20 spreads were counted), which was fewer than the number seen in the wild type at 4 hr (36.2 ± 5.8 , $n = 3$; Figure 2F). Thus, the *rtf1* mutation significantly decreased the steady number of Rad51 foci. These data suggest that the Paf1C component Rtf1 is necessary for the timely appearance of the correct number of Rad51 foci. The result that Rtf1 regulates DSB formation is consistent with the notion that Rtf1 is critical for H3K4 methylation. However, as will be described below, we found differential roles of Rtf1 and Set1 in DSB formation by a more detailed

comparative analysis of the *rtf1* single mutant and the *rtf1 set1* double mutant.

***Rtf1* and *Set1* play independent roles in regulating meiotic DSBs**

To obtain a better understanding of the effect of the *paf1C* mutation *rtf1* on DSB formation and to determine its relationship to the effects of a *set1* mutation, we employed the following three approaches to analyze DSBs: mapping of DSBs along the chromosomes using CHEF gel electrophoresis (on the background of *dmc1* and *rad50S* mutations, which block DSB turnover), analysis of DSBs at single chromosomal loci by conventional Southern blotting, and *Mre11* focus counting on the *rad50S* background. The combination of these methods revealed a unique role of *Rtf1* in meiotic DSB formation that is not shared with *Set1* and does not involve H3K4 methylation.

Mapping of DSBs on chromosomes on the *dmc1* background: We mapped DSBs along single chromosomes by separating chromosomes with CHEF gel electrophoresis followed by Southern blotting using a probe for the end of the chromosome. For accurate quantifications, we used the *dmc1* mutant background, in which DSBs are not repaired and accumulate instead (Bishop *et al.* 1992). We analyzed DSBs on chromosomes III (0.34 Mbp in length), V (0.58 Mbp), and XII (1.08 Mbp). In the *dmc1* mutant, each chromosome shows a unique region-specific capacity for DSB formation with hot and cold regions (Borde and de Massy 2013; de Massy 2013). As reported previously (Borde *et al.* 2009; Bani Ismail *et al.* 2014), the *set1 dmc1* double mutation reduces the frequency of DSBs along chromosomes III and XII, particularly on the left arm of chromosome III and in the region just to the right of the *CEN12* locus (Figure 3, A and B). Most of DSB-hot regions were downregulated and some regions were more prone for DSB formation in the *set1 dmc1* mutant than in the wild type (*dmc1*), as reported previously (Sollier *et al.* 2004; Borde *et al.* 2009). On the *dmc1* background, the *rtf1* mutation led to a slight, but significant reduction in the total number of DSBs on chromosomes III and XII (Figure 3C). Compared to the *dmc1* single mutant, the *rtf1 dmc1* double mutant had a decreased frequency of some DSBs in DSB-hot regions on both arms of chromosomes III and XII (shown by blue arrows in Figure 3). Importantly, the *rtf1 dmc1* double mutant showed a different distribution of DSBs on these chromosomes in comparison to the *set1 dmc1* mutant. DSBs in the *rtf1 dmc1* were decreased in some regions that were DSB hot in the wild type and also affected by the *set1* mutation. However, there were regions where DSB levels were apparently increased in the *rtf1 dmc1* double mutant relative to their number in the *dmc1* mutant (shown in red arrows). Increased levels of DSBs in these regions were not seen in the *set1 dmc1* mutant. Thus, the effect of the *rtf1* mutation on the DSB landscape is different from that of the *set1* mutation. Moreover, the DSB distribution in the *rtf1 set1 dmc1* triple mutant was similar to that in the *rtf1 dmc1* double mutant,

but different from the distribution in the *set1 dmc1* mutant. These findings suggest that *Rtf1* may function upstream of *Set1* in regulating the DSB landscape and may function in part independently of H3K4 methylation. The *rtf1 set1 dmc1* triple mutant exhibited a greater reduction in total DSBs than the decreases detected in the *set1 dmc1* or *rtf1 dmc1* double mutants, suggesting distinct contributions of *Set1* and *Rtf1* to the regulation of DSB frequency. Similar, albeit less drastic, effects of the *rtf1* mutation on DSB formation are seen on chromosome V (Figure S2A).

From the results of the DSB mapping, three conclusions may be drawn. First, *Rtf1* and *Set1* contribute to DSB patterning on chromosomes differentially, in a region-specific and chromosome-specific manner. Second, *Rtf1* functions upstream of *Set1* in regulating DSB patterning at least on some chromosomes. Third, *Rtf1* and *Set1* distinctly contribute to the regulation of DSB frequency in the absence of detectable H3K4 methylation marks.

Mapping of DSBs at individual chromosomal locus on the *dmc1* background: Using conventional Southern blotting, we also analyzed DSBs at the following representative loci: *YCR048W*, *GAT1*, and *PES4* (Figure 4A). At the *YCR048W* and *GAT1* loci, similar to our findings in the *set1 dmc1* mutant, the number of DSBs at 6 hr in the *rtf1 dmc1* double mutant decreased to ~30 and 40% of the levels seen in the *dmc1* mutant, respectively (Figure 4, B and C). Moreover, the *rtf1 set1 dmc1* triple mutant exhibited fewer DSBs at both loci compared to the levels observed in *rtf1 dmc1* or *set1 dmc1* double mutants. These findings support the notion of independent functions of *Rtf1* and *Set1* in the regulation of DSB formation. Previous studies indicated that the number of DSBs in some regions, *e.g.*, at the *PES4* locus, was increased in the *set1* mutant compared to the respective DSB frequency in the wild type (Borde *et al.* 2009; Sommermeyer *et al.* 2013). We also quantified DSBs at the *PES4* locus on the *dmc1* background and confirmed that at this location DSBs were more numerous in the *set1 dmc1* double mutant than in the *dmc1* mutant, consistent with the previous study (Borde *et al.* 2009). The *rtf1 dmc1* double mutant exhibited moderately increased levels of DSBs at this locus that were more frequent than the DSBs in the *dmc1* mutant, and similar to those in the *set1 dmc1* mutant. DSB levels in the *rtf1 set1 dmc1* triple mutant were similar to those detected in the *rtf1 dmc1* double mutant. These observations provided evidence that *Rtf1* and *Set1* repress DSB frequency at the *PES4* locus through a similar mechanism.

We also looked for a locus with a specific increase in DSBs specifically as a result of the *rtf1* mutation and identified the *ELO2* locus on chromosome III as a candidate. At this locus, the wild-type (*dmc1*) cells formed very few DSBs (bottom panels of Figure 4, B and C). On the other hand, DSBs were upregulated in the *rtf1 dmc1* mutant. This increase in the number of DSBs was less pronounced in the *set1 dmc1* mutant. These observations again support the hypothesis that *Rtf1* is of critical importance in regulating DSB formation, and that *Rtf1* has

Figure 3 Rtf1 and Set1 differently promote DSB formation. Distribution of DSBs along chromosomes XII (A) and III (B) was analyzed by PFGE followed by indirect labeling of one chromosome end using probes for *CHA1* and *AYT1*, respectively. Samples from meiotic time courses of the *dmc1* single (blue, MBY009/010), *rtf1 dmc1* double (red, SGY859/860), *set1 dmc1* double (green, SGY862/863), *set1 dot1 dmc1* triple (orange, SGY545/546), and *set1 rtf1 dmc1* triple (purple, SGY870/871) mutants were analyzed. On the left of the chromosome III panels, the approximate sizes of the chromosomes and the positions of the recombination hotspots, *HIS4-LEU2*, *ELO2*, and *THR4* are indicated. Blue arrows on the right side indicate DSBs, which are reduced by the *rtf1* mutation. Red arrows on the right side indicate DSB bands, which are increased by the *rtf1* mutation. Graphs below show the traces of the Phosphorimager signal at 8 hr. Pairwise comparison of traces is shown as indicated: the *dmc1* single, blue; *rtf1 dmc1*, red; *set1 dmc1*, green; *set1 dot1 dmc1*, orange; and *set1 rtf1 dmc1*, purple. Quantification of total DSBs at 8 hr for each chromosome by CHEF analysis of various strains (C) was carried out as described in *Materials and Methods*. Each plotted value represents the mean values $\pm$ SD from three independent time courses.

a repressive role. Moreover, at least at the *ELO2* locus, this repressive role is unlikely to be shared with *Set1*.

H3K4me marks are recognized by the PHD finger of the *Spp1* protein, a constituent of the COMPASS complex.

Spp1 binds to chromosome axes and also plays an additional role in the generation of DSBs (Acquaviva *et al.* 2013; Sommermeyer *et al.* 2013). We investigated the relationship between *Rtf1* and *Spp1* in regulating DSB

Figure 4 Rtf1 promotes the formation of DSBs in the absence of Set1. (A) Schematic representations of the *YCR048W*, *GAT1*, *PES4*, and *ELO2* loci. Sizes of DSB fragments are shown with lines below. (B) DSB formation at the above loci was determined in different strains by Southern blotting. The experiments were independently performed in triplicate and representative blots are shown. (C) The major DSB bands (DSB I) at the *YCR048W*, *GAT1*, *PES4*, and *ELO2* loci were quantified. The symbols represent the *dmc1* mutant (blue circles; MSY9/10), *rtf1 dmc1* mutant (red circles; SGY859/860), *set1 dmc1* mutant (green circles; SGY862/863), and *dmc1 set1 rtf1* mutant (purple circles; SGY870/871). Each plotted value represents the mean values $\pm$ SD from three independent time courses.

formation along chromosomes (Figure S3). CHEF-Southern for chromosomes III and XII revealed that the *rtf1* mutation affected the distribution of DSBs in the absence of *Spp1* in a similar way as in the absence of *Set1*. We also analyzed spore viability of *spp1 rtf1* double mutants. *spp1 rtf1* double mutants exhibited a frequency of survival of 87.9%, which is reduced relative to *spp1* single mutant (Figure S4A). These observations suggest that the *Set1*-independent role of *Rtf1* in DSB control is not mediated by the axis component *Spp1*. Like the *rtf1* and *set1* mutants (Figure 1C), the *spp1* single mutant showed a delay in the entry into MI (Figure S4B).

DSB frequency differs between *rtf1* and *dot1 set1* mutants

It is known that Paf1C promotes H3K79 methylation through its action on H2BK123 ubiquitination (Jaehning 2010). Our previous study showed that the *Dot1* histone H3K79 methyltransferase regulates DSB formation, particularly in the absence of *SET1* (Bani Ismail *et al.* 2014). Therefore, we hypothesized that some of the effects of *Rtf1* on the emergence of DSBs may be mediated by the action of *Dot1* in addition to that of *Set1*. To explore this possibility, we compared DSBs along chromosomes III and XII in *rtf1 dmc1*, *rtf1*

Figure 5 Mre11-focus formation in the *rad50S* mutant is reduced by the introduction of the *rtf1* mutation. (A) Distribution of DSBs along chromosomes III and XII was analyzed by PFGE followed by indirect labeling of one chromosome end using probes against *CHA1* and *AYT1*, respectively. Samples from meiotic time courses of strains described above were analyzed. (B) DSB profiles are shown as pairwise comparisons of the four strains. (C) Quantification of total DSBs by CHEF analysis of each strain in A. Details are described in *Materials and Methods*. Each plotted value represents the mean values from two independent time courses. (D) DSB formation at the *YCR048W* locus in different strains was determined by Southern blotting. Genomic DNA was digested with *Bgl*I. (E) DSB frequency was quantified by Southern blotting in the following strains: the *rad50S* mutant (blue, SGY83/84), *rtf1 rad50S* mutant (red, SGY107/108), *set1 rad50S* mutant (green, SGY89/90), and *set1 rtf1 rad50S* mutant (purple, SGY118/120). Each plotted value represents the mean values $\pm$ SD from three independent time courses. (F) Immunostaining analysis of Mre11 (green) in the *rad50S* mutant (SGY83/84), *rtf1 rad50S* mutant (SGY107/108), *set1 rad50S* mutant (SGY89/90), and *set1 rtf1 rad50S* mutant (SGY118/120) strains was carried out. Bar, 2 μ m. (G) Quantification of Mre11 foci in various mutants. For each focus-positive chromosome spread, the numbers of Mre11 foci at 8 hr were counted and plotted as shown. Error bars represent standard error of the mean (SEM; $n = 3$).

set1 dmc1, and *dot1 set1 dmc1* mutants (Figure 3). As reported previously (Bani Ismail *et al.* 2014), the *dot1 set1 dmc1* triple mutant had fewer DSBs on both chromosome III

and chromosome XII than the *set1 dmc1* double mutant. The right arm of chromosome III and the left arm of XII were particularly affected in the triple mutant. The pattern

seen in the *set1 dot1 dmc1* triple mutant is clearly different from that observed in the *rtf1 dmc1* double mutant. Furthermore, there were fewer DSBs in the *set1 dot1 dmc1* triple mutant than in the *rtf1 dmc1* double mutant. The difference was clearly seen in the right arm of chromosome III and in the left arm of chromosome XII. We conclude that *Rtf1* affects the DSB landscape irrespectively of the *Dot1* function, *i.e.*, independently of H3K79 methylation. Moreover, *Rtf1* may have a negative (repressive) role on DSB formation in the absence of two histone methylation marks, H3K4me and H3K79me. Previous results showed that the combination of the *set1* and *dot1* mutations can suppress *dmc1* arrest in prophase I (Bani Ismail *et al.* 2014). On the other hand, the *rtf1* mutation did not suppress the arrest (Figure S2B).

Mapping of DSBs in the *rad50S* background: To more precisely estimate changes in DSB levels, we also mapped DSBs on the background of the *rad50S* mutation, which accumulates unresected DSBs and thus yields a discrete Southern signal (Alani *et al.* 1990) (Figure 5, A and B). On chromosome III, the *rtf1 rad50S* mutant exhibited a reduced DSB frequency of 35.2% compared to the 47.2% of wild type (Figure 5C). DSB frequencies were more strongly reduced in the *set1 rad50S* mutant (22.1%) and the *rtf1 set1 rad50S* triple mutant showed more reduced DSB frequency than either double mutant, suggesting nonredundant roles of *Rtf1* and *Set1* in DSB frequency. This conclusion was confirmed using conventional Southern blotting to analyze DSBs at the *YCR048W* locus on chromosome III (Figure 5D). At this locus, *rtf1 rad50S* and *set1 rad50S* double mutants exhibited similarly reduced amounts of DSBs compared to the numbers detected in the wild type (28 and 13%, respectively at *YCR048W*; 8 hr; Figure 5E). Moreover, the *rtf1 set1 rad50S* triple mutant exhibited an even larger reduction of DSBs in comparison to the numbers seen in the *rad50S* mutant (4.3% at *YCR048W*; 8 hr time point). Unlike on the *dmc1* background, the DSB pattern on chromosome III was very similar between *rtf1 rad50S* and *set1 rad50S* double mutants, as well as the *rtf1 set1 rad50S* triple mutant (Figure 5B). This observation indicates similar effects of *rtf1* and *set1* on DSB patterning on chromosome III in the *rad50S* background. Notably, the DSB pattern on chromosome XII is different between *rtf1 rad50S* and *set1 rad50S* mutants, and the *rtf1 set1 rad50S* mutant exhibited a similar pattern to the *rtf1 rad50S* mutant. Thus, the contributions of *Rtf1* and *Set1* with respect to DSB pattern and frequency are different between the background of *dmc1* and *rad50S*, and also between chromosomes (see Discussion).

Counting *Mre11* foci on the *rad50S* background: Finally, as an independent measure of overall DSB levels, we performed immunostaining analysis of *Mre11*, which binds to DSB ends and shows focal staining on meiotic chromosomes of the *rad50S* background. The number of *Mre11* foci is proportional to that of DSBs in the genome (M. Shinohara, unpublished

results). We determined the average number of *Mre11* foci in the *rtf1 rad50S* and *set1 rad50S* double mutants (Figure 5F). In the *rad50S* single mutant, the average number of *Mre11* foci comprised 24.3 ± 4.6 (mean $\pm$ SE; $n = 3$) at 8 hr (Figure 5G). In the *rtf1 rad50S* and *set1 rad50S* double mutants that number was reduced to 10.2 ± 0.78 and 12.2 ± 0.32 , respectively. This finding was another confirmation of the hypothesis that *Rtf1* and *Set1* have a comparable contribution toward DSB formation. The *rtf1 set1 rad50S* triple mutant exhibited an even larger reduction of *Mre11* foci with the average number of 4.2 ± 0.7 , *i.e.*, much lower than the levels detected in the double mutants. These findings indicate that *Rtf1* and *Set1* independently contribute to the formation of *Mre11* foci, further supporting the notion that *Rtf1* and *Set1* have different functions in the regulation of DSB formation.

Genomic features of DSB redistribution in *rtf1* mutants

To specifically investigate the genomic features associated with the altered hotspot distribution in *rtf1* mutants, we determined hotspot patterns across the genome using a microarray-based approach (Blitzblau *et al.* 2007). ssDNA arrays measure the relative levels of ssDNA, which transiently accumulates at DSB sites during meiotic recombination. In the absence of *DMC1*, ssDNA is not turned over (Bishop *et al.* 1992), allowing for a cumulative measurement of meiotic DSB activity (Blitzblau *et al.* 2007; Buhler *et al.* 2007). Analysis of ssDNA profiles of *dmc1* and *rtf1 dmc1* mutants confirmed the redistribution of the DSB hotspots along chromosomes III, V, and XII (Figure 6A) seen in the Southern blot analysis (Figure 3 and Figure S2), and revealed similar changes across the entire genome (Figure S5A and File S1). Inspection of individual ssDNA peaks further indicated that the redistribution is largely a consequence of differential DSB activity at existing hotspots rather than the appearance of *de novo* hotspots.

To quantify these changes, we compiled a list of the strongest ssDNA peaks, representative of the strongest DSB hotspots, using a stringent significance cutoff. This analysis yielded 476 significant peaks in the *dmc1* mutant and 413 peaks in the *rtf1 dmc1* mutant (Figure 6B). The majority of the wild-type (*dmc1*) hotspots (279) were maintained in the absence of *RTF1*. In the *rtf1 dmc1* mutant, 197 peaks fell below the significance cutoff when compared to the *dmc1* mutant, while 134 new hotspot peaks became significant. Thus, the activity of a large fraction of hotspots is differentially affected by the loss of *RTF1*.

Based on these broad changes, we investigated whether specific chromosomal features are associated with these altered DSB patterns. No clear associations with larger scale chromosomal features, such as proximity to telomeres or centromeres, were observed (data not shown). We observed that hotspots gained in the absence of *RTF1* were somewhat enriched in regions far from replication origins (Figure 6C). This bias is statistically significant (Wilcoxon–Mann–Whitney test), but the functional importance of this effect is unclear,

Figure 6 Genome-wide mapping of ssDNAs in the *rtf1* mutant. (A) ssDNA distribution on chromosome III, V, and XII in the *dmc1* single (blue, MBY009/010), *rtf1 dmc1* double (red, SGY859/860). For chromosome V and XII, only a ~300-kb region of the left end of the chromosome is shown. Triangles indicate significant hotspots of ssDNA enrichment. (B) Venn diagram representing the overlap between significant hotspots in the *dmc1* single and *rtf1 dmc1* double mutants. Blue, the *dmc1* only; red *rtf1* only; purple common in *dmc1* and *rtf1 dmc1*. (C) Distribution of distances between DSB peaks and replication origins in different DSB classes. Distance between a DSB peak and the nearest origin was measured and plotted for each class. (D) Distribution of all significant hotspots in *dmc1* and *rtf1 dmc1* mutants relative to the position of transcripts for four classes: in genes or between tandem, divergent, or convergent transcripts. (E) Relationship of average nucleosome occupancy in each class was mapped relative to the distance from the DSB summits (in base pairs).

as we failed to observe a significant difference with respect to average replication timing (Figure S5B).

In wild type, the strongest hotspots preferentially occur in the promoter regions of divergent gene pairs (Blitzblau *et al.* 2007). As *RTF1* function is linked to transcriptional elongation (Mueller and Jaehning 2002; Squazzo *et al.* 2002), we wondered whether this promoter bias is altered

in the *rtf1 dmc1* mutants. We classified hotspot peaks based on their location with respect to gene bodies and whether they occurred in the intergenic regions between divergent, convergent, or tandem gene pairs (Figure 6D). These analyses revealed a significant redistribution of hotspots in the *rtf1 dmc1* mutants (chi-square test; $P = 0.007$), and showed that the hotspots most affected by the loss of *RTF1* are those

occurring in promoters at divergent genes. By contrast, hotspots between tandem or convergent gene pairs and in gene bodies remained largely unaffected. These data indicate that *RTF1* is not required for the overall promoter bias of meiotic DSB hotspots but that it plays a role in increasing DSB activity in the promoters of divergent gene pairs.

Given that *RTF1* regulates histone modifications (Figure 1A), we asked whether the changed hotspot activity observed in the *rtf1 dmc1* mutants was reflected in the local chromatin structure. Genome-wide meiotic nucleosome occupancy has been determined for wild-type cells (Pan *et al.* 2011). Therefore, we investigated whether hotspots with changed activity upon loss of *RTF1* differed in their wild-type nucleosome occupancy from hotspots that were unaffected (Figure 6E). To obtain the necessary DSB resolution for this analysis, we identified the precise sites of DSB formation underlying the ssDNA peaks by correlating them with available *Spo11* oligonucleotide data (Pan *et al.* 2011). This analysis revealed that hotspots that become more active in the *rtf1 dmc1* mutants have a higher chance to be occupied by nucleosomes in wild type. By contrast, DSBs maintained between the *dmc1* and *rtf1 dmc1* mutants or lost in the *rtf1 dmc1* mutant are largely depleted of nucleosomes in wild type. These observations suggest that suppression of DSB formation by *RTF1* is associated with increased local nucleosome occupancy, raising the intriguing possibility that *Rtf1*, and perhaps Paf1C, modulates DSB activity by controlling chromatin accessibility.

Discussion

Paf1C promotes DSB formation

Previous studies in budding yeast showed that methylation of two histone H3 lysines, H3K4 and H3K79, has a important role during meiotic DSB formation (Sollier *et al.* 2004; Borde *et al.* 2009; Bani Ismail *et al.* 2014). In particular, histone H3K4 methylation is a critical determinant of efficient DSB formation as well as selection of DSB sites (Borde *et al.* 2009; Acquaviva *et al.* 2013; Sommermeyer *et al.* 2013). However, in the absence of H3K4 methylation, DSB levels in the genome are only moderately reduced, while the overall distribution of DSBs along chromosomes, with some exceptions, remains largely unaffected (Borde *et al.* 2009). The complex effect of the H3K4-methylation deficit on DSB formation is apparent at the *PES4* locus that exhibits increased levels of DSBs in the absence of this methylation mark. Similar to the H3K4 methylation mutant, the *set1* mutant still shows a reasonable number of DSBs, harboring ~60% of DSBs detected in the wild type (Borde *et al.* 2009; Bani Ismail *et al.* 2014). Consistent with this mild DSB defect, the *set1* mutant and H3K4 methylation defective mutants showed high spore viability. This implies that H3K4 is not essential to DSB formation *per se*, but rather plays a regulatory role in efficient DSB formation, as proposed by Lam and Keeney (2015). Moreover, in the *set1* mutant, residual DSBs were nonrandomly distributed along the genome, suggesting the presence of additional determinants that shape the DSB landscape.

Indeed, we recently found that *Dot1* controls DSB frequency only in the absence of the *Set1*-dependent H3K4 methylation through the methylation of H3K79. However, the *set1 dot1* double mutant maintained a reasonable number of DSBs (40–50% of the levels seen in wild type) (Bani Ismail *et al.* 2014), indicating the existence of other determinants for DSB formation.

Initially, we were interested in the role of Paf1C in DSB formation since Paf1C is required for histone H2B ubiquitination, which, in turn, promotes both *Set1*-dependent H3K4 methylation and *Dot1*-dependent H3K79 methylation (Jaehning 2010). This functional link predicts that DSB defects seen in *paf1c* mutants should be very similar to those in the *set1 dot1* double mutant. However, although H3K4 methylation was abolished and H3K79 methylation strongly reduced in a mutant of the Paf1C component, *RTF1*, this mutant exhibited changes in DSB distribution and frequency that were different from those in the *set1 dot1* double mutant. These data suggest a role for *Rtf1*, and possibly for the whole Paf1C, in DSB formation that is distinct from the roles of *Set1* and *Dot1*. Moreover, the *set1 rtf1* double mutant had fewer DSBs than the *set1* or *rtf1* single mutants, consistent with a scenario whereby *Rtf1* regulates DSB formation in the absence of *Set1*, and, conversely, *Set1* promotes DSB formation in the absence of *Rtf1*. These roles of *Set1* and *Rtf1* are apparently independent of H3K4 methylation. The H3K4me-independent role of *Set1* can be explained by the recent finding that *Spp1*, a component of the *Set1*-containing complex, directly binds to *Mer2* (Acquaviva *et al.* 2013; Sommermeyer *et al.* 2013).

Paf1C shapes the DSB landscape

Importantly, the distribution of DSBs along a single chromosome in the *set1 dot1* double mutant is almost identical to that in the *set1* single mutant. This observation suggests the existence of additional factors determining DSB landscape on various chromosomes, particularly in some DSB-hot and DSB-cold regions in the absence of H3K4 and H3K79 methylation marks. In this study, we demonstrated that the DSB landscape in the *rtf1* mutant was altered in a different way compared to changes in DSB distribution in the *set1* and the *set1 dot1* mutants. Although overall DSB levels were reduced in the *rtf1* mutant, regions that displayed few DSBs in the wild type as well as in *set1* mutant cells, showed an increased capacity for DSB formation. Genome-wide ssDNA mapping identified 134 hotspots that show significantly increased DSB activities in the absence of *Rtf1*. This finding suggests a repressive role of *Rtf1*, and thus of the whole Paf1C, on the generation of DSBs in a region-specific manner. The formation of DSBs in these regions is independent of H3K4me. Therefore, *Rtf1* may play distinct roles in the control of DSB formation, positively affecting it through H3K4 methylation, as well as negatively regulating it. In some regions, Paf1C apparently promotes DSB formation through its role in H3K4 methylation. In other regions, however, this protein complex represses the emergence

of DSBs, possibly through mechanisms other than histone modification.

A possible scenario is that Paf1C modulates the dynamics or distribution of nucleosomes in promoter regions, as has been described for the *SER3* promoter (Pruneski *et al.* 2011). A redistribution of nucleosomes upon loss of Paf1C function may explain why hotspots that become more active in the *rtf1* mutant in fact appear occupied by nucleosomes in wild type. If this model applies, one would expect that nucleosome occupancy would be specifically reduced at these hotspots in the *rtf1* mutant. Conversely, the apparent loss of hotspots from divergent promoter regions may be the result of increased nucleosome occupancy at these loci in the *rtf1* mutant. A detailed analysis of nucleosome occupancy and dynamics in the *rtf1* mutant will help address this possibility. Moreover, any effects on nucleosome dynamics need not be direct, as Paf1C is mainly known for its function in mediating nucleosome reposition and histone modification during transcriptional elongation (Mueller and Jaehning 2002; Squazzo *et al.* 2002; Pruneski *et al.* 2011). Indeed, stronger DSB sites in the *set1* mutant were enriched in promoters of genes that were more highly transcribed in the *set1* mutant (Borde *et al.* 2009). A similar situation may apply for *rtf1* mutants, although at least in mitotic cells, *rtf1* deletion does not lead to strong changes in promoter usage (Penheiter *et al.* 2005).

An alternative, and not necessarily exclusive, model is that Paf1C may contribute to the formation of a particular chromosomal structure. During DSB formation, tethering of recombination hotspots through the 5'-region of genes is critical for the access of the hotspot to the Mer2 protein, a Spo11-activating protein on the axes (Acquaviva *et al.* 2013; Sommermeyer *et al.* 2013). One possibility is that Paf1C regulates tethering of DSB sites to the axes by changing the axis or loop structure (Kleckner 2006). Meiotic chromosome axes contain a number of proteins, including Hop1, Red1, Mek1, and the cohesion protein Rec8, whose mutations affect DSB landscape. Interestingly, the positioning of axis protein binding sites is strongly dependent on local transcription (Sun *et al.* 2015). One possibility is that Paf1C may control the assembly of these axis proteins on DNA molecules through its role in transcription, and, as a result, regulate the structure of the axis and tethered loops. To test this idea, we need to obtain more information on the regions exhibiting high levels of DSBs in the *rtf1* mutant on a genome-wide level and their relationships with other chromosomal proteins during meiosis.

In human, in addition to five conserved components, CDC73, CTR9, LEO1, PAF1, and RTF1, PAF1C contains one additional component, SKI8 (Kim *et al.* 2010). Interestingly, Ski8 in yeast is an essential factor for Spo11-mediated DSB formation (Arora *et al.* 2004). One possibility that needs to be investigated is that Paf1C may regulate the Ski8-dependent activity of Spo11. This is still an open question since there have been no reports about any functional relationship between Ski8 and Paf1C in yeast. Paf1C also promotes Set2-dependent H3K36 methylation (Jaehning 2010). It has previously been shown that the *set2* mutant increased DSBs

at the *HIS4* locus in the *rad50S* background (Merker *et al.* 2008). However, the role of this methylation mark in genome-wide DSB formation also remains to be elucidated.

Paf1C contributes to DSB formation differentially in *rad50S* and *dmc1* mutants

One of the surprising results of our study was a drastic difference of the effect of *rtf1* mutation on DSB frequency in the *rad50S* and *dmc1* mutants, which tend to accumulate unresected, protein-linked DSB ends and ssDNA molecules, respectively. During meiosis, DSB ends and ssDNA molecules activate checkpoint kinases Tel1^{ATM} and Mec1^{ATR}, correspondingly. It is well established that these two checkpoint effector kinases control DSB formation (Lange *et al.* 2011; Zhang *et al.* 2011; Farmer *et al.* 2012; Gray *et al.* 2013; Garcia *et al.* 2015). Particularly, it is known that Tel1^{ATM} promotes the repression of DSB formation through the phosphorylation of Rec114 (Carballo *et al.* 2013). We found the *rtf1* mutation reduced DSB levels much more on the *rad50S* background than on the *dmc1* background. This difference indicates a positive role of Rtf1 in DSB formation in conditions when Tel1^{ATM} is activated. It is possible that Paf1C mediates the Tel1^{ATM}-dependent feedback control of DSBs. Given that Dot1 is involved in Tel1^{ATM} signaling during meiosis (Ontoso *et al.* 2013), we propose a simple mechanism whereby Paf1C controls the Tel1^{ATM}-dependent regulation of DSBs indirectly through the Dot1-dependent H3K79 methylation. Alternatively, in the *rtf1* mutant, the normal DSB formation pathway is compromised and compensatory DSB formation depends on a Mec1-dependent pathway that is not activated in the *rad50S* mutant.

Since Tel1^{ATM} and Mec1^{ATR} are sequentially activated during meiotic recombination from unresected, protein-linked DSBs to resected DSBs, we hypothesize that the formation of early DSBs is likely to proceed initially under the control of Tel1^{ATM}, predominantly involving H3K4 methylation. In contrast, the formation of late DSBs, apparently under the principal control of Mec1^{ATR}, may use alternative cues for DSB generation, in addition to H3K4me. Collectively, these processes may affect the outcome of recombination as has been reported recently (Joshi *et al.* 2015).

Acknowledgments

We thank Doug Bishop, Susan Gasser, and Neil Hunter for discussions. We are also indebted to members of the Shinohara lab, especially Hisako Matsumoto and Ayaka Tokumura for their technical assistance. This work was supported by the Japanese Society for the Promotion of Science (JSPS) KAKENHI grants 22125001 and 22125002 to A.S., as well as by grants from the Takeda Science Foundation to A.S. M.S. was supported by JSPS through the Next Generation World-Leading Researchers program. This work was also supported in part by National Institutes of Health grant GM088248 and March Dimes research grant 6-FY13-105 to A.H. S.G. was supported by the Indian government as well as by the Institute for Protein Research.

S.K.G., H.S., M.S., A.H., and A.S. conceived and designed the experiments; S.K.G., H.S., and M.S. performed the experiments; S.K.G., M.S., and A.S. analyzed the data; N.J.P. and M.M.C. carried out the microarray experiments and analyzed the data; and S.K.G., M.S., N.J.P., A.H., and A.S. wrote the manuscript.

Literature Cited

- Acquaviva, L., L. Szekvolgyi, B. Dichtl, B. S. Dichtl, C. de La Roche Saint Andre *et al.*, 2013 The COMPASS subunit Spp1 links histone methylation to initiation of meiotic recombination. *Science* 339: 215–218.
- Alani, E., R. Padmore, and N. Kleckner, 1990 Analysis of wild-type and *rad50* mutants of yeast suggests an intimate relationship between meiotic chromosome synapsis and recombination. *Cell* 61: 419–436.
- Arora, C., K. Kee, S. Maleki, and S. Keeney, 2004 Antiviral protein Ski8 is a direct partner of Spo11 in meiotic DNA break formation, independent of its cytoplasmic role in RNA metabolism. *Mol. Cell* 13: 549–559.
- Bani Ismail, M., M. Shinohara, and A. Shinohara, 2014 Dot1-dependent histone H3K79 methylation promotes the formation of meiotic double-strand breaks in the absence of histone H3K4 methylation in budding yeast. *PLoS One* 9: e96648.
- Baudat, F., J. Buard, C. Grey, A. Fledel-Alon, C. Ober *et al.*, 2010 PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. *Science* 327: 836–840.
- Bishop, D. K., 1994 RecA homologs Dmc1 and Rad51 interact to form multiple nuclear complexes prior to meiotic chromosome synapsis. *Cell* 79: 1081–1092.
- Bishop, D. K., D. Park, L. Xu, and N. Kleckner, 1992 DMC1: a meiosis-specific yeast homolog of *E. coli* recA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell* 69: 439–456.
- Blat, Y., R. U. Protacio, N. Hunter, and N. Kleckner, 2002 Physical and functional interactions among basic chromosome organizational features govern early steps of meiotic chiasma formation. *Cell* 111: 791–802.
- Blitzblau, H. G., and A. Hochwagen, 2011 Genome-wide detection of meiotic DNA double-strand break hotspots using single-stranded DNA. *Methods Mol. Biol.* 745: 47–63.
- Blitzblau, H. G., G. W. Bell, J. Rodriguez, S. P. Bell, and A. Hochwagen, 2007 Mapping of meiotic single-stranded DNA reveals double-stranded-break hotspots near centromeres and telomeres. *Curr. Biol.* 17: 2003–2012.
- Blitzblau, H. G., C. S. Chan, A. Hochwagen, and S. P. Bell, 2012 Separation of DNA replication from the assembly of break-competent meiotic chromosomes. *PLoS Genet.* 8: e1002643.
- Borde, V., and B. de Massy, 2013 Programmed induction of DNA double strand breaks during meiosis: setting up communication between DNA and the chromosome structure. *Curr. Opin. Genet. Dev.* 23: 147–155.
- Borde, V., N. Robine, W. Lin, S. Bonfils, V. Geli *et al.*, 2009 Histone H3 lysine 4 trimethylation marks meiotic recombination initiation sites. *EMBO J.* 28: 99–111.
- Buhler, C., V. Borde, and M. Lichten, 2007 Mapping meiotic single-strand DNA reveals a new landscape of DNA double-strand breaks in *Saccharomyces cerevisiae*. *PLoS Biol.* 5: e324.
- Carballo, J. A., S. Panizza, M. E. Serrentino, A. L. Johnson, M. Geymonat *et al.*, 2013 Budding yeast ATM/ATR control meiotic double-strand break (DSB) Levels by down-regulating Rec114, an essential component of the DSB-machinery. *PLoS Genet.* 9: e1003545.
- Chu, S., and I. Herskowitz, 1998 Gametogenesis in yeast is regulated by a transcriptional cascade dependent on Ndt80. *Mol. Cell* 1: 685–696.
- de Massy, B., 2013 Initiation of meiotic recombination: How and where? Conservation and specificities among eukaryotes. *Annu. Rev. Genet.* 47: 563–599.
- Farmer, S., W. K. Leung, and H. Tsubouchi, 2011 Characterization of meiotic recombination initiation sites using pulsed-field gel electrophoresis. *Methods Mol. Biol.* 745: 33–45.
- Farmer, S., E. J. Hong, W. K. Leung, B. Argunhan, Y. Terentyev *et al.*, 2012 Budding yeast Pch2, a widely conserved meiotic protein, is involved in the initiation of meiotic recombination. *PLoS One* 7: e39724.
- Foreman, P. K., and R. W. Davis, 1996 CDP1, a novel *Saccharomyces cerevisiae* gene required for proper nuclear division and chromosome segregation. *Genetics* 144: 1387–1397.
- Garcia, V., S. Gray, R. M. Allison, T. J. Cooper and M. J. Neale, 2015 Tel1-mediated interference suppresses clustered meiotic double-strand-break formation. *Nature* 520: 114–118.
- Gray, S., R. M. Allison, V. Garcia, A. S. Goldman, and M. J. Neale, 2013 Positive regulation of meiotic DNA double-strand break formation by activation of the DNA damage checkpoint kinase Mec1(ATR). *Open Biol.* 3: 130019.
- Jaehning, J. A., 2010 The Paf1 complex: Platform or player in RNA polymerase II transcription? *Biochim. Biophys. Acta* 1799: 379–388.
- Joshi, N., M. S. Brown, D. K. Bishop, and G. V. Borner, 2015 Gradual implementation of the meiotic recombination program via checkpoint pathways controlled by global DSB levels. *Mol. Cell* 57: 797–811.
- Keeney, S., 2001 Mechanism and control of meiotic recombination initiation. *Curr. Top. Dev. Biol.* 52: 1–53.
- Keeney, S., C. N. Giroux, and N. Kleckner, 1997 Meiosis-specific DNA double-strand breaks are catalyzed by Spo11, a member of a widely conserved protein family. *Cell* 88: 375–384.
- Kim, J., M. Guermah, and R. G. Roeder, 2010 The human PAF1 complex acts in chromatin transcription elongation both independently and cooperatively with SII/TFIIS. *Cell* 140: 491–503.
- Kleckner, N., 2006 Chiasma formation: chromatin/axis interplay and the role(s) of the synaptonemal complex. *Chromosoma* 115: 175–194.
- Klein, F., P. Mahr, M. Galova, S. B. Buonomo, C. Michaelis *et al.*, 1999 A central role for cohesins in sister chromatid cohesion, formation of axial elements, and recombination during yeast meiosis. *Cell* 98: 91–103.
- Koch, C., P. Wollmann, M. Dahl, and F. Lottspeich, 1999 A role for Ctr9p and Paf1p in the regulation G1 cyclin expression in yeast. *Nucleic Acids Res.* 27: 2126–2134.
- Krogan, N. J., M. Kim, S. H. Ahn, G. Zhong, M. S. Kobor *et al.*, 2002 RNA polymerase II elongation factors of *Saccharomyces cerevisiae*: a targeted proteomics approach. *Mol. Cell. Biol.* 22: 6979–6992.
- Krogan, N. J., J. Dover, A. Wood, J. Schneider, J. Heidt *et al.*, 2003 The Paf1 complex is required for histone H3 methylation by COMPASS and Dot1p: linking transcriptional elongation to histone methylation. *Mol. Cell* 11: 721–729.
- Krogh, B. O., and L. S. Symington, 2004 Recombination proteins in yeast. *Annu. Rev. Genet.* 38: 233–271.
- Kugou, K., T. Fukuda, S. Yamada, M. Ito, H. Sasanuma *et al.*, 2009 Rec8 guides canonical Spo11 distribution along yeast meiotic chromosomes. *Mol. Biol. Cell* 20: 3064–3076.
- Lam, I., and S. Keeney, 2015 Mechanism and regulation of meiotic recombination initiation. *Cold Spring Harb. Perspect. Biol.* 7: a016634.
- Lange, J., J. Pan, F. Cole, M. P. Thelen, M. Jasin *et al.*, 2011 ATM controls meiotic double-strand-break formation. *Nature* 479: 237–240.

- Lichten, M., and B. de Massy, 2011 The impressionistic landscape of meiotic recombination. *Cell* 147: 267–270.
- Merker, J. D., M. Dominska, P. W. Greenwell, E. Rinella, D. C. Bouck *et al.*, 2008 The histone methylase Set2p and the histone deacetylase Rpd3p repress meiotic recombination at the *HIS4* meiotic recombination hotspot in *Saccharomyces cerevisiae*. *DNA Repair (Amst.)* 7: 1298–1308.
- Mueller, C. L., and J. A. Jaehning, 2002 Ctr9, Rtf1, and Leo1 are components of the Paf1/RNA polymerase II complex. *Mol. Cell. Biol.* 22: 1971–1980.
- Myers, S., R. Bowden, A. Tumian, R. E. Bontrop, C. Freeman *et al.*, 2010 Drive against hotspot motifs in primates implicates the PRDM9 gene in meiotic recombination. *Science* 327: 876–879.
- Ng, H. H., S. Dole, and K. Struhl, 2003 The Rtf1 component of the Paf1 transcriptional elongation complex is required for ubiquitination of histone H2B. *J. Biol. Chem.* 278: 33625–33628.
- Ontoso, D., I. Acosta, F. van Leeuwen, R. Freire, and P. A. San-Segundo, 2013 Dot1-dependent histone H3K79 methylation promotes activation of the Mek1 meiotic checkpoint effector kinase by regulating the Hop1 adaptor. *PLoS Genet.* 9: e1003262.
- Pan, J., M. Sasaki, R. Kniewel, H. Murakami, H. G. Blitzblau *et al.*, 2011 A hierarchical combination of factors shapes the genome-wide topography of yeast meiotic recombination initiation. *Cell* 144: 719–731.
- Panizza, S., M. A. Mendoza, M. Berlinger, L. Huang, A. Nicolas *et al.*, 2011 Spo11-accessory proteins link double-strand break sites to the chromosome axis in early meiotic recombination. *Cell* 146: 372–383.
- Parvanov, E. D., P. M. Petkov, and K. Paigen, 2010 Prdm9 controls activation of mammalian recombination hotspots. *Science* 327: 835.
- Penheiter, K. L., T. M. Washburn, S. E. Porter, M. G. Hoffman, and J. A. Jaehning, 2005 A posttranscriptional role for the yeast Paf1-RNA polymerase II complex is revealed by identification of primary targets. *Mol. Cell* 20: 213–223.
- Petronczki, M., M. F. Siomos, and K. Nasmyth, 2003 Un menage a quatre: the molecular biology of chromosome segregation in meiosis. *Cell* 112: 423–440.
- Pratto, F., K. Brick, P. Khil, F. Smagulova, G. V. Petukhova *et al.*, 2014 DNA recombination. Recombination initiation maps of individual human genomes. *Science* 346: 1256442.
- Pruneski, J. A., S. J. Hainer, K. O. Petrov, and J. A. Martens, 2011 The Paf1 complex represses *SER3* transcription in *Saccharomyces cerevisiae* by facilitating intergenic transcription-dependent nucleosome occupancy of the *SER3* promoter. *Eukaryot. Cell* 10: 1283–1294.
- Shi, X., A. Finkelstein, A. J. Wolf, P. A. Wade, Z. F. Burton *et al.*, 1996 Paf1p, an RNA polymerase II-associated factor in *Saccharomyces cerevisiae*, may have both positive and negative roles in transcription. *Mol. Cell. Biol.* 16: 669–676.
- Shilatfard, A., 2012 The COMPASS family of histone H3K4 methylases: mechanisms of regulation in development and disease pathogenesis. *Annu. Rev. Biochem.* 81: 65–95.
- Shinohara, A., H. Ogawa, and T. Ogawa, 1992 Rad51 protein involved in repair and recombination in *S. cerevisiae* is a RecA-like protein. *Cell* 69: 457–470.
- Shinohara, M., S. L. Gasior, D. K. Bishop, and A. Shinohara, 2000 Tid1/Rdh54 promotes colocalization of Rad51 and Dmc1 during meiotic recombination. *Proc. Natl. Acad. Sci. USA* 97: 10814–10819.
- Shinohara, M., K. Sakai, T. Ogawa, and A. Shinohara, 2003 The mitotic DNA damage checkpoint proteins Rad17 and Rad24 are required for repair of double-strand breaks during meiosis in yeast. *Genetics* 164: 855–865.
- Shinohara, M., S. D. Oh, N. Hunter, and A. Shinohara, 2008 Crossover assurance and crossover interference are distinctly regulated by the ZMM proteins during yeast meiosis. *Nat. Genet.* 40: 299–309.
- Smagulova, F., I. V. Gregoret, K. Brick, P. Khil, R. D. Camerini-Otero *et al.*, 2011 Genome-wide analysis reveals novel molecular features of mouse recombination hotspots. *Nature* 472: 375–378.
- Sollier, J., W. Lin, C. Soustelle, K. Suhre, A. Nicolas *et al.*, 2004 Set1 is required for meiotic S-phase onset, double-strand break formation and middle gene expression. *EMBO J.* 23: 1957–1967.
- Sommermeier, V., C. Beneut, E. Chaplais, M. E. Serrentino, and V. Borde, 2013 Spp1, a member of the Set1 Complex, promotes meiotic DSB formation in promoters by tethering histone H3K4 methylation sites to chromosome axes. *Mol. Cell* 49: 43–54.
- Squazzo, S. L., P. J. Costa, D. L. Lindstrom, K. E. Kumer, R. Simic *et al.*, 2002 The Paf1 complex physically and functionally associates with transcription elongation factors in vivo. *EMBO J.* 21: 1764–1774.
- Storlazzi, A., L. Xu, L. Cao, and N. Kleckner, 1995 Crossover and noncrossover recombination during meiosis: timing and pathway relationships. *Proc. Natl. Acad. Sci. USA* 92: 8512–8516.
- Sun, X., L. Huang, T. E. Markowitz, H. G. Blitzblau, D. Chen *et al.*, 2015 Transcription dynamically patterns the meiotic chromosome-axis interface. *eLife*: e07424.
- Symington, L. S., and J. Gautier, 2011 Double-strand break end resection and repair pathway choice. *Annu. Rev. Genet.* 45: 247–271.
- Wach, A., A. Brachat, R. Pohlmann, and P. Philippsen, 1994 New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast* 10: 1793–1808.
- Xu, L., B. M. Weiner, and N. Kleckner, 1997 Meiotic cells monitor the status of the interhomolog recombination complex. *Genes Dev.* 11: 106–118.
- Yamashita, K., M. Shinohara, and A. Shinohara, 2004 Rad6-Bre1-mediated histone H2B ubiquitylation modulates the formation of double-strand breaks during meiosis. *Proc. Natl. Acad. Sci. USA* 101: 11380–11385.
- Zhang, L., K. P. Kim, N. E. Kleckner, and A. Storlazzi, 2011 Meiotic double-strand breaks occur once per pair of (sister) chromatids and, via Mec1/ATR and Tel1/ATM, once per quartet of chromatids. *Proc. Natl. Acad. Sci. USA* 108: 20036–20041.
- Zhu, Z., S. Mori, H. Oshiumi, K. Matsuzaki, M. Shinohara *et al.*, 2010 Cyclin-dependent kinase promotes formation of the synaptonemal complex in yeast meiosis. *Genes Cells* 15: 1036–1050.

Communicating editor: N. Hunter

GENETICS

Supporting Information

www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.177287/-/DC1

The Double-Strand Break Landscape of Meiotic Chromosomes Is Shaped by the Paf1 Transcription Elongation Complex in *Saccharomyces cerevisiae*

Santosh K. Gothwal, Neem J. Patel, Meaghan M. Colletti, Hiroyuki Sasanuma, Miki Shinohara,
Andreas Hochwagen, and Akira Shinohara

Figure S1. The effect of *paf1C* mutations on various histone methylation marks. Levels of various histone methylation marks during meiosis. Levels of histone H3K4-me1, -me2 and -me3 as well as H3K79-me3 as well as α -tubulin in the wild type (NKY1543/1303), *rtf1* (SGY3/4), *leo1* (SGY5/6), and *cdc73* (SGY7/8) strains as a control were determined by western blotting. At each time point, cells were fixed with TCA and cell lysates were subject to the analysis. Representative images are shown.

A

B

Figure S2. Rtf1 promotes DSB formation in the absence of Set1. (A) Distribution of DSBs along Chromosome V was analyzed by PFGE followed by indirect labeling of one chromosome end using a probe for RMD6. Samples from meiotic time courses of the *dmc1* mutant (MBY009/010), *dmc1 rtf1* mutant (SGY859/860), *dmc1 set1* mutant (SGY862/863), and *dmc1 set1 rtf1* mutant (SGY870/871) were analyzed. Representative blots are shown. Pairwise comparison of traces are shown as indicated on the right. (B) Meiotic cell division I was analyzed by DAPI staining in *dmc1* (blue triangles; MBY009/010), and *dmc1 rtf1* (red triangles; SGY862/863) cells. At least 100 cells were counted for each time point. Each plotted value represents the mean values $\pm$ standard deviation (S.D.) from three independent time courses.

Figure S3. Rtf1 promotes the formation of DSBs in the absence of Spp1. Distribution of DSBs along Chromosomes III and XII was analyzed by PFGE followed by indirect labeling of one chromosome end using probes against *CHA1* and *AYT1*, respectively. Samples from meiotic time courses of the *dmc1* mutant (SGY225/226), *dmc1 rtf1* mutant (SGY227/228), *dmc1 spp1* mutant (SGY842/843), and *dmc1 spp1 rtf1* mutant (SGY847/848) were analyzed. Representative blots are shown. Pairwise comparison of traces are shown as indicated.

Figure S4. Rtf1 promotes the formation of DSBs in the absence of Spp1. (A) Spore viability of various strains was measured after dissecting tetrads. Spores were incubated at 30°C for 3 days. Each bar indicates the percentage of spore viability and number of total dissected tetrads (parentheses). *spp1* mutant, green, SGY837/838; *rtf1 spp1* mutant, purple, SGY846/848. (B) Meiotic cell division I was analyzed by DAPI staining in wild-type (blue circles; NKY1303/1543) and *spp1* (green triangles; SGY47/48) cells. At least 100 cells were counted for each time point. Each plotted value represents the mean values $\pm$ standard deviation (S.D.) from three independent time courses.

Figure S5. Genome-wide mapping of ss DNAs in the absence of Rtf1. (A) ssDNA distribution on chromosome I to XVI in the *dmc1* single (blue, MBY009/010), *rtf1 dmc1* double (red, SGY859/860). Triangles indicate statistically significant hotspots. (B) Relative replication timing of the three different hotspot classes. Wild-type replication timing of each hotspot was extracted from (Blitzblau et al., 2012) and was plotted as function of hotspot class.

Table S1. Strain list

MSY831	<i>MATa</i> , <i>ho::LYS2</i> , <i>lys2</i> , <i>ura3</i> , <i>leu2::hisG</i> , <i>trp1::hisG</i>
MSY833	<i>MATα</i> , <i>ho::LYS2</i> , <i>lys2</i> , <i>ura3</i> , <i>leu2::hisG</i> , <i>trp1::hisG</i>
NKY1303	<i>MATa</i> , <i>ho::LYS2</i> , <i>lys2</i> , <i>ura3</i> , <i>leu2::hisG</i> , <i>his4B-LEU2(MluI)</i> , <i>arg4-bgl</i>
NKY1543	<i>MATα</i> , <i>ho::LYS2</i> , <i>lys2</i> , <i>ura3</i> , <i>leu2::hisG</i> , <i>his4X-LEU2(BamHI)-URA3</i> , <i>arg4-nsp</i>
SGY3	NKY1303 with <i>rtf1::TRP1</i>
SGY4	NKY1543 with <i>rtf1::TRP1</i>
SGY5	NKY1303 with <i>cdc73::TRP1</i>
SGY6	NKY1543 with <i>cdc73::TRP1</i>
SGY7	NKY1303 with <i>leo1::TRP1</i>
SGY8	NKY1543 with <i>leo1::TRP1</i>
SGY19	NKY1303 with <i>set1::URA3</i>
SGY20	NKY1543 with <i>set1::URA3</i>
SGY47	NKY1303 with <i>set1::URA3</i> , <i>rtf1::TRP1</i>
SGY48	NKY1543 with <i>set1::URA3</i> , <i>rtf1::TRP1</i>
SGY51	NKY1303 with <i>set1::URA3</i> , <i>cdc73::TRP1</i>
SGY52	NKY1543 with <i>set1::URA3</i> , <i>cdc73::TRP1</i>
SGY83	NKY1303 with <i>rad50S::URA3</i>
SGY84	NKY1543 with <i>rad50S::URA3</i>
SGY870	NKY1303 with <i>set1::URA</i> , <i>rtf1::TRP1</i> , <i>dmc1::URA3</i>
SGY871	NKY1543 with <i>set1::URA</i> , <i>rtf1::TRP1</i> , <i>dmc1::URA3</i>
SGY862	NKY1303 with <i>set1::URA3</i> , <i>dmc1::URA3</i>
SGY863	NKY1543 with <i>set1::URA3</i> , <i>dmc1::URA3</i>
SGY854(MBY9)	NKY1303 with <i>dmc1::URA</i>
SGY855(MBY10)	NKY1543 with <i>dmc1::URA3</i>
SGY859	NKY1303 with <i>rtf1::TRP1</i> , <i>dmc1::URA3</i>
SGY860	NKY1543 with <i>rtf1::TRP1</i> , <i>dmc1::URA3</i>
SGY107	NKY1303 with <i>rtf1::TRP1</i> , <i>rad50S::URA3</i>
SGY108	NKY1543 with <i>rtf1::TRP1</i> , <i>rad50S::URA3</i>
SGY89	NKY1303 with <i>rad50S::URA3</i> , <i>set1::URA3</i>

SGY90	NKY1543 with <i>rad50S::URA3, set1::URA3</i>
SGY118	NKY1303 with <i>rad50S::URA3, set1::URA3, rtf1::TRP1</i>
SGY120	NKY1543 with <i>rad50S::URA3, set1::URA3, rtf1::TRP1</i>
SGY543	NKY1303 with <i>dmc1::URA3, set1::URA3, dot1::KANMX6</i>
(MBY 282)	
SGY544	NKY1543 with <i>dmc1::URA3, set1::URA3, dot1::KANMX6</i>
(MBY285)	
SGY225	MSY831 with <i>dmc1::URA3</i>
SGY226	MSY833 with <i>dmc1::URA3</i>
SGY227	MSY831 with <i>rtf1::TRP1, dmc1::URA3</i>
SGY228	MSY833 with <i>rtf1::TRP1, dmc1::URA3</i>
SGY837	MSY831 with <i>spp1::HYGMX6,</i>
SGY838	MSY833 with <i>spp1::HYGMX6,</i>
SGY842	MSY831 with <i>spp1::HYGMX6, dmc1::URA3</i>
SGY843	MSY833 with <i>spp1::HYGMX6, dmc1::URA3</i>
SGY846	MSY831 with <i>rtf1::TRP1, spp1::HYGMX6,</i>
SGY848	MSY833 with <i>rtf1::TRP1, spp1::HYGMX6,</i>
SGY847	MSY831 with <i>rtf1::TRP1, spp1::HYGMX6, dmc1::URA3</i>
SGY848	MSY833 with <i>rtf1::TRP1, spp1::HYGMX6, dmc1::URA3</i>

Table S2. Primer list

Strains	DNA sequence
<i>paf1::TRP1</i>	5'-CAGAAATGTATTTCAGTACAATAGAACAGTGCTCATAATAGTAT AAAGGGAGACATTACTATATATATAATATAGGAAGC 5'-CTTGTAAGTTTCCTTTTCTTCCTCTGGTTTTTGTTCAGTATG AACATATTATTAACGAGGCAAGTGCACAAACAATAC
<i>cdc73::TRP1</i>	5'-TCTTAATTCGGGGCGTTAAAAGAATAATAATTTGAGCAAGAAA CTGGTGAGACATTACTATATATATAATATAGGAAGC 5'-TTCAACGGTATCCTCTTGAAATAAGTTCCTTTTCTAAACTATG CCAGTATTATTAACGAGGCAAGTGCACAAACAATAC
<i>rtf1::TRP1</i>	5'-GAAGGATACCAAAGTTTGTAATTGTATTGCACTAATTTGTTGA GAGCACAGACATTACTATATATATAATATAGGAAGC 5'-ATTTGATATCCAATTCACCAACCATGCGCTCCAACCCACCAA GGCGCTATTATTAACGAGGCAAGTGCACAAACAATAC
<i>leo1::TRP1</i>	5'-CGTAAACTAGGTGCGTCCAATCAAAGTAATCCAATTAGATATA CTGGACAGACATTACTATATATATAATATAGGAAGC 5'-CCTCGTCTTCGTGTCCTCGATGACCGCAACCCTTCTTCTTTT TGTCTATTATTAACGAGGCAAGTGCACAAACAATAC
<i>ctr9::TRP1</i>	5'-ACTTCTGTGCAAAGTTCTAATTGTCTGGTCCATTTGTGTTGAG AGCAAGAGACATTACTATATATATAATATAGGAAGC 5'-CAATGATAATGATGATAACGACGGATTGTTCTAAGAAGAGGC GAGGGTATTATTAACGAGGCAAGTGCACAAACAATAC

File S1. Supplemental Data. ssDNA distribution on chromosome I to XVI in the *dmc1* single (blue, MBY009/010), *rtf1 dmc1* double (red, SGY859/860) mutants. Significant peaks in the *dmc1* single and *rtf1 dmc1* double mutant strains are shown in blue and red triangles, respectively.

position on chr 1 (kb)

position on chr 14 (kb)

position on chr 13 (kb)

position on chr 6 (kb)