Published in final edited form as:

Dev Sci. 2016 May ; 19(3): 452–468. doi:10.1111/desc.12312.

Rethinking the concepts of ‘local or global processors’: evidence from Williams syndrome, Down syndrome, and Autism Spectrum Disorders

Dean D’Souza¹, Rhonda Booth², Monica Connolly¹, Francesca Happé³, and Annette Karmiloff-Smith¹

¹Birkbeck Centre for Brain and Cognitive Development, University of London, UK

²Institute of Child Health, University College London, UK

³Social, Genetic and Developmental Psychiatry Research Centre, Institute of Psychiatry, King’s College London, UK

Abstract

Both Williams syndrome (WS) and Autism Spectrum Disorder (ASD) have been characterized as preferentially processing local information, whereas in Down syndrome (DS) the reported tendency is to process stimuli globally. We designed a cross-syndrome, cross-task comparison to reveal similarities and differences in local/global processing in these disorders. Our in-depth study compared local/global processing across modalities (auditory-verbal/visuo-spatial) and levels of processing (high/low) in the three syndromes. Despite claims in the literature, participants with ASD or WS failed to show a consistent local processing bias, while those with DS failed to show a reliable global processing bias. Depending on the nature of the stimuli and the task, both local and global processing biases were evident in all three neurodevelopmental disorders. These findings indicate that individuals with neurodevelopmental disorders cannot simply be characterized as local or global processors.

Introduction

The world is perceived as hierarchically organized. For instance, visual scenes comprise global percepts (e.g. trees) that are composed of local details (leaves). The ability to process and integrate information at both global and local levels makes it possible to build up context, discover structure, make classifications, and form generalizations over a wide range of contexts. This ability develops early. For instance, Slater and colleagues (1991) demonstrated that even neonates have the capacity to make local information (line segments) cohere into unified wholes (angles, shapes). However, in three neurodevelopmental disorders, namely, Williams syndrome (WS), Down syndrome (DS), and Autism Spectrum

Address for correspondence: Annette Karmiloff-Smith, Birkbeck Centre for Brain and Cognitive Development, University of London, 32 Torrington Square, London WC1N 7HX, UK; a.karmiloff-smith@bbk.ac.uk.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Data S1. Additional information on the relationship between Mental Age and task performance.

Disorder (ASD), local processing (also sometimes referred to as a piecemeal or detailed-focused information processing style) and global processing (also sometimes referred to as a holistic processing style) are purported to be dissociated (see Table 1 for a description of these neurodevelopmental disorders). In both WS and ASD, the processing of local detail is reported to predominate over the processing of global properties (e.g. Bellugi, Lichtenberger, Jones, Lai & St George, 2000; Brosnan, Scott, Fox & Pye, 2004; Farran, Jarrold & Gathercole, 2003; Happé & Frith, 2006; Mottron, Belleville & Ménard, 1999; Wang, Mottron, Peng, Berthiaume & Dawson, 2007). For example, children with ASD tend not to benefit from familiar gestalts and are thus slower at enumerating dots in canonical patterns (like dots on dice) than children with mild learning disabilities and mental-age matched controls (Jarrold & Russell, 1997). They appear to count each dot one after the other, rather than utilizing the global information of the canonical pattern. Individuals with WS also preferentially process local information. For example, on a drawing task, individuals with WS accurately represent the local detail of a picture but not the gestalt relationships among the local elements (Bellugi, Sabo & Vaid, 1988; Bellugi, Lichtenberger, Mills, Galaburda & Korenberg, 1999). Moreover, the local processing bias in WS and ASD may have a common underlying mechanism (namely, a bias in *attention*; Porter & Coltheart, 2006). In contrast, individuals with DS present with the opposite profile, that is, they often show impairments in processing local detail. For example, Bihrlé, Bellugi, Delis and Marks (1989) found a double dissociation between individuals with DS and WS. Participants were asked to copy a large global figure composed of smaller local forms (e.g. a 'D' made up of small 'Y's). While chronological-age and mental-age matched controls copied the figures accurately, individuals with WS tended to produce only the local forms of the figures, and individuals with DS only the overall global shape.

It has not hitherto been established how or why these biases emerge, although impairment of the dorsal visual pathway may account for some of them. In typical development, local-global processing increases in proficiency and changes over time. Converging paradigms indicate a developmental shift from an initial global processing bias in infancy to a local processing bias in early childhood (toddlers, preschoolers), with the global bias gradually returning from middle childhood onwards (in some domains as early as 6 years of age; Poirel, Mellet, Houdé & Pineau, 2011¹). Interestingly, the shift from local to global processing in the visual domain has been associated with a reduction of grey matter along the visual dorsal stream – specifically, the calcarine sulcus, lingual gyrus, and parietal cortex of the right hemisphere (Poirel *et al.*, 2008). This loss of grey matter might reflect selective specialization (through neural pruning) in visual processing of global information. It raises the possibility that a disorder of the dorsal stream might display problems with processing global information, disrupting the typical global-to-local-to-global shift. Empirical studies of individuals with WS (known as 'local processors'; see Pani, Mervis & Robinson, 1999, for discussion) provide evidence that this might be the case. For example, a functional Magnetic Resonance Imaging (fMRI) study showed that on global processing tasks, individuals with

¹See also Dukette & Stiles, 2001; Farroni, Valenza, Simion & Umiltà, 2000; Frick, Colombo & Allen, 2000; Ghim & Eimas, 1988; Hadad & Kimchi, 2006; Johnson & Aslin, 1995; Kellman, Spelke & Short, 1986; Kimchi, Hadad, Behrmann & Palmer, 2005; Kovács, 2000; Kovács, Kozma, Feher & Benedek, 1999; Kozma, Kovács & Benedek, 2001; Poirel, Mellet, Houdé & Pineau, 2008; Quinn & Bhatt, 2006; Quinn, Bhatt, Brush, Grimes & Sharpnack, 2002; Quinn, Burke & Rush, 1993; Quinn & Eimas, 1986.

WS were not only worse than TD controls in their behavioural responses, they also showed reduced activation of the dorsal stream pathway (Mobbs, Eckert, Menon, Mills, Korenberg *et al.*, 2007). Reduced grey matter in intraparietal sulcus and adjacent parietal areas has also been identified in WS (Meyer-Lindenberg, Kohn, Mervis, Kippenhan, Olsen *et al.*, 2004).

Dorsal stream deficits have also been reported in ASD (e.g. Atkinson, 2009; Bertone, Mottron, Jelenic & Faubert, 2003; Braddick, Atkinson & Wattam-Bell, 2003; Spencer, O'Brien, Riggs, Braddick, Atkinson *et al.*, 2000). For instance, individuals with ASD find it more difficult than TD controls to discriminate the direction of a field of coherently moving dots when a proportion of the dots are moving randomly (Milne, Swettenham, Hansen, Campbell, Jeffries *et al.*, 2002; Spencer *et al.*, 2000). However, because individuals with ASD use the same amount of contrast as TD controls to detect sinusoidal luminance manipulations of a Gaussian patch ('flicker detection'), it has been argued that ASD involves impairment in the processing of global motion information rather than a *general* dorsal deficit per se (Pellicano, Gibson, Maybery, Durkin & Badcock, 2005; see also Koldewyn, Whitney & Rivera, 2011). Thus, in the autism literature, three different (albeit underspecified) theories have been proposed to characterize and explain the bias: (1) the weak central coherence theory (WCC); (2) the enhanced perceptual functioning model (EPF); and (3) the hierarchization deficit hypothesis (HDH).

According to WCC, the local bias in ASD results from a detail-focused processing style that characterizes the disorder (Frith, 1989, 2003; Frith & Happé, 1994; Happé, 1999; Happé & Frith, 2006). This is consistent with evidence that individuals with autism show relatively enhanced performance on tasks that require participants to overcome global-to-local interference, e.g. on the embedded figures task (Jolliffe & Baron-Cohen, 1997; Shah & Frith, 1983), on visual search tasks (Happé, 1999), and in copying impossible figures (Mottron *et al.*, 1999). The theory is also consistent with evidence that individuals with ASD do not use context (e.g. meaning, semantic organization) to facilitate task performance to the same extent as typically developing (TD) controls. For example, in contrast to TD children, in children with ASD recall of randomly arranged word lists is not much improved when the words are rearranged into meaningful groupings (Hermelin & O'Connor, 1970).

In contrast to the WCC theory, more recent studies, based on the EPF model, suggest that individuals with ASD do process and use global information (e.g. Mottron *et al.*, 1999; Ozonoff, Strayer, McMahon & Filloux, 1994; Plaisted, Swettenham & Rees, 1999). Their local bias is thought to reflect enhanced functioning of early, low-level perceptual processes (Mottron, Dawson, Soulières, Hubert & Burack, 2006) and stimulus complexity (i.e. local details are often less complex than global properties, although this would not of course explain the global bias in DS; Bertone, Mottron, Jelenic & Faubert, 2005; Minshew & Goldstein, 1993, 1998; Minshew, Goldstein & Siegel, 1995; Minshew, Sweeney & Bauman, 1997). This is consistent with the finding that individuals with ASD show an atypically enhanced ability to detect minor modifications in their environment (Rimland, 1971).

The EPF model proposes that individuals with ASD do not necessarily have difficulty in perceiving global form, but they have an overdeveloped or 'over-specialized' (low-level) perceptual system that can disrupt higher-level cognition and behaviour, depending on the

requirements of the task (Motttron & Burack, 2001; Motttron *et al.*, 2006). Indeed, evidence in support of this hypothesis² has led to some modifications in the WCC model. The original theory (Frith, 1989) posited a *deficit* in global processing or in the drive to cohere information. The modified theory (Frith, 2003; Happé & Frith, 2006) now claims that WCC arises as a result of a local processing *bias* that can be overcome in tasks with explicit demands for global processing. Thus, two of the three main theories explain the dissociation of local and global processing in ASD as the result of either an overdeveloped low-level processing system or a detail-focused processing bias.

The third major theory is the HDH (Motttron & Belleville, 1993), according to which both local and global processing function normally in individuals with ASD. The abnormality is purported to exist in the *interaction* between the two levels.³ Specifically, the authors suggest that the local and global levels are not hierarchically distinguishable in individuals with ASD, who find it difficult to switch from one level of representation/abstraction to the other (Plaisted *et al.*, 1999). This is consistent with evidence that a fairly rigid trade-off between either local or global processing exists in children and adults with ASD, with individuals performing well on one level or the other but rarely both, whereas TD individuals switch back and forth easily between local and global processing (Happé & Booth, 2008).

Although these three theories have been applied to ASD, they can also be evaluated as explanations of reported hierarchical abnormalities or impaired local/global processing in WS and DS. A processing bias focused on detail (consistent with the WCC model) has also been used to explain the behaviour of individuals with WS (e.g. Bellugi, Marks, Bihrlé & Sabo, 1988; Bellugi *et al.*, 1999; Bihrlé *et al.*, 1989). Indeed, individuals with WS neither show global precedence (Deruelle, Schon, Rondan & Mancini, 2005) nor make use of context (Elsabbagh, Cohen & Karmiloff-Smith, 2010) on, for instance, auditory perception tasks. Furthermore, although there is little evidence consistent with the EPF model in individuals with WS, there are data pointing to a hierarchical deficit in WS (Pani *et al.*, 1999), similar to the HDH account of ASD. Pani and colleagues (1999) administered a visual search task to participants with WS and, after manipulating the stimuli in ways that more or less facilitated their grouping, found that individuals with WS do not have a problem with processing information at any particular level of organization. Instead, their problem is thought to lie in switching between levels.

In contrast to individuals with WS or ASD, a processing bias focused on global properties of stimuli has been claimed in individuals with DS, which might be interpreted as relatively

²See Bouvet, Simard-Meilleur, Paignon, Motttron & Donnadiou (2014); Kéita, Guy, Berthiaume, Motttron & Bertone (2014); Muth, Hönekopp & Falter (2014); Olu-Lafe, Liederman & Tager-Flusberg (2014); and Yamasaki, Maekawa, Takahashi, Fujita, Kamio *et al.* (2014), for more recent evidence.

³Whereas (weak) central coherence – local–global processing – used to be viewed as a continuum (e.g. Happé, 1999), the evidence suggests that it is underpinned by at least two independent mechanisms. For instance, unilateral lesions of the left cerebral hemisphere impair local processing, whereas those in the right hemisphere impair global processing (Robertson & Lamb, 1991). Functional neuroimaging evidence (e.g. Heinze, Hinrichs, Scholz, Burchert & Mangun, 1998) and developmental studies (e.g. Dukette & Stiles, 1996, 2001) also support this general finding. For example, although segmentation and integration skills are somewhat reciprocal, they can vary even in the same individual and develop along different time courses (Dukette & Stiles, 2001). Moreover, the fact that the perceptual relation between local and global levels is sensitive to variations such as number and relative size of the local elements weakens the underlying assumption that the global form and local elements map directly onto distinct perceptual units, differing only in level of globality (Kimchi, 1992).

strong (i.e. normal) central coherence (CC) rather than WCC. For example, in producing hierarchical figures in the Navon task, individuals with DS tend to produce the correct global form, often ignoring or making errors on the local details (Bihrlé *et al.*, 1989; Bellugi *et al.*, 1999; Bellugi, Bihrlé, Jeringan, Trauner & Doherty, 1990; Bellugi *et al.*, 2000).

In sum, much of the literature seems to point to dissociation between local and global information processing in individuals with WS, DS, or ASD. Those with WS or ASD are characterized as having problems with processing global information, whereas those with DS experience more difficulty with processing local information. The literature suggests that these phenomena arise from the reduced ability to process global information in WS, a detail-focused or enhanced low-level perceptual bias in ASD, and the reduced ability to focus on local information in DS. An inability to flexibly switch between levels has also been hypothesized for both WS and ASD. To test this idea (i.e. that local/global processing is dissociated in these neurodevelopmental disorders), a battery of tasks that tap local/global processing across modalities (visuo-spatial, verbal-auditory) and levels of processing (from low to high: perceptual, attentional, semantic, strategic) was administered to individuals with WS or DS, and their data were compared with data collected from individuals with ASD and TD controls using identical measures (Booth & Happé, 2010; Happé & Booth, 2008; see Table 2 for tasks, organized by modality and level of processing). In our view, it is crucial to study all three neurodevelopmental disorders within a single approach, because tasks and stimuli are often susceptible to differences in methodology. Specifically, individuals with WS (so-called 'local processors') were compared to individuals with DS (so-called 'global processors') matched on Chronological Age (CA) and Mental Age (MA), while participants with ASD were compared to CA- and MA-matched TD controls. The groups were matched in this way because it is problematic to match across all four groups appropriately because those with ASD were high functioning. If participants with WS/ASD fail to show a local bias relative to participants with DS and TD controls on these tasks, then the characterization of WS/ASD as local processors will need to be reconceptualized.

Methods

Participants

Twenty-one individuals with WS and 31 individuals with DS gave informed consent to participate in the study. These participants had been clinically diagnosed and/or tested respectively for microdeletion of the ELN gene via *fluorescence in situ hybridization (FISH)* or for full trisomy 21. Individuals with known health problems other than WS and DS (e.g. hearing and visual problems) were not contacted. One participant with DS turned out to be non-verbal and could not complete any of the tasks. One participant with WS was also excluded from the final analyses because although his facial dysmorphology suggested WS, his FISH test subsequently turned out to be negative.⁴ Data collected from the participants with WS and DS were compared with unpublished data (Happé & Booth, 2008) collected from 32 individuals with ASD and 31 controls. ASD diagnosis had been confirmed by a psychiatrist or paediatrician according to DSM-IV criteria (APA, 1994). To control for the

⁴This participant had only four of the 28 genes deleted in the WS Critical Region (Karmiloff-Smith, Broadbent, Farran, Longhi, D'Souza *et al.*, 2012).

effects of gender, individuals with ASD and their carefully matched controls were all male. This is because considerably more males than females are diagnosed with ASD (Jacquemont, Coe, Hersch, Duyzend, Krumm *et al.*, 2014). Gender was sufficiently balanced in the DS (40% males) and WS groups (45% males); no association was found between Group (DS, WS) and Gender (male, female), $\chi^2(1) = 0.12, p = .776$.⁵ Five participants with high functioning autism had comorbidities, four of whom attention-deficit/hyperactivity disorder (ADHD) and one attention-deficit disorder (ADD). However, these five participants did not significantly differ from the other individuals with ASD in age or IQ, all $t(23) < 1.30, p > .20$, and their exclusion did not affect group results. They were therefore included in the analyses.

Participants' verbal and visuo-spatial mental ages were obtained for purposes of comparison. All participants with WS and DS were administered the British Picture Vocabulary Scales (BPVS; Dunn, Dunn & Whetton, 1982) and the Block Design test (Wechsler, 1992), which were selected for their high reliability and validity estimates (Cyr & Brooker, 1984). Participants with ASD and TD participants were administered the WISC-III/WAIS-III (Wechsler, 1992, 1997). From these measures, an overall Mental Age was obtained for each participant (age equivalent scores were calculated using raw scores). Table 3 shows the mean chronological ages (CA), mental age (MA), verbal mental ages (VMA), and visuo-spatial mental ages (VSMA) for each group, not including the two cases (one WS, one DS) referred to earlier who were removed from all analyses.

CA and MA data were normally distributed in all groups ($Z_{Skewness} < 2$; Kolmogorov-Smirnov [K-S], $p > .05$). However, VMA and VSMA data failed tests of normality in the DS group (K-S, $p < .05$; $Z_{Skewness} > 2$). Therefore, VMA and VSMA in the WS and DS groups were logarithmically transformed (base 10) for analyses.

As expected from the literature, VMA was better in WS than in DS, $t(48) = 3.61, p = .001, d = 1.04$, while VSMA was worse in WS than in DS, $t(39) = 2.06, p = .046, d = 0.66$. This uneven cognitive profile is common in these clinical populations. Nevertheless, independent t -tests failed to detect significant differences between the WS and DS groups on CA, $t(48) = 1.29, p = .205$, or MA, $t(39) = 1.42, p = .164$. Also, independent-samples Mann-Whitney U tests revealed that the females were not significantly different from the males on CA, MA, VMA, or VSMA, either in the WS group or in the DS group (all, $p > .169$).

The ASD and Control groups did not differ on CA, $t(61) = 0.01, p = .993$, MA, $t(61) = 0.16, p = .909$, VMA, $t(61) = 0.02, p = .985$, or VSMA, $t(61) = 0.20, p = .839$.

Design

Eight tasks were administered to participants in a set order, consisting of two blocks lasting about 30 minutes each. A short 5-minute break was taken between the two blocks in order to prevent fatigue. Four tasks were presented using SuperLab Pro 2.0 and an Apple MacBook Pro with a 17" screen.

⁵None of the expected frequencies was less than five.

Participants were only allowed to attempt the tasks after they had successfully demonstrated in the practice trials that they had understood what was required of them. Between test phases, positive comments were made in order to encourage participants. During test phases, however, feedback was not given on participant responses. The testing procedure was identical to the one employed by Happé and Booth (2008) for the individuals with ASD and the controls. Of the eight tasks used in the original (unpublished) study (see Happé & Booth, 2008), we selected four tasks (Table 2), one at each level of processing (high, low) in each of the two modalities (auditory/verbal, visuospatial), on the basis that they are arguably purer measures of local/global processing than the other tasks, with fewer or no demands on memory or general ability.

Task 1: Phoneme Segmentation

In this low-level (verbal modality) task, participants were instructed to detect the presence of a target phoneme (/p/) within a nonword. For each nonword, the target phoneme (e.g. /p/) was presented either as the initial sound (e.g. *plo*), or as the medial or final sound (e.g. *lipod*, *vip*), or not at all (e.g. *dowen*). Because individuals discriminate phonemes by segmenting syllables or words (Savin & Bever, 1970), the discrimination of a target phoneme in a nonword can be used as a measure of local processing in the auditory domain. It was predicted that participants with a local bias would be just as quick at identifying medial or final targets as targets in the initial position, because they would be less influenced by context.

Materials—Forty-five nonwords were used: 15 in each of the following conditions: (1) target phoneme (/p/) presented as initial sound; (2) target (/p/) presented as medial or final sound;⁶ and (3) target absent. One-, two-, and three-syllable nonwords were present in each condition in equal numbers. While real words would have provided a stronger gestalt, nonwords were used to eliminate familiarity and to control for individual differences in word knowledge. The stimuli were pre-recorded by a British English speaker and played to participants via high definition speakers in order to eliminate visual cues and to ensure that the presentation of sounds was consistent across participants.

Procedure—Participants were asked to listen for a /p/ sound that would occur in some of the words, and press the 'Yes' button whenever they heard that sound and 'No' when they did not. The stimuli were presented in a fixed random order. Participants had unlimited time to respond, and could have each test item repeated back to them (no more than once). Reaction time was measured from the offset of each nonword.

Task 2: Sentence Completion

In this higher-level (verbal) task, participants were required to complete sentence stems such as '*Little boys grow up to be men and ...*', '*In the sea, there are fish and ...*'. Completions might be global (e.g. *dads / whales*) or local (e.g. *women / chips*). Participants with a detail-

⁶In the pilot study, no significant differences were found between the *medial* and *final* conditions (Happé & Booth, 2008). Therefore, these two conditions were collapsed in the present analyses.

focused processing bias were expected to provide more local completions than those without this bias.

Materials—Fifteen sentence stems (from Happé, Briskman & Frith, 2001; Booth & Happé, 2010) were read to participants; ten produced a local–global conflict for their completion, five (‘filler’ sentence stems) did not. The filler sentence stems were added to mask the nature of the task and make it more naturalistic. These control items were interspersed with test items in a set order for all participants.

Procedure—Participants were read a sentence stem and asked to say something that would finish off the sentence. Response time was recorded from the offset of the sentence stem to the onset of the answer. The maximum time allowed for each completion was 20 seconds. Two measures were obtained: (i) number of local completions; (ii) mean response time.

Task 3: Navon Similarity Judgment

Hierarchical figures (Figure 1) can be described and analysed on two levels: local and global (Navon, 1977). The local forms that comprise the global form of the figure are processed less quickly and less accurately in TD individuals than the global form itself (Navon, 2003). It is also harder to ignore the global form than the local elements (Navon, 2003). These observations gave rise to the *global-precedence hypothesis*, which predicts ‘global advantage’ and a ‘global-to-local interference effect’ in TD individuals (Kimchi, 1992).

We avoided the drawing version of the Navon, so as to control for motor planning differences across groups. Rather, we used a two-alternative forced-choice (similarity-judgment) version of the classic Navon task, predicting that individuals with a general local bias (WS, ASD) would demonstrate a preference for the local elements over the global form. Individuals with a general global bias (DS, Controls) were predicted to show the opposite effect.

Materials—The stimuli were hierarchical figures, namely, global letters composed of local letters (Figure 1). Twelve stimuli types were employed: Ah (i.e. the letter ‘A’ composed of small ‘h’ letters), An, Fh, Ft, Ha, Hf, Hn, Ht, Na, Nh, Tf, and Th. The letters – A, F, H, N, T – were selected on the basis of their similarity with each other in terms of their physical complexity and frequency in the English language (Solso & King, 1976). Each stimulus type was 45×55 mm in size and subtended at 6.3° when viewed from a distance of 50 cm. Local elements were 4×5 mm in size and subtended at 0.6° . The test trials ($N = 36$) consisted of these 12 stimuli types, each presented at one of three different levels of density: 4-, 5-, or 6-local elements wide.

Procedure—Participants were seated approximately 50 cm away from the 17" screen on which the stimuli were presented. The participant was then administered two practice trials and 36 test trials. In each trial, a red cross on a white background would appear at the centre of the top half of the screen for 1000 ms, and subsequently disappear for 150 ms. A hierarchical figure (the *standard*) would then appear where the red cross had been, for 250 ms. The standard figure would then disappear and immediately be replaced by two hierarchical figures (the *comparison* figures), positioned side by side on the bottom half of

the screen (Figure 2). One of the comparison figures would have the same global form as the standard figure but different local elements, whereas the other would have the same local elements as the standard but a different global form. Participants had to point to the figure that most looked like the standard. The trial would end once the participant had responded or after 4 s had elapsed. If the participant did not respond within the given time, an error message and a loud sound were presented, and the trial was repeated once.

The test trials were presented in a fixed random order. The side of presentation was counterbalanced, with each comparison type (global match, local match) presented equally in the left and right positions. Across trials density was varied, but within each trial, the figures were of the same density.

Task 4: Fragmented Pictures

The ability to conceptually integrate visual elements into a meaningful whole was investigated using a version of the Fragmented Picture-Completion Task (Snodgrass, Smith, Feenan & Corwin, 1987), in which participants are presented with fragmented pictures which are gradually completed in sequential steps, and asked to name what each picture represents as soon as they think they know. Because success on this task requires global processing at a visual-semantic level, it is predicted that individuals with WS or ASD will find it more difficult to cohere the elements than those with DS.

Although both the Fragmented Pictures task and the Navon Familiarity-Judgment task would involve at least some input from higher-level processing areas, the former was judged to be a 'higher level' task than the latter because it requires participants to stay on task and match meaningful stimuli in memory (e.g. an elephant) to meaningless (fragmented) stimuli; hence, it involves a wide range of higher-level processes (sustained attention, working memory, long-term memory, conflict resolution, switching, decision making, etc.). This is irrespective of whether the task is more or less demanding than the mainly perceptual Navon task.

Materials—Ten sets of picture sequences (from Snodgrass and colleagues' 1987 Fragmentation Picture-Completion Task) were used. Each picture sequence consisted of eight different pictures of an image that varied in degree of fragmentation by random deletion of pixels – from the most fragmented to the complete image (Figure 3). These were presented (one after the other) at the centre of a 17" inch screen using SuperLab Pro software. The images were selected for their familiarity amongst children (Cycowicz, Friedman, Rothstein & Snodgrass, 1997). Images that could be identified by an isolated detail (e.g. an eye) were deselected.

Procedure—Each fragmented image was presented to the participant one after the other for 5 seconds at a time, from the most fragmented through to the completed picture (1–8), or until the participant guessed what the image was supposed to represent in the picture sequence. The participants were instructed to tell the instructor as soon as they thought they knew what the image was. If they were correct, the experimenter would end the trial (picture sequence) and begin the next one. If they were incorrect, the trial continued. Response times were recorded.

Results

Although a total of 113 individuals took part in these experiments, data from 10 participants that were collected using the laptop were lost (2 WS, 8 DS). For each task, two independent sets of analyses were carried out: WS vs. DS, and ASD vs. TD controls. This was because the WS and DS groups, and the ASD and TD groups, were closely matched, but the former two groups were significantly different from the latter two on CA and MA. If the data were non-normal (i.e. if $Z_{Skewness} > 2$, or Kolmogorov-Smirnov $p < .05$), then they were log transformed (base 10). If the transformed data were non-normal (e.g. bimodal), then the untransformed data were analysed using the appropriate non-parametric test (e.g. Mann-Whitney).

Task 1: Phoneme Segmentation

Seven participants with DS found it difficult to detect the phoneme and subsequently withdrew from the experiment. Of the remaining participants, the WS and DS groups did not differ on CA, $t(25) = 0.20$, $p = .844$, or on MA, $t(21) = 0.18$, $p = .857$; the ASD and Control groups also did not differ on CA, $t(59) = 0.05$, $p = .959$, or on MA, $t(59) = 0.40$, $p = .690$. Sensitivity and response bias were calculated by subjecting correct and incorrect detections of the target to signal detection analysis. Seven individuals (3 DS, 2 WS, 1 ASD, 1 Control) had A' values (the non-parametric index of sensitivity: Grier, 1971) calculated across conditions of between .22 and .60, too close to (or below) chance to be deemed reliable, and their data were removed from analysis. The remaining A' values ($N = 89$: 16 WS; 12 DS; 31 ASD; 30 Controls) ranged from .62 to .96 ($M = .84$, $SD = .07$).

Accuracy—None of the groups demonstrated a response bias, nor did they differ on a non-parametric measure of bias, B' (ASD vs. Control, $U = 491.00$, $z = 0.38$, $p = .707$;⁷ WS vs. DS, $U = 95.00$, $z = -0.05$, $p = .982$ ⁸). Table 4 presents the mean number of correct judgments for each of the three conditions (initial, medial/final, absent) and the signal detection parameter of sensitivity (A') for each group.

The initial and medial/final data were non-normal in the ASD and Control groups, and subsequently log transformed. A 2 (Group: ASD, Control) $\times$ 2 (Position: Initial, Medial/Final) ANOVA revealed a main effect of Position, $F_{1,53} = 101.96$, $p < .000001$, $\eta^2 = .66$. No main effect of Group, $F_{1,53} = 2.17$, $p = .147$, and no interaction effect, $F_{1,53} = 0.46$, $p = .502$, were found. In other words, both groups were significantly more likely to identify correctly the target phoneme in the initial than medial/final condition, but individuals with ASD and controls did not differ on the number of correct judgments made.

Initial and medial/final data were also non-normal in the WS group. The data were therefore log transformed. A 2 (Group: WS, DS) $\times$ 2 (Position: Initial, Medial/Final) ANOVA revealed a main effect of Position, $F_{1,24} = 19.10$, $p = .0002$, $\eta^2 = .44$. No main effect of Group, $F_{1,24} = 0.52$, $p = .479$, and no interaction effect, $F_{1,24} = 0.01$, $p = .913$, were found. In other words, both groups (like the ASD and Control groups) were significantly more

⁷An independent t -test was also carried out: $t(59) = 0.23$, $p = .818$.

⁸An independent t -test was also carried out: $t(26) = 0.20$, $p = .840$.

likely to correctly identify the target phoneme in the initial than medial/final condition, but individuals with WS and DS did not differ on the number of correct judgments made.

However, a significant difference on A' was found between the ASD and Control groups, $U = 583.50$, $z = 2.28$, $p = .023$, $r = .30$, and between WS and DS groups, $U = 33.00$, $z = -2.72$, $p = .006$, $r = -.52$. This indicates that the ASD and DS groups were less able to discriminate between target absent and target present trials than the Control and WS groups, respectively.

Response time: ASD vs. Controls—A 2×2 mixed ANOVA with group (ASD, Control) as the between-subjects factor, and phoneme position (initial, medial/final) as the within-subject factor was used to examine the effect of phoneme position on response times. Response times were capped at 10 seconds (less than 2% of all cases were longer than 10 s) and logarithmically transformed to reduce positive skew. Three outliers (> 3.29 SD) were removed. There were no significant differences between the individuals with ASD and Controls on CA, $t(57) = 0.04$, $p = .971$, or MA, $t(57) = 0.03$, $p = .978$.

There was no main effect of group, $F_{1,57} = 0.06$, $p = .801$, which indicates that response times were similar across groups. There was a main effect of phoneme position, $F_{1,57} = 9.52$, $p = .003$, $\eta^2 = .14$, with participants taking longer to detect the target phoneme when it was presented in the medial/final position than in the initial position. There was also an interaction effect between group and phoneme position, $F_{1,57} = 6.31$, $p = .015$, $\eta^2 = .10$.

Paired samples t -tests were used to elucidate the interaction effect. The controls were significantly faster at detecting target phonemes when they were presented in the initial position than when in the medial or final positions, $t(28) = 5.20$, $p = .00002$, $d = 1.97$. However, the position of the phoneme did not differentiate the timing of detection in the ASD group, $t(29) = 0.35$, $p = .733$ (Figure 4).

Response time: WS vs. DS—A 2×2 mixed ANOVA with group (WS, DS) as the between-subjects factor, and phoneme position (initial, medial/final) as the within-subject factor, was used to examine the effect of phoneme position on response times. Response times were capped at 10 seconds (less than 2% of all cases were longer than 10 s) and logarithmically transformed to reduce positive skew. Three outliers (> 3.29 SD) were removed. The two groups did not significantly differ from each other on verbal MA, $t(25) = 1.73$, $p = .095$.

There was a main effect of Group, $F_{1,25} = 4.86$, $p = .037$, $\eta^2 = .16$. There was a main effect of position, $F_{1,25} = 6.71$, $p = .016$, $\eta^2 = .21$, and an interaction effect between group and phoneme position, $F_{1,25} = 4.89$, $p = .036$, $\eta^2 = .16$.

Paired samples t -tests were used to elucidate the interaction effect. Individuals with WS were significantly faster at detecting target phonemes when they were presented in the initial position than when in the medial or final positions, $t(15) = 3.96$, $p = .001$, $r = .71$. Notice that the effect in the WS group ($r = .71$) is similar in size to the one in the Control group ($r = .70$). However, the position of the phoneme did not differentiate the timing of detection in the DS group, $t(10) = 0.23$, $p = .823$ (Figure 5).

In sum, individuals with autism did not differ from controls on accuracy or response time, which suggests that they did not have a hierarchical deficit. In addition, context (viz. phoneme position) had no effect on reaction times in the ASD group (1.32 s vs. 1.36 s), whereas it clearly did with WS and Controls. Context also had no significant effect on reaction times in the DS group (2.59 s vs. 2.86 s). This may have been because they did not answer immediately as instructed. Indeed, they took twice as long to provide an answer and their reaction times were 2–3 times more variable. This may have attenuated the trend to respond more quickly to the target in the medial/final position than in the initial position. It is also important to note that data from only 12 participants with DS were analysed; seven found the task too difficult to complete.

Task 2: Sentence Completion

Individuals (2 WS; 2 DS; 2 ASD; 4 Controls) whose local completion score or mean response time (RT) was two standard deviations above or below the group mean were identified as outliers and removed from the analyses. The WS and DS groups did not significantly differ on CA, $t(29.15) = 1.12$, $p = .271$ (equal variances not assumed),⁹ or on MA, $t(37) = 1.55$, $p = .130$ (MA data were marginally non-normal in the DS group and thus MA data in both groups were log transformed). The ASD and Control groups did not significantly differ on CA, $t(61) = 0.01$, $p = .993$, or on MA, $t(61) = 0.06$, $p = .950$.

The local completions data failed tests of normality in the ASD and Control groups, but not in the WS or DS groups. Therefore, the local completion data were log transformed for the ASD vs. Control comparison. All groups produced more global completions than local completions ($p < .001$) (Table 5).

Individuals with ASD made significantly more local completions than controls, $t(50.11)^{10} = 3.15$, $p = .003$, $d = 0.89$. They also took longer to complete the sentence stems than controls, but this did not reach significance, $t(55) = 1.73$, $p = .089$.

Individuals with WS did not significantly differ from those with DS on local completions or mean RT, $t(39) = 0.14$, $p = .893$, $t(39) = 1.20$, $p = .238$.

Task 3: Navon Similarity Judgment

Some of the participants (10 WS, 10 DS) found the task too difficult to complete. Four outliers ($\pm 2 SD$) were also removed (1 WS, 1 DS, 2 Controls). The individuals with WS ($N = 7$) did not significantly differ from those with DS ($N = 10$) on CA, $t(17) = 0.63$, $p = .537$, or MA, $t(13) = 0.24$, $p = .817$. Also, those with ASD ($N = 31$) did not significantly differ from the controls ($N = 29$) on CA, $t(60) < 0.01$, $p = .998$, or MA, $t(60) = 0.12$, $p = .906$.

The proportional response data were skewed ($z = 1.86$, -1.02 ; $D(31) = 0.22$, $p = .001$) in the ASD group, but not in the Control group ($z < 1.96$, 1.96 ; K-S, $p = .066$). The data were therefore arcsine transformed. Individuals with ASD made significantly more local matches

⁹Levene's test, $F = 6.30$, $p = .016$. When equal variances are assumed: $t(42) = 1.19$, $p = .242$.

¹⁰The data failed Levene's test for equality of variance ($F = 4.88$, $p = .031$). Therefore equal variances were not assumed.

than controls, $t(36.98)^{11} = 2.89$, $p = .006$, $d = 0.95$. WS did not significantly differ from DS, $t(15) = 1.22$, $p = .242$.

All groups, except the DS group, made more global than local matches. In the DS group, individuals made more local matches ($M = 18.20$, $SD = 4.76$) than global matches ($M = 17.80$, $SD = 4.76$). One-sampled t -tests showed that the proportion did not significantly differ from 0.5 in the DS group, $t(9) = 0.13$, $p = .897$, but it did in the WS group, $t(6) = 2.79$, $p = .032$. In other words, the DS group made as many local matches as they did global matches (Figure 6) but, more importantly, individuals with DS made fewer global *relative to local* matches than those with WS.

Task 4: Fragmented Pictures

Seven participants with WS found the task too difficult to complete and withdrew from the experiment. The remaining WS and the DS groups did not differ on CA, $t(29) = 0.29$, $p = .778$, or MA, $t(22) = 1.09$, $p = .288$. The ASD and Control groups also did not differ on CA, $t(60) < 0.01$, $p = .998$, or MA, $t(60) = 0.12$, $p = .906$.

Total number of incorrect guesses—Six individuals (1 WS, 1 DS, 2 ASD, 2 TD) were identified as outliers (each made 5–18 incorrect guesses per item, which were two standard deviations above the group mean) and removed from the following analysis. The data were positively skewed in all four groups, and therefore log transformed. However, numbers of incorrect guesses did not significantly differ between the ASD and Control groups, $t(41) = 1.35$, $p = .184$, or between the WS and DS groups, $t(13.58) = 0.43$, $p = .671$ (Table 6).

Mean frame number for correct identification—One control was identified as an outlier (2.15 SD above the group mean) and removed from the following analysis. Contrary to expectations, the mean frame number for correct identification in the ASD group was not significantly different from the mean frame number in the control group, $t(59) = 1.61$, $p = .113$. However, it was significantly lower in the WS group than in the DS group, $t(30) = 3.54$, $p = .001$, $d = 1.29$.

Mean response time for correct identification—One individual with ASD was identified as an outlier (2.30 SD above the group mean) and removed from the following analysis. The ASD and Control groups did not significantly differ on response time for correct identification, $t(59) = 1.59$, $p = .117$.¹² Nor did individuals with WS take longer to identify a picture than DS, $t(30) = 0.26$, $p = .795$.

Total number of items identified at the 6th, 7th, or 8th frame—Participants were also compared on the number of pictures that were recognized only near their most complete form (i.e. either at the 6th, 7th, or 8th frame). One control was identified as an outlier (2.15 SD above the group mean) and removed from the following analysis. The number of pictures correctly identified at the 6th, 7th, or 8th frame was significantly higher in the ASD

¹¹Levene's, $F = 23.80$, $p < .001$, so equal variances were not assumed.

¹²However, if the outlier were included, then RT would – as predicted – be slower in the ASD than in the Control group, $t(60) = 1.86$, $p = .034$ (one-tailed).

group than in the control group, $t(59) = 2.49$, $p = .015$, $d = .65$. It is important to note that no control failed to identify a picture in its complete form (i.e. at the 8th frame).

The data in the WS and DS groups were negatively skewed. They were therefore reverse log transformed. The number of pictures correctly identified at the 6th, 7th, or 8th frame (i.e. *late*) was also significantly higher in the DS group than in the WS group, $t(29) = 4.01$, $p = .0004$, $d = 1.49$.

Discussion

The purpose of this study was to elucidate local/global processing by ascertaining whether more in-depth research across modalities and levels of processing could reveal differences or commonalities in integrative processing across the three neurodevelopmental disorders (WS, DS, ASD).

First and foremost, our study illustrates the importance of cross-syndrome comparisons, because had each neurodevelopmental group merely been compared with TD controls, we might have concluded that they could be characterized as showing either a local or global bias, as the literature has consistently maintained. However, a comparison between the disorders yields a far more complex picture (Table 7).

First, despite purported similarities in local/global processing in ASD and WS, neither (but especially not the WS group) demonstrated a general local bias. In a similar vein, a preference for global over local processing did not always materialize in the DS population; in some tasks, the opposite was true. For example, in the Phoneme Segmentation task, context (phoneme position) had no effect on DS/ASD whereas, like TD controls, it did on WS. This demonstrates that the local bias was not predominant in WS as far as an auditory task is concerned. By contrast, in the Fragmented Pictures task, individuals with ASD performed similarly to controls, while those with DS found it significantly more difficult than individuals with WS to integrate the fragmented visual elements into meaningful wholes. Furthermore, in the Sentence Completion task, those with DS displayed an equally high number of local completions as those with WS. This could have been because, despite the fact that the sentences were deliberately short, individuals with DS have poor verbal working memory so found it easier to maintain the local context in working memory than the global context. Yet in the Navon Similarity Judgment task, participants with ASD showed a reduced preference for global-level information compared with controls as expected, but those with DS demonstrated neither a global nor a local bias.

According to theory, a neat dissociation should have emerged, with individuals with WS/ASD showing a local bias on all four tasks, and individuals with DS displaying a global bias. But despite claims in the literature, neither WS nor ASD were consistently 'local processors' in their responses. No clear links could be identified between the participants' task performances and the cognitive atypicalities purported to underpin them. Nonetheless, individuals with ASD showed a local bias in both of the low-level tasks (Phoneme Segmentation, Navon), which hints that low-level attentional/perceptual tasks are more likely to capture local processing in ASD than higher-level tasks.

On the EPF account, individuals with ASD are expected to outperform the TD controls on low-level tasks. But in the Phoneme Segmentation task, the ASD group was less able to discriminate between target absent and target present trials than the controls.

Furthermore, the HDH predicts that individuals with a hierarchical deficit (namely, WS and ASD) should find tasks that require flexibility in shifting from one level to the next (e.g. the Phoneme Segmentation and Fragmented Pictures tasks) to be relatively difficult. Yet individuals with ASD showed no difficulty in detecting target phonemes in the Phoneme Segmentation task, and the WS group outperformed the DS group in the Fragmented Pictures task.

It should be noted that a large proportion of individuals with WS or DS withdrew from the Navon Familiarity-Judgment task. Although all the tasks were designed to be as easy to complete as possible, some participants clearly found that the stimuli in the Navon task were presented too quickly for them to decide which figure they thought was most like the target (the task was designed to be quick in order to capture participants' 'instinctive' preferences rather than higher-level explicit decision making). Thus it is possible that only (relatively) high functioning individuals completed this task, and that the results may have differed had low functioning individuals also been able to complete it.

Furthermore, a large proportion of individuals with DS could not complete the Phoneme Segmentation task (even though the nonwords were well articulated). This finding may reflect the difficulties that individuals with DS are reported to have in the auditory domain (e.g. Pueschel, Gallagher, Zartler & Pezzullo, 1987). Nevertheless, the remaining participants were not at floor. They were more likely than not to correctly identify the target phoneme, though phoneme position (context) had no significant effect.

In sum, both differences and commonalities emerged across the three neurodevelopmental disorders. Our findings thus indicate that individuals with WS and ASD should not be categorized as 'local processors'. It is at best over-simplistic; at worst, false. Moreover, it is likely that the emphasis on local *versus* global levels of information is actually a false dichotomy. Depending on task and stimulus conditions, individuals attend to either the parts, to the whole pattern, or to both (Dukette & Stiles, 1996, 2001; Tada & Stiles-Davis, 1989). It is possible that the differences between the groups have more to do with the quality of the participants' internal representations and the available operations that can be performed on them (Aslin & Smith, 1988). The ability to segment and cohere local-global information will be affected more by the nature and difficulty of the task than by any general local/global processing bias.

Interestingly, although individuals with ASD have difficulty switching from local to global processing, as do those with WS, those with ASD do not have difficulty switching from global to local (White, O'Reilly & Frith, 2009). Indeed, different computational processes are likely to underpin the local/global asymmetry found in these neurodevelopmental disorders. The local and global structures of a stimulus are defined by many factors and by the context in which the stimulus is situated. Moreover, differences in the ability to process the spatial frequency of stimuli may actually account for perceptual differences across

neurodevelopmental disorders. Indeed, there is evidence to suggest that individuals with DS have problems with processing high frequency spatial visual information, whereas those with WS have difficulty with processing low frequency spatial visual information (e.g. Leonard, Annaz, Karmiloff-Smith & Johnson, 2011). One advantage that a Spatial Frequency hypothesis has over theories discussed earlier is that it focuses on the stimulus, i.e. the source of information in the task, rather than difficult-to-define characteristics such as local/global processing. Another advantage of the Spatial Frequency hypothesis is that it is based on how the visual system actually functions. There is also a small but growing body of evidence that the hypothesis invoking variations in Spatial Frequency may also apply to other perceptual systems (e.g. auditory/speech perception; Ivry & Lebel, 1993), as well as higher-level processes (e.g. categorical and coordinate spatial relationships in memory; Kosslyn, 1987; Kosslyn, Koenig, Barret, Cave, Tang *et al.*, 1989). This could explain why individuals with DS performed so poorly (relative to those with WS) in the Fragmented Pictures task. It may also be useful for understanding another domain – face processing – that has been a topic of debate about local versus global processing, particularly regarding Williams syndrome (Deruelle, Mancini, Livet, Casse-Perrot & De Schonen, 1999; Deruelle, Rondan, Mancini & Livet, 2003; Karmiloff-Smith, Thomas, Annaz, Humphreys, Ewing *et al.*, 2004; Tager-Flusberg, Plesa-Skwerer, Faja & Joseph, 2003).

In conclusion, performances on tasks ‘that tap local/global processing’ have been claimed by many (but not all, Tager-Flusberg *et al.*, 2003) to reflect a local processing bias in WS and ASD, and a global processing bias in DS. However, our data do not support this hypothesis. Indeed, both a local and a global processing bias is apparent for some tasks and some stimuli in individuals across all three neurodevelopmental disorders, i.e. WS, DS, and ASD. Thus, individuals with these neurodevelopmental disorders cannot simply be characterized as having a local or global processing style. Rather, they can all process both local and global information, depending on the task, but they do so in different and atypical ways. We thus conclude that the use of ‘local’ or ‘global’ processors for explaining neurodevelopmental disorders needs to be reconceptualized and that a cross-syndrome/cross-task design is useful in identifying more subtle similarities and differences across such disorders than merely a comparison with TD controls.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We are very grateful for the contributions families have made towards this study. The study was funded by the Wellcome Trust, Economic and Social Research Council, and Williams Syndrome Foundation. The recruiting of participants was supported by the Williams Syndrome Foundation, the Down Syndrome Association, and Down Syndrome Education International.

References

American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4. Author; Washington, DC: 1994.

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4. Author; Washington, DC: 2013.
- Aslin RN, Smith LB. Perceptual development. *Annual Review of Psychology*. 1988; 39(1):435–473.
- Atkinson AP. Impaired recognition of emotions from body movements is associated with elevated motion coherence thresholds in autism spectrum disorders. *Neuropsychologia*. 2009; 47:3023–3029. [PubMed: 19500604]
- Bellugi U, Bihle A, Jeringan T, Trauner D, Doherty S. Neuropsychological, neurological, and neuroanatomical profile of Williams syndrome. *American Journal of Medical Genetics*. 1990; 6(Suppl.):115–125. [PubMed: 2144426]
- Bellugi U, Lichtenberger L, Jones W, Lai Z, St George M. The neurocognitive profile of Williams syndrome: a complex pattern of strengths and weaknesses. *Journal of Cognitive Neuroscience*. 2000; 12(Suppl. 1):7–29. [PubMed: 10953231]
- Bellugi U, Lichtenberger L, Mills D, Galaburda A, Korenberg J. Bridging cognition, brain, and molecular genetics: evidence from Williams syndrome. *Trends in Neurosciences*. 1999; 22:197–207. [PubMed: 10322491]
- Bellugi, U.; Marks, S.; Bihle, A.; Sabo, H. Dissociation between language and cognitive functions in Williams syndrome. In: Bishop, D.; Mogford, K., editors. *Language development in exceptional circumstances*. Psychology Press; Hove: 1988. p. 177-189.
- Bellugi, UL.; Sabo, H.; Vaid, J. Spatial deficits in children with Williams syndrome. In: Stile, J.; Davis, M.; Bellugi, U., editors. *Spatial cognition: Brain bases and development*. Lawrence Erlbaum Associates; Hillsdale, NJ: 1988. p. 273-297.
- Bertone A, Mottron L, Jelenic P, Faubert J. Motion perception in autism: a 'complex' issue. *Journal of Cognitive Neuroscience*. 2003; 25:218–225. [PubMed: 12676059]
- Bertone A, Mottron L, Jelenic P, Faubert J. Enhanced and diminished visuo-spatial information processing in Autism depends on stimulus complexity. *Brain*. 2005; 128:2430–2441. [PubMed: 15958508]
- Bihle AM, Bellugi U, Delis DC, Marks S. Seeing either the forest or the trees: dissociation in visuospatial processing. *Brain and Cognition*. 1989; 11(1):37–49. [PubMed: 2528973]
- Booth R, Happé F. 'Hunting with a knife and ... fork': examining central coherence in autism, attention deficit/hyperactivity disorder, and typical development with a linguistic task. *Journal of Experimental Child Psychology*. 2010; 107:377–393. [PubMed: 20655060]
- Bouvet L, Simard-Meilleur AA, Paignon A, Mottron L, Donnadieu S. Auditory local bias and reduced global interference in autism. *Cognition*. 2014; 131(3):367–372. [PubMed: 24637103]
- Braddick O, Atkinson J, Wattam-Bell J. Normal and anomalous development of visual motion processing: motion coherence and 'dorsal-stream vulnerability'. *Neuropsychologia*. 2003; 41(13):1769–1784. [PubMed: 14527540]
- Brosnan MJ, Scott FJ, Fox S, Pye J. Gestalt processing in autism: failure to process relationships and implications for contextual understanding. *Journal of Child Psychology and Psychiatry*. 2004; 45(3):459–469. [PubMed: 15055366]
- Cycowicz YM, Friedman D, Rothstein M, Snodgrass JG. Picture naming by young children: norms for name agreement, familiarity, and visual complexity. *Journal of Experimental Child Psychology*. 1997; 65(2):171–237. [PubMed: 9169209]
- Cyr J, Brooker BH. Use of appropriate formulas for selecting WAIS-R short forms. *Journal of Consulting and Clinical Psychology*. 1984; 52(5):903–905.
- Deruelle C, Mancini J, Livet MO, Casse-Perrot C, De Schonen S. Configural and local processing of faces in children with Williams syndrome. *Brain and Cognition*. 1999; 41(3):276–298. [PubMed: 10585239]
- Deruelle C, Rondan C, Mancini J, Livet M. Exploring face processing in Williams syndrome. *Cognition, Creier, Comportament*. 2003; 7:157–171.
- Deruelle C, Schon D, Rondan C, Mancini J. Global and local music perception in children with Williams syndrome. *NeuroReport*. 2005; 25:631–634. [PubMed: 15812322]
- Donnai D, Karmiloff-Smith A. Williams syndrome: from genotype through to the cognitive phenotype. *American Journal of Medical Genetics: Seminars in Medical Genetics*. 2000; 97(2):164–171. [PubMed: 11180224]

- Dukette D, Stiles J. Children's analysis of hierarchical patterns: evidence from a similarity judgment task. *Journal of Experimental Child Psychology*. 1996; 63(1):103–140. [PubMed: 8812036]
- Dukette D, Stiles J. The effects of stimulus density on children's analysis of hierarchical patterns. *Developmental Science*. 2001; 4:233–251.
- Dunn, LM.; Dunn, LM.; Whetton, C. *British Picture Vocabulary Scale*. Nfer-Nelson; Windsor: 1982.
- Elsabbagh M, Cohen H, Karmiloff-Smith A. Discovering structure in auditory input: evidence from Williams syndrome. *American Journal of Intellectual and Developmental Disability*. 2010; 115(2): 128–139.
- Farran EK, Jarrold C, Gathercole SE. Divided attention, selective attention and drawing: processing preferences in Williams syndrome are dependent on the task administered. *Neuropsychologia*. 2003; 41:676–687. [PubMed: 12591025]
- Farroni T, Valenza E, Simion F, Umiltà C. Configural processing at birth: evidence for perceptual organisation. *Perception*. 2000; 29(3):355–372. [PubMed: 10889944]
- Frick JE, Colombo J, Allen JR. Temporal sequence of global-local processing in 3-month-old infants. *Infancy*. 2000; 1(3):375–386.
- Frith, U. *Autism: Explaining the enigma*. Blackwell Publishing; Oxford: 1989.
- Frith, U. *Autism: Explaining the enigma*. 2. Blackwell Publishing; Oxford: 2003.
- Frith U, Happé F. Autism: beyond 'theory of mind'. *Cognition*. 1994; 50(1-3):115–132. [PubMed: 8039356]
- Ghim HR, Eimas PD. Global and local processing by 3- and 4-month-old infants. *Perception & Psychophysics*. 1988; 43(2):165–171. [PubMed: 3340515]
- Grier J. Nonparametric indexes for sensitivity and bias: computing formulas. *Psychological Bulletin*. 1971; 75(6):424–429. [PubMed: 5580548]
- Hadad BS, Kimchi R. Developmental trends in utilizing perceptual closure for grouping of shape: effects of spatial proximity and collinearity. *Perception & Psychophysics*. 2006; 68(8):1264–1273. [PubMed: 17378413]
- Happé F. Autism: cognitive deficit or cognitive style? *Trends in Cognitive Sciences*. 1999; 3(6):216–222. [PubMed: 10354574]
- Happé F, Booth R. The power of the positive: revisiting weak coherence in autism spectrum disorders. *Quarterly Journal of Experimental Psychology*. 2008; 61:50–63.
- Happé F, Briskman J, Frith U. Exploring the cognitive phenotype of autism: weak 'central coherence' in parents and siblings of children with autism: I. Experimental test. *Journal of Child Psychology and Psychiatry*. 2001; 42:299–307. [PubMed: 11321199]
- Happé F, Frith U. The weak coherence account: detailed-focused cognitive style in autism spectrum disorders. *Journal of Autism and Developmental Disorders*. 2006; 36:5–25. [PubMed: 16450045]
- Heinze HJ, Hinrichs H, Scholz M, Burchert W, Mangun GR. Neural mechanisms of global and local processing: a combined PET and ERP study. *Journal of Cognitive Neuroscience*. 1998; 10(4):485–498. [PubMed: 9712678]
- Hermelin, B.; O'Connor, N. *Psychological experiments with autistic children*. Pergamon; Oxford: 1970.
- Ivry RB, Leiby PC. Hemispheric differences in auditory perception are similar to those found in visual perception. *Psychological Science*. 1993; 4:41–45.
- Jacquemont S, Coe BP, Hersch M, Duyzend MH, Krumm N, et al. A higher mutational burden in females supports a 'female protective model' in neurodevelopmental disorders. *American Journal of Human Genetics*. 2014; 94(3):415–425. [PubMed: 24581740]
- Jarrold C, Baddeley AD, Phillips CE. Verbal short-term memory in Down syndrome: a problem of memory, audition, or speech? *Journal of Speech, Language, and Hearing Research*. 2002; 45:531–544.
- Jarrold C, Russell J. Counting abilities in autism: possible implications for central coherence theory. *Journal of Autism and Developmental Disorders*. 1997; 27:25–37. [PubMed: 9018580]
- Johnson SP, Aslin RN. Perception of object unity in 2-month-old infants. *Developmental Psychology*. 1995; 31(5):739–745.

- Jolliffe T, Baron-Cohen S. Are people with autism and Asperger syndrome faster than normal on the Embedded Figures Test? *Journal of Child Psychology and Psychiatry*. 1997; 38(5):527–534. [PubMed: 9255696]
- Karmiloff-Smith A, Broadbent H, Farran EK, Longhi E, D'Souza D, et al. Social cognition in Williams syndrome: genotype/phenotype insights from partial deletion patients. *Frontiers in Psychology*. 2012; 3(168):1–8. [PubMed: 22279440]
- Karmiloff-Smith A, D'Souza D, Dekker TM, Van Herwegen J, Xu F, et al. Genetic and environmental vulnerabilities in children with neurodevelopmental disorders. *Proceedings of the National Academy of Sciences, USA*. 2012; 109(Supplement 2):17261–17265.
- Karmiloff-Smith A, Thomas M, Annaz D, Humphreys K, Ewing S, et al. Exploring the Williams syndrome face processing debate: the importance of building developmental trajectories. *Journal of Child Psychology and Psychiatry*. 2004; 45:1258–1274. [PubMed: 15335346]
- Kéita L, Guy J, Berthiaume C, Mottron L, Bertone A. An early origin for detailed perception in Autism Spectrum Disorder: biased sensitivity for high-spatial frequency information. *Scientific Reports*. 2014;4.
- Kellman PJ, Spelke ES, Short KR. Infant perception of object unity from translatory motion in depth and vertical translation. *Child Development*. 1986; 57(1):72–86. [PubMed: 3948595]
- Kimchi R. Primacy of wholistic processing and global/local paradigm: a critical review. *Psychological Bulletin*. 1992; 112(1):24–38. [PubMed: 1529037]
- Kimchi R, Hadad B, Behrmann M, Palmer S. Microgenesis and ontogenesis of perceptual organization. *Psychological Science*. 2005; 16:282–290. [PubMed: 15828975]
- Koldewyn K, Whitney D, Rivera SM. Neural correlates of coherent and biological motion perception in autism. *Developmental Science*. 2011; 14(5):1075–1088. [PubMed: 21884323]
- Kosslyn SM. Seeing and imagining in the cerebral hemispheres: a computational approach. *Psychological Review*. 1987; 94:148–175. [PubMed: 3575583]
- Kosslyn SM, Koenig O, Barret A, Cave CB, Tang J, et al. Evidence for two types of spatial representations: hemispheric specialization for categorical and coordinate relations. *Journal of Experimental Psychology, Human Perception and Performance*. 1989; 15:723–735. [PubMed: 2531207]
- Kovács I. Human development of perceptual organization. *Vision Research*. 2000; 40(10):1301–1310. [PubMed: 10788641]
- Kovacs I, Kozma P, Feher A, Benedek G. Late maturation of visual spatial integration in humans. *Proceedings of the National Academy of Sciences, USA*. 1999; 96(21):12204–12209.
- Kozma P, Kovács I, Benedek G. Normal and abnormal development of visual functions in children. *Acta Biologica Szegediensis*. 2001; 45(1-4):23–42.
- Laws G, Gunn D. Phonological memory as a predictor of language comprehension in Down syndrome: a five-year follow-up study. *Journal of Child Psychology and Psychiatry*. 2004; 45(2):326–337. [PubMed: 14982246]
- Leonard H, Annaz D, Karmiloff-Smith A, Johnson MH. Developing spatial frequency biases for face recognition in Autism and Williams syndrome. *Journal of Autism and Developmental Disorders*. 2011; 41(7):968–973. [PubMed: 20945155]
- Meyer-Lindenberg A, Kohn P, Mervis CB, Kippenhan JS, Olsen RK, et al. Neural basis of genetically determined visuospatial construction deficit in Williams syndrome. *Neuron*. 2004; 43(5):623–631. [PubMed: 15339645]
- Milne E, Swettenham J, Hansen P, Campbell R, Jeffries H, et al. High motion coherence thresholds in children with autism. *Journal of Child Psychology and Psychiatry*. 2002; 43(2):255–263. [PubMed: 11902604]
- Minshew NJ, Goldstein G. Is autism an amnesic disorder? Evidence from the California Verbal Learning Test. *Neuropsychology*. 1993; 7:209–216.
- Minshew NJ, Goldstein G. Autism as a disorder of complex information processing. *Mental Retardation and Developmental Disabilities Research Reviews*. 1998; 4:129–136.
- Minshew NJ, Goldstein G, Siegel DJ. Speech and language in high functioning autistic individuals. *Neuropsychology*. 1995; 9:255–261.

- Minshew, NJ.; Sweeney, JA.; Bauman, ML. Neurological aspects of autism. In: Cohen, DJ.; Volkmar, FR., editors. *Handbook of autism and pervasive developmental disorders*. 2. John Wiley & Sons; New York: 1997. p. 344
- Mobbs D, Eckert MA, Menon V, Mills D, Korenberg J, et al. Reduced parietal and visual cortical activation during global processing in Williams syndrome. *Developmental Medicine & Child Neurology*. 2007; 49(6):433–438. [PubMed: 17518929]
- Mottron L, Belleville S. A study of perceptual analysis in a high-level autistic subject with exceptional graphic abilities. *Brain and Cognition*. 1993; 23(2):279–309. [PubMed: 8292330]
- Mottron L, Belleville S, Ménard E. Local bias in autistic subjects as evidenced by graphic tasks: perceptual hierarchisation or working memory deficit? *Journal of Child Psychology and Psychiatry*. 1999; 40(5):743–755. [PubMed: 10433408]
- Mottron, L.; Burack, JA. Enhanced perceptual functioning in the development of autism. In: Burack, JA.; Charman, T.; Yirmiya, N.; Zelazo, PR., editors. *The development of autism: Perspectives from theory and research*. Lawrence Erlbaum Associates; Mahwah, NJ: 2001. p. 131-148.
- Mottron L, Dawson M, Soulières I, Hubert B, Burack J. Enhanced perceptual functioning in Autism: an update, and eight principles of autistic perception. *Journal of Autism and Developmental Disorders*. 2006; 36(1):27–43. [PubMed: 16453071]
- Muth A, Hönekopp J, Falter CM. Visuo-spatial performance in autism: a meta-analysis. *Journal of Autism and Developmental Disorders*. 2014; 44(12):3245–3263. [PubMed: 25022252]
- Navon D. Forest before trees: the precedence of global features in visual perception. *Cognitive Psychology*. 1977; 9(3):353–383.
- Navon D. What does a compound letter tell the psychologist's mind? *Acta Psychologica*. 2003; 114(3): 273–309. [PubMed: 14670701]
- Olu-Lafe O, Liederman J, Tager-Flusberg H. Is the ability to integrate parts into wholes affected in autism spectrum disorder? *Journal of Autism and Developmental Disorders*. 2014; 44:2652–2660. [PubMed: 24737003]
- Ozonoff S, Strayer DL, McMahon WM, Filloux F. Executive function abilities in autism and Tourette syndrome: an information processing approach. *Journal of Child Psychology and Psychiatry*. 1994; 35(6):1015–1032. [PubMed: 7995842]
- Pani JR, Mervis CB, Robinson BF. Global spatial organization by individuals with Williams syndrome. *Psychological Science*. 1999; 10:453–458.
- Pellicano E, Gibson L, Maybery M, Durkin K, Badcock DR. Abnormal global processing along the dorsal visual pathway in autism: a possible mechanism for weak visuospatial coherence? *Neuropsychologia*. 2005; 43(7):1044–1053. [PubMed: 15769490]
- Plaisted KC, Swettenham J, Rees L. Children with autism show local precedence in a divided attention task and global precedence in a selective attention task. *Journal of Child Psychology and Psychiatry*. 1999; 40(5):733–742. [PubMed: 10433407]
- Poher BR. Williams-Beuren syndrome. *New England Journal of Medicine*. 2010; 362:239–252. [PubMed: 20089974]
- Poirel N, Mellet E, Houdé O, Pineau A. First came the trees, then the forest: developmental changes during childhood in the processing of visual local-global patterns according to the meaningfulness of the stimuli. *Developmental Psychology*. 2008; 44:245–253. [PubMed: 18194023]
- Poirel N, Simon G, Cassotti M, Leroux G, Percey G, et al. The shift from local to global visual processing in 6-year-old children is associated with grey matter loss. *PLoS ONE*. 2011; 6(6):e20879. [PubMed: 21687636]
- Porter MA, Coltheart M. Global and local processing in Williams syndrome, Autism, and Down syndrome: perception, attention, and construction. *Developmental Neuropsychology*. 2006; 30(3): 771–789. [PubMed: 17083292]
- Pueschel SM, Gallagher PL, Zartler AS, Pezzullo JC. Cognitive and learning processes in children with Down syndrome. *Research in Developmental Disabilities*. 1987; 8(1):21–37. [PubMed: 2958898]
- Quinn PC, Bhatt RS. Are some gestalt principles deployed more readily than others during early development? The case of lightness versus form similarity. *Journal of Experimental Psychology: Human Perception and Performance*. 2006; 32(5):1221. [PubMed: 17002533]

- Quinn PC, Bhatt RS, Brush D, Grimes A, Sharpnack H. Development of form similarity as a Gestalt grouping principle in infancy. *Psychological Science*. 2002; 13(4):320–328. [PubMed: 12137134]
- Quinn PC, Burke S, Rush A. Part-whole perception in early infancy: evidence for perceptual grouping produced by lightness similarity. *Infant Behavior and Development*. 1993; 16(1):19–42.
- Quinn PC, Eimas PD. Pattern-line effects and units of visual processing in infants. *Infant Behavior and Development*. 1986; 9(1):57–70.
- Rimland B. The differentiation of childhood psychosis. *Journal of Autism and Childhood Schizophrenia*. 1971; 1:161–174. [PubMed: 5172390]
- Robertson LC, Lamb MR. Neuropsychological contributions to theories of part/whole organization. *Cognitive Psychology*. 1991; 23(2):299–330. [PubMed: 2055002]
- Savin H, Bever T. The nonperceptual reality of the phoneme. *Journal of Verbal Learning and Verbal Behavior*. 1970; 9(3):295–302.
- Shah A, Frith U. An islet of ability in autistic children: a research note. *Journal of Child Psychology and Psychiatry*. 1983; 24(4):613–620. [PubMed: 6630333]
- Slater A, Mattock A, Brown E, Bremner JG. Form perception at birth: revisited. *Journal of Experimental Child Psychology*. 1991; 51(3):395–406. [PubMed: 2072083]
- Snodgrass JG, Smith B, Feenan K, Corwin J. Fragmenting pictures on the Apple Macintosh computer for experimental and clinical application. *Behavior Research Methods, Instruments and Computers*. 1987; 19(2):270–274.
- Solso RL, King JF. Frequency and versatility of letters in the English language. *Behavior Research Methods and Instrumentation*. 1976; 8(3):283–286.
- Spencer J, O'Brien J, Riggs K, Braddick O, Atkinson J, et al. Motion processing in autism: evidence for a dorsal stream deficiency. *NeuroReport*. 2000; 11(12):2765–2767. [PubMed: 10976959]
- Tada WL, Stiles-Davis J. Children's analysis of spatial patterns: an assessment of their 'errors' in copying geometric forms. *Cognitive Development*. 1989; 4(2):177–195.
- Tager-Flusberg H, Plesa-Skwerer D, Faja S, Joseph RM. People with Williams syndrome process faces holistically. *Cognition*. 2003; 89:11–24. [PubMed: 12893122]
- Wang L, Mottron L, Peng D, Berthiaume C, Dawson M. Local bias and local-to-global interference without global deficit: a robust finding in autism under various conditions of attention, exposure time, and visual angle. *Cognitive Neuropsychology*. 2007; 24(5):550–574. [PubMed: 18416507]
- Wechsler, D. Wechsler Intelligence Scale for Children. 3. The Psychological Corporation; London: 1992.
- Wechsler, D. Wechsler Adult Intelligence Scale. 3. The Psychological Corporation; London: 1997.
- White S, O'Reilly H, Frith U. Big heads, small details and autism. *Neuropsychologia*. 2009; 47:1274–1281. [PubMed: 19428391]
- Yamasaki, T.; Maekawa, T.; Takahashi, H.; Fujita, T.; Kamio, Y., et al. Electrophysiology of visual and auditory perception in autism spectrum disorders. In: Patel, VB.; Preedy, VR.; Martin, CR., editors. *Comprehensive guide to autism*. Springer; New York: 2014. p. 791-808.

Research highlights

- Individuals with neurodevelopmental disorders cannot simply be characterized as having a local or global processing style.
- Cross-syndrome/cross-task/cross-modality comparisons reveal why.

Figure 1. Examples of one stimuli type (Tf), presented at three different densities: (a), (b), and (c)

Figure 2. The Navon task: presentation of stimuli

Figure 3. Example stimuli from the Fragmented Picture Completion task

Figure 4. Mean response time (in seconds) for correct detection of the phoneme at initial and medial/final positions for the ASD and Control groups. Error bars represent standard error of the mean

Figure 5. Mean response time (in seconds) for correct detection of the phoneme at initial and medial/final positions for the DS and WS groups. Error bars represent standard error of the mean

Figure 6. Proportion of local matches (local / [local + global]) for each group. Error bars represent standard error of the mean

Table 1
A brief description of three neurodevelopmental disorders

Group	Description	References
Williams syndrome (WS)	A rare genetic disorder, caused by a hemizygotic microdeletion of approximately 1.6 Mb containing ~28 genes on chromosome 7 (7q11.23), and characterized by a wide range of physical, cognitive, and behavioural atypicalities, including an uneven cognitive profile with particularly weak visuo-spatial construction abilities.	Donnai & Karmiloff-Smith, 2000; Poher, 2010
Down syndrome (DS)	A disorder of neural development, caused by the trisomy of chromosome 21 and characterized by decelerated maturation (neoteny), incomplete morphogenesis, atavisms, and a number of physical, cognitive, and behavioural atypicalities, including language delay and poor verbal working memory.	Jarrold, Baddeley & Phillips, 2002; Laws & Gunn, 2004
Autism spectrum disorder (ASD)	A neurodevelopmental disorder, with no single known cause, characterized by impaired social interaction and communication, and by restricted and stereotyped patterns of behaviour and interests.	APA, 2013

Table 2
Tasks organized by modality and level of processing

	Auditory-Verbal	Visuo-spatial
Lower-level tasks	Phoneme Segmentation	Navon Similarity Judgment
Higher-level tasks	Sentence Completion	Fragmented Pictures

Table 3
Group means (SD) for chronological age (CA), mental age (MA), verbal mental age (VMA), and visuospatial mental age (VSMA)

Task	Group	N	CA in years (SD)	MA in years (SD)	VMA in years (SD)	VSMA in years (SD)
Overall	WS	20	22.7 (8.7)	8.4 ^a (2.6)	10.5 (3.7)	6.3 ^a (2.4)
	DS	30	25.3 (5.8)	7.4 ^a (2.1)	7.0 (3.1)	7.5 ^a (2.0)
	ASD	32	14.8 (2.3)	13.8 (3.4)	14.2 (3.9)	13.4 (3.4)
	Control	31	14.8 (2.4)	13.9 (2.9)	14.2 (3.4)	13.6 (3.0)
Phoneme Segmentation	WS	16	23.6 (9.1)	9.2 ^a (2.1)	11.5 (3.2)	7.0 ^a (2.3)
	DS	12	23.8 (6.3)	9.1 ^a (2.2)	9.0 (3.3)	8.0 ^a (1.0)
	ASD	31	14.8 (2.4)	14.0 (3.3)	14.4 (3.8)	13.6 (3.3)
	Control	30	14.9 (2.4)	14.1 (2.8)	14.5 (3.2)	13.7 (2.9)
Sentence Completion	WS	19	22.6 (9.0)	8.6 ^a (2.4)	10.8 (3.4)	6.4 ^a (2.4)
	DS	26	25.5 (5.8)	7.5 ^a (2.1)	7.5 (2.9)	7.5 ^a (2.1)
	ASD	32	14.8 (2.3)	13.8 (3.4)	14.2 (3.9)	13.4 (3.4)
	Control	31	14.8 (2.4)	13.9 (2.9)	14.2 (3.4)	13.6 (3.0)
Navon Similarity Judgment	WS	8	20.6 (8.7)	8.3 ^a (1.8)	10.6 (3.3)	6.0 ^a (1.8)
	DS	11	22.7 (4.7)	8.1 ^a (1.8)	8.3 (2.6)	7.6 ^a (1.2)
	ASD	31	14.8 (2.4)	13.8 (3.4)	14.2 (3.9)	13.4 (3.4)
	Control	31	14.8 (2.4)	13.9 (2.9)	14.2 (3.4)	13.6 (3.0)
Fragmented Pictures	WS	11	22.7 (8.7)	8.6 ^a (1.7)	11.1 (3.0)	6.1 ^a (1.7)
	DS	21	23.9 (5.6)	7.7 ^a (2.3)	7.4 (3.4)	7.4 ^a (1.2)
	ASD	31	14.8 (2.4)	13.8 (3.4)	14.2 (3.9)	13.4 (3.4)
	Control	31	14.8 (2.4)	13.9 (2.9)	14.2 (3.4)	13.6 (3.0)

^aThese are approximations, because some individuals were close to floor.

Table 4
Number of correct responses for each condition of the Phoneme Segmentation task (max = 15) and the overall sensitivity index, A' , for each group

Group	<i>N</i>	Initial (<i>SD</i>)	Medial/final (<i>SD</i>)	Absent (<i>SD</i>)	A'^a
WS	16	13.69 (1.92)	9.19 (4.12)	11.63 (2.42)	.86 (.06)
DS	12	12.17 (3.07)	7.67 (3.45)	10.25 (2.38)	.77 (.07)
ASD	31	13.26 (1.65)	9.39 (2.69)	11.58 (1.88)	.84 (.07)
Control	30	13.60 (1.35)	9.70 (2.98)	12.13 (1.85)	.87 (.07)

^a ASD < Control, $p = .023$; DS < WS, $p = .006$.

Table 5
Measures of local–global sentence completions

Group	N	MA	Local completions	Mean response time in sec
WS	17	8.64 (2.42)	1.41 (1.23)	3.44 (1.63)
DS	24	7.50 (2.06)	1.46 (0.98)	4.24 (2.38)
ASD	30	13.74 (3.69)	1.20 (1.24)	3.60 (1.95)
Control	27	13.80 (3.22)	0.37 (0.57)	2.81 (1.43)

Table 6
Fragmented Pictures task results by group overall: Mean (SD)

Group	N	Mean frame no. for correct identification (max = 8)	Mean RT for correct identification (s)	Total no. of items identified at 6th, 7th, or 8th frame (max = 10)
WS	11	5.8 (0.6)	26.1 (5.7)	5.3 (2.5)
DS	21	6.7 (0.7)	26.7 (5.7)	8.5 (1.6)
ASD	31	5.3 (0.5)	24.6 (2.7)	4.1 (1.7)
Control	31	5.1 (0.5)	23.4 (2.3)	3.0 (1.7)

Table 7
Summary of findings from the current cross-syndrome comparison, organized by level of processing (high/low) and modality (auditory-verbal/visuospatial)

	Auditory-verbal		Visuospatial	
Low	Phoneme Segmentation		Navon	
	WS (G)	ASD (L)	WS (G)	ASD (G)
	DS (L)	TD (G)	DS (L=G)	<TD (G)
High	Sentence Completion		Fragmented Pictures	
	WS (G)	ASD (G)	WS (G)	ASD (G)
	=DS (G)	<TD (G)	>DS (G)	=/<TD (G)

Note WS = Williams syndrome, DS = Down syndrome, ASD = Autism Spectrum Disorder, TD = typically developing controls, L = the group demonstrated a local bias, G = the group demonstrated a global bias