Published in final edited form as:

Brain Struct Funct. 2012 January ; 217(1): 115–125. doi:10.1007/s00429-011-0333-x.

Neuroanatomic changes and their association with cognitive decline in mild cognitive impairment: a meta-analysis

Thomas Nickl-Jockschat,

Department of Psychiatry, Psychotherapy and Psychosomatic, RWTH Aachen University, Pauwelsstr. 30, 52074 Aachen, Germany

JARA, Translational Brain Medicine, Aachen, Germany

Alexandra Kleiman,

JARA, Translational Brain Medicine, Aachen, Germany

Department of Neurology, RWTH Aachen University, Pauwelsstrasse 30, 52074 Aachen, Germany

Jörg B. Schulz,

JARA, Translational Brain Medicine, Aachen, Germany

Department of Neurology, RWTH Aachen University, Pauwelsstrasse 30, 52074 Aachen, Germany

Frank Schneider,

Department of Psychiatry, Psychotherapy and Psychosomatic, RWTH Aachen University, Pauwelsstr. 30, 52074 Aachen, Germany

JARA, Translational Brain Medicine, Aachen, Germany

Angela R. Laird,

Research Imaging Center, University of Texas Health Science Center, 4502 Medical Drive, San Antonio, TX 78229, USA

Peter T. Fox,

Research Imaging Center, University of Texas Health Science Center, 4502 Medical Drive, San Antonio, TX 78229, USA

Simon B. Eickhoff, and

Department of Psychiatry, Psychotherapy and Psychosomatic, RWTH Aachen University, Pauwelsstr. 30, 52074 Aachen, Germany

JARA, Translational Brain Medicine, Aachen, Germany

Institute of Neuroscience and Medicine, Research Center Jülich GmbH, Wilhelm-Johnen-Straße, 52428 Jülich, Germany

---

Correspondence to: Kathrin Reetz, kreetz@ukaachen.de.

Thomas Nickl-Jockschat and Alexandra Kleiman contributed equally to this work.

Electronic supplementary material The online version of this article (doi:10.1007/s00429-011-0333-x) contains supplementary material, which is available to authorized users.

Conflict of interest The authors declare that they have no conflict of interest.

Kathrin Reetz

JARA, Translational Brain Medicine, Aachen, Germany

Department of Neurology, RWTH Aachen University, Pauwelsstrasse 30, 52074 Aachen, Germany

Institute of Neuroscience and Medicine, Research Center Ju"lich GmbH, Wilhelm-Johnen-Stra"e, 52428 J"lich, Germany

Kathrin Reetz: kreetz@ukaachen.de

Abstract

Mild cognitive impairment (MCI) is an acquired syndrome characterised by cognitive decline not affecting activities of daily living. Using a quantitative meta-analytic approach, we aimed to identify consistent neuroanatomic correlates of MCI and how they are related to cognitive dysfunction. The meta-analysis enrolls 22 studies, involving 917 MCI (848 amnesic MCI) patients and 809 healthy controls. Only studies investigating local changes in grey matter and reporting whole-brain results in stereotactic coordinates were included and analysed using the activation likelihood estimation approach. Probabilistic cytoarchitectonic maps were used to compare the localization of the obtained significant effects to histological areas. A correlation between the probability of grey matter changes and cognitive performance of MCI patients was performed. In MCI patients, the meta-analysis revealed three significant clusters of convergent grey matter atrophy, which were mainly situated in the bilateral amygdala and hippocampus, extending to the left medial temporal pole and thalamus, as well as in the bilateral precuneus. A sub-analysis in only amnesic MCI revealed a similar pattern. A voxel-wise analysis revealed a correlation between grey matter reduction and cognitive decline in the right hippocampus and amygdala as well as in the left thalamus. This study provides convergent evidence of a distinct neuroanatomical pattern in MCI. The correlation analysis with cognitive-mnemonic decline further highlights the impact of limbic structures and the linkage with data from a functional neuroimaging database provides additional insight into underlying functions. Although different pathologies are underlying MCI, the observed neuroanatomical pattern of structural changes may reflect the common clinical denominator of cognitive impairment.

Keywords

Mild cognitive impairment; Meta-analysis; Voxel-based morphometry; Cognitive impairment; Activation likelihood estimation approach; Mini-mental state examination

Introduction

Mild cognitive impairment (MCI) is as an acquired syndrome defined as cognitive decline greater than expected for an individual's age and education but that does not interfere notably with activities of daily life (Gauthier et al. 2006). According to Petersen, amnesic MCI (aMCI) is a more specified term describing a subtype of MCI, in which there are memory complaint, objective memory impairment; essentially preserved general cognitive function, largely intact functional activities and no dementia (Petersen 2004). Given the current available morphometric MRI studies selecting MCI and aMCI patients according to

the older Petersen criteria (Petersen 2004; Petersen et al. 1999), it has to be noted, that results of this meta-analysis are based on these definitions. In the general elderly population, the prevalence of MCI is estimated between 3.2 and 19.3% (Ritchie et al. 2001). A meta-analysis in cohort studies revealed a yearly rate of progression of MCI to dementia of approximately 5–10% (Mitchell and Shiri-Feshki 2009). However, population-based studies have shown that up to 44% of patients presenting with MCI may return to normal cognitive-mnemonic functioning within a year (Ritchie et al. 2001; Ganguli et al. 2004). These findings underline that various, yet not fully understood factors may influence cognitive and mnemonic decline in an elderly population (Gauthier et al. 2006). Moreover, MCI is also supposed to originate from various aetiologies (Petersen et al. 2001) involving beginning neurodegenerative disorders but also for example vascular lesions, depression or diverse medical conditions. Thus, a method to distinguish between converters into dementia and non-converters has become one of the interests in MCI research.

Structural MRI studies in MCI patients have revealed grey matter loss in the hippocampus (Muller et al. 2007; Krasuski et al. 1998; Jack et al. 1999) but also have resulted in very heterogeneous findings concerning atrophy in several frontal, parietal and the occipitotemporal regions (Chetelat et al. 2002; Convit et al. 2000; Karas et al. 2004; Killiany et al. 2000; Visser et al. 1999). The question thus emerges whether common neuroanatomical substrates of this widely defined syndrome exist and how neurostructural findings correlate with neuropsychological markers that indicate cognitive-mnemonic decline. The aim of the present study was therefore:

1. to identify regions of consistent brain structure changes as reported in the current literature and relate these findings to cytoarchitectural areas,
2. to investigate whether the likelihood of finding regional atrophy correlates with neuropsychological markers that indicate cognitive-mnemonic decline in the respective studies, and
3. to assess the physiological functions of brain regions showing consistent atrophy.

Methods and materials

Literature search and selection

PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/>) and Brain-Map (<http://brainmap.org/>) databases were searched by using the following search string “smci + vbm”, “mci + voxel”, “mild + cognitive + impairment + voxel”, “mild + cognitive + impairment + vbm” to identify morphometric MRI studies investigating MCI. BrainMap provides a database that establishes structure–function correspondence in the brain through the combined application of experimental psychology, human neuroscience, and non-invasive neuroimaging (Laird et al. 2005b). Inclusion criteria for this meta-analysis were: (1) original peer-reviewed studies, (2) selection of MCI patients according to the Petersen criteria (Petersen 2004; Petersen et al. 1999), (3) quantitative automated MRI whole-brain grey matter assessment using voxel-based morphometry (VBM), (4) comparison of MCI patients to a matched healthy control group, (5) reporting of results as coordinates in stereotactic space (Talairach/MNI).

In total, 22 studies published between 2002 and 2010 fit these criteria and were included in our meta-analysis (see Table 1). Out of the 917 included MCI patients, 848 subjects were classified as amnesic MCI (aMCI). The remaining “non-amnesic” MCI subjects were subdivided in 36 subjects with multi-cognitive domain MCI (MCD-MCI) and 33 single-cognitive domain MCI (SCD-MCI) (Schmidt-Wilcke et al. 2009; Bell-McGinty et al. 2005; Pa et al. 2009; Jauhiainen et al. 2008). By now, there have been various re-definitions of (a) MCI and/or prodromal stages of Alzheimer’s disease (AD), but all of our included studies referred to the criteria according to Petersen (Petersen et al. 1999; Petersen 2004). These criteria are: (1) memory complaint usually corroborated by an informant, (2) objective memory impairment for age, (3) essentially preserved general cognitive function, (4) largely intact functional activities, (5) not demented. Overall, 130 MCI patients were reported to have converted to AD during follow-up, but separate VBM data for converters were only reported for three studies encompassing 41 subjects. None of the included studies reported increased grey matter changes in MCI patients compared to healthy controls.

Anatomic likelihood estimation meta-analysis procedure

The meta-analysis was carried out using a revised version (Eickhoff et al. 2009) of the activation likelihood estimation (ALE) approach for coordinate-based meta-analysis of neuroimaging results (Laird et al. 2005a; Turkeltaub et al. 2002). This algorithm aims at identifying areas showing a convergence of findings across studies, which is higher than expected under a spatially random spatial association. The key idea behind ALE is to treat the reported foci as centres of 3D Gaussian probability distributions reflecting the spatial uncertainty associated with each reported set of coordinates. The probabilities of all foci reported in a given study were combined for each voxel, resulting in a modelled anatomical effects map (paralleling the modelled activation maps in functional imaging meta-analyses). Taking the union across these yielded voxel-wise ALE scores describing the convergence of results at each particular location. To distinguish ‘true’ convergence between studies from random convergence, i.e., noise, ALE scores were compared to an empirical null-distribution reflecting a random spatial association between experiments. Hereby, a random-effects inference is invoked, focussing on inference on the above-chance convergence between studies, not clustering of foci within a particular study. Computationally, deriving this null-hypothesis involved sampling a voxel at random from each of the anatomical effects maps and taking the union of these values in the same manner as done for the (spatially contingent) voxels in the true analysis. The p value of a ‘true’ ALE was then given by the proportion of equal or higher values obtained under the null-distribution. The resulting non-parametric p values were transformed into Z scores and thresholded at a cluster level corrected threshold of $p < 0.05$.

Anatomical allocation by probabilistic maps

We used the SPM Anatomy Toolbox v1.5 (Eickhoff et al. 2007) to compare the localisation of the significant effects to histological areas as described in probabilistic cytoarchitectonic maps, which were summarised in a maximum probability map (MPM). The latter attributes each voxel of the reference space to the most likely cytoarchitectonic area or myeloarchitectonically defined fibre tract (Table 2) at this position and allows the definition of non-overlapping representations of all areas from a set of inevitably overlapping

probabilistic maps (Amunts and Zilles 2001; Eickhoff et al. 2006; Zilles et al. 2002; Eickhoff et al. 2005). For all clusters in regions that have not yet been histologically examined, we used the MPM of macroanatomical regions provided in the “Harvard-Oxford cortical and subcortical structural atlas” (Desikan et al. 2006; Makris et al. 2006).

Correlations between MMSE and likelihood of grey matter changes

The Mini-Mental State Examination (MMSE) (Folstein et al. 1975) is the most commonly used bedside test to screen for cognitive impairments. We aimed to investigate whether assessment of an MCI sample with lower MMSE scores increases the probability of finding grey matter reduction in a particular brain region. Thus, mean MMSE scores reported in the original studies were correlated on a voxel-wise level with the modelled anatomical effects values using a Spearman rank correlation ($p < 0.05$).

Functional analysis by Behavioural Domains

In order to determine structure–function relationships, we used the BrainMap database (Laird et al. 2005b) to assess the physiological functions of brain regions showing consistent atrophy in MCI. The BrainMap database archives peak coordinates of activations of functional imaging studies and their corresponding metadata (e.g. number of subjects, analysis technique, paradigm, cognitive domain, etc.). Metadata on the domain of the behavioural system is classified according to six main categories and their related subcategories: cognition, action, perception, emotion, interoception or pharmacology (a complete list of BrainMap’s behavioural domains can be found at <http://brainmap.org/scribe/>). To determine possible functional consequences of the structural changes, we analysed the behavioural domain metadata associated with each cluster from the ALE results to determine frequency of domain hits relative to its distribution across the whole brain (i.e. the entire database) as described before (Laird et al. 2009).

Results

Study characteristics

The mean age of subjects across the twenty-two VBM studies ranged from 65.4 to 77.9 years in the MCI group ($n = 917$ MCI patients, mean age 71.7 ± 3.2 (SD) years) and from 63 to 78.1 years in the control group ($n = 809$ healthy controls, mean age: 70.8 ± 4.4 (SD) years). Mean performance (within each study) in MMSE ranged from 23.5 to 28.8 points in MCI patients. The subgroup of aMCI comprised 848 subjects (mean age 72 ± 2.5 (SD) years; age range of 67.7–77.9) and 778 healthy controls (mean age 71.1 ± 4.2 (SD) years; age range 63.3–78.1). Mean performance in MMSE ranged from 23.1 to 28.7 points in aMCI patients. The demographic and clinical data are also summarised in Table 1.

Regions with significant grey matter differences between subjects with MCI and healthy controls

Across studies, we found in total three clusters of consistent grey matter reduction in MCI patients (Table 2, Fig. 1). The two largest clusters were situated in the medial temporal lobes. Analysis by cytoarchitectural probability showed that the largest cluster was mainly situated in the laterobasal part of the left amygdala, the left thalamus, the left medial

temporal pole and the cornu ammonis region of the left hippocampus. The second largest cluster in the contralateral temporal lobe was primarily located in the right laterobasal amygdala and extended to the fascia dentate region of the right hippocampus. The third largest cluster was to be found in the parietal precuneus broadening to the posterior cingulate cortex bilaterally, accentuated on the right side.

A sub-analysis in only aMCI subject compared to healthy controls revealed a similar pattern of brain structure changes with a slightly different quantity and size of clusters. The three clusters observed in the above-described analysis were almost identically found in this analysis again. However, the analysis in the aMCI subsample revealed two additional mid-sized clusters, one in the left superior temporal gyrus and the other one in the left thalamus (please see Supplementary Table 1; Supplementary Fig. 1).

Correlation of grey matter changes with MMSE performance

A voxel-wise analysis whether grey matter changes correlated with MMSE performance yielded three clusters indicating an increased probability of finding brain atrophy in these regions in patients with lower MMSE scores (Table 2, Fig. 2). The largest and the smallest cluster indicated at a correlation of grey matter volume reductions with MMSE performance in the right superficial parts of the amygdala and the right parahippocampal gyrus. The third cluster was primarily situated in the left thalamus. The MMSE correlation analysis in the aMCI group revealed one larger cluster in the right amygdala and one smaller cluster in the right thalamus (please see Supplementary Table 1; Supplementary Fig. 2).

Functional analysis by Behavioural Domains

Functional Behavioural Domain analysis was carried out for all clusters of convergence that indicated significant differences between MCI patients and healthy controls. Behavioural Domains that were most frequently represented in the clusters of convergence were “emotion” (bilateral hippocampus/amygdala, parietal precuneus, left thalamus), “cognition-memory” (bilateral hippocampus/amygdala, parietal precuneus), “cognition-language” (left hippocampus/amygdala) and “emotion-perception” (left thalamus).

Discussion

This quantitative meta-analysis, encompassing 22 wholebrain VBM studies with a total of 917 MCI patients and 809 healthy controls, is the first one presenting consistent regional brain atrophy, and clinical correlation with clinical cognitive-mnemonic performance as well as a description of structure–function correspondence in MCI. Our results demonstrate three significant clusters of convergence in MCI patients. These clusters were located bilaterally in the middle temporal lobe, including the hippocampus and the amygdala, and the parietal precuneus. This pattern was confirmed in a sub-analysis of only aMCI (848 out of 917 MCI subjects). Cognitive decline correlated with findings of brain atrophy in the bilateral hippocampus/amygdala region and the left thalamus. Behavioural Domain analysis found that physiologically these regions were mainly associated with emotion, cognition and memory.

The neuropathological substrate of the most common (amnesic) subtype of MCI, was shown to be the medial temporal lobe and concomitant pathologic abnormalities including argyrophilic grain disease, hippocampal sclerosis and vascular disease were found in *post mortem* studies (Petersen et al. 2006). The involvement of the medial temporal lobe in mnestic functions, especially encoding and retrieval of episodic and spatial memory, is well known and has been repeatedly demonstrated in functional neuroimaging, lesion studies and various animal models (Squire 1992; Sauvage et al. 2008; Fortin et al. 2004). Functionally, the observed structures are commonly regarded as key assemblage points in a widespread cerebral network of episodic memory encoding and retrieval that encompasses also ventrolateral temporal, medial and lateral parietal, and medial and lateral frontal isocortical brain (Dickerson and Eichenbaum 2010). Although there is yet no general consensus whether there is a specific topographic organisation in the medial temporal lobe that differentiates the functional neuroanatomy of memory encoding or retrieval, data derived from fMRI studies hint at differential activation in this brain region during respective paradigms. The hippocampus, located in the medial temporal lobe, belongs to the limbic system and is associated with various cortical regions. Together these structures are responsible for the formation of the long-term memory and spatial navigation. Specific activation of the hippocampal cornu ammonis (CA) areas was reported during episodic memory encoding, while retrieval related activation was mainly found in the subiculum (Zeineh et al. 2003). Item-based (e.g. item recollection, distinction between new and previously shown items) and associative memory processes also seem to involve different functional topographies. The hippocampal formation is primarily engaged in recollection or associative memories, while the entorhinal cortex was found to be activated for familiarity or for item-based non-associative memory (Davachi et al. 2003; Eichenbaum et al. 2007; Squire 2004). Spatial memory functions have been shown to be associated with posterior parahippocampal and perirhinal activity (Davachi et al. 2003; Maguire 1997; Stern et al. 1996). Mirroring these findings of the current literature on hippocampal involvement in episodic memory, our Behavioural Domain analysis of the clusters in the medial temporal lobes was able to verify this association over a large sample of studies. Remarkably, the larger of the two clusters situated in the cornu ammonis region of the hippocampus was on the right side. Lesion studies in humans and animals suggest a lateralization of hippocampal functions in spatial memory with positional memory and object-location memory more impaired by lesions of the right hippocampal formation (Cipolotti and Bird 2006; Kessels et al. 2001). Both memory systems have been shown to be impaired in MCI patients compared to normally ageing individuals (Kessels et al. 2010a), but were also able to validly discriminate MCI from AD patients, thus supporting the notion that tests enrolling visual and spatial features may have better diagnostic value than those relying on verbal memory (Kessels et al. 2010b). Recently, a VBM meta-analysis of MCI patients which converted to AD, proposed atrophy of the left medial temporal lobe as a neurostructural biomarker to predict conversion from amnesic MCI to AD (Ferreira et al. 2009). In this context, our metaanalysis findings of the hippocampal atrophy in patients with MCI further emphasise the important role of the hippocampus in MCI pathobiology.

The two largest clusters of convergence also comprised the amygdalae on either hemisphere. The role of the amygdalae in emotion processing and emotional memory has been

highlighted by functional imaging experiments and lesion studies in animal models (Maren and Fanselow 1995; Pape and Pare 2010; Roozendaal et al. 2009). Studies in humans have further demonstrated the essential role of the amygdala in the modulation of emotions, emotional memory processes and storage (Aldolphs 1999; Sarter and Markowitsch 1985; Fine and Blair 2000). On a cytoarchitectonic level, one of the most widely accepted subparcellation schemes distinguishes the superficial, the laterobasal and the centromedial part of the amygdala (Amunts et al. 2005; de Olmos and Heimer 1999). The basal and lateral nuclei of the amygdala receive projections from associative visual and auditory cortical regions, the orbital and the cingulate cortex and from the hippocampus and the entorhinal cortex. Besides projections to higher cortical regions, the efferences of the basolateral group are directed towards the centromedial complex of the amygdala. The centromedial amygdala itself sends efferences to the paraventricular and the lateral nuclei of the hypothalamus, the locus coeruleus, the periaqueductal grey region and the parabrachial nucleus, thus mediating autonomic responses to aversive stimuli (Pitkanen et al. 1997; Pitkanen et al. 1995). Given the functional role of the amygdala involving the storage of emotional memory contents, our findings in the meta-analysis emphasise the crucial role in mild cognitive impairment.

Interestingly, our meta-analysis and the correlation with the MMSE revealed clusters in the thalamus. From the neuropathological perspective, the limbic thalamus is strongly involved in AD (Braak and Braak 1991). Amyloid deposits and neurofibrillary tangles have been found in the thalamus (Braak and Braak 1991) as well as a significant loss of its grey matter in patients with MCI and AD (Karas et al. 2004). Functional neuroimaging studies underline the essential role in dynamical interactions of thalamocortical networks (Lopes da Silva 1991). In patients with MCI, thalamic dysfunctions have been reported, which likely affect the thalamocortical integrity (Cantero et al. 2009). These findings are not surprising given the role of the thalamus in declarative memory (de Rover et al. 2008; Van der Werf et al. 2003) and postulate a mild involvement of this subcortical structure in neurodegenerative processes.

The third cluster was mainly located in the posterior cingulate/precuneus region, accentuated on the left side in the MCI group. The precuneus located in the posteromedial portion of the parietal lobe, has a complex set of interconnections, namely multiple reciprocal and afferent/efferent cortical connections to the cingulate, the adjacent parietal and frontal cortex and subcortical projections to the thalamus, striatum, claustrum as well as the brainstem (Cavanna and Trimble 2006). According to the interwoven network, the precuneus most likely plays an important role in the modulation of self-processing, consciousness, episodic memory retrieval and visuo-spatial imagery (Cavanna and Trimble 2006). Alterations in the posterior cingulate/precuneus region in patients with MCI have not only been reported on the structural level (Lo et al. 2010; Derflinger et al. 2011) but also implications for the default mode network (DMN). The DMN includes the posterior cingulate extending into the precuneus, lateral parietal and medial prefrontal regions. Regional activity levels and network connectivity of resting brain function in MCI and AD feature additional default-related changes particularly in the posterior cingulate/precuneus and hippocampal region (Beason-Held 2011). Interestingly, these regions demonstrating the aberrant default network activity also overlap the anatomy of regions with highly amyloid burden in early AD (Klunk et al. 2004). Moreover, the posterior cingulate/precuneus region may play a role in cognitive

reserve, which has been claimed as a factor mitigating the clinical manifestation of AD (Serra et al. 2011).

In the present analysis, we observed several clusters in which the likelihood of finding differences to control populations depended on the patients' MMSE score, within the right hippocampal/amygdala region and the left thalamus. As already mentioned, MCI has to be regarded as a syndrome with underlying different diseases with distinct pathophysiologies under the main symptom of cognitive decline causing no inference with daily life functioning. Given the pathological heterogeneity, the further course of MCI is often difficult to estimate. In more detail, if MCI patients present in the further course a more severe cognitive decline (e.g. lower MMSE scores) they are more likely to convert to AD. In contrast, MCI patients showing no further reduction of cognitive function will potentially be clinical stable or even improve over time. Since progression to AD seems to be associated with marked deficits in spatial memory subsystems and lesions of the right hippocampal formation are known to be associated with congruent functional impairments, MCI patients on the verge of conversion to AD can be expected to display more atrophy in this region than other patient subgroups. Thus, these regions might provide a viable diagnostic tool for the identification of MCI patients at high risk to convert into dementia using structural neuroimaging. However, this interpretation has several caveats. Firstly, data of human lesion studies is heterogeneous by nature, especially in a small brain region with functionally diverse structures such as the medial temporal lobe. Due to the small sample of "non-amnesic" MCI subjects in our MCI group, a direct comparison between aMCI and "non-amnesic" MCI in order to detect specific regions for each group appeared not to be meaningful, but should be followed-up in future studies with well-balanced sufficient samples. Moreover, since MMSE scores are a rather unspecific measure of cognitive decline, correlations of brain structure changes with more specific memory tests will be a promising goal for future studies once available.

Whereas a previous VBM meta-analysis only found structural abnormalities in the right gyrus rectus, the left temporal pole/anterior superior temporal sulcus and the right amygdala (Schroeter et al. 2009), we found a bilateral involvement of the hippocampus and amygdala, the precuneus as well as of the left thalamus in patients with MCI. In this vein, the regions we found in our MCI group, the previous VBM meta-analysis by Schroeter mainly reported these findings in the AD group (Schroeter et al. 2009). Given the fact that the previous meta-analysis only comprised five MRI studies with 165 patients for the MCI group and nine MRI studies with 177 patients for the AD group, these observed differences might be due to the small sample of the previous meta-analysis.

As any meta-analysis technique, this approach is limited by the details of the primary research articles, including their definition of MCI and reported variables. Overall, most definitions of MCI relate to the concept of "cognitive decline greater than that for an individual's age or education level but that does not interfere notably with activities of daily life" (Gauthier et al. 2006; Dubois et al. 2007). However, clinical outcome can vary considerably due to differences in diagnostic entry criteria that may appear as minor (Dubois and Levy 2004; Gauthier et al. 2006). In consequence, also neuroanatomical findings might vary due to different entry criterion (Dubois and Levy 2004; Gauthier et al. 2006).

Finally and as already mentioned above only a relatively small number of published studies matched our inclusion criteria, we were therefore not able to perform separate analyses on clinical subgroups. Thus, more detailed analyses on structural findings pertaining to clinical subgroups of MCI will be an important topic for future research once additional whole-brain based VBM studies on this population are available.

Conclusion

In the current quantitative meta-analysis of structural neuroimaging data encompassing 917 MCI patients and 809 healthy controls, we identified common neuroanatomic correlates in the clinical syndrome of cognitive impairment. Firstly, we provide objective evidence that limbic and temporoparietal structures are consistently involved in pathobiology of MCI. Secondly, the correlation analysis with the clinical neuropsychological finding of cognitivemnemonic decline primarily highlights the impact of the hippocampus and amygdala in this clinical syndrome. Thirdly, we demonstrated using a neuroinformatics approach that these consistent regions of brain atrophy are functionally linked to “emotion” and “cognition”.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

ARL, PTF, and SBE were funded by the Human Brain Project (R01-MH074457) and the Initiative and Networking Fund of the Helmholtz Association within the Helmholtz Alliance on Systems Biology (Human Brain Model). JBS received research support from grants of the German Ministry of Education and Research (BMBF) for the projects GeneMove (01GM0503), NGFNplus and Competence Network Degenerative Dementias. SBE and KR were funded by the DFG (Deutsche Forschungsgemeinschaft) Translational Brain Research in Psychiatry and Neurology (DFG ZUK32/1).

References

- Aldolphs R. The human amygdala and emotion. *Prog Clin Neurosci.* 1999; 5:125–137.
- Amunts K, Zilles K. Advances in cytoarchitectonic mapping of the human cerebral cortex. *Neuroimaging Clin N Am.* 2001; 11(2):151–169. [PubMed: 11489732]
- Amunts K, Kedo O, Kindler M, Pieperhoff P, Mohlberg H, Shah NJ, Habel U, Schneider F, Zilles K. Cytoarchitectonic mapping of the human amygdala, hippocampal region and entorhinal cortex: intersubject variability and probability maps. *Anat Embryol (Berl).* 2005; 210(5–6):343–352. [PubMed: 16208455]
- Beason-Held LL. Dementia and the default mode. *Curr Alzheimer Res.* 2011 [Epub ahead of print].
- Bell-McGinty S, Lopez OL, Meltzer CC, Scanlon JM, Whyte EM, Dekosky ST, Becker JT. Differential cortical atrophy in subgroups of mild cognitive impairment. *Arch Neurol.* 2005; 62(9):1393–1397. [PubMed: 16157746]
- Braak H, Braak E. Alzheimer's disease affects limbic nuclei of the thalamus. *Acta Neuropathol.* 1991; 81(3):261–268. [PubMed: 1711755]
- Cantero JL, Atienza M, Gomez-Herrero G, Cruz-Vadell A, Gil-Neciga E, Rodriguez-Romero R, Garcia-Solis D. Functional integrity of thalamocortical circuits differentiates normal aging from mild cognitive impairment. *Hum Brain Mapp.* 2009; 30(12):3944–3957. [PubMed: 19449329]
- Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and behavioural correlates. *Brain.* 2006; 129(Pt 3):564–583. [PubMed: 16399806]

- Chetelat G, Desgranges B, De La Sayette V, Viader F, Eustache F, Baron JC. Mapping gray matter loss with voxel-based morphometry in mild cognitive impairment. *Neuroreport*. 2002; 13(15):1939–1943. [PubMed: 12395096]
- Cipolotti L, Bird CM. Amnesia and the hippocampus. *Curr Opin Neurol*. 2006; 19(6):593–598. [PubMed: 17102699]
- Convit A, de Asis J, de Leon MJ, Tarshish CY, De Santi S, Rusinek H. Atrophy of the medial occipitotemporal, inferior, and middle temporal gyri in non-demented elderly predict decline to Alzheimer's disease. *Neurobiol Aging*. 2000; 21(1):19–26. [PubMed: 10794844]
- Davachi L, Mitchell JP, Wagner AD. Multiple routes to memory: distinct medial temporal lobe processes build item and source memories. *Proc Natl Acad Sci USA*. 2003; 100(4):2157–2162. [PubMed: 12578977]
- de Olmos JS, Heimer L. The concepts of the ventral striatopallidal system and extended amygdala. *Ann N Y Acad Sci*. 1999; 877:1–32. [PubMed: 10415640]
- de Rover M, Petersson KM, van der Werf SP, Cools AR, Berger HJ, Fernandez G. Neural correlates of strategic memory retrieval: differentiating between spatial-associative and temporal-associative strategies. *Hum Brain Mapp*. 2008; 29(9):1068–1079. [PubMed: 17948888]
- Derflinger S, Sorg C, Gaser C, Myers N, Arsic M, Kurz A, Zimmer C, Wohlschläger A, Muhlau M. Grey-matter atrophy in Alzheimer's disease is asymmetric but not lateralized. *J Alzheimers Dis*. 2011 [Epub ahead of print].
- Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, Albert MS, Killiany RJ. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage*. 2006; 31(3):968–980. [PubMed: 16530430]
- Dickerson BC, Eichenbaum H. The episodic memory system: neurocircuitry and disorders. *Neuropsychopharmacology*. 2010; 35(1):86–104. [PubMed: 19776728]
- Dubois B, Levy R. Cognition, behavior and the frontal lobes. *Int Psychogeriatr*. 2004; 16(4):379–387. [PubMed: 15715355]
- Dubois B, Feldman HH, Jacova C, Dekosky ST, Barberger-Gateau P, Cummings J, Delacourte A, Galasko D, Gauthier S, Jicha G, Meguro K, O'Brien J, Pasquier F, Robert P, Rossor M, Salloway S, Stern Y, Visser PJ, Scheltens P. Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria. *Lancet Neurol*. 2007; 6(8):734–746. [PubMed: 17616482]
- Eichenbaum H, Yonelinas AP, Ranganath C. The medial temporal lobe and recognition memory. *Annu Rev Neurosci*. 2007; 30:123–152. [PubMed: 17417939]
- Eickhoff SB, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, Zilles K. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *Neuroimage*. 2005; 25(4):1325–1335. [PubMed: 15850749]
- Eickhoff SB, Heim S, Zilles K, Amunts K. Testing anatomically specified hypotheses in functional imaging using cytoarchitectonic maps. *Neuroimage*. 2006; 32(2):570–582. [PubMed: 16781166]
- Eickhoff SB, Paus T, Caspers S, Grosbras MH, Evans AC, Zilles K, Amunts K. Assignment of functional activations to probabilistic cytoarchitectonic areas revisited. *Neuroimage*. 2007; 36(3):511–521. [PubMed: 17499520]
- Eickhoff SB, Laird AR, Grefkes C, Wang LE, Zilles K, Fox PT. Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: a random-effects approach based on empirical estimates of spatial uncertainty. *Hum Brain Mapp*. 2009; 30(9):2907–2926. [PubMed: 19172646]
- Ferreira LK, Diniz BS, Forlenza OV, Busatto GF, Zanetti MV. Neurostructural predictors of Alzheimer's disease: a meta-analysis of VBM studies. *Neurobiol Aging*. 2009
- Fine C, Blair RJR. The cognitive and emotional effects of amygdala damage. *Neurocase*. 2000; 6:435–450.
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12(3):189–198. [PubMed: 1202204]
- Fortin NJ, Wright SP, Eichenbaum H. Recollection-like memory retrieval in rats is dependent on the hippocampus. *Nature*. 2004; 431(7005):188–191. [PubMed: 15356631]

- Ganguli M, Dodge HH, Shen C, DeKosky ST. Mild cognitive impairment, amnesic type: an epidemiologic study. *Neurology*. 2004; 63(1):115–121. [PubMed: 15249620]
- Gauthier S, Reisberg B, Zaudig M, Petersen RC, Ritchie K, Broich K, Belleville S, Brodaty H, Bennett D, Chertkow H, Cummings JL, de Leon M, Feldman H, Ganguli M, Hampel H, Scheltens P, Tierney MC, Whitehouse P, Winblad B. Mild cognitive impairment. *Lancet*. 2006; 367(9518): 1262–1270. [PubMed: 16631882]
- Jack CR Jr, Petersen RC, Xu YC, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Waring SC, Tangalos EG, Kokmen E. Prediction of AD with MRI-based hippocampal volume in mild cognitive impairment. *Neurology*. 1999; 52(7):1397–1403. [PubMed: 10227624]
- Jauhiainen AM, Kangasmaa T, Rusanen M, Niskanen E, Tervo S, Kivipelto M, Vanninen RL, Kuikka JT, Soininen H. Differential hypometabolism patterns according to mild cognitive impairment subtypes. *Dement Geriatr Cogn Disord*. 2008; 26(6):490–498. [PubMed: 18987469]
- Karas GB, Scheltens P, Rombouts SA, Visser PJ, van Schijndel RA, Fox NC, Barkhof F. Global and local gray matter loss in mild cognitive impairment and Alzheimer's disease. *Neuroimage*. 2004; 23(2):708–716. [PubMed: 15488420]
- Kessels RP, de Haan EH, Kappelle LJ, Postma A. Varieties of human spatial memory: a meta-analysis on the effects of hippocampal lesions. *Brain Res Brain Res Rev*. 2001; 35(3):295–303. [PubMed: 11423159]
- Kessels RP, Meulenbroek O, Fernandez G, Olde Rikkert MG. Spatial working memory in aging and mild cognitive impairment: effects of task load and contextual cueing. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn*. 2010a; 17(5):556–574. [PubMed: 20574862]
- Kessels RP, Rijken S, Joosten-Weyn Banningh LW, Van Schuylenborgh VANEN, Olde Rikkert MG. Categorical spatial memory in patients with mild cognitive impairment and Alzheimer dementia: positional versus object-location recall. *J Int Neuropsychol Soc*. 2010b; 16(1):200–204. [PubMed: 19883520]
- Killiany RJ, Gomez-Isla T, Moss M, Kikinis R, Sandor T, Jolesz F, Tanzi R, Jones K, Hyman BT, Albert MS. Use of structural magnetic resonance imaging to predict who will get Alzheimer's disease. *Ann Neurol*. 2000; 47(4):430–439. [PubMed: 10762153]
- Klunk WE, Engler H, Nordberg A, Wang Y, Blomqvist G, Holt DP, Bergstrom M, Savitcheva I, Huang GF, Estrada S, Ausen B, Debnath ML, Barletta J, Price JC, Sandell J, Lopresti BJ, Wall A, Koivisto P, Antoni G, Mathis CA, Langstrom B. Imaging brain amyloid in Alzheimer's disease with Pittsburgh Compound-B. *Ann Neurol*. 2004; 55(3):306–319. [PubMed: 14991808]
- Krasuski JS, Alexander GE, Horwitz B, Daly EM, Murphy DG, Rapoport SI, Schapiro MB. Volumes of medial temporal lobe structures in patients with Alzheimer's disease and mild cognitive impairment (and in healthy controls). *Biol Psychiatry*. 1998; 43(1):60–68. [PubMed: 9442345]
- Laird AR, Fox PM, Price CJ, Glahn DC, Uecker AM, Lancaster JL, Turkeltaub PE, Kochunov P, Fox PT. ALE meta-analysis: controlling the false discovery rate and performing statistical contrasts. *Hum Brain Mapp*. 2005a; 25(1):155–164. [PubMed: 15846811]
- Laird AR, Lancaster JL, Fox PT. BrainMap: the social evolution of a human brain mapping database. *Neuroinformatics*. 2005b; 3(1):65–78. [PubMed: 15897617]
- Laird AR, Eickhoff SB, Li K, Robin DA, Glahn DC, Fox PT. Investigating the functional heterogeneity of the default mode network using coordinate-based meta-analytic modeling. *J Neurosci*. 2009; 29(46):14496–14505. [PubMed: 19923283]
- Lo CY, Wang PN, Chou KH, Wang J, He Y, Lin CP. Diffusion tensor tractography reveals abnormal topological organization in structural cortical networks in Alzheimer's disease. *J Neurosci*. 2010; 30(50):16876–16885. [PubMed: 21159959]
- Lopes da Silva F. Neural mechanisms underlying brain waves: from neural membranes to networks. *Electroencephalogr Clin Neurophysiol*. 1991; 79(2):81–93. [PubMed: 1713832]
- Maguire EA. Hippocampal involvement in human topographical memory: evidence from functional imaging. *Philos Trans R Soc Lond B Biol Sci*. 1997; 352(1360):1475–1480. [PubMed: 9368936]
- Makris N, Goldstein JM, Kennedy D, Hodge SM, Caviness VS, Faraone SV, Tsuang MT, Seidman LJ. Decreased volume of left and total anterior insular lobule in schizophrenia. *Schizophr Res*. 2006; 83(2–3):155–171. [PubMed: 16448806]

- Maren S, Fanselow MS. Synaptic plasticity in the basolateral amygdala induced by hippocampal formation stimulation in vivo. *J Neurosci.* 1995; 15(11):7548–7564. [PubMed: 7472506]
- Mitchell AJ, Shiri-Feshki M. Rate of progression of mild cognitive impairment to dementia-meta-analysis of 41 robust inception cohort studies. *Acta Psychiatr Scand.* 2009; 119(4):252–265. [PubMed: 19236314]
- Muller MJ, Greverus D, Weibrich C, Dellani PR, Scheurich A, Stoeter P, Fellgiebel A. Diagnostic utility of hippocampal size and mean diffusivity in amnesic MCI. *Neurobiol Aging.* 2007; 28(3):398–403. [PubMed: 16529847]
- Pa J, Boxer A, Chao LL, Gazzaley A, Freeman K, Kramer J, Miller BL, Weiner MW, Neuhaus J, Johnson JK. Clinical-neuroimaging characteristics of dysexecutive mild cognitive impairment. *Ann Neurol.* 2009; 65(4):414–423. [PubMed: 19399879]
- Pape HC, Pare D. Plastic synaptic networks of the amygdala for the acquisition, expression, and extinction of conditioned fear. *Physiol Rev.* 2010; 90(2):419–463. [PubMed: 20393190]
- Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med.* 2004; 256(3):183–194. [PubMed: 15324362]
- Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild cognitive impairment: clinical characterization and outcome. *Arch Neurol.* 1999; 56(3):303–308. [PubMed: 10190820]
- Petersen RC, Doody R, Kurz A, Mohs RC, Morris JC, Rabins PV, Ritchie K, Rossor M, Thal L, Winblad B. Current concepts in mild cognitive impairment. *Arch Neurol.* 2001; 58(12):1985–1992. [PubMed: 11735772]
- Petersen RC, Parisi JE, Dickson DW, Johnson KA, Knopman DS, Boeve BF, Jicha GA, Ivnik RJ, Smith GE, Tangalos EG, Braak H, Kokmen E. Neuropathologic features of amnesic mild cognitive impairment. *Arch Neurol.* 2006; 63(5):665–672. [PubMed: 16682536]
- Pitkanen A, Stefanacci L, Farb CR, Go GG, LeDoux JE, Amaral DG. Intrinsic connections of the rat amygdaloid complex: projections originating in the lateral nucleus. *J Comp Neurol.* 1995; 356(2):288–310. [PubMed: 7629320]
- Pitkanen A, Savander V, LeDoux JE. Organization of intra-amygdaloid circuitries in the rat: an emerging framework for understanding functions of the amygdala. *Trends Neurosci.* 1997; 20(11):517–523. [PubMed: 9364666]
- Ritchie K, Artero S, Touchon J. Classification criteria for mild cognitive impairment: a population-based validation study. *Neurology.* 2001; 56(1):37–42. [PubMed: 11148233]
- Roozendaal B, McEwen BS, Chattarji S. Stress, memory and the amygdala. *Nat Rev Neurosci.* 2009; 10:423–433. [PubMed: 19469026]
- Sarter M, Markowitsch HJ. Involvement of the amygdala in learning and memory: a critical review, with emphasis on anatomical relations. *Behav Neurosci.* 1985; 99(2):342–380. [PubMed: 3939663]
- Sauvage MM, Fortin NJ, Owens CB, Yonelinas AP, Eichenbaum H. Recognition memory: opposite effects of hippocampal damage on recollection and familiarity. *Nat Neurosci.* 2008; 11(1):16–18. [PubMed: 18037884]
- Schmidt-Wilcke T, Poljansky S, Hierlmeier S, Hausner J, Ibach B. Memory performance correlates with gray matter density in the ento-/perirhinal cortex and posterior hippocampus in patients with mild cognitive impairment and healthy controls—a voxel based morphometry study. *Neuroimage.* 2009; 47(4):1914–1920. [PubMed: 19442751]
- Schroeter ML, Stein T, Maslowski N, Neumann J. Neural correlates of Alzheimer's disease and mild cognitive impairment: a systematic and quantitative meta-analysis involving 1351 patients. *Neuroimage.* 2009; 47(4):1196–1206. [PubMed: 19463961]
- Serra L, Cercignani M, Petrosini L, Basile B, Perri R, Fadda L, Spano B, Marra C, Giubilei F, Carlesimo GA, Caltagirone C, Bozzali M. Neuroanatomical correlates of cognitive reserve in Alzheimer disease. *Rejuvenation Res.* 2011; 14(2):143–151. [PubMed: 21204647]
- Squire LR. Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans. *Psychol Rev.* 1992; 99(2):195–231. [PubMed: 1594723]
- Squire LR. Memory systems of the brain: a brief history and current perspective. *Neurobiol Learn Mem.* 2004; 82(3):171–177. [PubMed: 15464402]

- Stern CE, Corkin S, Gonzalez RG, Guimaraes AR, Baker JR, Jennings PJ, Carr CA, Sugiura RM, Vedantham V, Rosen BR. The hippocampal formation participates in novel picture encoding: evidence from functional magnetic resonance imaging. *Proc Natl Acad Sci USA*. 1996; 93(16): 8660–8665. [PubMed: 8710927]
- Turkeltaub PE, Eden GF, Jones KM, Zeffiro TA. Meta-analysis of the functional neuroanatomy of single-word reading: method and validation. *Neuroimage*. 2002; 16(3 Pt 1):765–780. [PubMed: 12169260]
- Van der Werf YD, Jolles J, Witter MP, Uylings HB. Contributions of thalamic nuclei to declarative memory functioning. *Cortex*. 2003; 39(4–5):1047–1062. [PubMed: 14584566]
- Visser PJ, Scheltens P, Verhey FR, Schmand B, Launer LJ, Jolles J, Jonker C. Medial temporal lobe atrophy and memory dysfunction as predictors for dementia in subjects with mild cognitive impairment. *J Neurol*. 1999; 246(6):477–485. [PubMed: 10431775]
- Zeineh NM, Engel SA, Thompson PM, Bookheimer Sy. Dynamics of the hippocampus during encoding and retrieval of face-name pairs. *Science*. 2003; 299(5606):577–580. [PubMed: 12543980]
- Zilles K, Palomero-Gallagher N, Grefkes C, Scheperjans F, Boy C, Amunts K, Schleicher A. Architectonics of the human cerebral cortex and transmitter receptor fingerprints: reconciling functional neuroanatomy and neurochemistry. *Eur Neuropsychopharmacol*. 2002; 12(6):587–599. [PubMed: 12468022]

Fig. 1.

Convergent areas of grey matter loss in MCI patients compared to healthy controls. Grey matter changes were found bilaterally in the amygdala/hippocampus region extending to the thalamus and medial temporal pole on the left side and to the parahippocampal gyrus on the right side. The third cluster was found accentuated on the right side in the parietal precuneus broadening to the posterior cingulate cortex

Fig. 2.
Voxel-wise correlation analysis with the MMSE scores revealed clusters of convergence in the right hippocampal/amygala region and the left thalamus

Table 1

Demographic and clinical characteristics of the included VBM studies

Study	Subjects	Sex (male)	Age (mean $\pm$ SD)	MMSE (mean $\pm$ SD)	Scanner	Gaussian kernel width (mm)
Barbeau et al. Neuropsychologia, 2008	MCI (28[28] ^d)	12	69.3 $\pm$ 8.6	27.4 $\pm$ 1.4	1.5 T S	6
	HC (28)	15	63.3 $\pm$ 7.2			
Bell-McGinty et al. Arch Neurol, 2005	MCI (37[9])	17	71.9 $\pm$ 7.6	25.5 $\pm$ 3.2	1.5 T GE	8
	HC (47)	27	66.9 $\pm$ 7.3			
Bozzali et al. Neurology, 2006	MCI (22[22])	10	70.5 $\pm$ 10.5	25.3 $\pm$ 1.6	1.5 T S	12
	HC (20)	7	65.8 $\pm$ 6.8			
Caroli et al. J Neurol, 2007	MCI (23[23])	13	70.0 $\pm$ 5.5	26.9 $\pm$ 1.9	1TP	12
	HC (17)	8	69.0 $\pm$ 3.0			
Chetelat et al. Neuroreport, 2002	MCI (22[22])	10	71.0 $\pm$ 8.0	27.3 $\pm$ 1.5	1.5 T GE	12
	HC (22)	10	66.6 $\pm$ 7.2			
Hamalainen et al. Neuroimage, 2007	MCI (56[56])	15	72.4 $\pm$ 4.2	23.6 $\pm$ 2.5	1.5 T S	8
	HC (22)	11	72.9 $\pm$ 4.5			
Hirata et al. Neurosci Lett, 2005	MCI (30[30])	NS	70.6 $\pm$ 8.4	26.0 $\pm$ 1.5	1TS	12
	HC (41)	NS	71.1 $\pm$ 7.7			
Jauhiainen et al. DGCD, 2008	MCI (14[7])	7	76.6 $\pm$ 1.7	25.7 $\pm$ 0.8	1.5 T S	12
	HC (13)	7	74.4 $\pm$ 1.2			
Morbelli et al. Eur J Nucl Med Mol Imaging, 2009	MCI (20[20])	8	75.6 $\pm$ 5.6	27.8 $\pm$ 1.2	1.5 T P	8
	HC (12)	2	70.4 $\pm$ 7.3			
Pa et al. Ann Neurol, 2009	MCI (58[26])	33	65.9 $\pm$ 7.2	28.8 $\pm$ 1.3	1.5 T S	12
	HC (36)	13	64.8 $\pm$ 8.2			
Pennanen et al. JNNP, 2005	MCI (51[51])	17	72.0 $\pm$ 5.0	24.0 $\pm$ 2.0	1.5 T S	12
	HC (32)	13	74.0 $\pm$ 4.0			
Rami et al. Int J Geriatr Psychiatry, 2009	MCI (16[6])	4	72.9 $\pm$ 4.8	26.0 $\pm$ 2.0	1.5 T GE	10
	HC (27)	10	74.3 $\pm$ 5.3			
Risacher et al. Curr Alzheimer Res, 2009	MCI (339[339])	214	74.7 $\pm$ 0.7	26.9 $\pm$ 0.2	1.5 T GE, P, S	10
	HC (206)	107	76.0 $\pm$ 0.5			
Saykin et al. Neurology, 2006	MCI (40[40])	23	72.9 $\pm$ 7.1	27.2 $\pm$ 2.2	1.5 T GE	12

Study	Subjects	Sex (male)	Age (mean $\pm$ SD)	MMSE (mean $\pm$ SD)	Scanner	Gaussian kernel width (mm)
Shiino et al. Neuroimage, 2006	HC (40)	12	71.0 $\pm$ 5.1			
	MCI (20[20])	10	67.7 $\pm$ 9.0	26.8 $\pm$ 1.9	1.5 T GE	12
	HC (88)	40	68.7 $\pm$ 8.7			
Stoub et al. PNAS, 2006	MCI (40[40])	16	77.9 $\pm$ 7.5	27.2 $\pm$ 1.6	1.5 T GE	12
	HC (50)	15	78.1 $\pm$ 6.0			
	MCI (15[15])	9	73.3 $\pm$ 6.7	27.8 $\pm$ 1.8	3TS	12
Trivedi et al. Alzheimers Dement, 2006	HC (15)	9	73.6 $\pm$ 7.1			
	MCI (14[14])	6	71.8 $\pm$ 7.3	26.6 $\pm$ 0.3	3 T GE	8
	HC (14)	6	70.4 $\pm$ 5.8			
Schmidt-Wilcke et al. Neuroimage, 2009	MCI (18[16])	10	65.7 $\pm$ 7.2	n.a.	1.5 T S	10
	HC (18)	10	63.0 $\pm$ 10.7			
	MCI (10[10])	5	72.7 $\pm$ 5.3	26.3 $\pm$ 2.9	1.5 T GE	8
Bonekamp et al. Neuroreport, 2010	HC (20)	10	75.3 $\pm$ 4.8			
	MCI (25[25])	8	73.6 $\pm$ 7.0	27.5 $\pm$ 1.4	3TS	8
	HC (13)	5	75.0 $\pm$ 5.0			
Guedj et al. Eur J Nucl Med Mol Imaging, 2009	MCI (19[19])	9	69.9 $\pm$ 9.5	27.1 $\pm$ 1.1	1.5 T S	6
	HC (28)	14	66.7 $\pm$ 6.9			

MCI mild cognitive impairment, *GR* general electrics, *HC* healthy controls, *MMSE* mini-mental state examination, *n.a.* not available, *P* Philips, *S* Siemens, *SD* standard deviation, *T* Tesla, *S* Siemens

^a[aMCI]

Table 2
Main findings of grey matter reductions in MCI patients relative to healthy controls and correlation with MMSE performance

Cluster	k _E	Side	MNI coordinates			Region
			x	y	z	
MCI < HC						
Cluster 1	2407	L	-22	-8	-22	Amygdala (LB); hippocampus (CA); thalamus; medial temporal pole
Cluster 2	1984	R	24	-8	-20	Amygdala (LB); hippocampus (FD); parahippocampal gyrus; hippocampus (EC)
Cluster 3	269	L/R	2	-54	32	Parietal precuneus; dorsal cingulate cortex
Correlation with MMSE						
Cluster 1	120	R	18	-4	-18	Amygdala (SF)
Cluster 2	33	L	-16	-24	8	Thalamus
Cluster 3	28	R	32	-14	-38	Hippocampus (EC)

MCI mild cognitive impairment, HC healthy controls, GM grey matter, MMSE mini-mental state examination, MNI Montreal Neurological Institute, Lat. laterality, R right, L left, LB laterobasal, FD fascia dentate, CA cornu ammonis, SF superficial, EC entorhinal cortex