CORRESPONDENCE

Animal and Human Bite Wounds

by Prof. Dr. med. Karin Rothe, Prof. Dr. med. Michael Tsokos, and Prof. Dr. med. Werner Handrick in issue 25/2014

Disastrous Progression

Surgical treatment, which is of central importance, was explained to an unsatisfactory degree (1). One should urgently advise against wound irrigation using button cannulas or infusion catheters (2). If a bite wound is such that this idea arises then this is merely an indication for “real” surgical debridement in the operating room and under general anesthesia. Half-hearted and inconsistent primary treatments often result in disastrous disease progression. The indication for primary surgical debridement should be defined generously. In my own practice it has proved successful to operate on any bite wound as a matter of principle and to permit exceptions when there are good reasons for doing so. Every patient with a bite wound on their hand should immediately be referred to a center for hand surgery; for facial bite wounds, institutions for plastic surgery are available. Necrosectomy is never limited by the anatomical situation but is mandatory in every case. Necroses have to be removed, even from the face (3).

It goes without saying that the necrosectomy should be done subtly, with the help of magnifying spectacles, in order to spare healthy tissue. Correct necrosectomy is the basis for the best possible healing result without too much scarring (4). The authors are correct to point out that patients and physicians often underestimate the seriousness of smaller injuries; this is especially the case for bite wounds to the hands. The unspeakable “faffing about” with loop drains threaded through infected tissue, preferably on the hand, is unfortunately still a common phenomenon.

DOI: 10.3238/arztebl.2016.0116a

REFERENCES

1. Rothe K, Tsokos M, Handrick W: Animal and human bite wounds. *Dtsch Arztebl Int* 2015; 112: 433–43.
2. Lichte P, Kobbe P, Taeger G, Nast-Kolb D, Hierner R, Oberbeck R: Bissverletzungen der Hand. *Unfallchirurg* 2009; 112: 719–27.
3. Rui-feng C, Li-song H, Ji-bo Z, Li-qui W: Emergency treatment on facial laceration of dog bite wounds with immediate primary closure: a prospective randomized trial study. *BMC Emergency Medicine* 2013; 13: 2.
4. Vogt M: Diagnostik und Therapie von Bissverletzungen durch Hunde, Katzen und Menschen. *Dtsch Med Wochenschr* 2003; 128: 1059–63.

Dr. med. Burkhard Heitmann
Chirurg, Unfallchirurg, Handchirurg
BG Unfallklinik Duisburg GmbH
burkhard.heitmann@bg-klinikum-duisburg.de

Helpless Wishful Thinking

I wish to criticize the fact that wound irrigation with infusion catheters or button cannulas, which are inserted

into the full depth of the bite wound in order to remove foreign bodies and pathogens (1) is described as an apparently common method and in such a neutral fashion. Only “irrigation under pressure is inadvisable”. It would have been much better to request immediate surgical necrosectomy for any bite wound, if needed after referral to a specialized hospital (2), and to present any cannulation or irrigation as no longer recommended.

It amounts to helpless wishful thinking to believe that injecting fluids with a syringe will help expel anything from the wound—especially not foreign bodies or pathogens. Injecting fluids will naturally spread the infection into tissue spaces, tendon sheaths, joint capsules, and so on. Even those in favor of irrigation admit this risk, and they warn against it without having any idea how it might be prevented (3, 4). Surgical debridement has been confirmed as a sensible option, but irrigation never has. Almost all patients who present with purulent bite wounds after 3–4 days for the purposes of revision in a scenario of manifest infection have previously received irrigation.

Let us finally put a stop to half-hearted irrigation.

DOI: 10.3238/arztebl.2016.0116b

REFERENCES

1. Rothe K, Tsokos M, Handrick W: Animal and human bite wounds. *Dtsch Arztebl Int* 2015; 112: 433–43.
2. Rieck B: Hundebissverletzungen beim Kind. *Behandlung und Nachsorge. Monatsschrift Kinderheilkunde* 2005; 8: 789–95.
3. Goldstein EJC: Bite wounds and infection. *Clin Infect Dis* 1992; 14: 633–40.
4. Stucker FJ, Shaw GY, Boyd S, et al.: Management of animal and human bites in the head and neck. *Otolaryngol Head Neck Surg* 1990; 116: 789–93.

Prof. Dr. med. Bernd Rieck
Klinik für Plastische, Ästhetische und Handchirurgie
Hildesheim
Bernd.Rieck@helios-kliniken.de

In Reply:

The incidence of bite injuries has been increasing in tandem with the growing population of house pets, to the point where it has become a global public health problem. Views on the diagnostic evaluation and therapy of such injuries are divergent in the literature, and there are hardly any prospective controlled studies.

The discussion has focused on the risk of infection, which depends on the biting animal on the one hand and on the bite victim on the other hand. The timing of the initial visit to the doctor contributes vitally to the risk of infection. Timely presentation and correct wound care have helped lower the rates of infection notably. Bite wounds require interdisciplinary approaches (surgeon / pediatric surgeon, infectious disease specialist, clinical microbiologist, intensive care doctor if required, plastic surgeon, hand surgeon).

The article intended to point out the need for structured management. It is an undisputable fact that bite wounds are often underestimated and that only infected wounds reach the specialist. I thank Heitmann and Rieck for agreeing in their critiques of the article that optimal surgical management and specialist care are of crucial importance. Necrosectomy, mechanical reduction of the number of pathogens, and optimization of the microcirculation in the wound area are the cornerstones of surgical therapy.

I wish to comment on the criticisms of irrigation treatment only in as far as by reminding readers that cleaning and irrigating wounds are among the generally accepted measures of wound treatment, and that contaminated and infected wounds in particular benefit from lavage or irrigation for the purposes of reducing pathogens. Wound irrigation is obviously not a substitute for necrosectomy. The use of infusion catheters or button cannulas is probably unnecessary and should only be undertaken if the wound drainage is satisfactory. Unfortunately, to date no surgical specialty society has developed guidelines for the surgical management of acute wounds. Kramer et al suggest—from the

perspective of the German Society for Hospital Hygiene—combined management consisting of debridement and antiseptic lavage, depending the wound's age and type (1).

Our article was intended to encourage all practicing colleagues to refer patients with bite wounds early and liberally for specialist, and where required interdisciplinary, care, in order to minimize functional deficits as far as possible (2).

DOI: 10.3238/arztebl.2016.0116c

REFERENCES

1. Kramer A, Frank AM, Becker C, Hinz P: Prävention postoperativer Wundinfektionen im Rahmen der chirurgischen Versorgung von Bissverletzungen. GMS Krankenhhyg Interdisziplinär 2010; 5: 1–14.
2. Rothe K, Tsokos M, Handrick W: Animal and human bite wounds. Dtsch Arztebl Int 2015; 112: 433–43.

Prof. Dr. med. Karin Rothe
Charité – Universitätsmedizin Berlin
Klinik und Poliklinik für Kinderchirurgie
karin.rothe@charite.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.