Published in final edited form as:

Org Lett. 2015 November 6; 17(21): 5468–5471. doi:10.1021/acs.orglett.5b02832.

Highly Diastereoselective Michael Reactions Using β -Nitrocarbonyl Nucleophiles

Amber A. S. Gietter-Burch, Roxana E. Mitrut, and Donald A. Watson*

Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19716, USA

Abstract

We have discovered a highly diastereoselective Michael reaction of α -substituted, β -nitrocarbonyl compounds to deliver highly functionalized stereo-diads containing fully substituted nitrogen-bearing centers. Good to excellent yields and diastereoselectivities are observed. This transformation is tolerant of various types of carbonyl groups on the nucleophilic partner, as well as range of unsaturated electrophiles. Mechanistic investigations are consistent with internal hydrogen-bonding in the nitroalkane tautomer as the major factor in the control of diastereoselectivity in these transformations.

Graphical abstract

Nitroalkanes are highly versatile intermediates in organic synthesis.¹ These compounds are able to undergo a variety of carbon-carbon bond forming reactions such as arylations,² allylations,³ Henry reactions,¹ and conjugate additions.¹ Nitroalkanes also serve as starting materials for the installation of a variety of other functional groups, such as amines, ketones, and alkanes.¹

Recently, as a method for rapidly preparing complex nitroalkanes, our group has developed several copper-catalyzed procedures for the C-alkylation of nitroalkanes using simple alkyl halide electrophiles.^{4,5,6} In the course of these studies, we developed conditions for the preparation of β -nitrocarbonyl compounds via the alkylation of nitroalkanes with α -bromocarbonyls.^{7,8} Although these reactions proceed in excellent yields with good functional group tolerance, diastereoselectivity in the reactions was modest (ca. 66:33).

*Corresponding Author: dawatson@udel.edu.

Supporting Information

Experimental procedures, NMR studies, and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes

The authors declare no competing financial interest.

(1)

In an effort to demonstrate the utility of these products in downstream synthesis, we previously reported a single example using β -nitroamide **1** as the nucleophile in a Michael reaction with methyl acrylate (eq 1).⁷ To our surprise, despite that fact that a 63:37 diastereomeric mixture of the β -nitroamide was brought into the reaction, a single detectable diastereomer of product **3** emerged from the reaction in high yield. The relative stereochemistry of the product was determined by X-ray crystallography, after reduction to the corresponding amine. No examples of diastereoselective Michael additions of this type have been previously reported. In fact, very few alkylation reaction of β -nitrocarbonyl compounds of any type have ever been described.^{9,10}

We recognized these Michael adducts could potentially be of interest for the synthesis of complex nitrogen-containing molecules, as the products contain a fully substituted nitrogen-bearing stereo-center in a highly functionalized environment. As such, we chose to investigate the generality of this Michael reaction.

Herein we report that good to excellent levels of diastereoselectivity are observed using a range of β -nitrocarbonyl nucleophiles and several different classes of Michael acceptors. In addition, we briefly investigated the mechanism of this reaction, allowing us to offer a model to rationalize the observed diastereoselectivity.

Before exploring the generality of the transformation, we wanted to investigate the reaction conditions for the Michael reaction. To do so, the reaction of β -nitro-Weinreb amide **4** with excess (3 equiv) methyl acrylate (**2**) was investigated (Table 1). We were pleased to find that high yield and high diastereoselectivity were also observed in this system (entry 1), which was comparable to the previously investigated diethylamide substrate (**1**). We found that the reaction could also be run in other polar solvents such as acetonitrile (entry 2), but was less efficient in less polar solvents such as dichloromethane (entry 3). Acetonitrile also proved easier to remove from the products and was therefore selected for further studies. We also found the reaction to be relatively rapid, affording nearly quantitative yields in just one hour (entry 4). A variety of other mild organic and inorganic bases were also examined, but none were as effective as DBU (see Supporting Information). Finally, attempts to lower the equivalents of methyl acrylate used in the reaction led to lower yields (entry 5). In all cases, however, high diastereoselectivity (>95:5) was observed.

With the optimized conditions in hand, we began to examine the scope of the reaction with respect to the β -nitrocarbonyl nucleophile. A variety of carbonyl moieties were well tolerated under the reaction conditions (Scheme 1). The model Weinreb amide product **5** was isolated in 89% yield. Substrates bearing other tertiary amides (e.g., cyclic amide **6**)

also participated equally well in the Michael reaction, providing high yield and diastereoselectivity. Secondary amides (**7**) also participated, but with slightly eroded levels of diastereoselectivity.

β -Nitroesters can also be used in this method. Substrates bearing linear alkyl groups α to the ester were well tolerated, leading to high yielding, highly diastereoselective reactions (**8–10**). Branched α -substituents, however, resulted in both lower yield and selectivity, reflecting the greater steric demands of the nucleophile (**11**). Somewhat surprisingly, even β -nitroketones proved to be competent starting materials in the reaction, providing reasonable yield and diastereoselectivity (**12**).

In the case of the ketone substrate (as well as some esters) significant levels of byproducts resulting from denitration of the starting materials were observed under the standard reaction conditions. These alkenes presumably arise via the E1CB elimination of an equivalent of nitrous acid from these more acidic starting materials. This side reaction, was suppressed using the more sterically encumbered base 1,1,3,3-tetramethylguanidine (TMG) and lowering the reaction temperature (see Scheme 1 and 2). Notably, the denitration reaction appears to be sensitive to steric environment; no byproducts from the denitration of the Michael adducts have been observed.¹¹

We also examined the scope of the transformation with respect to Michael acceptors (Scheme 2). Other acrylate esters such as benzyl acrylate (**13** and **14**) afforded high yield and excellent diastereoselectivity.

Less electrophilic acrylamide derivatives (**15** and **16**) also afforded good yields, but required slightly elevated temperatures and longer reaction times (40 °C, 2 h). In addition, when using acrylamide derivatives, increasing the steric bulk at the alpha position of the β -nitrocarbonyl had a profound effect on the reactivity. When substitution was larger than methyl, the reaction did not proceed (**16** vs. **17**). Attempts to circumvent this steric limitation were unsuccessful; increasing the temperature afforded increased elimination of the nitro group from the starting material and variation of the base led to poor conversion. Vinyl ketones (**18**) and sulfones (**19**) proved to be excellent and highly selective substrates. Unsaturated nitriles also led to high yielding reactions with respectable levels of stereoselectivity (**20**) These products provide a variety of functional handles for downstream synthetic manipulation. With acrolein, good diastereoselectivity was observed (**21**); however, the formation of polymer byproducts severely suppressed the yield, even at cryogenic temperatures (–40 °C, 24 h).

We also briefly investigated more substituted Michael acceptors. Those incorporating internal cyclic alkenes, such as cyclopentenone, provided reasonable reactivity with non-sterically demanding β -nitrocarbonyls. However, stereocontrol in these systems was poor (**22**). With somewhat more sterically demanding nucleophiles, no product was observed (**23**). Likewise, acyclic Michael acceptors bearing either α - or β -substituents failed in the reaction. For example, product **24** was not observed.

As shown in Schemes 1 and 2, the reaction demonstrates excellent functional group tolerance. In addition to the functional groups described above, aryl chlorides (**7**), aryl bromides (**8**), silyl ethers and acyl protected alcohols (**10**), and distal esters (**14**) all afford excellent yields with high diastereoselectivity of the desired products.

We next sought to examine the origin of the diastereoselectivity observed in the Michael reaction. First, we established that the dr of the product is independent of that of the starting material by using diastereomerically-enriched¹² samples of β -nitroamide **4** (Scheme 3).

Next, we examined the thermodynamics of the reaction of DBU and β -nitroamide **4** in d₃-MeCN at rt (eq 2). No nitroanate anion, and only free DBU and **4**, were observed by ¹H NMR, even after extended reaction times. This suggests that any deprotonation event is endothermic under the reaction conditions.

(2)

Finally, to understand the kinetics of deprotonation of the β -nitrocarbonyl starting materials, we studied the epimerization of β -nitroamide **4** in the presence of DBU in d₃-MeCN at rt using diastereomerically enriched samples. The reactions were monitored by ¹H NMR. Independent of the diastereomeric ratio of the starting sample, compound **4** equilibrates to an approximate 60:40 ratio of diastereomers within 10 minutes (Scheme 4 and Supporting Information). This timeframe is much faster than the time course of the Michael reaction, which requires approximately 1 h to complete, suggesting that deprotonation is also rapid under these conditions.

Given these observations, a possible rationale for the observed diastereoselectivity is outlined in Figure 6. Rapid, reversible deprotonation of the nitroalkane could establish a small concentration of the highly nucleophilic tautomer **25**. Intramolecular hydrogen-bonding to the adjacent carbonyl would organize compound **25** as shown. From this intermediate, the Michael acceptor would be expected to react at the face away from the alkyl group alpha to the carbonyl, as shown. This model is consistent with the observed stereochemistry of the products.

This model is also consistent with the observation that the highest levels of diastereoselectivity are observed with the substrates bearing the most basic carbonyl groups (see Scheme 1 and 2). For example, while nearly all of the substrates examined react with high levels of diastereoselectivity, substrates bearing tertiary amides are slightly more selective than those with esters. This trend correlates well to the carbonyls ability to participate in hydrogen bonding.

In conclusion, we have demonstrated that Michael additions involving α -chiral, β -nitrocarbonyl derivatives as nucleophiles can be highly diastereoselective and efficient. This mild method works with a range of nucleophiles of this type, as well as a range of Michael-acceptors. In addition, the mildness of the method allows broad functional group tolerance. The products from these reactions are highly functionalized stereo-diads containing a fully-substituted, nitrogen-bearing stereocenter. These products should be of significant utility as building blocks in the synthesis of more complex molecules.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The University of Delaware, the University of Delaware Research Foundation, the Research Corp. Cottrell Scholars Program, and the NIH NIGMS (R01 GM102358) are gratefully acknowledged for support. A.A.S.G.-B. thanks the University of Delaware for a graduate fellowship. R.E.M. acknowledges a REACT Fellowship supported by the NSF (CAREER CHE-1254360). Data was acquired at UD on instruments obtained with the assistance of NSF and NIS funding (NSF CHE0421224, CHE1229234, and CHE0840401; NIH P20 GM103541, S10 RR02692, and S10 OD016267).

References

1. Ono, N. *The Nitro Group In Organic Synthesis*. John Wiley And Sons; New York: 2001.
2. (a) Dey C, Lindstedt E, Olofsson B. *Org Lett*. 2015; 17:4554. [PubMed: 26352796] (b) Vogl EM, Buchwald SL. *J Org Chem*. 2002; 67:106. [PubMed: 11777446] (c) Walvoord RR, Berritt S, Kozlowski MC. *Org Lett*. 2012; 14:4086. [PubMed: 22839593] (d) Walvoord RR, Kozlowski MC. *J Org Chem*. 2013; 78:8859. [PubMed: 23895411]
3. (a) Aleksandrowicz P, Piotrowska H, Sas W. *Tetrahedron*. 1982; 38:1321.(b) Maki K, Kanai M, Shibasaki M. *Tetrahedron*. 2007; 63:4250.(c) Trost BM, Surivet J-P. *Angew Chem Int Ed*. 2000; 39:3122.(d) Tsuji J, Yamada T, Minami I, Yuhara M, Nisar M, Shimizu I. *J Org Chem*. 1987; 52:2988.(e) Wade PA, Morrow SD, Hardinger SA. *J Org Chem*. 1982; 47:365.
4. Gildner PG, Gietter AAS, Cui D, Watson DA. *J Am Chem Soc*. 2012; 134:9942. [PubMed: 22691127]
5.
For another recent nitroalkane alkylation reaction, see:

Petruzziello D, Gualandi A, Grilli S, Cozzi PG. *Eur J Org Chem*. 2012; 2012:6697.
6.
For early papers on nitroalkane alkylation, see:

(a) Seebach D, Lehr F. *Angew Chem Int Ed Engl*. 1976; 15:505.(b) Seebach D, Henning R, Lehr F, Gonnermann J. *Tetrahedron Lett*. 1977; 18:1161.(c) Katritzky AR, De Ville G, Patel RC. *J Chem Soc, Chem Commun*. 1979; 602(d) Russell GA, Hershberger J, Owens K. *J Am Chem Soc*. 1979; 101:1312.(e) Katritzky AR, Kashmiri MA, De Ville GZ, Patel RC. *J Am Chem Soc*. 1983; 105:90.(f) Russell GA, Khanna RK. *Tetrahedron*. 1985; 41:4133.(g) Branchaud BP, Yu G-X. *Tetrahedron Lett*. 1988; 29:6545.
7. Gietter AAS, Gildner PG, Cinderella AP, Watson DA. *Org Lett*. 2014; 16:3166. [PubMed: 24870052]
8.
For other methods for preparing these compounds, see:

(a) Kornblum N, Boyd SD. J Am Chem Soc. 1970; 92:5784.(b) Miyakoshi T, Saito S, Kumanotani J. Chem Lett. 1981; 10:1677.(c) Russell GA, Ros F. J Am Chem Soc. 1982; 104:7349.(d) Kunetsky RA, Dilman AD, Tsvaygboym KP, Ioffe SL, Strelenko YA, Tartakovsky VA. Synthesis. 2003; 1339(e) Wilson JE, Casarez AD, MacMillan DWC. J Am Chem Soc. 2009; 131:11332. [PubMed: 19627154]

9.

For previous Micheal reactions, see:

(a) Alvarez-Ibarra C, Csáký AG, Gómez de la Oliva C. J Org Chem. 2000; 65:3544. [PubMed: 10843645] (b) Han Y, Liu Y, Rockenbauer A, Zweier JL, Durand Gg, Villamena FA. J Org Chem. 2009; 74:5369. [PubMed: 19530689]

10.

For Henry-type reactions, see:

(a) Sacher JR, Weinreb SM. Tetrahedron. 2011; 67:10203. [PubMed: 22125345] (b) Ji C-B, Liu Y-L, Cao Z-Y, Zhang Y-Y, Zhou J. Tetrahedron Lett. 2011; 52:6118.

11.

In some cases, we did observe a small amount (<5%) of a byproduct that results from a complex reaction sequence that includes denitration, Dieckmann condensation, and multiple alkylation steps. Re-subjection of isolated Michael adducts to the reaction conditions, however, did not result in this byproduct, suggesting that it is formed via a unique pathway that is independent of the Michael reaction. Details and characterization of the side product are provided in the Supporting Information.

12.

Obtained via chromatographic separation on silica gel.

Figure 6.
Possible Model for Observed Diastereoselectivity

Scheme 1.
Scope with Respect to β -Nitrocarbonyls

Scheme 2.
Scope with Respect to Michael Acceptor

Scheme 3.

Product Diastereomeric Ratio is Not a Function of Starting Material Stereochemistry

Scheme 4.
Epimerization Studies of **4**.

Table 1

Optimization of Reaction Conditions

<p>4, dr: 63:37</p>					
entry	solvent	time (h)	equiv 2	yield 5 (%) ^a	dr ^a
1	DMF	5	3	96	>95:5
2	MeCN	5	3	96	>95:5
3	CH ₂ Cl ₂	5	3	39	>95:5
4	MeCN	1	3	95	>95:5
5	MeCN	1	1.5	71	>95:5

^aYield and diastereoselectivity determined by NMR using 1,3,5-trimethoxybenzene as an internal standard.