Published in final edited form as:

Angew Chem Int Ed Engl. 2014 September 1; 53(36): 9462–9465. doi:10.1002/anie.201403607.

Catalytic Enantioselective Carboannulation with Allylsilanes

Nicolas R. Ball-Jones, Dr. Joseph J. Badillo, Ngon T. Tran, and Prof. Annaliese K. Franz
Department of Chemistry, University of California, One Shields Avenue, Davis, CA 95616 (USA)

Annaliese K. Franz: akfranz@ucdavis.edu

Abstract

The first catalytic asymmetric carboannulation with allylsilanes is presented. The enantioselective [3+2] annulation is catalyzed using a Sc(III)-indapybox complex with tetrakis-[3,5-bis(trifluoromethyl)phenyl]-borate (BARF) to enhance catalytic activity and control stereoselectivity. Functionalized cyclopentanes containing a quaternary carbon are derived from alkylidene oxindole, coumarin, and malonate substrates with high stereoselectivity. The enantioselective 1,4-conjugate addition and enantioselective lactone formation (via trapping of the β -silyl carbocation) is also described.

Keywords

asymmetric catalysis; cyclopentane; NaBARF; scandium; conjugate addition

Lewis acid-catalyzed conjugate addition and cyclization reactions of unsaturated carbonyl compounds are important routes for the synthesis of complex heterocycles and carbocycles. Despite significant advances in asymmetric catalysis,^[1,2] many conjugate addition and cyclization reactions using unsaturated carbonyl compounds still lack catalytic asymmetric variants. First reported by Knölker in 1990,^[3] the cyclopentane annulation of air- and moisture-stable allylsilane nucleophiles with unsaturated carbonyl compounds is a transformation where a catalytic asymmetric variant has eluded development.^[4-6] The challenge associated with controlling additions to unsaturated carbonyl compounds is highlighted by the fact that only one method has been reported for a catalytic enantioselective conjugate addition reaction using allylsilanes (Figure 1A).^[7] Herein, we report the first catalytic asymmetric annulation of allylsilanes with unsaturated carbonyl compounds (Figure 1B) to access cyclopentanes possessing up to three stereocenters, including a quaternary carbon spirocenter.

The reaction of alkylidene oxindole **1** with allylsilane **2a**^[8] offers a platform to study the reactivity and selectivity for the annulation reaction (Table 1). The selective formation of a cyclic product (e.g. **4**) or an allylation (Hosomi-Sakurai) product (e.g. **5**) is determined based on control of the β -silyl-stabilized carbocation intermediate (e.g. **3**).^[9,10] Selective formation of spirocyclopentane **4**, resulting from rearrangement of the β -silyl carbocation intermediate,

is a critical component of the successful development of this methodology. Our initial studies with chiral tin(IV),^[11] copper(II),^[7] and scandium(III)[5a,12] complexes afforded only low selectivity and/or low reactivity for the synthesis of spirocyclopentane **4a** (e.g. entries 1-4).^[13]

To identify a more effective catalyst, we investigated Sc(III)-pybox complexes with NaBARf [BARf = B(3,5-C₆H₃(CF₃)₂)₄] (Table 1, entries 5-9).^[14] A significant activating effect was observed for both ScCl₃- and Sc(OTf)₃-derived complexes, affording spirocyclopentane **4a**^[15] with high yield and up to 97:3 enantioselectivity (entries 5 and 6). Using other counterions such as NaSbF₆ and KPF₆, or additives such as TMSCl, did not enhance reactivity (Table S1). The amount of NaBARf was also observed to play a significant effect on the enantioselectivity where increasing the NaBARf from 5 to 50 mol % lowered the rate of the reaction and afforded racemic product **4a** (entry 7). It was also observed that the indapybox ligand plays a critical role in diastereocontrol: in the absence of ligand, the annulation proceeded with low diastereoselectivity (65:35 dr, entry 8). Evaluation of various ligands showed that (*R,S*)-indapybox (**L2**) affords with the highest diastereo- and enantioselectivity (Table S1).

The primary activating effect of NaBARf is attributed to formation of the cationic scandium complex.^[12] Formation of a cationic Sc(OTf)₂BARf•(*R,S*)-indapybox complex is supported by isolation of NaOTf from the precipitate (63% yield based on FAAS and ¹⁹F NMR analysis).^[16] The erosion of enantioselectivity with increasing NaBARf may be due to formation of Sc(BARf)₃ or Sc(BARf)₂X species; however, investigation of a preformed Sc(BARf)₃ species did not show comparable erosion (entry 9).^[17] Analysis of the Sc(OTf)₂BARf•(*R,S*)-indapybox complex using ¹H, ¹⁹F and ⁴⁵Sc NMR spectroscopy indicates a dynamic catalyst complex that is rapid and reversible on the NMR time scale. Based on literature precedent for carbocation stabilization, we also hypothesize that NaBARf can play a secondary role to facilitate formation of the transient β-silyl carbocation (e.g. **3**).^[18]

The Sc(III)-pybox/NaBARf system is also effective for the catalytic enantioselective 1,4-conjugate addition of allyltrimethyl-silane (**2b**) to afford allylation product **5** (eq 1). Even using a small silyl group such as TMS, these conditions still afford spirocyclopentane **4b**, which suggests that conditions using NaBARf suppress silyl elimination and/or promote 1,2-silyl migration of the β-silyl carbocation.

(1)

With this optimal catalyst system, we demonstrated that the spirocyclopentane annulation proceeds with excellent yields, diastereo- and enantioselectivity for a variety of alkylidene oxindole substrates (Table 2). Various ester (**1c-1g**) and nitrile (**1h**) substrates proceed efficiently. Phenyl-substituted alkylidene **1i** required higher catalyst loading and extended reaction times, affording spirocycle **4i** with high diastereoselectivity and moderate enantioselectivity. Chelating oxindoles containing urea and Cbz groups (**1j** and **1k**) also afford product **4** in excellent yield and enantioselectivity.^[19]

The NH spirooxindoles, such as **4l** and **4m**, can be accessed by simple deprotection of the N-Acyl oxindoles with KHCO_3 and H_2O_2 in high yield (80-88% yield over two steps). The silyl-substituted spirooxindoles can be oxidized using modified Tamao-Fleming conditions to generate hydroxy-substituted spirooxindoles (e.g. **6n**) with retention of stereochemistry.^[21]

Rate enhancement observed with ester-substituted alkylidene **1d** compared to phenyl-substituted alkylidene **1i** (45 min vs 4 d) suggests that the reaction is promoted by coordination of the proximal ester group with the transient β -silyl carbocation (e.g. **7**, Table 3). Using *tert*-butyl substituted oxindole **1f** leads to trapping of the transient β -silyl carbocation and isolation of lactone **8f**.^[15,22] The yield of lactone **8** is influenced by the use of chloride vs triflate salt (53 vs 18% yield, Table 3), which is attributed to Lewis acid initiated deprotection of the *tert*-butyl ester. The reaction of **1f** using excess NaBARF (60 mol %) promotes formation of allylation product **5f** with 95% conversion (in 48 h).

The enhanced catalytic activity with NaBARF and the reactivity trends observed with alkylidene oxindoles translate to malonate- and coumarin-derived electrophiles **9** and **11**, which broadens access to functionalized carbocycles containing a quaternary carbon (eqs 2 and 3).^[23] Allylsilane annulation followed by oxidation afforded carbocycle **10** in high diastereoselectivity, but with low enantioselectivity (eq 2). Coumarin-derived cyclopentane **11**, a carbocycle not previously accessed using [3+2] annulation methodology, can also be accessed with high diastereoselectivity (eq 3). While oxindole derivatives are highly enantioselective, malonate and coumarin derivatives eluded asymmetric induction.

(2)

(3)

In summary, we have developed the first catalytic asymmetric [3+2] annulation of unsaturated carbonyl compounds with allylsilanes, giving spirocyclopentane core structures in high yields, diastereo- and enantioselectivities. Significant ligand and counterion effects are observed to control the path of the transient β -silyl carbocation (annulation vs allylation vs lactone), as well as the diastereo- and enantioselectivity. The annulation methodology can be applied to malonate- and coumarin-derived electrophiles with high yields and diastereoselectivities. More detailed mechanistic and computational studies are underway to elucidate the factors controlling the stereoselectivity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank Prof. Dean Tantillo (UCD) and Dr. Gonzalo Jiménez-Osés (UCLA) for helpful discussions, and Prof. Mark Mascal and Dr. Gorkem Gunbas (UCD) for a generous donation of NaBARF. This research is supported by donors of the American Chemical Society Petroleum Research Fund and NIH/NIGMS (P41-GM0089153). A.K.F. acknowledges 3M Corporation for a Nontenured Faculty Award, N.R.B.-J. and N.T.T. acknowledge the Department of Education for a GAANN fellowship, J.J.B. acknowledges the NSF for a graduate research fellowship, and N.T.T. acknowledges UC Davis for a Bradford Borge Chemistry Fellowship.

References

- Howell GP. *Org Process Res Dev.* 2012; 16:1258.
- a) Christoffers J, Koripelly G, Rosiak A, Rössle M. *Synthesis.* 2007; 2007:1279.b) Evans DA, Rovis T, Johnson JS. *Pure Appl Chem.* 1999; 71:1407.
- Knölker HJ, Jones PG, Pannek JB. *SynLett.* 1990; 7:429.
- For diastereoselective examples, see: Danheiser RL, Dixon BR, Gleason RW. *J Org Chem.* 1992; 57:6094.; Panek JS, Jain NF. *J Org Chem.* 1993; 58:2345.; Knölker HJ, Foitzik N, Schmitt O. *Tetrahedron Lett.* 1999; 40:3557.; Knölker HJ, Jones PG, Graf R. *Synlett.* 1996; 11:1155..
- a) Hanhan NV, Ball-Jones NR, Tran NT, Franz AK. *Angew Chem.* 2012; 124:1013. For the catalytic asymmetric allylsilane [3+2] annulation with isatins, see: *Angew Chem Int Ed.* 2012; 51:989.b) Matsumura Y, Suzuki T, Sakakura A, Ishihara K. *Angew Chem.* 2014:n/a. During review of this manuscript, the first catalytic asymmetric allylsilane [4+2] annulation with β,γ -unsaturated α -ketoesters was reported, see: *Angew Chem Int Ed.* 2014:n/a.
- Allylsilane precursors have been used to generate trimethylenemethane nucleophile, see: Trost BM, Cramer N, Silverman SM. *J Am Chem Soc.* 2007; 129:12396. [PubMed: 17880222] .
- Shizuka M, Snapper ML. *Angew Chem.* 2008; 120:5127.*Angew Chem Int Ed.* 2008; 47:5049.

8. Organ MG, Dragan V, Miller M, Froese RDJ, Goddard JD. J Org Chem. 2000; 65:3666. [PubMed: 10864750]
9. Brook, MA. Silicon in Organic, Organometallic, and Polymer Chemistry. John Wiley & Sons; New York: 2000. p. 521-523.
10.
For reviews on allylsilane reactivity, see: Masse CE, Panek JS. Chem Rev. 1995; 95:1293.; Chabaud L, James P, Landais Y. Eur J Org Chem. 2004; 2004:3173.; Schmidt AW, Knölker HJ. Synlett. 2010; 15:2207..
11. Kieffer ME, Repka LM, Reisman SE. J Am Chem Soc. 2012; 134:5131. [PubMed: 22390403]
12. a) Evans DA, Wu J, Masse CE, MacMillan DWC. Org Lett. 2002; 4:3379. [PubMed: 12323023] b) Evans DA, Fandrick KR, Song HJ, Scheidt KA, Xu R. J Am Chem Soc. 2007; 129:10029. [PubMed: 17658808]
13.
Other Lewis acidic metal complexes (e.g. In, Bi, Zn, Cu) are ineffective as catalysts. See Tables S1-S4.
14. a) Nishida H, Takada N, Yoshimura M, Sonoda T, Kobayashi H. Bull Chem Soc Jpn. 1984; 57:2600.b) Krossing I, Raabe I. Angew Chem. 2004; 116:2116.Angew Chem Int Ed. 2004; 43:2066.
15.
The structure and absolute stereochemistry of **4m** was assigned and confirmed based on X-ray crystallography (anomalous dispersion method). The relative stereochemistry of lactone **8f** was assigned by X-ray analysis. Absolute stereochemistry of **8f** was assigned by analogy to structure **4m**. CCDC 958776 (**4m**) and CCDC 961076 (**8f**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
16. a) Brookhart M, Grant B, Volpe J. Organometallics. 1992; 11:3920.b) Arndtsen BA, Bergman RG. Science. 1995; 270:1970.
17.
See table S5 for details. Also see: France S, Shah MH, Weatherwax A, Wack H, Roth JP, Lectka T. J Am Chem Soc. 2005; 127:1206. [PubMed: 15669860] .
18.
For an example of NaBARf promoting formation of β -silyl carbocations, see: Lambert JB, Zhao Y, Wu H. J Org Chem. 1999; 64:2729. [PubMed: 11674342] . For an example of NaBARf allowing carbocation isolation, see: Bahr SR, Boudjouk P. J Org Chem. 1992; 57:5545..
19.
A chelating electrophile is required for enantiocontrol: *N*-methyl substrate **4o** afforded racemic product in 60% conversion.
20.
See Table S7 for details of deprotection conditions.
21. a) Peng ZH, Woerpel KA. Org Lett. 2001; 3:675. [PubMed: 11259034] b) Peng ZH, Woerpel KA. Org Lett. 2000; 2:1379. [PubMed: 10814452]
22.
For examples of heterocycle formation via trapping of carbocations, see: Yamazaki S, Ohmitsu K, Ohi K, Otsubo T, Moriyama K. Org Lett. 2005; 7:759. [PubMed: 15727434] ; Shupe BH, Allen EE, MacDonald JP, Wilson SO, Franz AK. Org Lett. 2013; 15:3218. [PubMed: 23758331] ; Ollevier T, Li Z. Adv Synth Cat. 2009; 351:3251..
- 23.

The relative stereochemistry of cyclopentanes 10 and 12 was assigned by analogy to literature reports, see: Knölker HJ, Foitzik N, Goesmann H, Graf R, Jones PG, Wanzl G. Chem Eur J. 1997; 3:538..

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

A. Snapper, 2008

Only report of a catalytic asymmetric Hosomi-Sakurai conjugate addition

B. This work

1st catalytic asymmetric [3+2] carboannulation with allylsilane

Figure 1. Catalytic asymmetric additions of allylsilanes to α,β -unsaturated carbonyl compounds

Table 1
Optimization of the allylsilane annulation with alkylidene oxindoles

entry	conditions	yield (%) ^[a]	dr ^[b]	er ^[c]
1	SnCl ₄ /(<i>S</i>)-BINOL (20 mol %), 24h	75 ^[b]	50:50 ^[d]	70:30
2	Cu(OTf) ₂ /(<i>S</i>)-PhBox/NaBARF (10 mol %), 168h	<5	nd	nd
3	ScCl ₃ /L2/NaSbF ₆ (10 mol %), 72h	<5	nd	nd
4	ScCl ₃ /L2/NaSbF ₆ , TMSCl ^[e]	<10 ^[b]	nd	nd
5	ScCl₃/L2/NaBARF	99	99:1	97:3
6	Sc(OTf) ₃ /L2/NaBARF	94	95:5	92:8
7	ScCl ₃ /L2 with 50 mol % NaBARF	73	95:5	56:44
8	ScCl ₃ /NaBARF	95 ^[b]	65:35	--
9	Sc(BArF) ₃ /L2 (20 mol %)	100 ^[b]	95:5	84:16

All reactions performed under argon with 3.0 equiv of allylsilane in a 1:1.1:1 ratio of ScCl₃(THF)₃, (*R,S*)-indapybox, and NaBARF for 1h at 5 mol % catalyst loading (unless otherwise noted).

^[a] isolated yield.

^[b] Determined using ¹H NMR spectroscopy.

^[c] Determined using chiral-phase HPLC.

^[d] Allylation product **5** was also observed.

^[e] 3 equivalents of TMSCl were used.

Table 2
Optimization of the allylsilane annulation with alkylidene oxindoles

4c 82% yield 97:3 dr 96:4 er	4d 92% yield 95:5 dr 98:2 er	4e 75% yield ^c 98:2 dr 98:2 er
4f 80% yield 85:10:5 dr 99.5:0.5 er	4g 95% yield 90:10 dr 98:2 er	4h 83% yield 86:7:7 dr 88:12 er
4i 72% yield ^d 92:8 dr 84:16 er	4j 97% yield 94:6 dr 99:1 er	4k 83% yield 99:1 dr 98:2 er
4l 80% yield ^e 97:3 dr 96:4 er	4m 88% yield ^e 95:5 dr 99:1 er	X-ray structure of 4m
4n 66% yield ^e 97:3 dr 93:7 er	6n 62% yield 97:3 dr 93:7 er	CsF; H ₂ O ₂ , KHCO ₃

Conditions: See Table 1, entry 5 (10 mol %).

[a] Isolated yield.

[b] Diastereo-meric and enantiomeric ratios determined as in Table 1.

[c] Using 20 mol % catalyst.

[d] Reaction run for 4 d using 20 mol % catalyst with 20 mol % TfOH.

[e] Yield over two steps (See Table S7).[20]

Table 3
Formation of lactone 8f via trapping of β -silyl carbocation

X	ratio (8f : 4f) ^[a]	yield of 8f (%) ^[b]	dr of 8f ^[c]	er of 8f ^[d]
OTf	63:37	53	44:17:19:20	85:15
Cl	19:81	18	21:47:23:9	94:6

^[a]Based on mass recovery.

^[b]Isolated yield.

^[c]Determined using ^1H NMR spectroscopy.

^[d]Determined using chiral-phase HPLC.