Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/jor

Brown-Sequard syndrome produced by calcified herniated cervical disc and posterior vertebral osteophyte: Case report

Dawei Guan^a, Guanjun Wang^a, Morgan Clare^b, Zhengda Kuang^{a,*}

^aDepartment of Spinal Surgery, General Hospital of Chinese People's Armed Police Forces, Beijing 100039, PR China

^bCollege of Medicine, University of Florida, Gainesville, FL 32610, USA

ARTICLE INFO

Article history:

Received 26 May 2015

Accepted 4 October 2015

Available online 29 October 2015

Keywords:

Brown-Sequard syndrome

Disc herniation

Calcification

Osteophyte

Subtotal vertebrectomy

ABSTRACT

Brown-Sequard syndrome (BSS) produced by cervical disc disorders has rarely been seen clinically and only 50 cases have been reported in English literatures. However, most of which have resulted from acute disc herniation. Here, we report a case of BSS produced by calcified herniated C4-C5 disc and posterior vertebral osteophyte, in which decompression through anterior approach was performed. This case revealed the potential of cervical spondylopathy leading to BSS in a chronic manner. Once the diagnosis is established, it is advisable to perform decompression as early as possible.

© 2015 Prof. PK Surendran Memorial Education Foundation. Published by Elsevier B.V. All rights reserved.

Brown-Sequard syndrome (BSS) produced by hemisection of the spinal cord will lead to ipsilateral motor weakness, proprioceptive and vibratory sensation loss and is associated with contralateral deficit in sensation of pain and temperature. BSS is most commonly caused by injuries and neoplasms in the cervical and thoracic region.^{1–5} BSS induced by disc disorders is rare; since the first published case in 1928, only 50 cases have been reported in English language literatures.^{6–10}

This case is the 51st published discogenic BSS, which is produced by calcified herniated C4-C5 disc and posterior vertebral osteophyte. Contrast to previous cases with acute progression, features of BSS took place 7 years after the onset of initial radicular symptom in a quite chronic manner.

This 53-year-old male patient had a 7-year history of mild pain and numbness in the left arm. During the 6 months before

admission, he had experienced progressive weakness in the left arm and leg, along with contralateral deficit in sensation of pain and temperature below T11.

Upon physical examination, features of BSS were shown, including ipsilateral weakness in the left arm and leg (MRC Grade 3/5), hypertonia and hyperreflexia in left lower extremity, as well as reduced contralateral sensation of pain and temperature below T11. In addition, Hoffmann sign, Babinski sign and ankle clonus test of the left side were positive.

CT scan showed calcifications of herniated C4-C5 disc and posterior vertebral osteophyte of C5 (Fig. 1). MRI showed a large central left-sided C4-C5 disc herniation, compressing the spinal cord, leading to increased signal intensity on T2WI (Fig. 2). MRI of thoracic spine showed no positive result, eliminating disorders in the thoracic region (Fig. 3).

* Corresponding author.

E-mail address: kuangzd@sina.com (Z. Kuang).

<http://dx.doi.org/10.1016/j.jor.2015.10.007>

0972-978X/© 2015 Prof. PK Surendran Memorial Education Foundation. Published by Elsevier B.V. All rights reserved.

Fig. 1 – CT. Calcification of left-sided herniated C4-C5 disc (A) and posterior vertebral osteophyte of C5 (B). To deal with the osteophyte, subtotal vertebrectomy and titanium mesh cages reconstruction were selected for decompression.

Subtotal vertebrectomy of C5 and reconstruction using titanium mesh cages filled with autogenous bone were performed through anterior approach (Hybrid, Stryker) (Fig. 4). At the time of hospital discharge, pain and numbness in the left arm were almost completely relieved, with the left limb muscle strength improved to MRC Grade 4/5. Additionally, the contralateral sensory disorder of pain and temperature was alleviated about 60%.

This patient underwent a 7-year progression from initial symptom to the occurrence of BSS, differing from the acute manner of previous cases whose clinical histories ranged from several days to the longest of 15 months.¹¹ The longstanding

history resulted in calcifications of the herniated disc and posterior vertebral osteophyte which firmly adhered to the dural sac, increasing the risk and difficulty of decompression (Fig. 1).

In this case, the contralateral sensory deficit occurred below T11, over 10 segments distal to the level of compression at C4-C5. The inconsistency was a characteristic of incomplete BSS, which is more common than complete BSS.^{12,13} Sensation of pain and temperature are transmitted through the spinothalamic tract, crossing the midline a few segments (1–3) above the entry. To distinguish the cause of contralateral sensory disorder, MRI of segments above the level of reduced sensation is needed.

Fig. 2 – Cervical MRI. The herniated C4-C5 disc compressed left side of the spinal cord (A), consistent to the weakness in left limb and sensory deficit in the right side. Increased signal intensity on T2 weighted image (B) indicated severe spinal cord injury, which may lead to poor prognosis.

Fig. 3 – Thoracic MRI. No positive result indicated that the contralateral sensory disorder resulted from compression in the cervical region.

To deal with posterior vertebral osteophyte, we selected subtotal vertebrectomy of C5 and reconstruction using titanium mesh cages filled with autogenous bone. In cases without disorders posterior to the vertebral body, anterior cervical discectomy and fusion (ACDF) is most frequently adopted and discectomy combined foraminotomy or laminectomy are adopted in 3 and 2 cases, respectively.^{7,14}

The patient wore a rigid cervical collar and received oral neural nutrition medicine for 2 months. Three months after the operation, no pain in the left arm was reported and muscle strength in the left limb improved to MRC Grade 4+/5 without limitation to daily activities. The contralateral sensory disorder was alleviated over 80% compared to the most severe condition.

1. Discussion

Cervical degeneration diseases, epidural hematomas, demyelinating diseases, and spinal cord herniation are rare causes of BSS.^{15–18} Including this case, only 51 cases of discogenic BSS have been published in English literatures. Nearly all of the cases were induced by cervical disc disorders, with only one exception resulting from thoracic disc herniation.¹⁰

In this case, the patient who underwent a 7-year cervical spondylopathy and 6-month BSS showed significant

Fig. 4 – X-ray after the operation. Subtotal vertebrectomy and titanium mesh cages reconstruction were performed through anterior approach. The cage was packed with autogenous spongy bone from removed C5 vertebra. Locking plate was implanted from C4-C6 for stabilization (Hybrid, Stryker).

improvement, but not complete relief in ipsilateral weakness and contralateral sensory deficit. However, in cases with short history, over half of the patients showed complete recovery.^{14,19} Because longstanding compression leads to poor prognosis, it is advised to perform the decompression as early as possible.

From this case, we should be alert to the potential of cervical spondylopathy resulting in BSS in a chronic manner, which may lead to poor prognosis. Close observation should be done and once the diagnosis of BSS is established, decompression should be performed as soon as possible.

Conflicts of interest

The authors have none to declare.

REFERENCES

1. Roth EJ, Park T, Pang T, Yarkony GM, Lee MY. Traumatic cervical Brown-Sequard and Brown-Sequard-plus syndromes: the spectrum of presentations and outcomes. *Paraplegia*. 1991;29:582–589.

2. Aryan HE, Farin A, Nakaji P, Imbesi SG, Abshire BB. Intramedullary spinal cord metastasis of lung adenocarcinoma presenting as Brown-Sequard syndrome. *Surg Neurol.* 2004;61:72–76.
3. Harris P. Stab wound of the back causing an acute subdural haematoma and a Brown-Sequard neurological syndrome. *Spinal Cord.* 2005;43:678–679.
4. Gill D, Ifthikharuddin S, Samkoff LM. Acute Brown-Sequard syndrome. *Arch Neurol.* 2004;61:131.
5. Lim E, Wong YS, Lo YL, Lim SH. Traumatic atypical Brown-Sequard syndrome: case report and literature review. *Clin Neurol Neurosurg.* 2003;105:143–145.
6. Antich PA, Sanjuan AC, Girvent FM, Simo JD. High cervical disc herniation and Brown-Sequard syndrome. A case report and review of the literature. *J Bone Joint Surg.* 1999;81:462–463.
7. Lee JK, Kim YS, Kim SH. Brown-Sequard syndrome produced by cervical disc herniation with complete neurologic recovery: report of three cases and review of the literature. *Spinal Cord.* 2007;45:744–748.
8. Rustagi T, Badve S, Maniar H, Parekh AN. Cervical disc herniation causing Brown-Sequard's syndrome: a case report and literature review. *Case Rep Orthop.* 2011;2011:943720.
9. Kim JT, Bong HJ, Chung DS, Park YS. Cervical disc herniation producing acute Brown-Sequard syndrome. *J Korean Neurosurg Soc.* 2009;45:312–314.
10. Witiw CD, Shamji MF. Brown-Sequard syndrome from herniation of a thoracic disc. *Can Med Assoc J.* 2014;186:1395.
11. Iwamura Y, Onari K, Kondo S, Inasaka R, Horii H. Cervical intradural disc herniation. *Spine.* 2001;26:698–702.
12. Kobayashi N, Asamoto S, Doi H, Sugiyama H. Brown-Sequard syndrome produced by cervical disc herniation: report of two cases and review of the literature. *Spine J.* 2003;3:530–533.
13. Mastronardi L, Ruggeri A. Cervical disc herniation producing Brown-Sequard syndrome: case report. *Spine.* 2004;29:E28–E31.
14. Abouhashem S, Ammar M, Barakat M, Abdelhameed E. Management of Brown-Sequard syndrome in cervical disc diseases. *Turk Neurosurg.* 2013;23:470–475.
15. Boukobza M, Guichard JP, Boissonet M, et al. Spinal epidural haematoma: report of 11 cases and review of the literature. *Neuroradiology.* 1994;36:456–459.
16. Bondurant FJ, Cotler HB, Kulkarni MV, McArdle CB, Harris Jr JH. Acute spinal cord injury. A study using physical examination and magnetic resonance imaging. *Spine.* 1990;15:161–168.
17. Kim JM, Oh SH, Kim KJ, Park SH, Park KS. Idiopathic spinal cord herniation as a treatable cause of progressive Brown-Sequard syndrome. *J Clin Neurol.* 2007;3:204–207.
18. Yokoyama K, Kawanishi M, Yamada M, Kuroiwa T. Cervical disc herniation manifesting as a Brown-Sequard syndrome. *J Neurosci Rural Pract.* 2012;3:182–183.
19. Sayer FT, Vitali AM, Low HL, Paquette S, Honey CR. Brown-Sequard syndrome produced by C3–C4 cervical disc herniation: a case report and review of the literature. *Spine.* 2008;33:E279–E282.