Published in final edited form as:

Curr Opin Immunol. 2016 April ; 39: 82–89. doi:10.1016/j.coi.2016.01.007.

Development and Maturation of Natural Killer Cells

Theresa L. Geiger^{1,2} and Joseph C. Sun^{1,2,3}

¹Louis V. Gerstner, Jr. Graduate School of Biomedical Sciences, Memorial Sloan Kettering Cancer Center, New York, NY 10065

²Immunology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065

³Department of Immunology and Microbial Pathogenesis, Weill Cornell Medical College, New York, NY 10065

Abstract

Natural killer (NK) cells are innate lymphocytes that are critical for host protection against pathogens and cancer due to their ability to rapidly release inflammatory cytokines and kill infected or transformed cells. In the 40 years since their initial discovery, much has been learned about how this important cellular lineage develops and functions. We now know that NK cells are the founding members of an expanded family of lymphocyte known as innate lymphoid cells (ILC). Furthermore, we have recently discovered that NK cells can possess features of adaptive immunity such as antigen specificity and long-lived memory responses. Here we will review our current understanding of the molecular mechanisms driving development of NK cells from the common lymphoid progenitor (CLP) to mature NK cells, and from activated effectors to long-lived memory NK cells.

Introduction

NK cells, like B and T cells, are a lymphocyte lineage derived from the CLP [1], and like B cells, are thought to develop primarily in the bone marrow [2], although other sites of development, such as the liver and thymus, have also been proposed (reviewed in [3]). However, unlike the antigen receptors of B and T cells, NK cell receptors are germ line encoded and do not require gene rearrangement by RAG recombinase [4], though recent work has suggested that RAG plays an unexpected cell-intrinsic role in NK cell development [5]. NK cells also undergo an “education” process during development where they acquire the ability to recognize lack of self MHC class I, or “missing-self”, a feature that facilitates their surveillance of target cells that have down-regulated MHC class I during infection or malignancy [6]. NK cells rely on both cytokines and transcription factors to promote and control their development. Cytokine signaling from interleukin (IL)-15 is

Corresponding author: Joseph C. Sun, PhD, Memorial Sloan Kettering Cancer Center, 408 East 69th Street, ZRC-1402, New York, NY 10065, Phone: 646-888-3228, Fax: 646-422-0452, sunj@mskcc.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

critical for the development of NK cells and is required throughout their lifetime [7,8]. Transcription factors such as Nfil3 and PU.1 are necessary for development of early NK cell progenitors [9-12], whereas Id2, Tox, and others are important later in development [13-15]. Eomes and T-bet are among factors that then control the final stages of NK cell maturation [16,17]. In the periphery, the activation and differentiation of NK cells are regulated by a plethora of transcription factors mediating distinct effector functions. This review will outline current knowledge about the stages of NK cell development and the factors driving each stage.

Stages of NK Cell Development and Differentiation

The CLP is characterized by expression of IL-7R α (CD127), c-kit (CD117), Sca-1, and Flt-3 (CD135), as well as the lack of common lineage markers such as CD3, CD4, CD8, CD19, Ter119, Gr-1 and NK1.1 (Figure 1) [1]. From the CLP, cells develop into NK cell precursors (NKP), which are defined by expression of the IL-15 receptor β chain (CD122), and lack of common lineage markers, including the NK cell markers NK1.1 and DX5 (CD49b) (Figure 1) [2]. This NKP population has been further refined based on the co-expression of CD27 and CD244, with the majority of these cells also expressing IL-7R α [18]. An intermediate population between the CLP and NKP termed “pre-NKP” has also recently been defined as lineage negative, CD244⁺ c-kit^{low} IL-7R α ⁺ Flt-3⁻ and CD122⁻ [18,19]. However, recent work suggests that this population is heterogeneous, composed of true NK-committed precursors as well as PLZF- and α 4 β 7 integrin-expressing ILC precursors (ILCP) (Figure 1) [20]. A precursor of this pre-NKP population also capable of producing all ILC lineages (including NK cells) has recently been identified by expression of the transcription factor Tcf-1 [21]. From the CD122⁺IL-7R α ⁺ NKP stage, cells develop into immature NK (iNK) cells, which lose expression of IL-7R α and acquire expression of NK1.1 but do not yet express CD49b (Figure 1) [2]. As immature NK cells gain expression of CD11b, CD43, Ly49 receptors, and CD49b (DX5), they also gain functional competence in cytotoxicity and production of interferon (IFN)- γ [22], and egress from the bone marrow.

The peripheral NK cell pool can be delineated by expression of CD27, with CD27^{lo/-} NK cells being more cytotoxic and producing more cytokines than CD27^{high} NK cells [23]. These mature peripheral NK cell populations have more recently been further refined into four stages of maturation, defined by sequential upregulation of CD11b expression followed by downregulation of CD27, with the most immature NK cells being CD27⁻CD11b⁻ and the most mature NK cells being CD27⁻CD11b⁺ [24]. During viral infection or pro-inflammatory cytokine exposure, mature peripheral NK cells can differentiate into effector and long-lived memory NK cells (reviewed in [25]). During the CD8⁺ T cell response to viral infection, at least two different effector cell populations are thought to be generated: KLRG1^{hi} short-lived effector cells (SLECs) and KLRG1^{lo} memory precursor effector cells (MPECs) [26]. Recent evidence suggests that a similar paradigm exists in the resting NK cell pool, with virus-specific KLRG1⁻ NK cells exhibiting a greater capacity to generate memory NK cells than their KLRG1⁺ counterparts [27]. In accordance with this finding, another recent study found that RAG expression during NK cell ontogeny was correlated with lower expression of KLRG1 and a greater memory potential [5].

Transcriptional Control of Early NK Cell Development

Lineage commitment to either an adaptive or innate lymphocyte cell fate is determined by a complex network of transcription factors (Figure 2). For example, Notch signaling through the ligands Jagged1 and Jagged2 preferentially drives NK cell development from the CLP [28-30], whereas delta-like ligands (DLL) promote T cell development [31]. Moreover, thymocytes can be diverted into an NK cell-like fate if the Notch1-dependent transcription factor Bcl11b is ablated during T cell development [32-34], suggesting active suppression of the NK cell fate. Similarly, early B cell factor 1 (Ebf1) and Pax5 promote the B cell fate by suppressing expression of ILC and T-cell promoting transcription factors Notch1, Tcf-1, Gata3, and Id2 [35]. Even within the innate lymphocyte lineages, differential expression of specific transcription factors give rise to distinct cell fates. For example, although both NK cells and non-NK cell “helper” ILCs require the transcription factors Id2 [13,36,37] and Nfil3 [9,10,38-42] for their development, only the helper ILC lineages require Gata3 for development [43-45]. These differential requirements are consistent with recent studies indicating that ILCs are not derived from the same CD122⁺ precursor as NK cells, but rather arise from an IL-7R α ⁺, α 4 β 7⁺, Id2-expressing precursor, referred to as the common helper innate lymphoid cell precursor or “CHILP” (reviewed in [46]).

Nfil3 (also known as E4BP4) is a critical factor in NK cell lineage commitment. Originally identified as a circadian clock gene [47], Nfil3 is widely expressed in many hematopoietic and non-hematopoietic cells, and is expressed as early as the CLP stage in developing lymphocytes [48]. Early studies in Nfil3-deficient mice revealed a specific loss of NK cells, whereas numbers of B cells, CD4⁺, and CD8⁺ T cells were normal [9-11]. Later studies revealed that Nfil3 expression is only required in developing NK cells through the NKP stage; conditional deletion during the iNK stage does not impact NK cell numbers, cytokine production, or response to viral infection [49]. Nfil3 expression is thought to be driven by IL-15, as IL-15 induces Nfil3 expression in NK cells and ectopic Nfil3 expression can partially rescue NK cell development *in vitro* in the absence of IL-15 signaling [9,50]. IL-15 is thought to drive Nfil3 expression through the kinase PDK1 and its downstream target mTOR [50] and mice with an NK cell specific deletion of mTOR have a block in maturation of bone marrow NK cells and a severe lack of NK cells in peripheral organs [51]. However, several recent studies have shown that certain tissue resident NK cells may be Nfil3-independent [52-54].

B cells, T cells, and most ILCs require IL-7 for their development, whereas NK cells [37,55,56] and type 1 ILCs [57] do not. The first step towards an NK cell fate coincides with gain of CD122 and loss of IL-7R α , reflecting a shift from IL-7 to IL-15 dependence. IL-15 signals through the transcription factor STAT5, and thus mice lacking Stat5b are deficient in NK cells [58]. Deleting Stat5a/b in NKP also results in complete lack of NK cells [59]. Similarly, the transcription factor Runx3 and its binding partner Cbfb can promote CD122 expression, and NKP deficient in these factors fail to produce peripheral NK cells in fetal liver chimeras [60]. An NKp46⁺ cell-specific deletion of Runx3 also results in a lack of peripheral NK cells [61]. T-bet and Eomes have also been shown to cooperate to promote expression of CD122 and mice lacking both these transcription factors are deficient in NK cells as well as memory CD8⁺ T cells [62].

PU.1, a member of the Ets family of transcription factors (reviewed in [63]), is important in the development of T and B cells, monocytes, dendritic cells, and granulocytes (reviewed in [64]). Although PU.1-deficient fetal liver cells are able to generate NK cells, chimeric mice have reduced numbers of NKP and immature NK cells [12]. PU.1-deficient NK cells also have increased expression of another Ets family factor, Ets-1, a finding that has led to suggestions that Ets-1 can compensate for lack of PU.1 in driving the NK cell lineage [12]. Although Ets-1 is likely expressed prior to the NKP stage, it appears to be required at a later point in development than PU.1. Peripheral NK cells are severely decreased in Ets-1-deficient mice [65] due to an arrest at the NKP stage [66]. Furthermore, residual NK cells in Ets-1-deficient mice were refractory when stimulated through activating receptors but hyperresponsive to proinflammatory cytokines, suggesting chronic basal stimulation [66]. Ets-1 was shown to promote other transcription factors critical in NK cell development, including Id2 and T-bet [66]. Similarly, mice lacking the Ets family member Mef have decreased NK cell numbers, as well as a defect in IFN- γ secretion and perforin expression [67].

Transcription Factors Governing NK Cell Maturation

In addition to the early role for Id family transcription factors in suppressing the adaptive lymphocyte fate while promoting innate lymphocyte development, these factors are also important later in the development of NK cells. Id2-deficient mice have a cell-intrinsic lack of peripheral NK cells [13] that was found to be due to an arrest at the iNK stage [14], indicating that Id2 is important in the transition from immature to mature NK cell. Both Id2 and Id3 are expressed in NKP, and Id2 continues to be expressed in NK cells through the mature NK cell stage. In addition, both Id2 and Id3 can promote NK cell development in culture [68,69] and Id2 is thought to be downstream of Nfil3, as ectopic Id2 expression can rescue NK cell development in Nfil3 deficient progenitors [9,48].

The T-box family of transcription factors is critical in several aspects of NK cell development and maturation. One family member, T-bet, is thought to regulate expression of SIP5, a receptor required for NK cell egress out of lymph nodes and bone marrow [70]. T-bet-deficient mice lack mature NK cells in the bone marrow and periphery, exhibiting an arrest at the iNK stage during development [16]. A more recent study suggests that T-bet stabilizes an immature (TRAIL⁺DX5⁻) NK cell state and that loss of T-bet results in higher expression of Eomes, another T-box transcription factor [17]. Eomes is required for transition to the mature (DX5⁺) NK cell stage and acquisition of Ly49 receptors [17]. However, there is some question as to whether this TRAIL⁺ population truly consists of immature NK cells or whether it represents a distinct lineage of ILC1 (reviewed in [3,71]). Loss of T-bet also results in reduced expression of another transcription factor, Blimp1, which is similarly required for progression from the iNK stage [72]. The transcription factor Gata3 is thought to promote T-bet expression, as Gata3-deficient NK cells have reduced T-bet expression and are immature in phenotype similar to T-bet-deficient NK cells [73]. Conversely, the forkhead box family transcription factors Foxo1 and Foxo3 suppress NK cell maturation by repressing T-bet [74]. Foxo3-deficient mice have normal numbers of NK cells but an increase in KLRG1⁺ NK cells, suggesting a role for Foxo3 in suppressing terminal maturation of NK cells [75].

The transcription factor Tox is required for transition from the iNK to mNK stage, as Tox-deficient mice have a severe defect in mature NK cells but no defect in immature NK cells or NKP [15]. Tox is believed to be downstream of Nfil3, as transduction of Nfil3-deficient bone marrow with Tox-expressing retroviruses was able to rescue NK cell and ILC development in recipient mice [42]. Aiolos, an Ikaros family member, is also required for terminal maturation of NK cells; Aiolos-deficient mice have an accumulation of CD27⁺CD11b⁻ and CD27⁺CD11b⁺ NK cells but a loss of the most mature CD27⁻CD11b⁺ NK cell subset [76]. Irf2 is similarly required for promoting mature NK cells, as Irf2-deficient mice maintain normal numbers of immature NK cells in the bone marrow but lack the most mature CD11b⁺ subset and circulating NK cells [77,78]. Runx3, important in promoting CD122 expression [60], has likewise been shown to be important late in NK cell development, promoting the expression of Ly49 receptors, CD11b, and CD43 [79].

Regulation of Effector NK Cell Responses and Memory Formation

The STAT family of transcription factors contains members that are phosphorylated downstream of pro-inflammatory cytokine receptors and form homo- or hetero-dimers that translocate to the nucleus to induce gene transcription (reviewed in [80]). During viral infection, type I IFNs and downstream STAT1 have been shown to enhance NK cell cytotoxicity (Figure 3) [81,82], and shield activated NK cells from cell death via an NKG2D-dependent fratricide mechanism [83]. IL-12 and downstream STAT4 are required for NK cell production of IFN- γ [81,82,84], and NK cell expansion and memory generation after mouse cytomegalovirus (MCMV) infection [85]. IL-33, IL-18, and MyD88 are also important for optimal expansion during viral infection, but are not required for memory cell formation [86,87]. In addition, IL-12, IL-18, and type I IFNs together drive expression of the transcription factor Zbtb32, which promotes NK cell proliferation after MCMV infection by antagonizing Blimp-1 [88]. The aryl hydrocarbon receptor (AhR) is another nuclear factor required for optimal cytotoxicity of NK cells [89]. Similarly, NK cells deficient in either of the transcription factors C/EBP or MITF have reduced cytotoxicity and IFN- γ secretion [90-93]. MITF may regulate cytotoxicity through interactions with MEF and PU.1 at the perforin promoter [67,94].

Control of the apoptosis pathway is thought to be involved in regulating NK cell memory formation, as the anti-apoptotic molecule Bcl-2 is downregulated in NK cells following MCMV infection [88,95]. The pro-apoptotic factor Bim controls the formation of memory NK cells, with Bim-deficient NK cells failing to contract normally following MCMV infection and displaying lower levels of memory-associated cell surface markers [95]. A recent study found that NK cells accumulate damaged mitochondria after MCMV infection, and that a small subset that cleared these mitochondria by autophagy preferentially survived to form memory NK cells, a process dependent on BNIP3 and BNIP3L [96]. Additional factors that may be specifically required for memory formation remain to be elucidated.

Concluding Remarks

As is evident by the number of groundbreaking studies discussed in this review, the development of NK cells is a highly dynamic process and new discoveries occur each week. Still, there is so much we do not yet understand. Future work will elucidate the mechanisms

regulating the various factors described above, how the factors interact with each other, and how they might be involved in disease processes.

Acknowledgements

We thank members of the Sun lab for helpful discussions. In particular we thank Aimee Beaulieu, Clair Geary, and Tim O'Sullivan, who read and commented on this review. We apologize to those whose work we were unable to discuss due to space limitations. T.L.G. is supported by a fellowship from the National Institute of Allergy and Infectious Diseases (F31 AI114019). J.C.S. is supported by the Searle Scholars Program, the Cancer Research Institute, and National Institutes of Health grants AI085034 and AI100874. Our laboratory is also supported by National Institutes of Health/National Cancer Institute Cancer Center Support Grant (CCSG) P30CA008748.

References

1. Kondo M, Weissman IL, Akashi K. Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell*. 1997; 91:661–672. [PubMed: 9393859]
2. Rosmaraki EE, Douagi I, Roth C, Colucci F, Cumano A, Di Santo JP. Identification of committed NK cell progenitors in adult murine bone marrow. *Eur J Immunol*. 2001; 31:1900–1909. [PubMed: 11433387]
3. Sojka DK, Tian Z, Yokoyama WM. Tissue-resident natural killer cells and their potential diversity. *Semin Immunol*. 2014; 26:127–131. [PubMed: 24548893]
4. Lanier LL, Phillips JH, Hackett J, Tutt M, Kumar V. Natural killer cells: definition of a cell type rather than a function. *J Immunol*. 1986; 137:2735–2739. [PubMed: 3489775]
- 5●●. Karo JM, Schatz DG, Sun JC. The RAG recombinase dictates functional heterogeneity and cellular fitness in natural killer cells. *Cell*. 2014; 159:94–107. [PubMed: 25259923] [Shows that NK cells with a history of RAG expression are more adept at generating memory NK cells.]
6. Orr MT, Lanier LL. Natural killer cell education and tolerance. *Cell*. 2010; 142:847–856. [PubMed: 20850008]
7. Di Santo JP. Natural killer cell developmental pathways: a question of balance. *Annu Rev Immunol*. 2006; 24:257–286. [PubMed: 16551250]
8. Yokoyama WM, Kim S, French AR. The dynamic life of natural killer cells. *Annu Rev Immunol*. 2004; 22:405–429. [PubMed: 15032583]
9. Gascoyne DM, Long E, Veiga-Fernandes H, de Boer J, Williams O, Seddon B, Coles M, Kioussis D, Brady HJM. The basic leucine zipper transcription factor E4BP4 is essential for natural killer cell development. *Nat Immunol*. 2009; 10:1118–1124. [PubMed: 19749763]
10. Kamizono S, Duncan GS, Seidel MG, Morimoto A, Hamada K, Grosveld G, Akashi K, Lind EF, Haight JP, Ohashi PS, et al. Nfil3/E4bp4 is required for the development and maturation of NK cells in vivo. *J Exp Med*. 2009; 206:2977–2986. [PubMed: 19995955]
11. Kashiwada M, Levy DM, McKeag L, Murray K, Schröder AJ, Canfield SM, Traver G, Rothman PB. IL-4-induced transcription factor NFIL3/E4BP4 controls IgE class switching. *Proc Natl Acad Sci USA*. 2010; 107:821–826. [PubMed: 20080759]
12. Colucci F, Samson SI, DeKoter RP, Lantz O, Singh H, Di Santo JP. Differential requirement for the transcription factor PU.1 in the generation of natural killer cells versus B and T cells. *Blood*. 2001; 97:2625–2632. [PubMed: 11313251]
13. Yokota Y, Mansouri A, Mori S, Sugawara S, Adachi S, Nishikawa S, Gruss P. Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2. *Nature*. 1999; 397:702–706. [PubMed: 10067894]
14. Boos MD, Yokota Y, Eberl G, Kee BL. Mature natural killer cell and lymphoid tissue-inducing cell development requires Id2-mediated suppression of E protein activity. *J Exp Med*. 2007; 204:1119–1130. [PubMed: 17452521]
15. Aliahmad P, la Torre de B, Kaye J. Shared dependence on the DNA-binding factor TOX for the development of lymphoid tissue-inducer cell and NK cell lineages. *Nat Immunol*. 2010; 11:945–952. [PubMed: 20818394]

16. Townsend MJ, Weinmann AS, Matsuda JL, Salomon R, Farnham PJ, Biron CA, Gapin L, Glimcher LH. T-bet regulates the terminal maturation and homeostasis of NK and Valpha14i NKT cells. *Immunity*. 2004; 20:477–494. [PubMed: 15084276]
17. Gordon SM, Chaix J, Rupp LJ, Wu J, Madera S, Sun JC, Lindsten T, Reiner SL. The transcription factors T-bet and Eomes control key checkpoints of natural killer cell maturation. *Immunity*. 2012; 36:55–67. [PubMed: 22261438]
18. Fathman JW, Bhattacharya D, Inlay MA, Seita J, Karsunky H, Weissman IL. Identification of the earliest natural killer cell-committed progenitor in murine bone marrow. *Blood*. 2011; 118:5439–5447. [PubMed: 21931117]
19. Carotta S, Pang SHM, Nutt SL, Belz GT. Identification of the earliest NK-cell precursor in the mouse BM. *Blood*. 2011; 117:5449–5452. [PubMed: 21422472]
- 20●●. Constantinides MG, Gudjonson H, McDonald BD, Ishizuka IE, Verhoef PA, Dinner AR, Bendelac A. PLZF expression maps the early stages of ILC1 lineage development. *Proc Natl Acad Sci USA*. 2015; 112:5123–5128. [PubMed: 25838284] [Maps the divergence of NK cell and ILC1 development using PLZF expression.]
- 21●●. Yang Q, Li F, Harly C, Xing S, Ye L, Xia X, Wang H, Wang X, Yu S, Zhou X, et al. TCF-1 upregulation identifies early innate lymphoid progenitors in the bone marrow. *Nat Immunol*. 2015; 16:1044–1050. [PubMed: 26280998] [Identifies an early ILC/NK cell progenitor defined by Tcf-1 expression.]
22. Kim S, Iizuka K, Kang H-SP, Dokun A, French AR, Greco S, Yokoyama WM. In vivo developmental stages in murine natural killer cell maturation. *Nat Immunol*. 2002; 3:523–528. [PubMed: 12006976]
23. Hayakawa Y, Smyth MJ. CD27 dissects mature NK cells into two subsets with distinct responsiveness and migratory capacity. *J Immunol*. 2006; 176:1517–1524. [PubMed: 16424180]
24. Chiossone L, Chaix J, Fuséri N, Roth C, Vivier E, Walzer T. Maturation of mouse NK cells is a 4-stage developmental program. *Blood*. 2009; 113:5488–5496. [PubMed: 19234143]
25. Min-Oo G, Kamimura Y, Hendricks DW, Nabekura T, Lanier LL. Natural killer cells: walking three paths down memory lane. *Trends Immunol*. 2013; 34:251–258. [PubMed: 23499559]
26. Kaech SM, Wherry EJ. Heterogeneity and cell-fate decisions in effector and memory CD8+ T cell differentiation during viral infection. *Immunity*. 2007; 27:393–405. [PubMed: 17892848]
- 27●●. Kamimura Y, Lanier LL. Homeostatic control of memory cell progenitors in the natural killer cell lineage. *Cell Rep*. 2015; 10:280–291. [PubMed: 25578733] [Establishes a paradigm for identifying memory NK cell precursors similar to that in CD8 T cells, based on low expression of KLRG1.]
28. DeHart SL, Heikens MJ, Tsai S. Jagged2 promotes the development of natural killer cells and the establishment of functional natural killer cell lines. *Blood*. 2005; 105:3521–3527. [PubMed: 15650053]
29. Jaleco AC, Neves H, Hooijberg E, Gameiro P, Clode N, Haury M, Henrique D, Parreira L. Differential effects of Notch ligands Delta-1 and Jagged-1 in human lymphoid differentiation. *J Exp Med*. 2001; 194:991–1002. [PubMed: 11581320]
30. Lehar SM, Dooley J, Farr AG, Bevan MJ. Notch ligands Delta 1 and Jagged1 transmit distinct signals to T-cell precursors. *Blood*. 2005; 105:1440–1447. [PubMed: 15486060]
31. Maillard I, Fang T, Pear WS. Regulation of lymphoid development, differentiation, and function by the Notch pathway. *Annu Rev Immunol*. 2005; 23:945–974. [PubMed: 15771590]
32. Ikawa T, Hirose S, Masuda K, Kakugawa K, Satoh R, Shibano-Satoh A, Kominami R, Katsura Y, Kawamoto H. An essential developmental checkpoint for production of the T cell lineage. *Science*. 2010; 329:93–96. [PubMed: 20595615]
33. Li L, Leid M, Rothenberg EV. An early T cell lineage commitment checkpoint dependent on the transcription factor Bcl11b. *Science*. 2010; 329:89–93. [PubMed: 20595614]
34. Li P, Burke S, Wang J, Chen X, Ortiz M, Lee S-C, Lu D, Campos L, Goulding D, Ng BL, et al. Reprogramming of T cells to natural killer-like cells upon Bcl11b deletion. *Science*. 2010; 329:85–89. [PubMed: 20538915]
35. De Obaldia ME, Bhandoola A. Transcriptional regulation of innate and adaptive lymphocyte lineages. *Annu Rev Immunol*. 2015; 33:607–642. [PubMed: 25665079]

36. Moro K, Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H, Furusawa J-I, Ohtani M, Fujii H, Koyasu S. Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)Sca-1(+) lymphoid cells. *Nature*. 2010; 463:540–544. [PubMed: 20023630]
37. Satoh-Takayama N, Lesjean-Pottier S, Vieira P, Sawa S, Eberl G, Vosschenrich CAJ, Di Santo JP. IL-7 and IL-15 independently program the differentiation of intestinal CD3-NKp46+ cell subsets from Id2-dependent precursors. *J Exp Med*. 2010; 207:273–280. [PubMed: 20142427]
38. Kashiwada M, Pham N-LL, Pewe LL, Harty JT, Rothman PB. NFIL3/E4BP4 is a key transcription factor for CD8 α^+ dendritic cell development. *Blood*. 2011; 117:6193–6197. [PubMed: 21474667]
- 39●. Geiger TL, Abt MC, Gasteiger G, Firth MA, O'Connor MH, Geary CD, O'Sullivan TE, van den Brink MR, Pamer EG, Hanash AM, et al. Nfil3 is crucial for development of innate lymphoid cells and host protection against intestinal pathogens. *J Exp Med*. 2014; 211:1723–1731. [PubMed: 25113970] [With Refs. [40●, 41●, 42●] identifies Nfil3 as required for the development of all ILC lineages.]
- 40●. Seillet C, Rankin LC, Groom JR, Mielke LA, Tellier J, Chopin M, Huntington ND, Belz GT, Carotta S. Nfil3 is required for the development of all innate lymphoid cell subsets. *J Exp Med*. 2014; 211:1733–1740. [PubMed: 25092873] [See annotation to Ref. [39●].]
- 41●. Xu W, Domingues RG, Fonseca-Pereira D, Ferreira M, Ribeiro H, Lopez-Lastra S, Motomura Y, Moreira-Santos L, Bihl F, Braud V, et al. NFIL3 orchestrates the emergence of common helper innate lymphoid cell precursors. *Cell Rep*. 2015; 10:2043–2054. [PubMed: 25801035] [See annotation to Ref. [39●].]
- 42●. Yu X, Wang Y, Deng M, Li Y, Ruhn KA, Zhang CC, Hooper LV. The basic leucine zipper transcription factor NFIL3 directs the development of a common innate lymphoid cell precursor. *eLife*. 2014;3. [See annotation to Ref. [39●].]
- 43●●. Yagi R, Zhong C, Northrup DL, Yu F, Bouladoux N, Spencer S, Hu G, Barron L, Sharma S, Nakayama T, et al. The transcription factor GATA3 is critical for the development of all IL-7R α -expressing innate lymphoid cells. *Immunity*. 2014; 40:378–388. [PubMed: 24631153] [With Ref. [44●] demonstrates that Gata3 is required for development and maintenance [45●] of helper ILC lineages but not NK cells.]
- 44●. Serafini N, Klein Wolterink RGJ, Satoh-Takayama N, Xu W, Vosschenrich CAJ, Hendriks RW, Di Santo JP. Gata3 drives development of ROR γ t+ group 3 innate lymphoid cells. *J Exp Med*. 2014; 211:199–208. [PubMed: 24419270] [See annotation to Ref. [43●●].]
- 45●. Zhong C, Cui K, Wilhelm C, Hu G, Mao K, Belkaid Y, Zhao K, Zhu J. Group 3 innate lymphoid cells continuously require the transcription factor GATA-3 after commitment. *Nat Immunol*. 2015 doi:10.1038/ni.3318. [See annotation to Ref. [43●●].]
46. Serafini N, Vosschenrich CAJ, Di Santo JP. Transcriptional regulation of innate lymphoid cell fate. *Nat Rev Immunol*. 2015; 15:415–428. [PubMed: 26065585]
47. Mitsui S, Yamaguchi S, Matsuo T, Ishida Y, Okamura H. Antagonistic role of E4BP4 and PAR proteins in the circadian oscillatory mechanism. *Genes Dev*. 2001; 15:995–1006. [PubMed: 11316793]
- 48●●. Male V, Nisoli I, Kostrzewski T, Allan DSJ, Carlyle JR, Lord GM, Wack A, Brady HJM. The transcription factor E4bp4/Nfil3 controls commitment to the NK lineage and directly regulates Eomes and Id2 expression. *J Exp Med*. 2014; 211:635–642. [PubMed: 24663216] [Identifies potential downstream targets of Nfil3 and along with Refs. [49●, 50●●] refines its role in NK cell development.]
- 49●. Firth MA, Madera S, Beaulieu AM, Gasteiger G, Castillo EF, Schluns KS, Kubo M, Rothman PB, Vivier E, Sun JC. Nfil3-independent lineage maintenance and antiviral response of natural killer cells. *J Exp Med*. 2013; 210:2981–2990. [PubMed: 24277151] [Shows that mature NK cells do not require Nfil3 for maintenance or function.]
- 50●●. Yang M, Li D, Chang Z, Yang Z, Tian Z, Dong Z. PDK1 orchestrates early NK cell development through induction of E4BP4 expression and maintenance of IL-15 responsiveness. *J Exp Med*. 2015; 212:253–265. [PubMed: 25624444] [Identifies PDK1 and mTOR as driving Nfil3 expression downstream of IL-15. This pathway is further refined in ref. [51●●].]
- 51●●. Marçais A, Cherfils-Vicini J, Viant C, Degouve S, Viel S, Fenis A, Rabilloud J, Mayol K, Tavares A, Bienvenu J, et al. The metabolic checkpoint kinase mTOR is essential for IL-15

- signaling during the development and activation of NK cells. *Nat Immunol.* 2014; 15:749–757. [PubMed: 24973821] [See annotation to Ref. [50].]
- 52●. Sojka DK, Plougastel-Douglas B, Yang L, Pak-Wittel MA, Artyomov MN, Ivanova Y, Zhong C, Chase JM, Rothman PB, Yu J, et al. Tissue-resident natural killer (NK) cells are cell lineages distinct from thymic and conventional splenic NK cells. *eLife.* 2014; 3:e01659. [PubMed: 24714492] [With Refs. [53●, 54●], describes Nfil3-independent tissue-resident NK cells.]
- 53●. Cortez VS, Fuchs A, Cella M, Gilfillan S, Colonna M. Cutting edge: Salivary gland NK cells develop independently of Nfil3 in steady-state. *J Immunol.* 2014; 192:4487–4491. [PubMed: 24740507] [See annotation to Ref. [52●].]
- 54●. Seillet C, Huntington ND, Gangatirkar P, Axelsson E, Minnich M, Brady HJM, Busslinger M, Smyth MJ, Belz GT, Carotta S. Differential requirement for Nfil3 during NK cell development. *J Immunol.* 2014; 192:2667–2676. [PubMed: 24532575] [See annotation to Ref. [52●].]
55. Freedden-Jeffry von U, Vieira P, Lucian LA, McNeil T, Burdach SE, Murray R. Lymphopenia in interleukin (IL)-7 gene-deleted mice identifies IL-7 as a nonredundant cytokine. *J Exp Med.* 1995; 181:1519–1526. [PubMed: 7699333]
56. Vonarbourg C, Mortha A, Bui VL, Hernandez PP, Kiss EA, Hoyler T, Flach M, Bengsch B, Thimme R, Hölscher C, et al. Regulated expression of nuclear receptor RORγt confers distinct functional fates to NK cell receptor-expressing RORγt(+) innate lymphocytes. *Immunity.* 2010; 33:736–751. [PubMed: 21093318]
- 57●. Klose CSN, Flach M, Möhle L, Rogell L, Hoyler T, Ebert K, Fabiunke C, Pfeifer D, Sexl V, Fonseca-Pereira D, et al. Differentiation of type 1 ILCs from a common progenitor to all helper-like innate lymphoid cell lineages. *Cell.* 2014; 157:340–356. [PubMed: 24725403] [Identifies the common helper innate lymphoid cell precursor (CHILP), which gives rise to helper ILCs but not NK cells.]
58. Imada K, Bloom ET, Nakajima H, Horvath-Arcidiacono JA, Udy GB, Davey HW, Leonard WJ. Stat5b is essential for natural killer cell-mediated proliferation and cytolytic activity. *J Exp Med.* 1998; 188:2067–2074. [PubMed: 9841920]
59. Eckelhart E, Warsch W, Zebedin E, Simma O, Stoiber D, Kolbe T, Rülcke T, Mueller M, Casanova E, Sexl V. A novel Ncr1-Cre mouse reveals the essential role of STAT5 for NK-cell survival and development. *Blood.* 2011; 117:1565–1573. [PubMed: 21127177]
60. Guo Y, Maillard I, Chakraborti S, Rothenberg EV, Speck NA. Core binding factors are necessary for natural killer cell development and cooperate with Notch signaling during T-cell specification. *Blood.* 2008; 112:480–492. [PubMed: 18390836]
- 61●. Ebihara T, Song C, Ryu SH, Plougastel-Douglas B, Yang L, Levanon D, Groner Y, Bern MD, Stappenbeck TS, Colonna M, et al. Runx3 specifies lineage commitment of innate lymphoid cells. *Nat Immunol.* 2015 doi:10.1038/ni.3272. [Identifies Runx3 and Cbfb as necessary for development of NK cells, ILC1, and ILC3.]
62. Intlekofer AM, Takemoto N, Wherry EJ, Longworth SA, Northrup JT, Palanivel VR, Mullen AC, Gasink CR, Kaech SM, Miller JD, et al. Effector and memory CD8+ T cell fate coupled by T-bet and eomesodermin. *Nat Immunol.* 2005; 6:1236–1244. [PubMed: 16273099]
63. Hollenhorst PC, McIntosh LP, Graves BJ. Genomic and biochemical insights into the specificity of ETS transcription factors. *Annu Rev Biochem.* 2011; 80:437–471. [PubMed: 21548782]
64. Carotta S, Wu L, Nutt SL. Surprising new roles for PU.1 in the adaptive immune response. *Immunol Rev.* 2010; 238:63–75. [PubMed: 20969585]
65. Barton K, Muthusamy N, Fischer C, Ting CN, Walunas TL, Lanier LL, Leiden JM. The Ets-1 transcription factor is required for the development of natural killer cells in mice. *Immunity.* 1998; 9:555–563. [PubMed: 9806641]
66. Ramirez K, Chandler KJ, Spaulding C, Zandi S, Sigvardsson M, Graves BJ, Kee BL. Gene deregulation and chronic activation in natural killer cells deficient in the transcription factor ETS1. *Immunity.* 2012; 36:921–932. [PubMed: 22608498]
67. Lacorazza HD, Miyazaki Y, Di Cristofano A, Deblasio A, Hedvat C, Zhang J, Cordon-Cardo C, Mao S, Pandolfi PP, Nimer SD. The ETS protein MEF plays a critical role in perforin gene expression and the development of natural killer and NK-T cells. *Immunity.* 2002; 17:437–449. [PubMed: 12387738]

68. Heemskerk MH, Blom B, Nolan G, Stegmann AP, Bakker AQ, Weijer K, Res PC, Spits H. Inhibition of T cell and promotion of natural killer cell development by the dominant negative helix loop helix factor Id3. *J Exp Med.* 1997; 186:1597–1602. [PubMed: 9348318]
69. Schotte R, Dontje W, Nagasawa M, Yasuda Y, Bakker AQ, Spits H, Blom B. Synergy between IL-15 and Id2 promotes the expansion of human NK progenitor cells, which can be counteracted by the E protein HEB required to drive T cell development. *J Immunol.* 2010; 184:6670–6679. [PubMed: 20483740]
70. Jenne CN, Enders A, Rivera R, Watson SR, Bankovich AJ, Pereira JP, Xu Y, Roots CM, Beilke JN, Banerjee A, et al. T-bet-dependent S1P5 expression in NK cells promotes egress from lymph nodes and bone marrow. *J Exp Med.* 2009; 206:2469–2481. [PubMed: 19808259]
71. O'Sullivan TE, Sun JC, Lanier LL. Natural Killer Cell Memory. *Immunity.* 2015; 43:634–645. [PubMed: 26488815]
72. Kallies A, Carotta S, Huntington ND, Bernard NJ, Tarlinton DM, Smyth MJ, Nutt SL. A role for Blimp1 in the transcriptional network controlling natural killer cell maturation. *Blood.* 2011; 117:1869–1879. [PubMed: 21131593]
73. Samson SI, Richard O, Tavian M, Ranson T, Vosshenrich CAJ, Colucci F, Buer J, Grosveld F, Godin I, Di Santo JP. GATA-3 promotes maturation, IFN-gamma production, and liver-specific homing of NK cells. *Immunity.* 2003; 19:701–711. [PubMed: 14614857]
- 74●●. Deng Y, Kerdiles Y, Chu J, Yuan S, Wang Y, Chen X, Mao H, Zhang L, Zhang J, Hughes T, et al. Transcription Factor Foxo1 Is a Negative Regulator of Natural Killer Cell Maturation and Function. *Immunity.* 2015; 42:457–470. [PubMed: 25769609] [Identifies Foxo1 as a transcription factor that suppresses NK cell maturation by suppressing T-bet.]
75. Huntington ND, Puthalakath H, Gunn P, Naik E, Michalak EM, Smyth MJ, Tabarias H, Degli-Esposti MA, Dewson G, Willis SN, et al. Interleukin 15-mediated survival of natural killer cells is determined by interactions among Bim, Noxa and Mcl-1. *Nat Immunol.* 2007; 8:856–863. [PubMed: 17618288]
- 76●●. Holmes ML, Huntington ND, Thong RPL, Brady J, Hayakawa Y, Andoniou CE, Fleming P, Shi W, Smyth GK, Degli-Esposti MA, et al. Peripheral natural killer cell maturation depends on the transcription factor Aiolos. *EMBO J.* 2014; 33:2721–2734. [PubMed: 25319415] [Identifies Aiolos as a transcription factor required for NK cell maturation.]
77. Lohoff M, Duncan GS, Ferrick D, Mittrücker HW, Bischof S, Prechtel S, Rölinghoff M, Schmitt E, Pahl A, Mak TW. Deficiency in the transcription factor interferon regulatory factor (IRF)-2 leads to severely compromised development of natural killer and T helper type 1 cells. *J Exp Med.* 2000; 192:325–336. [PubMed: 10934221]
78. Taki S, Nakajima S, Ichikawa E, Saito T, Hida S. IFN regulatory factor-2 deficiency revealed a novel checkpoint critical for the generation of peripheral NK cells. *J Immunol.* 2005; 174:6005–6012. [PubMed: 15879093]
79. Ohno S-I, Sato T, Kohu K, Takeda K, Okumura K, Satake M, Habu S. Runx proteins are involved in regulation of CD122, Ly49 family and IFN-gamma expression during NK cell differentiation. *Int Immunol.* 2008; 20:71–79. [PubMed: 18003603]
80. O'Shea JJ, Schwartz DM, Villarino AV, Gadina M, McInnes IB, Laurence A. The JAK-STAT pathway: impact on human disease and therapeutic intervention. *Annu Rev Med.* 2015; 66:311–328. [PubMed: 25587654]
81. Orange JS, Biron CA. Characterization of early IL-12, IFN- α beta, and TNF effects on antiviral state and NK cell responses during murine cytomegalovirus infection. *J Immunol.* 1996; 156:4746–4756. [PubMed: 8648121]
82. Nguyen KB, Salazar-Mather TP, Dalod MY, Van Deusen JB, Wei X-Q, Liew FY, Caligiuri MA, Durbin JE, Biron CA. Coordinated and distinct roles for IFN- α beta, IL-12, and IL-15 regulation of NK cell responses to viral infection. *J Immunol.* 2002; 169:4279–4287. [PubMed: 12370359]
- 83●. Madera S, Rapp M, Firth MA, Beilke JN, Lanier LL, Sun JC. Type I IFN promotes NK cell expansion during viral infection by protecting NK cells against fratricide. *J Exp Med.* 2016 In press. [Identifies a role for type I IFN signaling in protecting activated NK cells from NKG2D-induced fratricide.]

84. Thierfelder WE, van Deursen JM, Yamamoto K, Tripp RA, Sarawar SR, Carson RT, Sangster MY, Vignali DA, Doherty PC, Grosveld GC, et al. Requirement for Stat4 in interleukin-12-mediated responses of natural killer and T cells. *Nature*. 1996; 382:171–174. [PubMed: 8700208]
85. Sun JC, Madera S, Bezman NA, Beilke JN, Kaplan MH, Lanier LL. Proinflammatory cytokine signaling required for the generation of natural killer cell memory. *J Exp Med*. 2012; 209:947–954. [PubMed: 22493516]
- 86●. Madera S, Sun JC. Cutting edge: stage-specific requirement of IL-18 for antiviral NK cell expansion. *J Immunol*. 2015; 194:1408–1412. [PubMed: 25589075] [Identifies the IL-18/MyD88 signaling axis as promoting optimal expansion of NK cells after MCMV infection.]
- 87●. Nabekura T, Girard J-P, Lanier LL. IL-33 Receptor ST2 Amplifies the Expansion of NK Cells and Enhances Host Defense during Mouse Cytomegalovirus Infection. *J Immunol*. 2015; 194:5948–5952. [PubMed: 25926677] [Identifies IL-33 as important for NK cell expansion, but not memory formation, after MCMV infection.]
- 88●. Beaulieu AM, Zawislak CL, Nakayama T, Sun JC. The transcription factor Zbtb32 controls the proliferative burst of virus-specific natural killer cells responding to infection. *Nat Immunol*. 2014; 15:546–553. [PubMed: 24747678] [Identifies Zbtb32 as a transcription factor important for antigen-specific proliferation of NK cells.]
89. Shin JH, Zhang L, Murillo-Sauca O, Kim J, Kohrt HEK, Bui JD, Sunwoo JB. Modulation of natural killer cell antitumor activity by the aryl hydrocarbon receptor. *Proc Natl Acad Sci USA*. 2013; 110:12391–12396. [PubMed: 23836658]
90. Kaisho T, Tsutsui H, Tanaka T, Tsujimura T, Takeda K, Kawai T, Yoshida N, Nakanishi K, Akira S. Impairment of natural killer cytotoxic activity and interferon gamma production in CCAAT/enhancer binding protein gamma-deficient mice. *J Exp Med*. 1999; 190:1573–1582. [PubMed: 10587348]
91. Seaman WE, Gindhart TD, Greenspan JS, Blackman MA, Talal N. Natural killer cells, bone, and the bone marrow: studies in estrogen-treated mice and in congenitally osteopetrotic (mi/mi) mice. *J Immunol*. 1979; 122:2541–2547. [PubMed: 376736]
92. Ito A, Kataoka TR, Kim DK, Koma Y, Lee YM, Kitamura Y. Inhibitory effect on natural killer activity of microphthalmia transcription factor encoded by the mutant mi allele of mice. *Blood*. 2001; 97:2075–2083. [PubMed: 11264174]
93. Kataoka TR, Komazawa N, Oboki K, Morii E, Nakano T. Reduced expression of IL-12 receptor beta2 and IL-18 receptor alpha genes in natural killer cells and macrophages derived from B6-mi/mi mice. *Lab Invest*. 2005; 85:146–153. [PubMed: 15492754]
94. Hesslein DGT, Lanier LL. Transcriptional control of natural killer cell development and function. *Adv Immunol*. 2011; 109:45–85. [PubMed: 21569912]
- 95●. Min-Oo G, Bezman NA, Madera S, Sun JC, Lanier LL. Proapoptotic Bim regulates antigen-specific NK cell contraction and the generation of the memory NK cell pool after cytomegalovirus infection. *J Exp Med*. 2014; 211:1289–1296. [PubMed: 24958849] [Shows that the apoptosis pathway regulates formation of memory NK cells.]
- 96●. O'Sullivan TE, Johnson LR, Kang HH, Sun JC. BNIP3- and BNIP3L-Mediated Mitophagy Promotes the Generation of Natural Killer Cell Memory. *Immunity*. 2015; 43:331–342. [PubMed: 26253785] [Identifies autophagy as necessary for optimal generation of memory NK cells.]

Highlights

- How do NK cells diverge from other innate lymphocyte cell fates?
- Transcription factors key in early NK cell development include Nfil3, Ets family.
- Transcription factors critical for NK cell maturation include Id2, T-bet, Eomes.
- Activating cytokines, STATs control NK cell cytotoxicity and IFN- γ production.
- Apoptosis and autophagy regulate development of memory NK cells.

Figure 1. Stages of NK Cell Development
NK cells are derived from the CLP, which differentiates into a heterogeneous pre-NKP/ILCP population distinguished from the NKP by its expression of IL-7R and lack of CD122 expression. From the NKP, cells begin to express NK cell markers NK1.1 and NKp46, and as they further mature they acquire expression of DX5 (CD49b) and CD11b while losing expression of CD27. As NK cells mature they also gain functional competence, expressing lytic molecules and cytokines such as IFN- γ . Cell surface proteins are color coded by the stage in which they are first expressed. Loss of a specific cell surface marker after a given stage is indicated by parentheses in the stage immediately following.

Figure 2. Transcriptional Control of the NK Cell Lineage

A complex network of transcription factors governs the decision to adopt an innate or adaptive lymphocyte fate from the CLP. A simplified list of factors promoting the helper ILC, T cell, and B cell fates is shown. From the CLP, the indicated transcription factors drive cells to become NKP, immature NK cells (iNK) and mature NK cells (mNK). Factors promoting transition from the iNK to mNK stage are listed in two columns based on whether they are thought to be important earlier (left) or later (right) in NK cell maturation. Transcription factors are placed based upon where a defect or development arrest is seen in the knockout mouse when possible. Expression of the factors may occur before the indicated stage.

Figure 3. Regulation of NK Cell Function and Memory

Activated NK cells can secrete lytic molecules and IFN- γ . They can also proliferate in response to specific antigens and contract to form long-lived “memory” NK cells. Different cytokines (blue), transcription factors (purple), and other molecules (red) govern each of these distinct NK cell functions.