CORRESPONDENCE

Rehabilitation in Patients With Coronary Heart Disease: Participation and Its Effect on Prognosis

by Prof. Dr. med. Axel Schlitt, Patricia Wischmann, Prof. Dr. rer. nat. habil. Andreas Wienke, Florian Hoepfner, Frank Noack, Prof. Dr. med. Rolf-Edgar Silber, and Prof. Dr. med. Karl Werdan in issue 31–32/2015

Two Very Different Patient Populations

Any study addressing this issue must be seen as positive since valid studies on the subject are still missing even though patients have undergone short-term comprehensive cardiac rehabilitation for decades (1). With regard to the number of subjects and the length of the observation period, this study could have provided reliable data on mortality. However, the two patient populations—patients who underwent rehabilitation and patients who did not undergo rehabilitation—differed in variables known to influence the prognosis of coronary heart disease to such an extent that one could not expect the study to produce a credible result with regard to mortality. The markedly uneven distribution of risk factors and the obvious differences in coronary artery bypass graft surgery (CABG) and percutaneous coronary intervention (PCI) between the two groups indicate that two very different patient populations were evaluated. This issue cannot be resolved by multivariate analysis.

The discussion section conveys the impression that this study adds to the series of controlled studies all arriving at the conclusion that patients who underwent rehabilitation have a survival advantage. Although the three studies from Germany referred to in the article had a controlled design, they were randomized based on the intake of omega-3 fatty acids (OMEGA study) and the use of transtelephonic ECGs (registry study) or recruited from a registry (ACOS registry), but not randomized based on participation in cardiac rehabilitation. Therefore, rehabilitation was not evaluated in a controlled fashion.

Since in Germany health-insured persons are legally entitled to participate in rehabilitation, randomization for rehabilitation is not possible. However, comprehensive case-control studies are feasible from a scientific perspective. In the age of evidence-based medicine, the German statutory pension insurance scheme (*Deutsche Rentenversicherung*) or rather the linked German Society of Rehabilitation Science (DGRW, *Deutsche Gesellschaft für Rehabilitationswissenschaften*) is therefore called upon to present valid studies on the efficacy of rehabilitation. Unfortunately, the following statement still describes the present situation: “Scientific studies have not succeeded in demonstrating that rehabilitation increases the actual retirement age, lowers the need for nursing care and reduces morbidity and mortality“ (2).

DOI: 10.3238/arztebl.2016.0147a

REFERENCES

- Schlitt A, Wischmann P, Wienke A, Hoepfner F, Noack F, Silber RE, Werdan K: Rehabilitation in patients with coronary heart disease—participation and its effect on prognosis. *Dtsch Arztebl Int* 2015; 112: 527–34.
- Wirth A, Klein G, Leptin H-J: Bessere Vernetzung notwendig. *Dtsch Arztebl* 2010; 25: A1253–6.

Prof. Dr. med. Alfred Wirth
wirthbr@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Unrealistic View

Schlitt et al. report an enormous reduction of the overall mortality of 93% in patients with coronary heart disease who participated in cardiac rehabilitation (1). A meta-analysis cited by the authors found only a decline in overall mortality among cardiac rehabilitation participants of 13%.

The value of Schlitt et al.’s study is that it highlights the variation in cardiac rehabilitation participation rates among the various German Federal States and thus points towards another potential cause of regional differences in cardiac mortality. However, we think that the number needed to treat of 4.6, which can be estimated from the study data, is unrealistic. We suspect that the authors have overestimated the actual impact of cardiac rehabilitation on overall mortality:

- Immortal time bias may have had an impact. All patients who underwent cardiac rehabilitation had at least survived until the start of the rehabilitation program. One reason for an overestimate of the effect of cardiac rehabilitation could be that in the rehab group waiting times until the start of the rehabilitation were included in the effect estimates even though no deaths could be observed during these times.
- In the rehab group, 0.2% of the patients with coronary heart disease died of non-cardiac causes compared with 7.3% in the non-rehab group which indicates that the patient groups differed by health status right from the start. While data were adjusted above all for cardiovascular factors, no adjustment was made for pre-existing diseases and for indicators of health status relevant to the approval of cardiac rehabilitation, such as the Barthel index and nursing care level.
- For two subgroups of the patient population, there is reason to assume that patients did either not start with the rehabilitation program or that the rehab group application processing was stopped because of a particularly poor health status. If, based on the intention-to-treat principle, these two subgroups were assigned to the rehab group, the effect estimate would move closer to 1.

DOI: 10.3238/arztebl.2016.0147b

REFERENCES

- Schlitt A, Wischmann P, Wienke A, Hoepfner F, Noack F, Silber RE, Werdan K: Rehabilitation in patients with coronary heart disease—participation and its effect on prognosis. *Dtsch Arztebl Int* 2015; 112: 527–34.

Dr. rer. nat. Dr. rer. san. Bernd Kowall

Prof. Dr. med. Andreas Stang, MPH

Institute of Medical Informatics, Biometry and Epidemiology,
University Hospital of Essen, Germany
Bernd.Kowall@uk-essen.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We would like to thank the authors of the letters for their comments. However, we think that in the multiple regression analysis performed, adjustments were made for all confounders measured so that the difference between the groups was eliminated. A prerequisite for meaningful results from multiple analysis is to collect data on all potential confounders. Thus, the problem with observational studies is not related to the observed covariables but to all non-observed covariables for which differences between the groups could exist. In this regard we would refer here to the second part of our reply.

We reject the criticism with regard to percutaneous coronary intervention (PCI)/coronary artery bypass graft surgery (CABG). PCIs were more common in the non-rehab group (62.7% versus 52.5%). Table 2 shows a tendency towards lower mortality among patients with PCI. This means that the non-rehab group appears to be at an advantage in the initial situation. This is similar for the variable CABG.

We would like to express our thanks for addressing the issue that it is not feasible to conduct a randomized study because in Germany, every patient with severe cardiac disease has the legal right to undergo rehabilitation pursuant to the indication catalogue of the funding organizations.

Re 1) In response to the criticism of our colleagues Kowall/Stang, we have investigated how many patients actually died during the first two weeks in which cardiac rehabilitation following an acute inpatient hospital stay must be started. The analysis revealed that two patients died during this time. For the sake of

completeness, we also analyzed weeks 3 and 4 (one patient died in each week). Thus, early mortality does not appear to have an impact on the results presented by us.

Re 2) Here, we have to agree. No data on Barthel index or nursing care level were collected. These have to be regarded as potential confounders that do have an impact on mortality. Because of these, the advantage of participating in cardiac rehabilitation may have been overestimated.

Re 3) The first subgroup consists of patients who did not undergo the rehabilitation program despite the approval of their rehabilitation application. Here, it is possible that the (deteriorated) health status was the reason why the patient did not participate in the rehabilitation program. Thus, the increased mortality in this group may be explained by this factor.

The second subgroup comprises patients whose rehabilitation applications were either not approved or not completely processed, for unknown reasons. Here, one can also speculate that a deteriorated health status may have played a role; thus, mortality could actually be higher in this subgroup, too.

We would like to thank the authors of the two letters. However, in our opinion the data presented in our study show that patients benefit from cardiac rehabilitation and that eligible patients in the Federal State of Saxony-Anhalt do not make enough use of this rehabilitation offering.

DOI: 10.3238/arztebl.2016.0148

REFERENCES

- Schlitt A, Wischmann P, Wienke A, Hoepfner F, Noack F, Silber RE, Werdan K: Rehabilitation in patients with coronary heart disease—participation and its effect on prognosis. *Dtsch Arztebl Int* 2015; 112: 527–34.

Corresponding author:

Prof. Dr. med. habil. Axel Schlitt, MHA

Paracelsus-Harz-Klinikum Bad Suderode, Quedlinburg, Germany
prof.dr.axel.schlitt@paracelsus-kliniken.de

Conflict of interest statement

Prof. Schlitt has received consultancy fees (Advisory Board) from Boehringer Ingelheim. He has received lecture fees and reimbursement of conference fees and travel expenses from Sanofi-Aventis, BMS, Pfizer, Novartis, Servier, Boehringer Ingelheim, MSD, and Bayer AG. He has received study funding (third-party funds) from GSK, Sanofi-Aventis, Mitsubishi, Endotis, Bayer AG, Boehringer Ingelheim, Novartis, Actelion, and BMS.