Published in final edited form as:

Obstet Gynecol Surv. 2015 December ; 70(12): 773–779. doi:10.1097/OGX.0000000000000256.

Prenatal stress, prematurity and asthma

Brock Medsker, MD, MS¹, Erick Forno, MD, MPH², Hyagriv Simhan, MD, MSc³, and Juan C. Celedón, MD, DrPH²

¹Division of Newborn Medicine, Department of Pediatrics, Children's Hospital of Pittsburgh of UPMC, University of Pittsburgh, Pittsburgh, PA

²Division of Pulmonary Medicine, Allergy and Immunology, Department of Pediatrics, Children's Hospital of Pittsburgh of UPMC, University of Pittsburgh, Pittsburgh, PA

³Division of Maternal-Fetal Medicine, Magee Women's Hospital, Department of Obstetrics, Gynecology and Reproductive Sciences, University of Pittsburgh, Pittsburgh, PA

Abstract

Asthma is the most common chronic disease of childhood, affecting millions of children in the U.S. and worldwide. Prematurity is a risk factor for asthma, and certain ethnic or racial minorities such as Puerto Ricans and non-Hispanic Blacks are disproportionately affected by both prematurity and asthma. In this review, we examine current evidence to support maternal psychosocial stress as a putative link between prematurity and asthma, while also focusing on disruption of the hypothalamic-pituitary-adrenal (HPA) axis and immune responses as potential underlying mechanisms for stress-induced “premature asthma”. Prenatal stress may not only cause abnormalities in the HPA axis but also epigenetic changes in the fetal glucocorticoid receptor gene (*NR3C1*), leading to impaired glucocorticoid metabolism. Moreover, maternal stress can alter fetal cytokine balance, favoring Th2 (allergic) immune responses characteristic of atopic asthma: IL-6, which has been associated with premature labor, can promote Th2 responses by stimulating production of IL-4 and IL-13. Given a link among stress, prematurity, and asthma, future research should include birth cohorts aimed at confirming and better characterizing “premature asthma”. If confirmed, clinical trials of prenatal maternal stress reduction would be warranted to reduce the burden of these common co-morbidities. While awaiting the results of such studies, sound policies to prevent domestic and community violence (e.g. from firearms) are justified, not only by public safety but also by growing evidence of detrimental effects of violence-induced stress on psychiatric and somatic health.

INTRODUCTION

Asthma is a major public health problem and the most common chronic disease of childhood [1]. In the United States (U.S.), the burden of asthma is unevenly distributed among racial/ethnic groups, with Puerto Ricans and non-Hispanic Blacks being more heavily affected than non-Hispanic whites or Mexican Americans [2].

Preterm birth is frequent in the U.S., and prematurity-related complications may explain ~15% of the global mortality in children younger than 5 years [3]. Prematurity is also a risk factor for childhood asthma. In a meta-analysis of 19 studies published over a 40-year period, premature children of all ages had 37% greater odds of asthma than children born at term (95% confidence interval [CI] for adjusted odds ratio [aOR]=1.30–1.43)[4]. Prematurity may partly explain racial or ethnic disparities in asthma, as pre-term birth is more common in Puerto Ricans (13.7%) or non-Hispanic Blacks (17.1%) than in non-Hispanic whites (10.8%) or Mexican Americans (11.3%)[5].

A potential explanation for the high prevalence of both prematurity and asthma in certain minority groups is maternal psychosocial stress. Maternal stress or anxiety during pregnancy has been associated with prematurity [6] and asthma or morbidity from asthma [7]. To date, no study has directly examined whether the link between maternal stress and childhood asthma is mediated by the effect of stress on prematurity rather than being independent of such effects (Figure 1).

Maternal stress leads to excess glucocorticoid production, which can impair the development of the hypothalamic-pituitary-adrenal (HPA) axis and fetal growth, while also causing heritable epigenetic changes [8]. Moreover, glucocorticoid-regulated genes are key to fetal lung development, especially during the first and second trimester of pregnancy [9]. In this hypothesis-generating review, we will discuss some potential mechanisms of stress-related asthma in premature children.

GLUCOCORTICOIDS AND THE HPA AXIS

In humans, the HPA axis is regulated through negative feedback mechanisms. After being released from the periventricular nucleus of the hypothalamus, corticotropin-releasing hormone (CRH) binds to CRH-receptors in the anterior pituitary gland to stimulate the synthesis of pro-opiomelanocortin (POMC), which is then cleaved to produce adrenocorticotrophic hormone (ACTH). ACTH is next released from the pituitary gland into the bloodstream, where it travels to the adrenal glands and triggers the synthesis of cortisol. Cortisol can then down-regulate the HPA axis by binding to glucocorticoid receptors (GR) and mineralocorticoid receptors (MR) in the hippocampus, periventricular hypothalamus and the pituitary gland.

The role of glucocorticoids in fetal programming has been recently reviewed [10, 11]. During pregnancy, the fetus is exposed to endogenous glucocorticoids from the maternal circulation and, potentially, to exogenous glucocorticoids used to treat relatively common (e.g. pre-term labor, maternal asthma or auto-immune disorders) or uncommon (e.g. congenital adrenal hyperplasia) conditions. This is important, as prenatal exposure to increased levels of endogenous (e.g. due to increased stress) or exogenous glucocorticoids can cause lasting alterations in the fetal and neonatal HPA axis [10].

THE PLACENTA AND THE HPA AXIS

The placenta influences the maternal-fetal HPA axis in two ways. Firstly, the placenta is a second source of CRH production. CRH is produced and released from the placenta through

positive feedback in response to both maternal and fetal cortisol levels [12, 13]. This differs from the negative feedback effect of cortisol on the maternal-fetal HPA axis. The overall result is an exponential increase in CRH levels in late gestation, which has been postulated to trigger parturition [14]. Secondly, the placenta serves as a buffer for glucocorticoids passing from mother to fetus due to the presence of 11 β -hydroxysteroid dehydrogenase type-2 (HSD11B2), an enzyme that converts cortisol into less active forms. HSD11B2 levels can in turn be up or down-regulated by environmental cues or maternal stress [15].

Acute and chronic stress has been shown to differentially affect expression of the gene for HSD11B2 (*HSD11B2*) in murine models and in humans. In pregnant women, increased levels of HSD11B2 have been found in the placenta in response to acute stress [16], and chronic stress (assessed by questionnaires [17]) is associated with down-regulation of *HSD11B2*, which can lead to excess cortisol and lower neonatal birth weight [15]. Stress could alter regulation of HSD11B2 through increased methylation of the gene promoter [18] or through inflammatory mediators [19].

PRENATAL STRESS AND PREMATURITY

The relation between prenatal stress and prematurity is complex and likely influenced by timing and degree of maternal perceived stress, type of stressor [20] [21] and maternal coping mechanisms, and maternal race or ethnicity. For example, exposure to an earthquake (a traumatic event) has been associated with preterm birth only when such exposure occurred during the first trimester of pregnancy [22]. Similarly, a mother's previous history of stressful events and development of successful mechanisms to cope with stress may be more important to the pathogenesis of preterm birth than the number of stressful events *per se* [23]. Moreover, race or ethnicity may modify the effect of stress on immune responses relevant to preterm birth. In a study of 78 pregnant and non-pregnant women, serum IL-6 levels measured two hours after exposure to an acute stressor were 46% higher in African Americans than in whites (95% CI=8%-81%)[24].

Maternal stress may directly lead to prematurity through effects on the maternal-fetal HPA axis and immune responses, or indirectly by promoting unhealthy behaviors that increase the risk of preterm delivery (e.g. smoking, substance abuse, or poor dietary habits) [14].

Maternal stress leads to increased cortisol levels, which can in turn lead to increased placental CRH and prostaglandin production, signaling the start of labor. Increased serum CRH at 33-weeks of pregnancy has been associated with a threefold increased risk of spontaneous pre-term birth (95% CI for adjusted relative risk= 1.2–9.4)[13]. Of interest, CRH levels were highest in African American women who reported higher levels of cumulative stress, racial discrimination and community violence [25].

Alternatively, stress can lead to impaired cellular immune responses and impaired suppression of pro-inflammatory cytokines [26]. For example, maternal stress could enhance susceptibility to genitourinary infections such as bacterial vaginosis (BV), ultimately leading to increased risk of premature delivery [27]. Prevalence of BV among pregnant women is estimated at 16–49%, with 2 to 3.3 times higher prevalence in African Americans than whites [27]. Racial disparities also exist in the cytokine response to genitourinary infections,

with African Americans having increased IL-1 β , TNF- α and decreased soluble TNF receptor responses resulting in an imbalance of pro-inflammatory cytokines [28]. Moreover, racial disparities could be amplified when the cytokine milieu associated with under-recognized infections such as sub-clinical microbial invasion of the amniotic cavity (MIAC) is considered. In a previous study, amniotic fluid cultures showed MIAC in 12.8% of women in preterm labor with intact membranes, and in 32.4% of women with preterm pre-labor rupture of membranes (pPROM) [29]. Of note, a separate study showed that cases of *Ureaplasma* infection detected by PCR alone demonstrated increased IL-6 in amniotic fluid, as well as histologically graded chorio-amnionitis and funisitis equal to culture-positive cases [30]. As a corollary, amniotic fluid IL-6 is utilized as a bedside test for detection of MIAC, achieving a positive-predictive value of 82% [31].

Whether due to infection or stress, certain cytokines play an important role in the physiology of preterm labor by triggering prostaglandin (PG) synthesis. Disruptions in the balance of pro-inflammatory and anti-inflammatory cytokines can initiate early steps in parturition, including prostaglandin production and cervical ripening [19]. This cytokine imbalance may be more pronounced in African American women. Fetal membranes from African American mothers have been shown to produce higher concentrations of IL-1 β , MMP-9, CRH and prostaglandin-E2 [28], suggesting a more intense response to inciting events such as infection or stress.

PRENATAL STRESS AND ASTHMA

Prenatal stress has been associated with asthma or wheeze (a key symptom of asthma) in several studies [7, 32–38]. Similar to prematurity, factors or exposures that may confound or modify the putative effects of prenatal stress on asthma include socioeconomic status and access to health care [39], *in utero* environmental tobacco smoke [39], intensity of stressful exposures or perceived maternal stress [36], maternal obesity [33], maternal race or ethnicity [39], air pollution [40], and parental history of asthma or allergies [32, 41].

There is conflicting or insufficient evidence of a critical “time window” for detrimental effects of maternal stress during pregnancy [33, 35, 37], an interaction between prenatal stress and the neonate’s sex [36], or the relative independent contributions of pre- and post-natal stress [32, 33, 35] to the pathogenesis of childhood asthma. Because a significant proportion of children who experience wheeze before age 4 years have transient wheeze but no asthma, a key limitation of most (but not all [34]) published studies is the relatively short duration of follow-up after birth, which precludes a confident diagnosis of asthma.

Several mechanisms could explain the influence of prenatal stress on the infant’s susceptibility to asthma. Firstly, prenatal stress may alter fetal cytokine balance, favoring persistence of enhanced Th2 immune responses (characterized by increased production of IL-4 and IL-13, and total IgE) in early childhood. Following exposure to prenatal stress, cord blood mononuclear cells produce IL-8 and TNF- α in response to a microbial challenge, as well as IL-13 in response to dust-mite allergen [42]. IL-6 may further promote Th2 differentiation by increasing IL-4 production by CD4+ T cells, as well as by stimulating production of IL-13 [43]. Infants born to mothers with high prenatal stress also have higher

cord blood IgE than those born to mothers with lower stress levels, perhaps due to IL-4 promoter isotype switching [38].

Secondly, perinatal stress influences the glucocorticoid receptor system, which may lead to persistent abnormalities in the HPA axis. Offspring of mothers who experience depression [44] or are exposed to major stressful events (e.g. war [45] or intimate partner violence [46]) have increased methylation of the gene for the glucocorticoid receptor (*NR3C1*) in cord blood [44] or peripheral [46, 47] mononuclear cells. Infants with increased *NR3C1* methylation of cord-blood mononuclear cells have significantly higher cortisol levels after experimental stress at 3 months of life [44], as well as decreased sensitivity to dexamethasone suppression [45]. Findings in rats further suggest that disrupted mother-infant bonding may cause post-natal methylation of *NR3C1* in premature infants, which could persist into adulthood [48]. This effect has been demonstrated in preterm infants, who have higher methylation rates of *NR3C1* in peripheral blood than term infants [49].

Thirdly, maternal stress may alter immune responses through effects on the HPA axis. In murine models, maternal stress has been shown to increase eosinophils in broncho-alveolar lavage fluid in the offspring [50]. In another model, prenatal stress was associated with increased plasma corticosterone (the murine equivalent of cortisol) and IL-13 following aero-allergen challenge [51]. Of interest, both the corticosterone and IL-13 responses were suppressed by inhibition of glucocorticoid synthesis or the glucocorticoid receptors [50, 51].

Post-natal stress may amplify the detrimental effects of pre-natal stress on asthma, and separating the relative contribution of such effects to the development of asthma can be very difficult [7]. For example, children or adolescents exposed to traumatic experiences have increased methylation of *NR3C1* [52] and are at risk for post-traumatic stress disorder (PTSD), a condition associated with increased risk of asthma [53].

PREMATURITY AND ASTHMA

Prematurity, which occurs in ~11% of births worldwide, has been consistently associated with childhood asthma. In one illustrative study, birth at <32 weeks gestation was associated with 2.8 times increased odds of wheezing [54]. Other factors correlated with prematurity, such as chronic lung disease, need for mechanical ventilation and corticosteroid use have also been linked to asthma [55]. The link between prematurity and asthma has been shown to decrease after school age [4], and thus “premature asthma” may be an independent phenotype. Moreover, we recently reported that prematurity is associated with fivefold increased odds of atopic or allergic asthma but is not significantly associated with non-atopic asthma in school-aged children [56], suggesting that both structural changes in the lung and abnormal immune responses are required for prematurity-induced asthma in later childhood (the “two-hit hypothesis”).

In 1989, Strachan et al first hypothesized that the rising prevalence of allergic diseases is due to reduced exposure to microbial infections due to declining family size and improved hygienic conditions (the “hygiene hypothesis”)[57]. Since then, a growing body of literature suggests that changes in the neonatal gut flora leads to atopy and atopic diseases such as asthma (a “modified hygiene hypothesis”). Although full characterization or “correction” of

gut flora abnormalities leading to atopy remains elusive in humans [58, 59], it is reasonable to postulate that prematurity leads to persistence of predominantly Th2 immune responses after birth through changes in the gut flora (broadly characterized by reduction in anaerobes such as *Bifidobacterium* species, and increased number of aerobic bacteria such as *E. coli* and *Clostridium difficile*). Such changes can in turn be explained by perinatal factors including caesarean delivery [60, 61] and antibiotic use [60, 62]. Alternatively, birth by caesarean delivery may alter neonatal immune responses in premature infants due to the absence of labor [61].

Premature infants have increased salivary levels of pro-inflammatory cytokines such as IL-6, IL-8 and IL-12 [63], which could play a role in neonatal morbidity. IL-6 is used to operationally define fetal inflammatory response syndrome (FIRS), a condition associated with increased risk of respiratory distress syndrome, sepsis, bronchopulmonary dysplasia, intraventricular hemorrhage, and necrotizing enterocolitis [64]. Both IL-6 and IL-8 are increased after cesarean delivery of full term infants [65], which may be relevant due to modestly increased asthma risk among infants delivered by cesarean section [66]. IL-6 is associated with increased airway resistance [67] and acts on a subset of effector memory CD8+ T-cells to enhance production of Th2 cytokines [68]. IL-8, a pro-inflammatory cytokine influenced by ozone and particulate matter, acts as a potent chemoattractant of neutrophils leading to airway inflammation [69]. Thus, beyond the developmental deficiencies associated with preterm birth, cytokine profile alterations may constitute a risk marker of wheeze or asthma.

DISCUSSION AND FUTURE DIRECTIONS

Prenatal stress has been associated with asthma in children born at term [70], and thus not all cases of “stress-induced asthma” could be explained by prematurity. Similarly, stress may lead to prematurity but no asthma. Although not formally studied, it is also plausible that: 1) prematurity is associated with asthma in children exposed to low levels of maternal (prenatal) stress, and 2) stress leads to prematurity-induced asthma (“premature asthma”) (Figure 1). Given marked detrimental effects of prematurity and asthma on the health of children worldwide, identifying modifiable risk factors for one or both conditions could have major public health implications.

Maternal stress can lead to elevations of cortisol and cytokine imbalance, leading to early parturition signals. The resulting pro-inflammatory environment can result in intrauterine growth restriction (IUGR), a condition associated with fetal hypoxemia, hypoglycemia, and increased catecholamine and cortisol concentrations. High cortisol levels may alter the transcription of GR-regulated genes, which –together with IUGR– could affect lung development [9] [71]. Elevated fetal cortisol is also inversely related to surfactant protein A and B mRNA expression [72], potentially leading to structural lung changes that predispose to post-natal asthma, particularly in neonates with an atopic predisposition (either through heredity or an altered HPA axis). Stress-induced changes in the post-natal HPA axis, including increased production of intrinsic glucocorticoids, may lead to alterations in the production of pro-inflammatory cytokines (e.g. IL-1 β , TNF- α , or IL-6) [42], as well as “classic” Th2 or T regulatory cytokines relevant to asthma (e.g. IL-4 or IL-13) [63]. For

example, IL-6 not only influences the development of prematurity but has been shown to promote CD4+ T cell-mediated production of IL-4 and IL-13 [43].

We have presented evidence that perinatal stress may cause changes in glucocorticoid and cytokine production, leading to preterm labor; in turn, changes in immune responses and lung development due to both stress and prematurity place these children at increased risk of asthma (Figure 1). Large birth cohorts with adequate data on prenatal stress and prematurity, as well as sufficient duration of follow up (to assess the development of “true asthma”), are needed to elucidate which proportion of stress-induced asthma is mediated by prematurity. If the presence and importance of stress-induced “premature asthma” are confirmed, clinical trials of prenatal stress reduction to prevent prematurity and asthma would be justifiable.

The U.S. has the highest rate of firearm-related deaths of all industrialized countries [73]. The recent and highly publicized shootings in Newtown (CT), Aurora (CO), Oak Creek (WI), Tucson (AZ) and Charleston (SC) continue to raise concerns about harmful effects of firearms. For example, we and others have shown that Puerto Ricans are not only exposed to violence in their communities [2, 74, 75] but may be particularly susceptible to developing psychosocial stress after such exposure [76, 77]. A growing body of evidence suggests that stress resulting from exposure to violence (e.g. firearm-related [78]) or heavy trauma (e.g. domestic abuse [79]) partly explains the high burden of asthma in Puerto Ricans [75] [79] and other economically disadvantaged groups [7, 80].

Even though health policies cannot be based solely on existing data on both prematurity and asthma, there is ample justification for policies designed to prevent domestic abuse and firearm violence. Fatal and nonfatal firearm violence not only poses a serious threat to public safety, but has detrimental effects on psychiatric and somatic health in general, and on anxiety disorders (e.g. PTSD) and asthma in particular. The time for sound policies on gun control in the U.S. is long overdue.

Acknowledgments

Funding

This work was supported by grants HL079966 and HL117191 from the U.S. National Institutes of Health (NIH), and by the Heinz Endowments. Dr. Forno’s contribution was supported by NIH grant HD052892.

References

1. WHO. Asthma. 2013. [cited 2014 December, 14]; WHO Asthma Fact Sheet]. Available from: <http://www.who.int/mediacentre/factsheets/fs307/en/>
2. Rosser FJ, et al. Asthma in Hispanics. An 8-year update. *Am J Respir Crit Care Med*. 2014; 189(11):1316–27. [PubMed: 24881937]
3. Liu L, et al. Global, regional, and national causes of child mortality in 2000–2013, with projections to inform post-2015 priorities: an updated systematic analysis. *The Lancet*. 2014
4. Jaakkola JJ, et al. Preterm delivery and asthma: a systematic review and meta-analysis. *J Allergy Clin Immunol*. 2006; 118(4):823–30. [PubMed: 17030233]
5. Martin JA, et al. Births: final data for 2010. *Natl Vital Stat Rep*. 2012; 61(1):1–72.
6. Shapiro GD, et al. Psychosocial stress in pregnancy and preterm birth: associations and mechanisms. *J Perinat Med*. 2013; 41(6):631–45. [PubMed: 24216160]

7. Yonas MA, Lange NE, Celedon JC. Psychosocial stress and asthma morbidity. *Curr Opin Allergy Clin Immunol*. 2012; 12(2):202–10. [PubMed: 22266773]
8. Reynolds RM. Glucocorticoid excess and the developmental origins of disease: Two decades of the testing the hypothesis. *Psychoneuroendocrinology*. 2013; 38:1–11. [PubMed: 22998948]
9. Sharma S, et al. Glucocorticoid genes and the developmental origins of asthma susceptibility and treatment response. *American Journal of Respiratory Cell and Molecular Biology*. 2014
10. Moisiadis VG, Matthews SG. Glucocorticoids and fetal programming part 1: Outcomes. *Nat Rev Endocrinol*. 2014; 10(7):391–402. [PubMed: 24863382]
11. Moisiadis VG, Matthews SG. Glucocorticoids and fetal programming part 2: Mechanisms. *Nat Rev Endocrinol*. 2014; 10(7):403–411. [PubMed: 24863383]
12. Rich-Edwards JW, Grizzard TA. Psychosocial stress and neuroendocrine mechanisms in preterm delivery. *Am J Obstet Gynecol*. 2005; 192(5 Suppl):S30–5. [PubMed: 15891710]
13. Wadhwa PD, et al. Placental corticotropin-releasing hormone (CRH), spontaneous preterm birth, and fetal growth restriction: a prospective investigation. *Am J Obstet Gynecol*. 2004; 191(4): 1063–9. [PubMed: 15507922]
14. Beydoun H, Saftlas AF. Physical and mental health outcomes of prenatal maternal stress in human and animal studies: a review of recent evidence. *Paediatr Perinat Epidemiol*. 2008; 22(5):438–66. [PubMed: 18782252]
15. Welberg LA, Thirivikraman K, Plotsky PM. Chronic maternal stress inhibits the capacity to up-regulate placental 11 β -hydroxysteroid dehydrogenase type 2 activity. *Journal of Endocrinology*. 2005; 186:R7–R12. [PubMed: 16135661]
16. Ghaemmaghami P, et al. The association between the acute psychobiological stress response in second trimester pregnant women, amniotic fluid glucocorticoids, and neonatal birth outcome. *Developmental Psychobiology*. 2014; 56:734–747. [PubMed: 23775363]
17. O'Donnell K, et al. Maternal prenatal anxiety and downregulation of placental 11 β -HSD2. *Psychoneuroendocrinology*. 2012; 37:818–826. [PubMed: 22001010]
18. Alikhani-Koopaei R, et al. Epigenetic regulation of 11 β -hydroxysteroid dehydrogenase type 2 expression. *Journal of Clinical Investigation*. 2004; 114:1146–1157. [PubMed: 15489962]
19. Christiaens I, et al. Inflammatory processes in preterm and term parturition. *Journal of Reproductive Immunology*. 2008; 79:50–57. [PubMed: 18550178]
20. Barrios YV, et al. Risk of spontaneous preterm birth in relation to maternal experience of serious life events during pregnancy. *Int J Womens Health*. 2014; 6:249–57. [PubMed: 24591850]
21. Khashan AS, et al. Reduced infant birthweight consequent upon maternal exposure to severe life events. *Psychosom Med*. 2008; 70(6):688–94. [PubMed: 18606728]
22. Glynn L, et al. When stress happens matters: Effects of earthquake timing on stress responsivity in pregnancy. *American Journal of Obstetrics and Gynecology*. 2001; 184:637–642. [PubMed: 11262465]
23. Hobel CJ, Goldstein A, Barrett ES. Psychosocial stress and pregnancy outcomes. *Clinical Obstetrics and Gynecology*. 2008; 51(2):333–348. [PubMed: 18463464]
24. Christian LM, et al. Stress-induced inflammatory responses in women: effects of race and pregnancy. *Psychosomatic Medicine*. 2013; 75:658–669. [PubMed: 23873713]
25. Tse AC, et al. Cumulative stress and maternal prenatal corticotropin-releasing hormone in an urban U.S. cohort. *Psychoneuroendocrinology*. 2012; 37(7):970–9. [PubMed: 22154479]
26. Segerstrom S, Miller GE. Psychological Stress and the Human Immune System: A Meta-Analytic Study of 30 Years of Inquiry. *Psychological Bulletin*. 2004; 130(4):601–630. [PubMed: 15250815]
27. Culhane J, Rauh V, Goldenberg R. Stress, Bacterial Vaginosis, and the Role of Immune Processes. *Current Infectious Disease Reports*. 2006; 8:459–464. [PubMed: 17064639]
28. Menon R, et al. An overview of racial disparities in preterm birth rates: caused by infection or inflammatory response? *Acta Obstetrica et Gynaecologica Scandinavica*. 2011; 90:1325–1331.
29. Romero R, et al. The preterm parturition syndrome. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2006; 113(Suppl 3):17–42. [PubMed: 17206962]

30. Yoon B, et al. The clinical significance of detecting *Ureaplasma urealyticum* by the polymerase chain reaction in the amniotic fluid of patients with preterm labor. *American Journal of Obstetrics and Gynecology*. 2003; 189:919–924. [PubMed: 14586326]
31. Kacerovsky MMI, Hornychova H, et al. Bedside assessment of amniotic fluid interleukin-6 in preterm prelabor rupture of membranes. *American Journal of Obstetrics and Gynecology*. 2014; 211:385.e1–9. [PubMed: 24705131]
32. Mathilda Chiu YH, et al. Prenatal and postnatal maternal stress and wheeze in urban children: effect of maternal sensitization. *Am J Respir Crit Care Med*. 2012; 186(2):147–54. [PubMed: 22582161]
33. Wright RJ, et al. Disrupted prenatal maternal cortisol, maternal obesity, and childhood wheeze. Insights into prenatal programming. *Am J Respir Crit Care Med*. 2013; 187(11):1186–93. [PubMed: 23590260]
34. Khashan AS, et al. Prenatal stress and risk of asthma hospitalization in the offspring: a Swedish population-based study. *Psychosom Med*. 2012; 74(6):635–41. [PubMed: 22753636]
35. Guxens M, et al. Parental psychological distress during pregnancy and wheezing in preschool children: the Generation R Study. *J Allergy Clin Immunol*. 2014; 133(1):59–67. e1–12. [PubMed: 23777854]
36. Turcotte-Tremblay AM, et al. Prenatal maternal stress predicts childhood asthma in girls: project ice storm. *Biomed Res Int*. 2014; 2014:201717. [PubMed: 24895550]
37. de Marco R, et al. Foetal exposure to maternal stressful events increases the risk of having asthma and atopic diseases in childhood. *Pediatr Allergy Immunol*. 2012; 23(8):724–9. [PubMed: 22957808]
38. Peters JL, et al. Prenatal negative life events increases cord blood IgE: interactions with dust mite allergen and maternal atopy. *Allergy*. 2012; 67(4):545–51. [PubMed: 22309645]
39. Quon BS, Goss CH. Maternal stress: a cause of childhood asthma? *Am J Respir Crit Care Med*. 2012; 186(2):116–7. [PubMed: 22798410]
40. Chiu YH, et al. Effects of prenatal community violence and ambient air pollution on childhood wheeze in an urban population. *J Allergy Clin Immunol*. 2013
41. Wright RJ, et al. Parental stress as a predictor of wheezing in infancy: a prospective birth-cohort study. *Am J Respir Crit Care Med*. 2002; 165(3):358–65. [PubMed: 11818321]
42. Wright RJ, et al. Prenatal maternal stress and cord blood innate and adaptive cytokine responses in an inner-city cohort. *Am J Respir Crit Care Med*. 2010; 182(1):25–33. [PubMed: 20194818]
43. Rincon M, Irvin C. Role of IL-6 in Asthma and Other Inflammatory Pulmonary Diseases. *International Journal of Biological Sciences*. 2012; 8(9):1281–1290. [PubMed: 23136556]
44. Oberlander TF, et al. Prenatal exposure to maternal depression, neonatal methylation of human glucocorticoid receptor gene (NR3C1) and infant cortisol stress responses. *Epigenetics*. 2008; 3(2):97–106. [PubMed: 18536531]
45. Yehuda R, et al. Influences of maternal and paternal PTSD on epigenetic regulation of the glucocorticoid receptor gene in holocaust survivor offspring. *American Journal of Psychiatry*. 2014; 171:872–880. [PubMed: 24832930]
46. Radtke KM, et al. Transgenerational impact of intimate partner violence on methylation in the promoter of the glucocorticoid receptor. *Transl Psychiatry*. 2011; 1:e21. [PubMed: 22832523]
47. Yehuda R, et al. Lower Methylation of Glucocorticoid Receptor Gene Promoter 1 in Peripheral Blood of Veterans with Posttraumatic Stress Disorder. *Biol Psychiatry*. 2014
48. Weaver IC, et al. Epigenetic programming by maternal behavior. *Nat Neurosci*. 2004; 7(8):847–54. [PubMed: 15220929]
49. Kantake M, et al. Postnatal epigenetic modification of glucocorticoid receptor gene in preterm infants: a prospective cohort study. *BMJ Open*. 2014; 4(7):e005318.
50. Lim R, Fedulov AV, Kobzik L. Maternal stress during pregnancy increases neonatal allergy susceptibility: Role of glucocorticoids. *Am J Physiol Lung Cell Mol Physiol*. 2014; 307(2):L141–8. [PubMed: 24838749]
51. Okuyama K, et al. The involvement of glucocorticoids in psychological stress-induced exacerbations of experimental allergic asthma. *International Archives of Allergy and Immunology*. 2014; 163:297–306. [PubMed: 24776388]

52. van der Knaap LJ, et al. Glucocorticoid receptor gene (NR3C1) methylation following stressful events between birth and adolescence. The TRAILS study. *Transl Psychiatry*. 2014; 4:e381. [PubMed: 24713862]
53. Goodwin RD, Fischer ME, Goldberg J. A twin study of post-traumatic stress disorder symptoms and asthma. *American Journal of Respiratory and Critical Care Medicine*. 2007; 176:983–987. [PubMed: 17702964]
54. Been JV, et al. Preterm birth and childhood wheezing disorders: a systematic review and meta-analysis. *PLoS Med*. 2014; 11(1):e1001596. [PubMed: 24492409]
55. Grischkan J, et al. Variation in childhood asthma among former preterm infants. *J Pediatr*. 2004; 144(3):321–6. [PubMed: 15001935]
56. Rosas-Salazar C, et al. Prematurity, atopy, and childhood asthma in Puerto Ricans. *J Allergy Clin Immunol*. 2014; 133(2):357–62. [PubMed: 24139607]
57. Strachan DP. Hay fever, hygiene, and household size. *Bmj*. 1989; 299(6710):1259–60. [PubMed: 2513902]
58. Ly NP, et al. Gut microbiota, probiotics, and vitamin D: interrelated exposures influencing allergy, asthma, and obesity? *J Allergy Clin Immunol*. 2011; 127(5):1087–94. quiz 1095–6. [PubMed: 21419479]
59. West CE, et al. The gut microbiota and inflammatory noncommunicable diseases: associations and potentials for gut microbiota therapies. *J Allergy Clin Immunol*. 2015; 135(1):3–13. quiz 14. [PubMed: 25567038]
60. Penders J, et al. Factors Influencing the Composition of the Intestinal Microbiota in Early Infancy. *Pediatrics*. 2006; 118(2):511–521. [PubMed: 16882802]
61. Pistiner M, et al. Birth by cesarean section, allergic rhinitis, and allergic sensitization among children with a parental history of atopy. *J Allergy Clin Immunol*. 2008; 122(2):274–9. [PubMed: 18571710]
62. Arboleya S, et al. Intestinal microbiota development in preterm neonates and effect of perinatal antibiotics. *J Pediatr*. 2015; 166(3):538–44. [PubMed: 25444008]
63. Talarico Sesso M, et al. Prospective evaluation of cytokine in saliva of preterm and fullterm neonates. *Immunobiology*. 2014; 219:830–835. [PubMed: 25155058]
64. Hofer N, et al. The fetal inflammatory response syndrome is a risk factor for morbidity in preterm neonates. *Am J Obstet Gynecol*. 2013; 209(6):542.e1–542.e11. [PubMed: 23994220]
65. Tadaki H, et al. Association of cord blood cytokine levels with wheezy infants in the first year of life. *Pediatr Allergy Immunol*. 2009; 20(3):227–33. [PubMed: 19438981]
66. Huang L, et al. Is elective cesarean section associated with a higher risk of asthma? A meta-analysis. *J Asthma*. 2015; 52(1):16–25. [PubMed: 25162303]
67. Rubini A. IL-6 increases airway resistance in the rat. *Cytokine*. 2010; 51(3):266–73. [PubMed: 20643563]
68. Lee N, et al. IL-6 receptor alpha defines effector memory CD8+ T cells producing Th2 cytokines and expanding in asthma. *Am J Respir Crit Care Med*. 2014; 190(12):1383–94. [PubMed: 25390970]
69. Kafoury RM, Kelley J. Ozone enhances diesel exhaust particles (DEP)-induced interleukin-8 (IL-8) gene expression in human airway epithelial cells through activation of nuclear factors-kappaB (NF-kappaB) and IL-6 (NF-IL6). *Int J Environ Res Public Health*. 2005; 2(3–4):403–10. [PubMed: 16819095]
70. Wright RJ, Gold DR. Population Disparities in Asthma. *Annual Review of Public Health*. 2005; 26:89–113.
71. Rozance PJ, et al. Intrauterine growth restriction decreases pulmonary alveolar and vessel growth and causes pulmonary artery endothelial cell dysfunction in vitro in fetal sheep. *American Journal of Physiology Lung Cellular and Molecular Physiology*. 2011; 2011:L860–L871.
72. Orgeig S, et al. Intrauterine growth restriction delays surfactant protein maturation in the sheep fetus. *American Journal of Physiology Lung Cellular and Molecular Physiology*. 2010; 298:L575–L583.
73. Hahn RA, et al. Firearms laws and the reduction of violence: a systematic review. *Am J Prev Med*. 2005; 28(2 Suppl 1):40–71. [PubMed: 15698747]

74. Martinez-Taboas A, et al. Prevalence and victimization correlates of pathological dissociation in a community sample of youths. *J Trauma Stress*. 2006; 19(4):439–48. [PubMed: 16929499]
75. Chen W, et al. ADCYAP1R1 and asthma in Puerto Rican children. *Am J Respir Crit Care Med*. 2013; 187(6):584–8. [PubMed: 23328528]
76. Galea S, et al. Hispanic ethnicity and post-traumatic stress disorder after a disaster: evidence from a general population survey after September 11, 2001. *Ann Epidemiol*. 2004; 14(8):520–31. [PubMed: 15350950]
77. Ortega AN, Rosenheck R. Posttraumatic stress disorder among Hispanic Vietnam veterans. *Am J Psychiatry*. 2000; 157(4):615–9. [PubMed: 10739422]
78. Ramratnam S, et al. Exposure to gun violence and asthma among children in Puerto Rico. *Respir Med*. 2015 In Press.
79. Cohen RT, et al. Violence, abuse, and asthma in Puerto Rican children. *Am J Respir Crit Care Med*. 2008; 178(5):453–9. [PubMed: 18556622]
80. Coogan PF, et al. Abuse during childhood and adolescence and risk of adult-onset asthma in African American women. *J Allergy Clin Immunol*. 2013; 131(4):1058–63. [PubMed: 23219171]

Learning Objectives

After completing this activity, the learner will be better able to

Understand the quality and limitation of the current evidence for a link between prenatal stress and prematurity or asthma.

Compare and contrast the feedback mechanisms for cortisol regulation at the brain and placenta in response to stress.

Demonstrate an understanding of racial differences in hormonal and cytokine response, and how this may apply to increased risk of preterm birth among certain groups.

Examine potential mechanisms by which alterations in the HPA axis and immune responses can contribute to preterm birth and asthma.

FIG. 1.
Potential mechanisms for the links among prenatal stress, prematurity and asthma.