Published in final edited form as:

Thromb Res. 2015 November ; 136(5): 907–910. doi:10.1016/j.thromres.2015.09.003.

Impact of tornadoes on hospital admissions for acute cardiovascular events

Federico Silva-Palacios^a, Ana Isabel Casanegra^a, Alan Shapiro^e, Minh Phan^c, Beau Hawkins^b, Ji Li^d, Julie Stoner^d, and Alfonso Tafur^a

^aVascular Medicine Program, Cardiovascular Section, Department of Medicine, University Of Oklahoma, United States

^bCardiovascular Section, Department of Medicine, University of Oklahoma, United States

^cHematology and Oncology Section, Department of Medicine, University of Oklahoma, United States

^dDepartment of Biostatistics and Epidemiology, College of Public Health, University of Oklahoma, United States

^eSchool of Meteorology, University of Oklahoma, United States

Abstract

Background—There is a paucity of data describing cardiovascular events after tornado outbreaks. We proposed to study the effects of tornadoes on the incidence of cardiovascular events at a tertiary care institution.

Population and methods—Hospital admission records from a single center situated in a tornado-prone area three months before and after a 2013 tornado outbreak were abstracted. To control for seasonal variation, we also abstracted data from the same period of the prior year (control). Hospital admissions for cardiovascular events (CVEs) including acute myocardial infarction, stroke and venous thromboembolism (VTE) were summated by zip codes, and compared by time period.

Results—There were 22,607 admissions analyzed, of which 6,705 (30%), 7,980 (35%), and 7,922 (35%) were during the pre-tornado, post-tornado, and control time frames, respectively. There were 344 CVE in the controls, 317 CVE in pre-tornado and 364 CVEs in post tornado periods. There was no difference in the prevalence of CVE during the post-tornado season compared with the control (PPR = 1.05 95% CI: 0.91 to 1.21, $p = 0.50$) or the pre-tornado season (PPR = 0.96, 95% CI: 0.83 to 1.21, $p = 0.63$).

Conclusion—In conclusion, tornado outbreaks did not increase the prevalence of cardiovascular events. In contrast to the effect of hurricanes, implementation of a healthcare policy change directed toward the early treatment and prevention of cardiovascular events after tornadoes does not seem warranted.

1. Background

Inevitable and often unpredictable natural disasters can have preventable health effects in impacted communities. For instance, after the 2011 EF5 tornado in Joplin, Missouri, there

was an increase in cases of mucormycosis [1]. Tornado strength is measured using the Enhanced Fujita (EF) scale, which measures 28 types of structural damage and estimates speeds. An EF 4 tornado is consistent with winds of 166–200mph (267–322 km/h) and severe structural damage. An EF 5 tornado is consistent with winds of more than 200 mph (N322 km/h) and devastating structural damage [2].

Health effects of other natural disasters have been studied. Hurricane Katrina was one of the most devastating natural disasters in the history of the United States. It produced catastrophic damage with an estimated cost of 75 billion dollars in the New Orleans area and along the Mississippi coast in 2005 with a death toll of 1833 [3]. Other natural events might not be as violent but can be deadlier, the heat wave of 2003 in Europe caused 70,000 deaths [4].

An observational study of 1795 adults that lived in this area 6 months before and 6–16 months after the hurricane showed an increase in glycosylated hemoglobin levels [5]. Similarly, there was a 3 fold increase in admissions for myocardial infarction incidence after the hurricane, an effect that persisted even 6 years after the event, thought to be related to various factors including population shifts, alterations in the health care system, and the effects of chronic stress and associated behaviors [6]. Reports of rates of post-traumatic stress disorder following tornadoes vary from 2% to 59% [7].

Tornado outbreaks are among some of the most destructive weather phenomena. Based on data from 1996–2007, tornadoes in the United States cause annual property damages exceeding \$1 billion, annual fatalities of 63.3 persons and annual injuries of 999 persons [8]. Almost 100,000 applications for federal disaster assistance have been submitted as a result of tornados between 1970–1980 [7]. In the United States, May tends to be the peak month for tornado occurrence in the Southern plains states, while the upper Midwest and Great Lakes region suffer the highest frequency in the summer months. The Moore, Oklahoma tornado was part of a tornado outbreak that occurred during the afternoon and evening hours of May 20, 2013, and was the last day of a three-day stretch of significant severe weather from May 18–20, 2013. This event also produced the most deadly and devastating tornado of the year for Oklahoma and the United States. With an approximately 14 miles long path, the tornado touched down in Newcastle and dissipated east of Moore. Peak width of damage path was approximately 1.1 mile. Peak damage rating: EF-5. [9]. On May 31st, 2013 there were two tornadoes; one sampled by mobile radar and also the widest known tornado on record. This tornado was known as the “El Reno Tornado” produced extensive crop and property damage and a total of 8 motorists were killed by the tornado as it overtook them along U.S. Interstate Highway 40 [10]. Several tornados touched down in the El Reno/South OKC area during the late afternoon of 31 May 2013. An EF3 tornado touched down approximately 8.3 miles west-southwest of El Reno and traveled south of El Reno before turning northeast and crossing I-40. The tornado traveled approximately 16.2 miles before dissipating 4 mi east-southeast of El Reno. The tornado grew to become the widest on record (2.6 mile wide) and had winds exceeding 296mph. The surprisingly low EF rating for this tornado (it was originally rated EF5 but was eventually downgraded by the National Weather Service to EF3) was because the tornado traveled over open terrain during much of its

lifespan and did not affect many structures (EF ratings are based on structural damage, not wind speed).

There is paucity of data describing cardiovascular events after tornado outbreaks. We proposed to study the effects of tornadoes on the incidence of cardiovascular events (CVE) during the Oklahoma tornado outbreaks of 2013.

2. Population and methods

2.1. Institution

The University of Oklahoma Medical Center is the only level one trauma center in the state [11] and the biggest hospital in the tornado areas by bed count and admissions [12]. The hospital in the city of Moore was destroyed by one of the tornadoes, and patients were diverted to our center [13]. Oklahoma is one of the top 17 states in US that has the highest average tornadoes per 10,000 mile² over 20 years [3] and is at the center of the so called “tornado alley” because is an area of disproportionate high incidence of tornadoes [14].

Medical records for all hospital admissions during the three months before and after the tornado outbreak of 2013 (pre-tornado 2/20/2013–5/19/2013), (post-tornado 5/20/2013–8/31/2013) were abstracted. To control for seasonal variation, we also abstracted data from the same period of the prior year (control, 5/20/2012–8/31/2012). There were no high magnitude tornadoes in 2012 in the corresponding geographic areas.

Hospital admissions for coronary artery disease, stroke and VTE were identified through a hospital wide ICD-9 code using a previously described and validated method [15]. Specific codes are available in Appendix A. The admissions were analyzed by zip codes affected by the tornado, identified with data from the National Weather Service with the assistance of the School of Meteorology of the University of Oklahoma. Age, gender, readmission rate were retrospectively abstracted for all the included patients.

To evaluate the possibility of a rather immediate effect of the Tornado on CVEs, specially CAD, rather than over 3 months, the same analysis was conducted with only 1 month after tornado.

3. Statistics

A univariate generalized linear model, using a log link to estimate the prevalence proportion ratio (PPR), was fit to compare the prevalence proportion of a CVE event among the three tornado time frames (post-tornado vs control, post- vs pre-tornado). Multivariate regression models were used to test the interaction between time period and gender or zip code group on the prevalence proportion of CVE. To explore the potential short-term impact of the tornado events, a one-month window relative to the tornado time period was also considered. A two-sided 0.05 alpha level was used to define statistical significance.

4. Results

There were 22,607 admissions analyzed, of which 6,705 (30%), 7,980 (35%), and 7,922 (35%) were during the pre-tornado, post-tornado, and control time frames, respectively. There were 344 (4.3%) CVE in the controls, 317 (4.7%) CVE in pre-tornado and 364 (4.6%) CVEs in post tornado periods. The prevalence of CVEs subtypes among all the patients was: 1.3%–1.5% for VTE, 1%–1.2% for CAD, and 2%–2.2% for stroke (Table 1).

There was no difference in the prevalence of CVE during the post tornado season compared with the control (PPR = 1.05 95% CI: 0.91 to 1.21, $p = 0.50$) or the pre-tornado season (PPR = 0.96, 95% CI: 0.83 to 1.21, $p = 0.63$).

When adjusted for gender, there was no significant association between the prevalence of a CVE diagnosis and time period (overall 2° of freedom test: $p = 0.55$).

The prevalence of CVEs was lower among those living in El Reno/South OKC compared to the other zip code areas; this difference was seen consistently across the three time periods. Within each zip code stratum, the prevalence of CVE was similarly distributed among the three time periods (Table 2). Due to the non-significant interaction term between time period and zip code ($p = 0.89$) we only tested the main effects. There was no significant association between the prevalence of a CVE diagnosis and time period (overall test, $p = 0.54$). Zip code was significantly associated with the odds of CVE diagnosis ($p < 0.0001$). Patients in El Reno/South OKC area had a 41% lower prevalence of a CVE diagnosis than in the other areas (PPR = 0.59, 95% CI: 0.47 to 0.75) and a 42% lower odds of a CVE diagnosis than patients in the Newcastle/Moore area (PPR = 0.58, 95% CI: 0.41 to 0.84), while there was no significant difference between patients in the Newcastle/Moore area versus other areas ($p = 0.91$).

During the follow up, 10% of the patients had recurrent visits for any indication. The percentage of patients with recurrent hospital stays and a diagnosis code of CVE was consistent among the time points (Table 3).

To evaluate the possibility of an immediate effect of the Tornado on CVEs, specially CAD, rather than over 3 months, the same analysis was conducted using a one-month time frame relative to the tornado events. The prevalence of CVE subtypes was not different among the three time periods: 0.9%–1.6% for VTE, 0.9%–1.5% for CAD, and 2.1%–2.3% for stroke. CAD in particular there were 19 events in controls (1.3% of all admits in Time period), 19 events in pre tornado (0.9% of all admits in time period) and 36 events in post tornado (1.5% of all admits in time period). There was no difference in the prevalence of CVE during the post-tornado season compared with the control (PPR = 0.96, 95% CI: 0.74 to 1.24, $p = 0.77$) or the pre-tornado season (PPR = 1.12, 95% CI: 0.85 to 1.47 $p = 0.43$) when using one-month time windows.

5. Discussion

There was no difference in cardiovascular event admissions secondary to tornadic activity. This included CAD, VTE and stroke admissions. These findings remained despite

controlling for zip code, gender and readmissions. The need to intensify preventive health care efforts specific to CVE does not seem necessary when anticipating this type of natural disaster.

Others have found an effect of natural disasters on the occurrence of CVE. These effects may persist for at least 6 years following a particular event and may be related to various factors including population shifts, alterations in the health care system, and the effects of chronic stress and associated behavior [6,16].

Proposed hypothesis for increased cardiovascular events after natural disasters have been inferred based on a study showing increased rates of alcohol and tobacco use among survivors of natural disaster. 209 adult survivors of Hurricane Katrina were interviewed 2 to 9 months after the Hurricane. Results revealed that survivors were smoking cigarettes, consuming alcohol, and experiencing alcohol-related problems at a substantially higher rate than expected based on pre-Hurricane prevalence data. Results also suggested that certain psychosocial factors were associated with participants' substance use and misuse following the Hurricane [17]. Another study has shown association of higher cardiovascular events in populations with high rates of smoking, alcohol consumption and sedentarism [18]. The link between cramped conditions and VTE has been recognized since World War II, when people sitting for hours at a time in air raid shelters developed the condition [19].

There is a chance that the discrepancy with our results is due to the nature and duration of different natural disasters. It is possible that we did not see the same result in our study here because of the high degree of preparedness in our tornado-prone area. There is a tornado siren testing conducted in communities once a week at noon so the population is familiar with the tornado warning system [20]. The average lead-time for tornado warning was 5 min in the early 1990's; in contrast the inhabitants of Moore, Oklahoma had a 36-minute lead time for the tornado event under study in this project [21]. Tornadoes occur frequently in the studied area, and are much more common than events such as major hurricanes which rarely impact the same coastal area within a 2- to 3-year time period. As such, the perceived emotional and physical burdens may be less in these tornado-prone areas that are subjected to such frequent events, and acute cardiovascular risk may be thereby mitigated.

Our study has several strengths. This report is among the first and largest studies to assess cardiovascular events related to tornadoes. Our dataset has undergone comprehensive statistical analysis and involves a large population. Another uniqueness of our study is through the inclusion of a scenario where two different F4 tornadoes affected the same area in the same month, which had a presumably incremental effect on tornado associated physical and psychosocial damage.

Some weaknesses must be addressed. This is a single center study; however validation of our results in a larger scale is unlikely due to the sporadic and geographically limited occurrence of natural disasters. Long-term impact has not been analyzed, but we expect the main effect would have been seen during the initial 3 months post tornado given the temporality of the associated stress. There is a chance that the events were concentrated more near the outbreak, even though our hospital was the main trauma center of the area and

only 11 miles from the outbreak zone, and our center was responsible for consuming the bulk of the trauma associated with this event. This is a cardiovascular analysis, we did not delve into other outcomes previously studied related to natural disasters, such as asthma, COPD and diabetes. We could not control for severity of events. In conclusion, tornado outbreaks did not increase the prevalence of cardiovascular events. In contrast to the effect of hurricanes, implementation of a healthcare policy change for early treatment and prevention of cardiovascular events after tornadoes does not seem warranted.

Acknowledgments

Partial funding provided by National Institutes of Health, National Institute of General Medical Sciences [Grant 1 U54GM104938; JS, JL].

Appendix A. ICD 9 codes used

Codes used were:

- a. Overall CVE: 451.0 to 451.99, 453.0, 453.40 to 453.52, 453.72 to 453.79, 453.82 to 453.89, 415.1 to 415.19, 410.0 to 410.99, 411.1, 430, 431, 432.0 to 432.99, 433.0 to 433.91, 434.0 to 434.91, 435.0 to 435.99, 436
- b. VTE (DVT & PE): 451.0 to 451.99, 453.0, 453.40 to 453.52, 453.72 to 453.79, 453.82 to 453.89, 415.1 to 415.19
- c. CAD [410.7 are NSTEMI, rest STEMI]: 410.0 to 410.99, 411.1
- d. Stroke (CVE) [includes ischemic and hemorrhagic]: 430, 431, 432.0 to 432.99, 433.0 to 433.91, 434.0 to 434.91, 435.0 to 435.99, 436.

References

1. Neblett Fanfair R, Benedict K, Bos J, Bennett SD, Lo YC, Adebajo T, et al. Necrotizing cutaneous mucormycosis after a tornado in Joplin, Missouri, in 2011. *N. Engl. J. Med.* 2012; 367(23):2214–2225. [PubMed: 23215557]
2. National Wind Institute. Enhanced Fujita Scale Report 2013. Available from: <http://www.depts.ttu.edu/nwi/Pubs/FScale/Fujita.php>.
3. National Centers for Environmental Information. [cited 2014 Sept 4] Historical Records and trends 2013. Available from: <http://www.ncdc.noaa.gov/climate-information/extreme-events/us-tornado-climatology/trends>
4. Robine JM, Cheung SL, Le Roy S, Van Oyen H, Griffiths C, Michel JP, et al. Death toll exceeded 70,000 in Europe during the summer of 2003. *C. R. Biol.* 2008; 331(2):171–178. [PubMed: 18241810]
5. Fonseca VA, Smith H, Kuhadiya N, Leger SM, Yau CL, Reynolds K, et al. Impact of a natural disaster on diabetes: exacerbation of disparities and long-term consequences. *Diabetes Care.* 2009; 32(9):1632–1638. [PubMed: 19542210]
6. Peters MN, Moscona JC, Katz MJ, Deandrade KB, Quevedo HC, Tiwari S, et al. Natural disasters and myocardial infarction: the six years after Hurricane Katrina. *Mayo Clinic Proc. Mayo Clin.* 2014; 89(4):472–477.
7. Bohonos JJ, Hogan DE. The medical impact of tornadoes in North America. *J. Emerg. Med.* 1999; 17(1):67–73. [PubMed: 9950391]
8. Simmons KM, Sutter D. Economic and societal impacts of tornadoes. *Amer. Meteorol. Soc., Boston, MA.* 2011:282.

9. National Weather Service Weather Forecast Office. [updated 5/6/2014; cited 2014 Sept 4] The Tornado Outbreak of May 20, 2013. Available from: <http://www.srh.noaa.gov/oun/?n=events-20130520>
10. National Weather Service Weather Forecast Office. [updated 5/6/2014; cited 2014 Sept 4] The May 31–June 1, 2013 Tornado and Flash Flooding Event 2013. Available from: <http://www.srh.noaa.gov/oun/?n=events-20130531>
11. American College of Surgeons. Verified Trauma Centers 2014. Available from: <https://http://www.facs.org/search/trauma-centers?state=OK&distance=50>.
12. American Hospital Association. Data Viewer Quick Report: American Hospital Association. 2014 Available from: <http://www.ahadataviewer.com/quickreport/>.
13. Norman Regional Hospital System. Norman Regional Moore Update 2014. Available from: http://www.normanregional.com/en/newsroom.html?news_id=305.
14. National Centers for Environmental Information. [cited 2014 Sept 4] Tornado Alley 2013. Available from: <http://www.ncdc.noaa.gov/climate-information/extremeevents/us-tornado-climatology/tornado-alley>
15. Birman-Deych E, Waterman AD, Yan Y, Nilasena DS, Radford MJ, Gage BF. Accuracy of ICD-9-CM codes for identifying cardiovascular and stroke risk factors. Med. Care. 2005; 43(5):480–485. [PubMed: 15838413]
16. Aoki T, Fukumoto Y, Yasuda S, Sakata Y, Ito K, Takahashi J, et al. The Great East Japan Earthquake Disaster and cardiovascular diseases. Eur. Heart J. 2012; 33(22):2796–2803. [PubMed: 22930461]
17. Flory K, Hankin BL, Kloos B, Cheely C, Turecki G. Alcohol and cigarette use and misuse among Hurricane Katrina survivors: psychosocial risk and protective factors. Subst. Use Misuse. 2009; 44(12):1711–1724. [PubMed: 19895302]
18. Hamer M, Molloy GJ, Stamatakis E. Psychological distress as a risk factor for cardiovascular events: pathophysiological and behavioral mechanisms. J. Am. Coll. Cardiol. 2008; 52(25):2156–2162. [PubMed: 19095133]
19. Simpson K. Shelter Deaths from Pulmonary Embolism. The Lancet. 1940; 236(6120):744.
20. City of Oklahoma City. [cited 2014 9 Sept 2014] Outdoor Warning Sirens. Available from: <http://www.okc.gov/sirens/>
21. National Oceanic and Atmospheric Administration. [updated 5/6/2014; cited 2014 Sept 4] Tornado Forecastings and Warnings 2013. Available from: http://www.noaa.gov/factsheets/newversion/Tornadoes_web_version_final.pdf

Table 1
Frequency of patients who had CVE/CVE subtypes events across 3 time periods

Type	Time Period								
	Control			Pre-tornado			Post-tornado		
	Number of event	% of patients among all the patients (N=7922)	% of patients among CVD patients (N=344)	Number of event	% of patients among all the patients (N=6705)	% of patients among CVD patients (N=317)	Number of event	% of patients among all the patients (N=7980)	% of patients among CVD patients (N=364)
1. Overall CVD	344	4.3	100.0	317	4.7	100.0	364	4.6	100.0
2. VTE	100	1.3	29.1	91	1.4	28.7	118	1.5	32.4
3. CAD	78	1.0	22.7	83	1.2	26.2	95	1.2	26.1
4. Stroke	170	2.1	49.4	148	2.2	46.7	157	2.0	43.1

CVE: cardiovascular events, VTE: venous thromboembolism, CAD: coronary artery disease.

Table 2

		Time Period								
		Control			Pre-tornado			Post-tornado		
Type	Zip code group	Number of event	Total N	% among patients in zip code stratum	Number of event	Total N	% among patients in zip code stratum	Number of event	Total N	% among patients in zip code stratum
1. Overall CVE	Newcastle/Moore	13	305	4.3	18	307	5.9	14	319	4.4
	EL Reno/South OKC	25	894	2.8	23	737	3.1	23	889	2.6
	Others	306	6723	4.6	276	5661	4.9	327	6772	4.8
2. VTE	Newcastle/Moore	.	.	.	6	307	2.0	5	319	1.6
	EL Reno/South OKC	9	894	1.0	7	737	0.9	5	889	0.6
	Others	91	6723	1.4	78	5661	1.4	108	6772	1.6
3. CAD	Newcastle/Moore	2	305	0.7	2	307	0.7	3	319	0.9
	EL Reno/South OKC	7	894	0.8	5	737	0.7	5	889	0.6
	Others	69	6723	1.0	76	5661	1.3	87	6772	1.3
4. Stroke	Newcastle/Moore	11	305	3.6	10	307	3.3	6	319	1.9
	EL Reno/South OKC	9	894	1.0	12	737	1.6	13	889	1.5
	Others	150	6723	2.2	126	5661	2.2	138	6772	2.0

Table 3

Percent of patients who had recurrent CVE

Type	Time Period					
	Control		Pre-tornado		Post-tornado	
	Number of patient with recurrent event	% of patient (denominator N=7922)	Number of patient with recurrent event	% of patient (denominator N=7980)	Number of patient with recurrent event	% of patient (denominator N=6705)
1. Overall CVE	15	0.189%	13	0.163%	15	0.224%
2. VTE	4	0.050%	6	0.075%	5	0.075%
3. CAD	2	0.025%	3	0.038%	4	0.060%
4. Stroke	8	0.101%	3	0.038%	5	0.075%

CVE: cardiovascular events, VTE: venous thromboembolism, CAD: coronary artery disease.