Published in final edited form as:

J Am Chem Soc. 2015 September 23; 137(37): 11876–11879. doi:10.1021/jacs.5b06740.

α -Arylation of Saturated Azacycles and *N*-Methylamines via Palladium(II)-catalyzed C(sp³)-H Coupling

Jillian E. Spangler[†], Yoshihisa Kobayashi[§], Pritha Verma[†], Dong-Hui Wang[†], and Jin-Quan Yu^{†,*}

[†]Department of Chemistry, The Scripps Research Institute, 10550 N. Torrey Pines Road, La Jolla, California 92037

[§]Medicinal Chemistry, Eisai Product Creation Systems, Eisai Co., Ltd, 5-1-3 Tokodai, Tsukuba-shi, Ibaraki 300-2635, Japan

Abstract

Pd(II)-catalyzed α -C(sp³)-H arylation of pyrrolidines, piperidines, azepanes and *N*-methyl amines with arylboronic acids has been developed for the first time. This transformation is applicable to both a wide array of pyrrolidines and boronic acids, including heteroaromatic boronic acids. A diastereoselective one-pot hetero-diarylation of pyrrolidines is also achieved.

Graphical Abstract

The prevalence of cyclic saturated amines in bioactive natural products and pharmaceutical compounds has led to significant interest in the functionalization of sp³ C-H bonds adjacent to nitrogen.¹ Important progress has been made in the direct arylation of cyclic amines through α -anion², α -radical,^{3–5} and α -cation⁶ pathways. Extensive efforts at Merck led to the development of an elegant procedure for the enantioselective arylation of *N*-Boc-pyrrolidine via an asymmetric lithiation/Negishi coupling.^{2d} In contrast, direct α -arylation through catalytic transition metal-catalyzed C(sp³)-H bond activation remains underdeveloped.^{7–11} To date, all pioneering studies on metal-catalyzed arylation employ low-valent ruthenium(0) catalysts, although the high reaction temperatures (120–150 °C) required for these transformations often leads to over-arylation and poor stereocontrol.

Corresponding Author: yu200@scripps.edu.

Supporting Information Available: Experimental procedures and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

These methods also use a heterocyclic directing group that requires multiple steps to remove.

In 2006, our group disclosed a method for the palladium-catalyzed C(sp³)-H oxidation of *N*-methylcarbamates.¹² Our early efforts in this field, in conjunction with the high value of α -aryl cyclic amines, prompted us to pursue the development of a method for palladium-catalyzed α -C(sp³)-H arylation of cyclic amines. Herein we report the discovery of a method for the direct Pd(II)-catalyzed α -C(sp³)-H cross-coupling of pyrrolidines, piperidines and azepanes with both aryl and heteroaryl boronic acids, a rare example of the coupling of methylene C-H bonds with organometallic reagents. The excellent monoselectivity of this reaction also enables a one-pot sequential diastereoselective diarylation.

Although our initial efforts to achieve Pd-catalyzed α -C(sp³)-H arylation with carbamate or heterocyclic directing groups were generally unsuccessful, we were inspired by a 1981 report of the cyclopalladation of *N*-alkylthioamides¹³ to explore the α -C(sp³)-H arylation of thioamides. As such, we were pleased to find that the palladacycle **2** can be formed as an air-stable yellow solid upon heating of thioamide **1** with stoichiometric Pd(PhCN)₂Cl₂ (Scheme 1). We further discovered that palladacycle **2** readily undergoes efficient cross-coupling with phenyl boronic acids at 80°C in *tert*-amyl alcohol in the presence of a mild base and stoichiometric 1,4-benzoquinone (1,4-BQ) to provide 2-phenyl pyrrolidine **3** in 78% yield. This reaction can be conducted without exclusion of oxygen with no decrease in yield. Consistent with our early discovery of Pd(II)-catalyzed cross-coupling of C-H bonds with organotin reagents,¹⁴ 1,4-BQ is essential as a promoter for reductive elimination; no product formation is observed in the absence of 1,4-BQ.

Upon further reaction development, we found that the formation of palladacycle **2** and subsequent cross-coupling with phenylboronic acid can be rendered catalytic by treatment of pyrrolidine **1** with 10 mol% Pd(TFA)₂ catalyst and 1,4-BQ (Table 1). We were pleased to find that arylation of the pyrrolidine is highly mono-selective (>20:1 mono:diarylation). It is interesting to note that no functionalization is observed at the terminal methyl groups of the thioamide (**4**), despite significant precedent for the preferential activation of primary C-H bonds over secondary C-H bonds.¹⁵ It is also surprising that our workhorse catalyst, Pd(OAc)₂, gives negligible amount of product (Table 1, entries 1–6). 1,4-BQ is also uniquely successful as the oxidant in this transformation due to facile oxidation of the thioamide with the other oxidants that we examined, including Ag(I) salts and peroxides; the resultant amide is an unreactive byproduct in this directed α -arylation.

With these optimized reaction conditions in hand, we next evaluated the scope of the arylboronic acids that can participate in this transformation. As shown in Table 2, this C-H arylation protocol is tolerant of a broad range of coupling partners, including both electron-rich (Table 2, entries **5a–f**, 51–78% yield) and electron-poor (Table 2, entries **5g–o**, 70–80% yield) boronic acids. However, the use of electron-rich boronic acids does lead to slight increase in the formation of the corresponding diarylated product (4–8%). We were pleased to find that *para*-, *meta*-, and even sterically hindered *ortho*-substituted boronic acids (**5a**, **5k**, 51–70% yield) can all be effectively coupled in this transformation. This reaction is tolerant

of a range of functional groups, including ketones (**5n**, **5o**, 72–80% yield), amides (**5f**, 78% yield), ethers (**5e**, **5g**, 75–76% yield), and aryl halides (**5h–l**, 70–79% yield).

The relatively efficient coordination of the thioamide directing group with Pd(II) catalysts led us to speculate that this coupling reaction would be compatible with heteroarylboronic acids, which are generally challenging coupling partners in palladium-catalyzed C(sp³)-H functionalization reactions due to their propensity to coordinate to the palladium catalyst. As such, we were pleased to find that a broad range of heteroarylboronic acids react in this transformation to provide α -heteroaryl pyrrolidines (Table 3). Electron-rich benzofuran- and indole-containing boronic acids are particularly good coupling partners and provide the corresponding pyrrolidine products in excellent yields (**6a–c**, 58–82% yield). A variety of substituted pyridinylboronic acids can be effectively coupled in good yield with complete mono-selectivity (**6d–h**, 51–76% yield). However, the coupling of unsubstituted pyridinylboronic acids was not successful under current conditions. Presumably, the pyridyl outcompetes the thioamide for coordinating with Pd catalyst. Similar phenomenon have been observed in asymmetric hydrogenation of 2-pyridyl cyclic imines.¹⁶

We next turned our attention to the scope of pyrrolidines that can be used in this transformation. As shown in Table 4, 3-substituted (**8a–8e**, 51–99% yield) and 2-substituted (**8g–l**, 54–86% yield) pyrrolidines couple efficiently in this transformation with moderate to high levels of diastereoselectivity. In all cases arylation occurs selectively at the less hindered α -methylene with preferential formation of the *trans*-diastereoisomer.¹⁷ An array of heteroatom-substituted pyrrolidines are tolerated in this transformation (**8b,c,e,h**, 51–87% yield). A (3,3,0)-bicyclic pyrrolidine is also an effective substrate in this transformation, providing the desired product in 92% yield (**8f**). An array of 2,5-diaryl pyrrolidines can be synthesized using this protocol with extremely high levels of diastereoselectivity (**8i–l**, 55–86% yield, >20:1 dr). In accordance with our previous observation that electron-rich boronic acids lead to higher levels of diarylation, aryl pyrrolidines **8i,j** (76–86% yield) showed higher reactivity than aryl pyrrolidines **8k,l** (55–57% yield).

We subsequently found that this arylation can be extrapolated into a consecutive one-pot hetero-diarylation of pyrrolidines to provide 2,5-diarylated pyrrolidine products (Table 5). All of these reactions proceed with complete diastereoselectivity (>20:1 dr). Notably, this one-pot procedure does not require the addition of a second batch of palladium catalyst. We observed that the use of a more mono-selective electron-deficient boronic acid in the first arylation step provides higher yields of the hetero-diarylated product with negligible formation of the undesired homo-diarylated byproducts.

In examining the reactivity of other *N,N*-dialkylthioamides in our α -C(sp³)-H arylation reaction, we discovered that *N*-methyl thioamides are also competent substrates in this transformation. As shown in Table 6, a variety of *N*-methyl thioamides can be arylated to provide the corresponding benzyl thioamides in good yield (**11a–e**, 68–94% yield). The reaction of *N*-methyl thioamides with both electron-rich and electron-deficient heteroarylboronic acids can also be achieved (**11f–h**, 77–99% yield). In all cases the reaction is completely regioselective; no methylene C(sp³)-H arylation is observed. Surprisingly, *N*-

methyl anilines also undergo regioselective α -C(sp³)-H arylation (**11i-k**, 51–75% yield) without reacting at the *ortho*-positions of the aryl group.

We also explored the application of our methodology to the arylation of larger azacycles. As shown in Scheme 2, azepane **12** undergoes arylation in good yield under standard reaction conditions (**13**, 68% yield). In contrast, the arylation of piperidine **14a** proceeds in poor yield, even with addition of a large excess of 1,4-BQ (**15a**, 13% yield). Further studies demonstrated that, while palladacycle formation from **14a** is facile, the rate reductive elimination is significantly slower than reductive elimination from complex **2**. As such, we hoped to optimize this transformation via the use of a more bulky thioamide directing group in order to promote reductive elimination.¹⁸ We were pleased to find that the use of a the bulky 2,2-diethylbutanoic acid directing group (**14b**) provides the corresponding product in excellent yield without formation of any diarylated byproduct (**15b**, 92% yield).

Lastly, these arylated products can be readily deprotected by cleavage of the thioamide to the corresponding amine with methyl lithium at 0 °C in good yield after protection as the Boc-carbamate **16** (73% yield, Scheme 3).¹⁹ Alternatively, the thioamide **1** can be converted to amide **17** by oxidation with silver(II)-salts in nearly quantitative yield.

In conclusion, we have demonstrated the α -arylation of saturated azacycles and *N*-methyl amines via Pd(II)-catalyzed C(sp³) H coupling with boronic acids. This method allows for highly monoselective arylation as well as sequential diastereoselective di-arylation of pyrrolidines. The successful C–H coupling of pyrrolidines with aryl boronic acids also demonstrates the first example of Pd-catalyzed cross-coupling of methylene C–H bonds with organometallic reagents.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge The Scripps Research Institute, Eisai Co., Ltd, and the NIH (NIGMS, 2R01GM084019) for financial support. We thank Dr. Richard Clark from Eisai for helpful discussions.

References

1. For reviews see: Campos KR. Chem Soc Rev. 2007; 36:1069–1084. [PubMed: 17576475] Mitchell EA, Peschiulli A, Lefevre N, Meerpoel L, Maes BUW. Chem–Eur J. 2012; 18:10092–10142. [PubMed: 22829434]
2. a) Beak P, Kerrick ST, Wu S, Chu J. J Am Chem Soc. 1994; 116:3231–3239. b) Dieter RK, ShengJian L. Tetrahedron Lett. 1995; 36:3613–3616. c) Dieter RK. J Org Chem. 1997; 62:7726–7735. d) Campos KR, Klapars A, Waldman JH, Dormer PG, Chen CY. J Am Chem Soc. 2006; 128:3538–3539. [PubMed: 16536525] e) Klapars A, Campos KR, Waldman JH, Zewge D, Dormer PG, Chen CY. J Org Chem. 2008; 73:4986–4993. [PubMed: 18507444] f) Barker G, O'Brien P, Campos KR. Org Lett. 2010; 12:4176–4179. [PubMed: 20718476] g) Barker G, McGrath JL, Klapars A, Stead D, Zhou G, Campos KR, O'Brien P. J Org Chem. 2011; 76:5936–5953. [PubMed: 21714542] h) Seel S, Thaler T, Takatsu K, Zhang C, Zipse H, Straub BF, Mayer P, Knochel P. J Am

- Chem Soc. 2011; 133:4774–4777. [PubMed: 21388211] i) Beng TK, Gawley RE. *Org Lett*. 2011; 13:394–397. [PubMed: 21174392] j) Millet A, Larini P, Clot E, Baudoin O. *Chem Sci*. 2013; 4:2241–2247.
3. a) Murahashi SI. *Angew Chem, Int Ed*. 1995; 34:2443–2465. b) Li Z, Li CJ. *J Am Chem Soc*. 2004; 126:11810–11811. [PubMed: 15382913] c) Li Z, Li CJ. *J Am Chem Soc*. 2005; 127:3672–3673. [PubMed: 15771482] d) Catino AJ, Nichols JM, Nettles BJ, Doyle MP. *J Am Chem Soc*. 2006; 128:5648–5649. [PubMed: 16637627] e) Yoshikai N, Mieczkowski A, Matsumoto A, Illies L, Nakamura E. *J Am Chem Soc*. 2010; 132:5568–5569. [PubMed: 20361736] f) Ratnikov MO, Xu X, Doyle MP. *J Am Chem Soc*. 2013; 135:9475–9479. [PubMed: 23734676]
4.
For C–H coupling of tetrahydrofuran with arylboronic acids via radical abstraction, see: Liu D, Liu C, Li H, Lei A. *Angew Chem Int Ed*. 2013; 52:4453–4456.
5.
For α -arylation of amines via photoredox, see: McNally A, Prier CK, MacMillan DWC. *Science*. 2011; 334:1114–1117. [PubMed: 22116882] Singh A, Arora A, Weaver JE. *Org Lett*. 2013; 15:5390–5393. [PubMed: 24098895] Prier CK, MacMillan DWC. *Chem Sci*. 2014; 5:4173–4178. [PubMed: 26236461] Beatty JW, Stephenson CRJ. *Acc Chem Res*. 2015; 48:1474–1484. [PubMed: 25951291]
6. a) Baslé O, Li CJ. *Org Lett*. 2008; 10:3661–3663. [PubMed: 18662002] b) Dai C, Meschini F, Narayanam JMR, Stephenson CRJ. *J Am Chem Soc*. 2012; 77:4425–4431. c) Girard AS, Knauber T, Li CJ. *Angew Chem Int Ed*. 2014; 53:74–100.
7. a) Pastine SJ, Brikov DV, Sames D. *J Am Chem Soc*. 2006; 128:14220–14221. [PubMed: 17076471] b) Prokopcová H, Bergman SD, Aelvoet K, Smout V, Herrebout W, Van der Veken B, Meerpoel L, Maes BUW. *Chem–Eur J*. 2010; 16:13063–13067. [PubMed: 20981669] c) Peschiulli A, Smout V, Storr TE, Mitchell EA, Eliás Z, Herrebout W, Berthelot D, Meerpoel L, Maes BUW. *Chem–Eur J*. 2013; 19:10378–10387. [PubMed: 23780756] d) Dastbaravardeh N, Schnürch M, Mihovilovic MD. *Org Lett*. 2012; 14:1930–1933. [PubMed: 22449256] e) Schwarz MC, Dastbaravardeh N, Kirchner K, Schnürch M, Mihovilovic MD. *Monatsh Chem*. 2013; 144:539–552. [PubMed: 26166880] f) Phani Kumar NY, Jeyachandran R, Ackermann L. *J Org Chem*. 2013; 78:4145–4152. [PubMed: 23574572]
8.
For Ta-catalyzed hydroaminoalkylation see: Herzon SB, Hartwig JF. *J Am Chem Soc*. 2007; 129:6690–6691. [PubMed: 17474747] Herzon SB, Hartwig JF. *J Am Chem Soc*. 2008; 130:14940–14941. [PubMed: 18937477] Eisenberger P, Ayinla RO, Lauzon JMP, Schafer LL. *Angew Chem, Int Ed*. 2009; 48:8361–8365. Payne PR, Garcia P, Eisenberger P, Yim JCH, Schafer LL. *Org Lett*. 2013; 15:2182–2185. [PubMed: 23600625] Garcia P, Lau YY, Perry MR, Schafer LL. *Angew Chem, Int Ed*. 2013; 52:9144–9148. Chong E, Brandt JW, Schafer LL. *J Am Chem Soc*. 2014; 136:10898–10901. [PubMed: 25041474]
9.
For references on Ru/Ir-catalyzed hydroaminoalkylation see: Jun CH, Hwang DC, Na SJ. *Chem Commun*. 1998:1405–1406. Chatani N, Asaumi T, Yorimitsu S, Ikeda T, Kakiuchi F, Murai S. *J Am Chem Soc*. 2001; 123:10935–10941. [PubMed: 11686697] Pan S, Endo K, Shibata T. *Org Lett*. 2011; 113:4692–4695. [PubMed: 21812393] Bergman SD, Storr TE, Prokopcová H, Aelvoet K, Diels G, Meerpoel L, Maes BUW. *Chem Eur J*. 2012; 18:10393–10398. [PubMed: 22786664] Schmitt DC, Lee J, Dechert-Schmitt AM, Yamaguchi E, Krische MJ. *Chem Commun*. 2013; 49:6096–6098. Schinkel M, Wang L, Bielefeld K, Ackermann L. *Org Lett*. 2014; 16:1876–1879. [PubMed: 24635222]
- 10.

- For references on Ir-catalyzed alkenylation of amides see: Tsuchikama D, Kasagawa M, Endo K, Shibata T. *Org Lett*. 2009; 11:1821–1823. [PubMed: 19354323] Pan S, Matsuo Y, Endo K, Shibata T. *Tetrahedron*. 2012; 68:9009–9015.
- 11.
- For references on the Rh/Ir-catalyzed borylation of amines and amides see: Kawamorita S, Miyazaki T, Iwai T, Ohmiya H, Sawamura M. *J Am Chem Soc*. 2012; 134:12924–12927. [PubMed: 22816772] Li Q, Liskey CW, Hartwig JF. *J Am Chem Soc*. 2014; 136:8755–8765. [PubMed: 24836159]
12. Wang DH, Hao XS, Wu DF, Yu JQ. *Org Lett*. 2006; 8:3387–3390. [PubMed: 16836412]
- 13.
- Tamaru Y, Kagotani M, Yoshida Z. *Angew Chem Int Ed Engl*. 1981; 20:980–981. Dunina VV, Golovan EB, Kazakova EI, Potapov GP, Beletskaya IP. *Metalloorg Khim*. 1991; 4:1391–1396. Nojima Y, Nonoyama M, Nakajima K. *Polyhedron*. 1996; 15:3795–3809. Dunina VV, Gorunova ON, Kuznetsova ED, Turubanova EI, Livantsov MV, Grishin YK, Kuzmina LG, Churakov AV. *Russ Chem Bull, Int Ed*. 2006; 55:2193–2211. For direct Pd insertion into α -C–H bonds of amines: c) Lu CC, Peters JC. *J Am Chem Soc*. 2004; 126:15818–15832. [PubMed: 15571407]
14. Chen X, Li JJ, Hao XS, Goodhue CE, Yu JQ. *J Am Chem Soc*. 2006; 128:78–79. [PubMed: 16390130]
- 15.
- Recent reviews of transition-metal-catalyzed C(sp³) H functionalization see: Daugulis O, Do HQ, Shabashov D. *Acc Chem Res*. 2009; 42:1074–1086. [PubMed: 19552413] Jazzar R, Hitce J, Renaudat A, Sofack-Kreutzer J, Baudoin O. *Chem–Eur J*. 2010; 16:2654–2672. [PubMed: 20143359] Lyons TW, Sanford MS. *Chem Rev*. 2010; 110:1147–1169. [PubMed: 20078038] Wasa M, Engle KM, Yu JQ. *Isr J Chem*. 2010; 50:605–616. [PubMed: 21552359] Baudoin O. *Chem Soc Rev*. 2011; 40:4902–4911. [PubMed: 21505712]
16. Guo C, Sun DW, Yang S, Mao SJ, Xu XH, Zhu SF, Zhou QL. *J Am Chem Soc*. 2015; 137:90–93. [PubMed: 25548865]
- 17.
- Stereochemistry was confirmed by single crystal X-ray crystallography of compound **9a**. This crystal structure was deposited in the Cambridge Crystallographic Data Center (CCDC-1408481). See supporting information.
18. Mann G, Shelby Q, Roy AH, Hartwig JF. *Organometallics*. 2003; 22:2775–2789. Culkin DA, Hartwig JF. *Organometallics*. 2004; 23:3398–3416.
19. a) Lubosch W, Seebach D. *Helv Chim Acta*. 1980; 63:102–116. b) Hodgson DM, Kloesges J. *Angew Chem Int Ed*. 2010; 49:2900–2903. c) Hodgson DM, Pearson CI, Thompson AL. *J Org Chem*. 2013; 78:1098–1106. [PubMed: 23305091]

Scheme 1.
Stoichiometric α -Arylation of Pyrrolidine

Scheme 2.
Arylation of Larger Azacycles

Scheme 3.
Deprotection of Thioamide

Table 1Optimization of the α -arylation of amines^{a,b}

entry	deviation from standard conditions	yield(%)
1	none	82
2	10 mol% PdCl ₂ (PhCN) ₂	63
3	10 mol% PdCl ₂	57
4	10 mol% Pd(OAc) ₂	7
5	10 mol% Pd(OTf) ₂ ·MeCN	22
6	10 mol% allylPdCl ₂	37
7	noBQ	0
8	0.5 eq BQ	34
9	1.5 eq BQ	61
10	with 2.0 eq PhBPin	8
11	with 2.0 eq PhBF ₃ K	0

^aReaction conditions: thioamide **1** (0.2 mmol, 1.0 eq), arylboronic acid (0.4 mmol, 2.0 eq), Pd(TFA)₂ (0.02 mmol, 0.1 eq), 1,4-BQ (0.22 mmol, 1.1 eq), KHCO₃ (0.4 mmol, 2.0 eq), t-AmylOH (4.0 mL), 100 °C, 4 h;

^bThe yield was determined by ¹H NMR analysis of the crude product using mesitylene as the internal standard.

Table 2

α-Arylation of Pyrrolidine: Boronic Acid Scope^{a,b}

^a Reaction conditions: thioamide **1** (0.2 mmol, 1.0 eq), arylboronic acid (0.4 mmol, 2.0 eq), Pd(TFA)₂ (0.02 mmol, 0.1 eq), 1,4-BQ (0.22 mmol, 1.1 eq), KHCO₃ (0.4 mmol, 2.0 eq), *t*-AmylOH (4.0 mL), 100 °C, 4 h.

^b Yield after isolation by chromatography.

^c 1.2 eq 1,4-BQ.

^d 1.4 eq 1,4-BQ.

Table 3

Heteroaryl Boronic Acid Scope^{a,b}

^aReaction conditions: thioamide **1** (0.2 mmol, 1.0 eq), arylboronic acid (0.4 mmol, 2.0 eq), Pd(TFA)₂ (0.02 mmol, 0.1 eq), 1,4-BQ (0.22 mmol, 1.1 eq), KHCO₃ (0.4 mmol, 2.0 eq), *t*-AmylOH (4.0 mL), 100 °C, 4 h.

^bYield after isolation by chromatography.

^c3 eq 1,4-BQ.

Table 4

Pyrrolidine Substrate Scope^{a,b}

^aReaction conditions: thioamide **7** (0.2 mmol, 1.0 eq), arylboronic acid (0.4 mmol, 2.0 eq), Pd(TFA)₂ (0.02 mmol, 0.1 eq), 1,4-BQ (0.4 mmol, 2.0 eq), KHCO₃ (0.4 mmol, 2.0 eq), *t*-AmylOH (4.0 mL), 100 °C, 4 h.

^bYields after isolation by chromatography.

^c0.15 eq Pd(TFA)₂ (0.03 mmol).

^d 1.4 eq 1,4-BQ (0.28 mmol), 0.15 eq Pd(TFA)₂ (0.03 mmol).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 5

One-pot Hetero-diarylation of Pyrrolidines^{a,b}

^aReaction conditions: thioamide **1** (0.2 mmol, 1.0 eq), arylboronic acid-1 (0.4 mmol, 2.0 eq), Pd(TFA)₂ (0.02 mmol, 0.1 eq), 1,4-BQ (0.28 mmol, 1.4 eq), KHCO₃ (0.4 mmol, 2.0 eq), *t*-AmylOH (4.0 mL), 100 °C, 4 h then arylboronic acid-2 (0.4 mmol, 2.0 eq), 1,4-BQ (0.4 mmol, 2.0 eq), 12 h.

^bYield after isolation by chromatography, dr determined by ¹H NMR.

Table 6

Arylation of *N*-Methylamines^{a,b}

^aReaction conditions: thioamide **10** (0.2 mmol, 1.0 eq), arylboronic acid (0.4 mmol, 2.0 eq), Pd(TFA)₂ (0.02 mmol, 0.1 eq), 1,4-BQ (0.4 mmol, 2.0 eq), KHCO₃ (0.4 mmol, 2.0 eq), *t*-AmylOH (4.0 mL), 100 °C, 4 h.

^bYield after isolation by chromatography.

^c1.0 eq 1,4-BQ (0.2 mmol).