Association between QTc of patients with schizophrenia and five genetic polymorphisms of *GSTZ1* and *XRCC1*

Mostafa Saadat

Correspondence to

Professor Mostafa Saadat,
Department of Biology, College
of Sciences, and Institute of
Biotechnology, Shiraz
University, Shiraz 71454, Iran;
saadat@susc.ac.ir,
msaadat41@yahoo.com

Received 7 September 2013

Accepted 4 November 2013

ABSTRACT

Background Several antipsychotic agents are known to prolong the QT interval. A study was undertaken to find the possible influence of polymorphisms of *GSTZ1* (MIM: 603758) and *XRCC1* (MIM: 194360) on the rate-corrected QT interval (QTc) in patients with schizophrenia.

Methods The study was carried out on 48 inpatients with schizophrenia. The patients were diagnosed with chronic schizophrenia according to structured clinical interviews using SCID-I (clinician version) to confirm and document DSM-IV diagnosis. Measurements of the QT and RR intervals were recorded using a magnifying grid on lead II. The QTc was calculated according to Bazett's formula. A PCR-based method was used to determine the *GSTZ1* and *XRCC1* genotypes.

Results Statistical analysis showed that there was no association between the study polymorphisms of *GSTZ1* and *XRCC1* and QTc.

Conclusions *GSTZ1* is not associated with QTc in patients treated with antipsychotic drugs.

INTRODUCTION

Genetic factors are involved in the QT interval length on the ECG.^{1–2} However, it is known that the rate-corrected QT interval (QTc) is influenced by various environmental parameters including medication.^{3–5} A number of drugs including several antipsychotic agents are known to prolong the QT interval in a dose-dependent manner.^{3–4}

The human *GSTZ1* (a member of GST ζ ; MIM: 603758) was discovered by a bioinformatics approach and identified in human expressed sequence tag databases.^{6–7} In mice and humans, GST ζ is expressed in many tissues at a low level.⁸ It has been reported that, in mice, *Gstz1* deficiency resulted in the generation of a constant level of oxidative stress.⁸ Several *GSTZ1* variant sequences have been identified in humans^{8–9}; two non-synonymous polymorphisms at nucleotide positions 94 (Glu32Lys; rs7975) and 124 (Gly42Arg; rs7972) have been reported¹⁰ and a G-1002A polymorphism in the promoter region of *GSTZ1* has also been identified.⁹

The x-ray repair cross-complementing 1 protein (XRCC1; MIM: 194360) plays an important role in DNA single-strand break repair in cells. Two polymorphisms at codons 194 (Arg194Trp; rs 1799782) and 399 (Arg399Gln; rs 25487) have been reported in human XRCC1.¹¹

Associations between the abovementioned polymorphisms and both schizophrenia and bipolar disorder have been reported.^{12–16} It has also been

shown that the QTc interval in patients with schizophrenia is correlated with the *GSTT1* polymorphism.¹⁷ In the present study the association between QTc in patients with schizophrenia and polymorphisms of *XRCC1* and *GSTZ1* was investigated.

MATERIALS AND METHODS

This study was performed in Shiraz, southern Iran. Forty-eight inpatients with schizophrenia from Ibn-Sina and Razi Hospitals, Shiraz University of Medical Sciences of mean $\pm$ SD age 43.5 $\pm$ 9.4 years participated in the study. The patients were diagnosed as having chronic schizophrenia according to clinical interview using SCID-I (clinician version) to confirm and document a diagnosis of DSM-IV, as described previously.¹⁸ The patients were receiving conventional antipsychotic drugs such as perphenazine, trifluoperazine, chlorpromazine, thioridazine and haloperidol. None were receiving antidepressants.

Measurements of the QT and RR intervals were recorded using a magnifying grid on lead II because the T wave is often well-defined in this lead. Two independent observers who were blinded to the genotypes of the patients determined the duration of the QT interval, which was recorded for three consecutive beats through lead II. The QTc was calculated using Bazett's formula in which the QT interval is adjusted for heart rate by dividing it by the square root of the RR interval.¹⁹

Immediately after blood collection, whole blood was stored at -20°C until use. The PCR conditions for determining *XRCC1* and *GSTZ1* genotypes and laboratory quality control were the same as those reported previously.^{9–10–12–14}

The Kolmogorov-Smirnov test was applied in order to show the normal distribution of QTc. To evaluate an association between the suggested independent variables and QTc, the independent Student t test and/or one-way analysis of variance were used. Statistical analysis was performed using the Statistical Package for Social Sciences V.11.5 (SPSS, Chicago, Illinois, USA).

RESULTS

The mean $\pm$ SD QTc in our patients was 505 $\pm$ 45 ms, which is significantly increased compared with normal controls. This value is exceedingly high given that 500 ms is the threshold to stop medications due to the substantial risk of a torsades des pointes arrhythmia about this value. Using the Kolmogorov-Smirnov test, the QTc showed a normal distribution ($Z=0.690$, $p=0.728$).

To cite: Saadat M. *Heart Asia* 2014;6:8–9.
doi:10.1136/heartasia-2013-010424

Table 1 Mean QTc interval (in ms) in patients with schizophrenia stratified according to their *XRCC1* and *GSTZ1* genotypes

Polymorphisms	Mean (ms)	SD	n	t (df=46) F (df=2, 45)	p Value
Polymorphisms of <i>XRCC1</i>					
Arg194Trp					
Arg/Arg	504	48	38	0.155	0.877
Arg/Trp	507	29	10		
Arg399Gln					
Arg/Arg	485	39	11	1.374	0.264
Arg/Gln	510	46	30		
Gln/Gln	514	46	7		
Polymorphisms of <i>GSTZ1</i>					
G-1002A					
GG	502	40	30	0.264	0.769
GA	507	56	16		
AA	524	9	2		
Glu32Lys					
Glu/Glu	500	40	28	0.816	0.449
Glu/Lys	516	51	16		
Lys/Lys	491	51	4		
Gly42Arg					
Gly/Gly	504	46	45	0.178	0.859
Gly/Arg	509	26	3		

The mean $\pm$ SD QTc in our patients according to their genotypes of *XRCC1* and *GSTZ1* polymorphisms are shown in table 1. Statistical analysis using the independent Student t test and/or one way analysis of variance showed that there was no association between the polymorphisms of *GSTZ1* and *XRCC1* and QTc.

DISCUSSION

We have previously found that the *GSTT1* polymorphism was associated with the risk of schizophrenia¹⁸ and QTc in these patients.¹⁷ The present results indicate that *GSTZ1*, which is another detoxifying glutathione S-transferase (GST) enzyme, is not associated with QTc in patients treated with antipsychotic drugs.

As ethnicity may influence the observed associations in multi-factorial traits,²⁰ replication of this study in other countries is recommended. The main limitations of our study are the small sample size, lack of stratification of patients according to drug intake and the lack of measurement of plasma concentrations of antipsychotic drugs and/or their active metabolites.

Acknowledgements The author is indebted to the participants for their close cooperation. The author would like to acknowledge Dr Hassan Farrashbandi for referring the patients and Mrs Fatemeh Mobayen for her assistance during the course of this study.

Funding This study was supported by Shiraz University.

Competing interests None.

Patient consent Obtained.

Ethics approval Ethics approval was obtained from the Institutional Review Board of Shiraz University.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Russell MW, Law I, Sholinsky P, *et al.* Heritability of ECG measurements in adults male twins. *J Electrocardiol* 1998;30:64–8.
- Carter N, Snieder H, Jeffery S, *et al.* QT-interval in twins. *J Hum Hypertens* 2001;14:389–90.
- Glassman AH, Bigger JT. Antipsychotic drugs: prolonged QTc interval, torsade de points, and sudden death. *Am J Psychiatry* 2001;158:1774–82.
- Elming H, Sonne J, Lublin HKF. The importance of the QT interval: a review of the literature. *Acta Psychiatrica Scand* 2003;107:96–101.
- Kaab S, Hintergeer M, Nabauer M, *et al.* Sotalol testing unmasks altered repolarization in patients with suspected acquired long-QT-syndrome: a case-control pilot study using i.v sotalol. *Eur Heart J* 2003;24:649–57.
- Board PG, Baker RT, Chelvanayagam G, *et al.* Zeta, a novel class of glutathione transferases in a range of species from plants to humans. *Biochem J* 1997;328:929–35.
- Board PG, Chelvanayagam G, Jermini LS, *et al.* Identification of novel glutathione transferases and polymorphic variants by expressed sequence tag database analysis. *Drug Metab Dispos* 2001;29:544–7.
- Blackburn AC, Matthaei KI, Lim C, *et al.* Deficiency of glutathione transferase zeta causes oxidative stress and activation of antioxidant response pathways. *Mol Pharmacol* 2006;69:650–7.
- Fang YY, Kashkarov U, Anders MW, *et al.* Polymorphisms in the human glutathione transferase zeta promoter. *Pharmacogenet Genomics* 2006;16:307–13.
- Blackburn AC, Tzeng HF, Anders MW, *et al.* Discovery of a functional polymorphism in human glutathione transferase zeta by expressed sequence tag database analysis. *Pharmacogenetics* 2000;10:49–57.
- Goode EL, Ulrich CM, Potter JD. Polymorphisms in DNA repair genes and associations with cancer risk. *Cancer Epidemiol Biomarkers Prev* 2002;11:1513–30.
- Saadat M, Pakyari N, Farrashbandi H. Genetic polymorphism in the DNA repair gene *XRCC1* and susceptibility to schizophrenia. *Psychiatry Res* 2008;157:241–5.
- Saadat M, Mohammadynejad P, Ghanizadeh A, *et al.* Genetic polymorphisms (at codons 194 and 399) in the DNA repair gene *XRCC1* and susceptibility to bipolar disorder. *Psychiatry Res* 2012;198:171.
- Derakhshandeh S, Saadat I, Farrashbandi H, *et al.* Association between genetic polymorphism of *XRCC1* Arg194Trp and risk of schizophrenia. *Psychiatry Res* 2009;169:186.
- Nafissi S, Saadat I, Farrashbandi H, *et al.* Association between three genetic polymorphisms of glutathione S-transferase Z1 (*GSTZ1*) and susceptibility to schizophrenia. *Psychiatry Res* 2011;187:314–15.
- Rezaei Z, Saadat I, Saadat M. Association between three genetic polymorphisms of glutathione S-transferase Z1 (*GSTZ1*) and susceptibility to bipolar disorder. *Psychiatry Res* 2012;198:166–8.
- Bahaoddini A, Farrashbandi H, Saadat M. Genetic polymorphism of glutathione S-transferase T1 (*GSTT1*) and QT-interval in schizophrenia patients. *J Mol Neurosci* 2009;38:173–7.
- Saadat M, Mobayen F, Farrashbandi H. Genetic polymorphism of glutathione S-transferase T1: a candidate genetic modifier of individual susceptibility to schizophrenia. *Psychiatry Res* 2007;153:87–91.
- Bazett H. An analysis of the time relationship of electrocardiograms. *Heart* 1920;7:353–70.
- Saadat M, Ansari-Lari M. Polymorphism of *XRCC1* (at codon 399) and susceptibility to breast cancer, a meta-analysis of the literatures. *Breast Cancer Res Treat* 2009;115:137–44.